

**Interactions between  
cirrus and embedded  
contrails**

K. Gierens

# Selected topics on interactions between cirrus clouds and embedded contrails

**K. Gierens**

Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre,  
Oberpfaffenhofen, Germany

Received: 10 September 2012 – Accepted: 14 September 2012 – Published:  
24 September 2012

Correspondence to: K. Gierens (klaus.gierens@dlr.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Persistent contrails and natural cirrus clouds often coexist in the upper troposphere and contrails can be embedded within cirrus clouds. The present paper deals with some questions regarding the interaction of cirrus clouds and embedded contrails. I have selected only questions that can be answered by analytical means. I find that (1) the emission index for water vapour is only slightly changed when an aircraft crosses a cirrus cloud, (2) that contrail formation is not affected by an ambient cirrus, (3) that cirrus ice crystals entrained into the trailing wing tip vortex do not efficiently retard the sublimation of contrail ice crystals, and (4) that cirrus can start to dissolve an embedded contrail after a couple of hours by aggregation.

## 1 Introduction

Although contrails and cirrus clouds do not have the same formation conditions, they often occur together in the same regions because the condition for cirrus formation, ice supersaturation, is also required for contrail persistence. Indeed, sensitive observations with lidar instruments that are capable to detect optically very thin cirrus, including sub-visible cirrus, suggest that contrails mostly occur together with cirrus (Sassen, 1997; Immler et al., 2008; Iwabuchi et al., 2012). Further, airborne lidar observations often show backscatter peaks within cirrus clouds that can be interpreted as embedded contrails.

A recent study of air traffic effects in the North Atlantic on cirrus (and contrail) coverage and outgoing longwave radiation (OLR, Schumann and Graf, 2012) showed an effect on OLR that was too large to be explainable by the additional cloud coverage due to contrails (1 to 2 %) alone. Instead one must assume that aircraft flying through cirrus clouds produce contrails and further effects within these natural clouds which eventually manifests itself as a significant change in OLR. These findings are consistent with results from radiation transfer calculations (Schumann et al., 2012).

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Interactions between  
cirrus and embedded  
contrails**K. Gierens

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

These results warrant to start studies concerning the interaction between cirrus clouds and embedded contrails using numerical models and analytical calculations where possible. To my knowledge, contrails have so far been modelled mainly as isolated objects and numerical simulations of the development of contrails within cirrus clouds have only recently been performed for the first time (Gierens and Spichtinger, 2012). In the present paper I will present some simple analytical calculations on certain aspects of contrail development within cirrus clouds. Generally the questions treated are of the kind “is process P important for the interaction or not?” and these questions can often be treated considering and comparing various time scales.

The questions treated are the following:

- How much do ice crystals sucked through an engine change the water vapour emission index?
- Is contrail formation within a cirrus affected by the cirrus?
- Can cirrus crystals mixed into the downward travelling wing vortex enhance the available water mass such that the sublimation of contrail ice is efficiently retarded?
- What is the role of aggregation between cirrus and contrail ice crystals?

These questions are dealt with in several sections. A summary is given at the end and since quite a number of calculations are performed I provide a nomenclature section as an appendix.

## 2 How much do ice crystals sucked through an engine change the water vapour emission index?

An aircraft flying through a cirrus cloud sucks ice crystals together with air through its engines. The ice sublimates and the resulting water vapour at engine exit adds to the

water vapour resulting from burning kerosene; thus an enhanced emission index for water vapour results which could affect contrail formation. How large is this effect?

Let us assume a fairly large ice mixing ratio of  $q_c = 10^{-3.5}$ . This value corresponds to the maximum mixing ratio measured in 28 research flights (see Schiller et al., 2008, their Fig. 2 and the second entry in their Table 2). An aircraft engine typically has an air fuel ratio  $\dot{m}_{\text{air}}/\dot{m}_{\text{fuel}} \approx 70$ , and the emission index of water vapour for kerosene is  $\text{EI}_{\text{H}_2\text{O}} = 1.25$ . With these values we find the ratio of water vapour emission at engine exit of sublimated ice vs. emission from kerosene combustion is  $(\dot{m}_{\text{air}}q_c)/(\dot{m}_{\text{fuel}}\text{EI}_{\text{H}_2\text{O}})$ . Even with the assumed large ice mixing ratio this ratio is merely of the order percent. Thus the contribution of sublimated cirrus ice on the water vapour emission index is generally negligible.

### 3 Is contrail formation within a cirrus affected by the cirrus?

How could a pre-existing cirrus affect contrail formation? When ambient air (i.e. cirrus air) is mixed into the exhaust plume, the plume temperature decreases and the relative humidity in the plume increases. Contrail formation requires that the relative humidity increases up to water saturation (Schmidt–Appleman criterion, see Schumann, 1996); if water saturation is reached, the water that is condensed on the exhaust particles, mainly on soot particles which serve as freezing nuclei, freezes and a contrail results. If this happens within a cirrus, ice crystals are mixed into the plume and start to grow as soon as the plume air is supersaturated with respect to ice, thereby consuming the excess vapour. Thus we have two competing processes affecting the plume supersaturation, the cooling due to mixing caused by jet dynamics and the growing ice crystals mixed into the plume from the ambient cirrus. The latter process adds latent heat to the plume but this effect is negligible.

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The temporal change of the saturation ratio  $S(t) = e(t)/e^*[T(t)]$  in an exhaust plume is given as

$$\begin{aligned} \frac{dS}{dt} &= \frac{e^*[T(t)] \frac{de}{dt} - e(t) \frac{de^*}{dT} \frac{dT}{dt}}{e^*[T(t)]^2} \\ &= \frac{1}{e^*[T(t)]} \frac{de}{dt} - S(t) \frac{L}{R_v T^2} \frac{dT}{dt}. \end{aligned} \quad (1)$$

The symbols are listed in the nomenclature section below. The second rhs term describes the change of the saturation vapour pressure due to jet cooling with a rate  $dT/dt$  which I prescribe as a function  $C(t)$  that is determined by the entrainment of ambient air into the hot plume air. Since the latent heat release is a negligible contribution to this term, it is not affected by the presence of an ambient cirrus and I do not consider it further.

The first rhs term describes change of  $S$  due to the change of the vapour pressure  $e$ . This term has a jet dynamical and a microphysical component as follows:

$$\frac{de}{dt} = GC(t) - c(t), \quad (2)$$

where  $G$  is the change in vapour pressure per change in temperature of the isobarically expanding plume (cf. Schumann, 1996):

$$G = \frac{c_p \rho E I_{H_2O}}{\epsilon Q (1 - \eta)}. \quad (3)$$

The condensation function  $c(t)$  for condensation on pre-existing cirrus ice crystals can be approximated following Gierens (2003) as

$$c(t) = \frac{e - e^*}{\tau_g} \quad (4)$$

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with the crystal growth time scale  $\tau_g$  (see, e.g. Kärcher and Lohmann, 2002; Gierens, 2003). In order to see whether an ambient cirrus affects contrail formation it suffices to compare the two  $S$ -change rates  $GC(t)/e^*$  and  $c(t)/e^*$  or their inverses which can be interpreted as the respective time scales.  $C(t)$  can be obtained from an explicit consideration of the entrainment of ambient air into the jet where we have (Schumann, 1996)

$$c_p(T_{\text{jet}} - T_{\text{cirrus}}) \approx \frac{Q(1 - \eta)}{N(t)}. \quad (5)$$

$N(t)$  is the plume dilution ratio, indicating that at plume age  $t$  the jet is composed of one unit exhaust air and  $N - 1$  units of ambient cirrus air. In the equation above I simply considered  $c_p$  to be a constant and I neglected the enthalpy of ambient air against that of the hot exhaust gases which are reasonable approximations.  $N(t)$  can be approximated from engine exit up to almost 3 h plume age as a power law (Schumann et al., 1998):

$$N(t) = N_0(t/t_0)^\gamma \quad (6)$$

where  $N_0 = 7000$  when  $t_0 = 1$  s.  $\gamma = 0.8$  was successfully used by Schumann et al. (1998) to fit a large number of in-situ measurements. The cooling rate is then:

$$C(t) \equiv \frac{dT_{\text{jet}}}{dt} = -\frac{Q(1 - \eta)}{c_p \frac{\gamma}{N(t)t}} \quad (7)$$

Setting everything together and inserting the expression for  $G$  we find for the dynamic time scale in the jet (of course the absolute value has to be taken):

$$\tau_{\text{jet}} = \frac{\epsilon e^* N(t) t}{E I_{\text{H}_2\text{O}} \rho \gamma} = 0.62 \frac{e^*}{\rho} N(t) t. \quad (8)$$

**Interactions between cirrus and embedded contrails**

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In order to find a typical order of magnitude for  $\tau_{\text{jet}}$  we can take  $t = 1$  s,  $N = N_0 = 7000$  and  $e^*/p = \mathcal{O}(10^{-3})$  which gives a timescale of a few seconds.

The condensation time scale can be obtained using expressions from Gierens (2003):

$$\tau_{\text{g}}^{-1} = 1.4 \times 10^{-6} T^{1.61} p^{-1} (se^*)^{1/3} \mathcal{N}_{\text{cir}}^{2/3}, \quad (9)$$

which can be inserted into Eq. (4) and gives the condensation timescale as

$$\tau_{\text{cond}} = \tau_{\text{g}}/s \quad (10)$$

with  $s = S - 1$ .

Figure 1 shows the ratio  $\tau_{\text{cond}}/\tau_{\text{jet}}$  as a function of cirrus ice crystal number concentration and plume age within the first second of the plume and for various ambient conditions. In order to yield a maximum effect of the ambient cirrus I have assumed water saturation for the value of  $s$ . In spite of this, the ratio is much larger than one for realistic cirrus conditions ( $\mathcal{N}_{\text{cir}} < 10 \text{ cm}^{-3}$ ). In microphysical terms: the cirrus crystals grow too slowly to efficiently halt the increase of plume supersaturation due to the cooling. This means that contrail formation is not affected by the presence of an ambient cirrus cloud. A ratio of unity between the two time scales is only reached at very high crystal concentrations,  $\mathcal{N} = \mathcal{O}(10^4 \dots 10^5) \text{ cm}^{-3}$ . Such high concentrations are perhaps observed in the contrails themselves. Thus it is eventually the growing contrail ice that is able to compete with the dynamic jet cooling for affecting plume supersaturation. The number of particles that can freeze at sufficient supersaturations in a plume is much higher than the number of soot particles (which usually nucleate ice first). If there were not enough soot for letting  $\tau_{\text{cond}}/\tau_{\text{jet}} \rightarrow 1$  other particles would freeze until eventually  $\tau_{\text{cond}}/\tau_{\text{jet}} = 1$  (see also Kärcher and Yu, 2009).

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



#### 4 Can cirrus crystals mixed into the downward travelling wing vortex enhance the available water mass such that the sublimation of contrail ice is efficiently retarded?

During the jet phase, cirrus ice crystals are entrained into the exhaust plume as enthalpy and water vapour are mixed in. Once the vortex pair is completely rolled up and starts to travel downwards, both contrail ice and cirrus ice crystals experience the adiabatic heating in the plume and start to sublime as soon as the in-plume relative humidity decreases below ice saturation. Now, if the contrail ice is alone present, a certain fraction (depending mainly on initial supersaturation and temperature) of the contrail ice crystals eventually sublimates completely (Unterstrasser and Sölch, 2010). When cirrus ice is present, this process could be slowed down because the large cirrus crystals could supply water vapour without sublimating completely and the added water vapour helps to keep the relative humidity closer to ice saturation than without the cirrus crystals. In the following I estimate whether this can be an important process.

It suffices to compare the sublimation time scales for the two ice populations, the contrail ice and the cirrus ice. Let  $\mathcal{M}$  be the ice water concentration (IWC). Then, for a certain crystal mass distribution we can invoke the Koenig-parameterisation for crystal growth/sublimation (Koenig, 1971) in the following form (Eq. 45 of Spichtinger and Gierens, 2009, without corrections for ventilation and kinetic effects):

$$\frac{d\mathcal{M}}{dt} = a\mu_b, \quad (11)$$

where  $a, b$  are parameters depending mainly on temperature and saturation ratio ( $a < 0$  for  $S < 1$ ), and where  $\mu_b$  is the moment of order  $b$  of the crystal's mass probability density function (pdf).  $\mathcal{M}$  itself is proportional to the first moment,  $\mathcal{M} \propto \mu_1$ , and the sublimation timescale is then

$$\tau = \frac{\mathcal{M}}{|d\mathcal{M}/dt|} \propto \frac{\mu_1}{\mu_b}. \quad (12)$$

### Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





The moments  $\mu_k$  are given in Spichtinger and Gierens (2009, Eq. 24) for a log-normal pdf, and these will be used here as well:

$$\mu_k = \mathcal{N} \bar{m}^k r_0^{k(k-1)/2}, \quad (13)$$

with crystal number density  $\mathcal{N}$  (the pdf is normalised to the number concentration of ice crystals), mean mass  $\bar{m}$  and a parameter  $r_0$  that quantifies the width of the pdf. With this information we can now compute the ratio of the sublimation time scales for contrail and cirrus ice:

$$\begin{aligned} \frac{\tau_{\text{cir}}}{\tau_{\text{con}}} &= \frac{\mathcal{N}_{\text{cir}} \bar{m}_{\text{cir}} r_{0,\text{cir}}}{\mathcal{N}_{\text{con}} \bar{m}_{\text{con}} r_{0,\text{con}}} \cdot \frac{\mathcal{N}_{\text{con}} \bar{m}_{\text{con}}^b r_{0,\text{con}}^{b(b-1)/2}}{\mathcal{N}_{\text{cir}} \bar{m}_{\text{cir}}^b r_{0,\text{cir}}^{b(b-1)/2}} \\ &= \frac{\bar{m}_{\text{cir}}^{1-b} r_{0,\text{cir}}^{1-b(b-1)/2}}{\bar{m}_{\text{con}}^{1-b} r_{0,\text{con}}^{1-b(b-1)/2}}. \end{aligned} \quad (14)$$

The width parameters  $r_0$  of cirrus and contrail pdfs are similar (values of 2 to 4 are typical) such that their ratio is of the order unity. The value of  $b$  is of the order 1/3 to 1/2, such that also the exponent  $1 - b(b - 1)/2$  approximates unity; the factors involving the width parameters can thus be safely neglected. What remains is the ratio of the mean crystal masses, viz.:

$$\frac{\tau_{\text{cir}}}{\tau_{\text{con}}} \approx \left( \frac{\bar{m}_{\text{cir}}}{\bar{m}_{\text{con}}} \right)^{(1-b)}. \quad (15)$$

This ratio is typically of the order 100. Thus the effect of cirrus ice in the downward travelling plume to halt the sublimation of the contrail ice is usually small or even negligible. This result is surprising if we compare the total ice mass concentrations in the plume:  $\mathcal{M}_{\text{con}}$  is typically of the order 1 to 10  $\text{mg m}^{-3}$  (Schumann, 2002, Fig. 11.5) while

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that of a cirrus can reach a few  $10 \text{ mgm}^{-3}$  (e.g. Schiller et al., 2008). Although there can be more cirrus ice in the contrail vortex than contrail ice itself, the reaction of the cirrus ice to changing saturation conditions is much too slow to retard the sublimation of the contrail ice when subsaturation is reached in the adiabatically heated plume.

## 5 What is the role of aggregation between cirrus and contrail ice crystals?

After the end of the vortex phase the contrail develops further within the cirrus cloud. There is generally a complex interaction between the two ice classes, depending on ambient conditions, on the state of the cirrus, on the composition of the cirrus ice crystals (heterogeneously vs. homogeneously formed ice crystals) and so on. Thus numerical simulations are generally required to capture all these possible interactions. But one can at least estimate the role of aggregation between cirrus and contrail ice crystals. In particular if the contrail is located in the lower part of a cirrus layer it is possible that large cirrus crystals falling from above into the contrail collect contrail ice crystals, thereby diminishing the contrail signature in the perturbed cloud.

I derive the expression for the aggregation time scale starting from the master equation for aggregation (cf. Kienast-Sjögren et al., 2012). Assuming that contrail crystals are depleted by aggregation with cirrus crystals and neglecting aggregation between contrail crystals, the master equation takes the simple form:

$$\frac{\partial n(m_{\text{con}}, t)}{\partial t} = - \int_0^{\infty} K(m_{\text{con}}, m_{\text{cir}}) n(m_{\text{con}}, t) n(m_{\text{cir}}, t) dm_{\text{cir}}. \quad (16)$$

Here,  $n(m, t)$  is a number density per infinitesimal mass interval. Using the normalized pdf  $f(m, t) = n(m, t)/\mathcal{N}$  and integrating both sides over all contrail crystal masses we

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



obtain:

$$\frac{\partial \mathcal{N}_{\text{con}}(t)}{\partial t} = -\mathcal{N}_{\text{con}} \mathcal{N}_{\text{cir}} \times \int_0^{\infty} \int_0^{\infty} K(m_{\text{con}}, m_{\text{cir}}) f(m_{\text{con}}, t) f(m_{\text{cir}}, t) dm_{\text{cir}} dm_{\text{con}}. \quad (17)$$

The double integral can be conceived as the expectation value of  $K$  given the mass pdfs,  $\langle K \rangle$ . The aggregation time scale is then

$$\tau_{\text{aggr}} = \frac{\mathcal{N}_{\text{con}}}{|\partial \mathcal{N}_{\text{con}} / \partial t|} = (\mathcal{N}_{\text{cir}} \langle K \rangle)^{-1}, \quad (18)$$

For the aggregation kernel I assume the following expression (Field and Heymsfield, 2003; Westbrook et al., 2004):

$$K = \pi(R_{\text{con}}^2 + R_{\text{cir}}^2) v_t E_a, \quad (19)$$

where  $v_t$  is the fall speed of the cirrus crystals. The fall speed of the contrail crystals has been neglected and that is the reason why I could neglect aggregation between contrail ice crystals in the master equation.  $E_a$  is the collision efficiency. As generally  $R_{\text{con}} < R_{\text{cir}}$  we may simplify the analysis by also neglecting  $R_{\text{con}}^2$  in the expression for  $K$ . The corresponding integral for  $\langle K \rangle$  then reduces to a single integral over the cirrus crystal masses. The terminal velocity is usually taken proportional to  $R_{\text{cir}}^y$ :  $v_t = v_0 (R_{\text{cir}}/R_0)^y$ .

Using this we have

$$\tau_{\text{aggr}} = (\mathcal{N}_{\text{cir}} \langle R_{\text{cir}}^{2+y} \rangle v_0 \pi E_a)^{-1}. \quad (20)$$

$y$  is in the range 0.7 to 1.3 depending on crystal size and habit (e.g. Heymsfield and laquinta, 2000, Eq. A29), but for crystals with maximum dimension smaller than about 0.5 mm (which are much more abundant than larger ones),  $y$  is about 1.25, and the prefactor  $v_0$  is of the order  $22\,700 \text{ ms}^{-1}$  for a unit radius  $R_0 = 1 \text{ m}$ . With this parameterisation an ice crystal of  $R = 100 \text{ }\mu\text{m}$  would fall with a speed of  $23 \text{ cm s}^{-1}$ . For a lognormal

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



radius pdf we can estimate  $\langle R_{\text{cir}}^{3.25} \rangle \approx 2\bar{R}^{3.25}$ . With  $E_a \approx 0.7$  we can write the following approximation:

$$\tau_{\text{aggr}} \approx 4 \times 10^{-5} \mathcal{N}_{\text{cir}}^{-1} \bar{R}_{\text{cir}}^{-3.25}. \quad (21)$$

As shown in Fig. 2, this aggregation time scale varies between approximately 1 s for an unrealistically high density of large crystals (the upper right corner in the figure) and several hours. Heymsfield and McFarquhar (2002, Fig. 4.6) present concentrations of large crystals with  $R_{\text{cir}} > 50 \mu\text{m}$ . These have typically concentrations of 1 to  $10 \text{ l}^{-1}$ . Thus, aggregation timescales are typically a couple of hours, similar to timescales needed for contrails to spatially expand several kilometres. Thus we can expect that cirrus clouds after some hours start to remove the ice of embedded contrails (in particular contrails at cloud bottom) via aggregation.

## 6 Summary

As a prooemium to more elaborate numerical studies (Gierens and Spichtinger, 2012), I have treated several questions regarding the interaction between cirrus and embedded contrails. Only questions that can be answered by analytical means, mainly by comparison of certain timescales, have been selected. The questions and their answers are:

Q: How much do ice crystals sucked through an engine change the water vapour emission index?

A: The emission index of water vapour changes by a few percent at most when an aircraft crosses a cirrus cloud.

Q: Is contrail formation within a cirrus affected by the cirrus?

A: No. The contrail forms as if no cirrus were present.

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Q: Can cirrus crystals mixed into the downward travelling wing vortex enhance the available water mass such that the sublimation of contrail ice is efficiently retarded?

A: Hardly, although the ice mass provided by the cirrus can be larger than the ice mass of the contrail itself. As the concentration of contrail crystals is several orders of magnitude larger than the concentration of cirrus crystals entrained in the plume, the cirrus crystals react slowly on changing saturation conditions and therefore have hardly any effect on the evolution of the saturation ratio within the plume, which is driven mainly by the vortex dynamics.

Q: What is the role of aggregation between cirrus and contrail ice crystals?

A: After a few hours it is possible that sedimenting cirrus crystals collect a significant fraction of contrail crystals (mainly for contrails embedded in the lower part of a cirrus), and thus that the contrail dissolution is enhanced by aggregation with cirrus crystals.

There are many more interesting questions regarding contrails embedded in cirrus clouds. To answer them is generally beyond the possibilities of analytical estimation. Numerical simulations are then required which are the topic of another paper.

## Nomenclature

- a* prefactor for crystal growth parameterisation
- b* exponent for crystal growth parameterisation
- c* condensation rate in terms of partial pressure ( $\text{Pa s}^{-1}$ )
- $c_p$  specific heat of air ( $1005.7 \text{ J kg}^{-1} \text{ K}^{-1}$ )
- C* cooling rate due to jet expansion ( $\text{K s}^{-1}$ )
- e* partial pressure of water vapour (Pa)

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$e^*$	saturation (with respect to ice) partial pressure of water vapour (Pa)
$E_a$	collision and collection efficiency ( $\approx 0.7$ )
$EI_{\text{H}_2\text{O}}$	emission index of water vapour (1.25)
$f$	normalised probability density function for crystal masses ( $\text{kg}^{-1}$ )
$G$	$de/dT$ in an expanding plume ( $\text{PaK}^{-1}$ )
$K$	aggregation kernel ( $\text{m}^3 \text{s}^{-1}$ )
$L$	latent heat of sublimation ( $2.8352 \times 10^6 \text{ Jkg}^{-1}$ )
$\mathcal{M}$	ice water concentration (IWC, $\text{kgm}^{-3}$ )
$\dot{m}_{\text{air/fuel}}$	air and fuel flow rates through an engine ( $\text{kg s}^{-1}$ )
$m$	crystal mass (kg)
$\bar{m}$	mean mass of an ice crystal pdf (kg)
$n$	number density of ice crystals per infinitesimal mass interval ( $\text{m}^{-3} \text{kg}^{-1}$ )
$\mathcal{N}$	number density of ice crystals ( $\text{m}^{-3}$ )
$N, N_0$	plume dilution ratio, at unit time
$p$	pressure (Pa)
$Q$	calorific value of kerosene ( $43.2 \text{ MJkg}^{-1}$ )
$q_c$	ice mixing ratio ( $\text{kg ice}(\text{kg air})^{-1}$ )
$r_0$	width parameter of crystal size distribution
$R$	crystal “radius” (m)
$\bar{R}$	mean crystal radius (m)
$R_v$	gas constant of water vapour ( $461.5 \text{ Jkg}^{-1} \text{ K}^{-1}$ )
$s$	supersaturation with respect to ice ( $S - 1$ )
$S$	ice saturation ratio
$t$	time or plume age (s)
$T, T_x$	various temperatures (K or $^{\circ}\text{C}$ )
$v_t$	terminal fall speed of ice crystals ( $\text{ms}^{-1}$ )
$y$	exponent in fall speed to radius relation
$\gamma$	exponent for plume dilution parameterization

## Interactions between cirrus and embedded contrails

K. Gierens

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


$\epsilon$  ratio of molar masses of water and air (0.622)  
 $\eta$  overall propulsion efficiency of an aircraft ( $\approx 0.3$ )  
 $\tau_x$  various process time scales (s)

*Acknowledgements.* I thank Ulrich Schumann for critically reading a preliminary version of this manuscript. The work contributes to the DLR program “Emissionen und Klimawirkung des Luftverkehrs”.

## References

- 5 Field, P. and Heymsfield, A.: Aggregation and scaling of ice crystal size distributions, *J. Atmos. Sci.*, 56, 544–560, 2003. 25247
- Gierens, K.: On the transition between heterogeneous and homogeneous freezing, *Atmos. Chem. Phys.*, 3, 437–446, doi:10.5194/acp-3-437-2003, 2003. 25241, 25242, 25243
- 10 Heymsfield, A. and Iaquinta, J.: Cirrus crystal terminal velocities, *J. Atmos. Sci.*, 57, 916–938, 2000. 25247
- Heymsfield, A. and McFarquhar, G.: Mid-latitude and tropical cirrus: microphysical properties, in: *Cirrus*, edited by: Lynch, D., Sassen, K., Starr, D., and Stephens, G., Oxford University Press, New York, USA, 78–101, 2002. 25248
- 15 Immler, F., Treffeisen, R., Engelbart, D., Krüger, K., and Schrems, O.: Cirrus, contrails, and ice supersaturated regions in high pressure systems at northern mid latitudes, *Atmos. Chem. Phys.*, 8, 1689–1699, doi:10.5194/acp-8-1689-2008, 2008. 25238
- Iwabuchi, H., Yang, P., Liou, K., and Minnis, P.: Physical and optical properties of persistent contrails: climatology and interpretation, *J. Geophys. Res.*, 117, D06215, doi:10.1029/2011JD017020, 2012. 25238
- 20 Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: homogeneous freezing of supercooled aerosols, *J. Geophys. Res.*, 107, 4698, doi:10.1029/2001JD000470, 2002. 25242
- Kärcher, B. and Yu, F.: Role of aircraft soot emissions in contrail formation, *Geophys. Res. Lett.*, 36, L01804, doi:10.1029/2008GL036649, 2009. 25243

## Interactions between cirrus and embedded contrails

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kienast-Sjögren, E., Spichtinger, P., and Gierens, K.: Formulation and test of an ice aggregation scheme for two-moment bulk microphysics schemes, *Atmos. Chem. Phys. Discuss.*, 12, 23975–24009, 2012,

<http://www.atmos-chem-phys-discuss.net/12/23975/2012/>. 25246

- 5 Koenig, L.: Numerical modeling of ice deposition, *J. Atmos. Sci.*, 28, 226–237, 1971. 25244
- Sassen, K.: Contrail-cirrus and their potential for regional climate change, *B. Am. Meteorol. Soc.*, 78, 1885–1903, 1997. 25238
- Schiller, C., Krämer, M., Afchine, A., Spelten, N., and Sitnikov, N.: Ice water content of Arctic, midlatitude, and tropical cirrus, *J. Geophys. Res.*, 113, D24208, doi:10.1029/2008JD010342, 2008. 25240, 25246
- 10 Schumann, U.: On conditions for contrail formation from aircraft exhausts, *Meteorol. Z.*, 5, 4–23, 1996. 25240, 25241, 25242
- Schumann, U.: Contrail cirrus, in: *Cirrus*, edited by: Lynch, D., Sassen, K., Starr, D., and Stephens, G., Oxford University Press, chapter 11, 231–255, Oxford, New York, 2002. 25245
- 15 Schumann, U. and Graf, K.: Aviation-induced cirrus and radiation changes at diurnal timescales, *J. Geophys. Res.*, submitted, 2012. 25238
- Schumann, U., Schlager, H., Arnold, F., Baumann, R., Haschberger, P., and Klemm, O.: Dilution of aircraft exhaust plumes at cruise altitudes, *Atmos. Environ.*, 32, 3097–3103, 1998. 25242
- Schumann, U., Mayer, B., Graf, K., and Mannstein, H.: A parametric radiative forcing model for contrail cirrus, *J. Appl. Meteorol. Climatol.*, 51, 1391–1406, 2012. 25238
- 20 Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1a: Model description and validation, *Atmos. Chem. Phys.*, 9, 685–706, doi:10.5194/acp-9-685-2009, 2009. 25244, 25245
- Unterstrasser, S. and Sölch, I.: Study of contrail microphysics in the vortex phase with a Lagrangian particle tracking model, *Atmos. Chem. Phys.*, 10, 10003–10015, doi:10.5194/acp-10-10003-2010, 2010. 25244
- 25 Westbrook, C., Ball, R., Field, P., and Heymsfield, A.: Theory of growth by differential sedimentation, with application to snowflake formation, *Phys. Rev. E*, 70, 021403, 1–7, 2004. 25247

**Interactions between  
cirrus and embedded  
contrails**

K. Gierens

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Interactions between cirrus and embedded contrails

K. Gierens



**Fig. 1.** Isolines of  $\tau_{\text{cond}}/\tau_{\text{jet}}$ , the ratio of the condensation time scale to the jet expansion time scale vs. plume age during contrail formation (first second) and vs. number concentration of ice crystals in the ambient cirrus for various conditions: **(a)**  $T = -40^\circ\text{C}$ ,  $p = 300$  hPa, **(b)**  $T = -50^\circ\text{C}$ ,  $p = 250$  hPa, **(c)**  $T = -60^\circ\text{C}$ ,  $p = 200$  hPa. Note that crystal number concentrations for cirrus clouds exceeding  $10\text{ cm}^{-3}$  are unrealistic. On the contrary, number concentrations in young contrails are of the order  $10^4$  to  $10^5\text{ cm}^{-3}$ . Hence, contrail formation within a cirrus cloud is generally not affected by the cirrus ice crystals (ratio  $\gg 1$ ) but the ice crystals forming and growing in the exhaust can affect the evolution of the supersaturation almost as strongly as the expansion process does.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Interactions between cirrus and embedded contrails

K. Gierens



**Fig. 2.** Aggregation time scale for the removal of contrail ice crystals by sedimenting cirrus ice crystals as a function of the mean cirrus crystal radius  $\bar{R}_{\text{cir}}$  and the cirrus crystal number concentration  $\mathcal{N}_{\text{cir}}$ . Note that only parameter combinations below the main diagonal can be considered realistic in cirrus clouds; thus aggregation timescales are typically of the order hours.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

