

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani, A. Stohl, and S. Eckhardt

1-NILU, Norwegian Institute for Air Research, 2027, Kjeller, Norway

Received: 7 June 2012 – Accepted: 11 August 2012 – Published: 5 October 2012

Correspondence to: M. Cassiani (mc@nilu.no)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Abstract

Megacities are extreme examples of the continuously growing urbanization of human population that pose (new) challenges to the environment and human health at a local scale. However, because of their size megacities also have larger-scale effects and more research is needed to quantify their regional and global scale impacts. We performed a study of the characteristics of plumes dispersing from a group of thirty-six of world's megacities using the Lagrangian particle model FLEXPART and focusing on black carbon (BC) emissions during the years 2003–2005. BC was selected since it is representative of combustion-related emissions and it has a significant role as a short-lived climate forcer. Based on the BC emissions two tracers were modeled; a purely passive tracer and one subject to wet and dry deposition. These tracers allowed us to investigate the role of deposition processes in determining the impact of megacities' pollutant plumes. The particles composing the plumes have been sampled in space and time. The time sampling allowed us to investigate the evolution of the plume from its release up to 48 days after emission and to generalize our results for any substance decaying with a time scale sufficiently shorter than the time window of 48 days. The physical characteristics of the time averaged plume have been investigated and this showed that although local conditions are important, overall the latitude of the city is the main factor influencing both the local and the regional-to-global dispersion of the megacities' plumes. We also repeated the calculations of some of the regional-pollution-potential metrics previously proposed by Lawrence et al. (2007), thus extending their results to a depositing scalar and retaining the evolution in time for all the plumes. Noteworthy our results agreed well with the previous results despite being obtained using a totally different modeling framework. For the environmental impact on a global scale we focused on the export of mass from the megacities to the sensitive polar regions. We found that the sole city of Saint Petersburg contributes more to the lower troposphere pollution and deposition in the Arctic than the whole ensemble of Asian megacities. In general this study showed that the pollution of urban origin in the

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lower troposphere of the Arctic is mainly generated by northern European sources. We also found that the deposition of BC in the Antarctic due to megacities is comparable to the emissions generated by local shipping activities. Finally multiplying population and ground level concentration maps, we found that the exposure of human population to megacities pollution occurs mainly inside the city boundaries and this is especially true if deposition is accounted for. However, some exceptions exist (Beijing, Tianjin, Karachi) where the impact on population outside city boundary is larger than that inside city boundary.

1 Introduction

More than 50 % of the human population is living in urban areas (United Nations Population Fund report unit <http://www.unfpa.org/swp/2007/english/introduction.html>) and this fraction is still increasing, although urban areas currently cover only 0.5 % of the Earth's land area (Schneider et al., 2009). Extreme examples of urbanization are the so-called megacities with a population of several million of people; ten millions are often used as a threshold to define a megacity (Gurjar and Lelieveld, 2005). This large concentration of human activity causes unprecedented pressure on the environment including the quality of air both inside the city (e.g. Gurjar et al., 2008) and on a regional and global scale. Some case studies (e.g. Guttikunda et al., 2003; Stohl et al., 2003; Real et al., 2008; Mena-Carrasco et al., 2009; Voss et al., 2010) have investigated regional and long range transport of megacity plumes and the associated chemical processes, but more must be done to quantify the relative importance of their local versus large scale impacts.

In a relatively recent paper Lawrence et al. (2007, called L07 in the remainder of the paper) investigated the differences and common features in the regional dispersion characteristics of plumes from a collection of world's largest cities. In that work three artificial tracers, with a characteristic decaying time of 1, 10 and 100 days were released and transported using a 3-D global Eulerian chemistry transport model. This study

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

gave significant information on the common regional features of dispersion from these megacities and identified cases where the dispersion was dominated by local conditions. It is well known that for studying dispersion characteristics of plumes originating from a relatively localized source, a modeling approach based on 3-D Lagrangian particle models is a viable alternative to Eulerian models. Lagrangian models minimize, if not completely erase, artificial numerical diffusivity effects and can account for various sub-grid scale processes in an accurate way. Based on these premises we decided to use the Lagrangian particle model FLEXPART (Stohl et al., 2005) for simulating the dispersion from a group of world's megacities during the year 2003–2005, as established within the framework of the European Union project MEGAPOLI (Megacities: Emissions, urban, regional and Global Atmospheric POLLution and climate effects, and Integrated tools for assessment and mitigation, <http://megapoli.info/>). Our investigation has a similar scope as the study of L07 but uses a Lagrangian particle model with a 4-D sampling grid. The extension into the time domain, allowed the use of additional metrics to describe the dispersion process of the plumes. Moreover, we specifically investigate long range transport into the sensitive polar regions and into the upper layer of the troposphere and the stratosphere. Finally, we answer some of the question pointed out by L07 by investigating the behavior of a passive tracer versus an aerosol-like tracer subject to removal processes by dry and wet deposition, and inferring information about population exposure by combining modeled near ground concentration and population maps.

In the next section a brief description of the FLEXPART model, the emission inventory and the meteorological input data will be given. In Sect. 3 the mean transport characteristics of the megacity plumes will be discussed. In Sect. 4 some metrics originally proposed by L07 will be calculated and compared with their original results. The contribution of megacities to pollution in the polar regions will be investigated in Sect. 5. In Sect. 6, a metric of the large scale versus local impacts of a megacity on human population will be defined.

2 Methods

2.1 Model description

For this study, the Lagrangian particle dispersion model FLEXPART (Stohl et al., 2005) was run for the period November 2002 to December 2005. FLEXPART is an extensively validated transport model (e.g. Stohl et al., 1998) and the code and documentation are freely available (see <http://transport.nilu.no/flexpart/>). It is worth to remind here that: (i) FLEXPART uses the well mixed criteria, introduced by Thomson (1987), to define stochastic differential equations with a Gaussian velocity distribution to simulate the effect of planetary boundary layer turbulence on particle dispersion and accounts for the vertical air density gradient (Stohl and Thomson, 1999), (ii) it includes a horizontal mesoscale meandering parameterization through an additional Langevin equation for the mesoscale velocity, (iii) it simulates convective exchanges using the scheme of Emanuel and Zivkovic-Rothman (1999), and (iv) the model includes dry/wet deposition processes. Thanks to the Lagrangian particle representation the model does not suffer of significant numerical diffusivity and the grid is used only to extract information from the particle field which in general preserves very well any flow structures generated during the advection and dispersion process. This last characteristic and the superior accuracy in the representation of sub-grid dispersion processes differentiates this class of models from Eulerian (fully grid based) chemistry transport models, and makes FLEXPART particularly suitable to study dispersion from localized sources in the planetary boundary layer.

The model is driven here by operational meteorological data from the European Centre for Medium-Range Weather Forecasts (ECMWF) retrieved with a horizontal resolution of $1^\circ \times 1^\circ$ and with 60 vertical levels in a hybrid coordinate system and a time resolution of 3h. The model output grid has a horizontal resolution of $1^\circ \times 1^\circ$ and 25 vertical levels ranging from 100 m in the layer closest to the ground to 3 km at an elevation above ground of 27–30 km. The grid extends also in the time space thus partitioning the particles among 10 time (age) classes based on the time elapsed since the particle

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

release, i.e. the time of emission. The age-classes duration used are expressed in days as, $0 \leq t < 1$, $1 \leq t < 3$, $3 \leq t < 6$, $6 \leq t < 9$, $9 \leq t < 14$, $14 \leq t < 20$, $20 \leq t < 27$, $27 \leq t < 34$, $34 \leq t < 41$, $41 \leq t < 48$. Note that, because of their different duration, the mass content of a perfectly conserved tracer differs for the different age classes. When a particle reaches 48 days of elapsed time it is removed from the simulation, therefore even the conserved scalar (BC_{tr}, see later) has a maximum time of residence in the atmosphere of 48 days. The model output is stored every 5 days as 5-day average concentration $c(x, y, z, t)$ or, written in a discretized, form $c_{ijk,it}$ where it is used as index for the age and i, j, k are indices for the longitude, latitude and elevation above terrain level (a.t.l.).

Aerosols are removed in FLEXPART by dry and wet deposition. The dry deposition is simulated according to a resistance approach (Wesely and Hicks, 1977) considering the aerodynamic resistance, the quasilaminar sublayer resistance, and gravitational settling (Slinn, 1982). For the aerosols considered in our simulations (see below), we assumed a logarithmic diameter distribution with geometric mean diameter 2.5×10^{-7} m and geometric standard deviation 1.25, and a density of 1.4×10^3 kg m⁻³. Wet deposition distinguishes between in-cloud and below-cloud scavenging (McMahon and Denison, 1979; Hertel et al., 1995). Scavenging coefficients are determined primarily as a function of cloud liquid water content and precipitation rate, respectively. Notice that FLEXPART has no aerosol processing scheme and details such as the conversion of some aerosols from an initially hydrophobic to a hydrophilic state (Vignati et al., 2010) are ignored.

2.2 Megacity emissions and deposition time scales

The emission inventory is extracted from the original Representative Concentration Pathway (RCP) 8.5 emission scenario for the year 2005 developed by the MESSAGE (Model for Energy Supply Strategy Alternatives and their General Environmental impact) modeling team and IIASA (Integrated Assessment Framework at the International Institute for Applied Systems Analysis), Austria (see <http://www.iiasa.ac.at> and

Riahi et al., 2007). The global emission data used here has a $1^\circ \times 1^\circ$ resolution and was obtained from the original data set at $0.5^\circ \times 0.5^\circ$ by T. M. Butler (personal communication, 2010) and provided to the MEGAPOLI project along with a mask defining the megacities' extensions. Any simulated megacity occupies one or more grid cells of this emission data set as listed in Table 1 and illustrated in Fig. 1 and it is possible to see that both single-grid-cell as well as multiple-grid-cell megacities are included. From the RCP 8.5 data set, the emissions of several compounds can be extracted.

A first estimate on the relative importance of megacity emissions on a global scale can be deduced immediately from Fig. 2, where the cumulative emission contribution for several chemical species (black carbon, BC; carbon monoxide, CO; nitrogen oxides, NO_x; sulfur dioxide, SO₂) for all the megacities up to a given degree of latitude is scaled with the total cumulative species emission (including non-megacity emission) to the same degree of latitude. For example the total cumulated emission, starting from the

South Pole and up to a specific latitude i_{lat} , could be written as, $CE_{i_{lat}} = \sum_{j=-90}^{i_{lat}} \sum_{i=1}^{360} E(i, j)$, where $E(i, j)$ is the emission in grid cell i, j . Similarly, the cumulative contribution for all the cities starting from the South Pole to latitude, i_{lat} , could be written as,

$$CEMC_{i_{lat}} = \sum_{i_{city}=1}^{N_{city}} \sum_{j=-90}^{i_{lat}} \sum_{i=1}^{360} E(i, j) \delta(j - i_{lat}(i_{city})) \delta(i - lon(i_{city}))$$

N_{city} is the number of cities considered and δ is a function with a value of one within the city limits and zero outside.

The data have been separated in Northern and Southern Hemisphere (NH and SH) and the cumulative contribution is calculated starting from the North and South Poles respectively. No data are shown south of the latitude of 35° S and north of the latitude of 60° N since there are no megacities. Further, the data are separated into autumn/winter emissions, and spring/summer emissions. It can be seen that the 35° S latitude megacities in the SH contribute more to total emissions than their counterparts

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in the NH. In the NH, there is a strong seasonal variation at higher latitudes especially for CO and BC, which is due to intense open biomass burning in spring and summer, so that megacity emissions contribute less in summer. In the autumn and winter months, a single city (St. Petersburg) accounts for almost 10 % of the total emissions north of about 60° N for the BC considered in this study. It can be of interest to note that adding to the list of cities an emitting area extending from a lower left corner of (59° N, 10° E) to an upper right corner (61° N, 26° E) the emission fraction goes up to 36 % of the total emission over 60° N for BC in autumn and winter. This area contains five large urban areas in the Baltic and Scandinavian countries: Tallinn (Estonia), Helsinki (Finland), Stockholm and Uppsala (Sweden), and Oslo (Norway). Since on the scale of this study these cities are geographically close to the investigated megacity of St. Petersburg, the results obtained for this latter city could be of guidance for the behavior of emissions that in the autumn and winter contribute 36 % of the total world emissions north of 60° N. These cities' contribution is lower in the summer but considering that during these months the concentrations of BC in the Arctic lower atmosphere are lower due to stronger deposition processes (e.g. Hirdman et al., 2010), the winter peak of these urban emissions could be significant.

The further discussion in this paper will be based only on the emissions of one compound, BC, which we consider as representative for all combustion-related emissions. This compound will be used for two different scalar tracers: one considered as a conserved tracer (BC_{tr}) and not subject to any removal process and another one (BC_{dp}) with dry and wet deposition processes included in the simulations.

For studying the impact of pollution on human population we used the GPWv3 data set of the Center for International Earth Science Information Network (CIESIN), Columbia University; available at <http://sedac.ciesin.columbia.edu/gpw> (January 2011). The population of a city was extracted, from the GPWv3 data set, using the same megacity mask used for the emission; the resulting population for the cities is listed in Table 1.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Before proceeding with a detailed analysis of the plume characteristics it is useful to explore the effects of the modeled wet and dry deposition processes on the suspended mass in the atmosphere as a function of time after emission (“age”). The deposition processes reduce the mass content significantly with age but differently for the different cities considered. In most figures in this paper, megacities have been divided in NH and SH megacities and the results of these two groups will be in general plotted separately to improve the clarity and readability of the figures.

Figure 3a, b shows the three year averaged concentration (mid December 2002 to mid December 2005, see Sect. 3) of BC_{dp} normalized for that of BC_{tr} in any age class on a logarithmic scale, while Fig. 3c, d shows the same ratio on a linear scale. This indicates how dry and wet deposition processes deplete the atmospheric mass of BC_{dp} with time. From Fig. 3c, d it is apparent how initially deposition processes are more active for European and tropical cities. However, after a few days the plumes originating from megacities located at midlatitudes in Asia are those characterized by the higher deposition rates, related to strong scavenging in the North Pacific stormtrack (Fig. 3a, b). For the last few age classes, European and North American plumes have again higher deposition rates. This complex behavior reflects the different three dimensional distributions of the plumes originating from different geographical regions. For very large ages, all the plumes should converge towards the same deposition rates typical of the Northern and Southern Hemisphere, while initially the deposition rate is strongly influenced by the local processes close to the source. This can be better appreciated defining two time scales obtained by fitting an exponential decay to the data on a logarithmic scale, T_{10} (based on Fig. 3a, b) and to the data on a linear scale, T_{li} (based on Fig. 3c, d). These time scales are reported in Table 1. T_{10} is an estimate of the average e-folding deposition time scale for the 48 days time window and the source location. T_{li} gives a good idea on how fast the mass emitted from a specific source is depleted from the plume and it strongly depends on the local processes in the boundary layer close to the source. T_{li} varies between one and ten days while T_{10} varies between about six

and nine days, but with most of the cases between about six and eight days, indicating the approximate global annual e-folding lifetime of the BCdp aerosols in FLEXPART.

An important feature of including age classes in the analysis is that many of the results reported here involving a mass assigned to the age classes, can be extended to consider the effect of a superimposed decay process (e.g. radioactive decay, or chemical destruction) just applying, a-posteriori, the decay – e.g. exponential characterized by an assigned time scale. This is true for both BCtr and BCdp. In the first case the superimposed decay is applied to a substance not undergoing deposition, in the second case to a substance undergoing wet and dry deposition.

3 Dispersion characteristics of transported plume

In this section we start the discussion about the megacity plumes with an analysis of their physical dispersion characteristics both horizontally and vertically. To characterize the horizontal dispersion of the plumes, the model output was first averaged over the time period of interest (e.g. three winter seasons or all three years)

$$\tilde{c}(x, y, z, t) = \frac{1}{N} \sum_{i=1}^N c_i(x, y, z, t)$$

Here N indicates the number of 5 days output intervals in the period of interest, x , y , z are the three space dimensions, and t indicates the elapsed time from the release (i.e. the age). From this time averaged concentration a vertically integrated concentration field was introduced,

$$\tilde{c}_z(x, y, t) = \sum_{k=1}^{Nz} \tilde{c}(x, y, z_k, t) \Delta z_k$$

Here Δz_k indicates the vertical grid level extension and Nz ($= 25$) is the number of vertical levels in the FLEXPART output grid. From this field an advection measure was

26360

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



obtained as the average distance of the mass from the source location, $\bar{d}(t)$, in the following called transport distance,

$$\bar{d}(t) = \frac{\sum_{i=1}^{360} \sum_{j=1}^{180} d_{ij}(\tilde{c}_z(t))_{ij} \Delta_{ij}}{\sum_{i=1}^{360} \sum_{j=1}^{180} (\tilde{c}_z(t))_{ij} \Delta_{ij}}$$

where d_{ij} is used to indicate the distance, on the Earth sphere, of the center of the grid point with indices (i, j) from the source position. Since some sources are composed of more than one single grid cell element, the source position was defined as the grid element containing the maximum ground level concentration averaged over the extension of the first age class (i.e. $0 < t < 1$ day). Δ_{ij} is the area of a grid element with indices (i, j) , and $(\tilde{c}_z)_{ij}$ is the concentration associated with the grid element. It should be noticed again that, since the grid extends also in the time dimension, any of these measures is defined as a function of age for the intervals defined above. In a similar manner, the averaged vertical position, $\bar{h}(t)$, of the horizontally integrated concentration $(\tilde{c}_{xy})_k$ as a function of time can be introduced, where k is used for the vertical grid cell index.

The results for the distance from the source location, $\bar{d}(t)$, for BCtr are shown in Fig. 4a and c, respectively for the NH and the SH and those for BCdp are shown in Fig. 4b and d. The smaller left panels are a zoom into the age classes 0–1 and 1–3 days and use a logarithmic scale for the plot. For BCtr in the NH, it can be seen that initially the plumes subject to the fastest advection are those emitted at higher latitude and associated therefore with European and North American megacities, while the plumes emitted by tropical or sub-tropical megacities have the slowest initial transport. However, the situation changes, and for elapsed time greater than 10 days all the higher latitude cities show comparatively short transport distances. On the contrary, the tropical and sub-tropical megacity plumes then become associated with the largest

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



transport distances. Asian megacities of intermediate latitude (blue symbols) have in general intermediate characteristics. The reason for this latitude-dependent behavior is influenced by three factors: (1) the plumes are transported preferentially zonally before spreading meridionally due to latitudinal transport barriers, and the length of a circle of latitude is largest at the equator, (2) cities at lower latitudes have a larger asymptotic distance possible for a uniform scalar distribution on the hemisphere and (3) the tropics are characterized by lower near-surface wind speeds but a high frequency of deep convection, so that the dispersion is at first slower than for higher-latitude cities but after a few days significant amounts of tracer in the tropics are transported by deep convection to the upper troposphere, where winds are faster. This is illustrated in Fig. 5a, c, which shows that tropical cities and particularly south-east Asian cities (e.g. Manila) are associated with a very fast increase of the plume average elevation.

The zoom figures for the horizontal transport behavior between 1–3 days (Fig. 4), show that in the NH the cities with the initial fastest plume transport are London, New-York, Osaka and Tokyo. These are all cities close to the ocean which is therefore an important factor in determining a fast early phase of plume transport. Among the European megacities the Po-valley shows a unique behavior since the tracer is trapped there by mountains on three sides, which strongly limits the early phase of transport. One consequence of this is the high pollution levels usually registered in this area (e.g. Bigi et al., 2012). However, overall in the NH the megacity with the slowest horizontal transport up to 10 days is the tropical city of Lagos which also has severe air pollution problems (Baumbach et al., 1995).

Removal processes lead to a reduction of BC mass in the atmosphere. The lifetime T_{li} of the BCdp tracer for the various megacities ranges from 1.0 days for Jakarta to 10.5 days for Cairo (Table 1), reflecting the very different efficiencies of wet scavenging in tropical and desert regions. Removal processes also change the transport characteristics of the BC plumes and these changes are sometimes difficult to understand (Fig. 4b, d). For example, the preferential removal of mass during upward transport associated with precipitation means that initially the vertical transport of BCdp is slower

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

compared to BCtr (Fig. 5). As a consequence, initially the horizontal transport is dominated by the slower low-level winds, leading also to shorter horizontal transport distances of BCdp compared to BCtr, see left panels in Fig. 4. After a few days, however, due to preferential wet scavenging in the lower troposphere versus lack of precipitation in the upper troposphere and lower stratosphere, for the remaining BCdp the fraction located in the upper troposphere and stratosphere increases. Thus, the centroid altitude of the BCdp tracer increases throughout all remaining age classes (Fig. 5b, d) and the horizontal transport distance also catches up with that of BCtr (Fig. 4b). This is summarized in Fig. 6a, b which shows the ratio, for BCdp versus BCtr, of the horizontal

transport distances ($D(t) = \sum_{i=1}^{N_{\text{cities}}} \bar{d}_i(t) / N_{\text{cities}}$) and of the average megacities plume

altitude ($H(t) = \sum_{i=1}^{N_{\text{cities}}} \bar{h}_i(t) / N_{\text{cities}}$) for all cities. Notice that these results for $H(t)$ do not

imply that a larger fraction of the total emissions can reach the upper troposphere for BCdp than for BCtr. However, after some time a larger relative fraction of the remaining BCdp than of the remaining BCtr resides in the upper troposphere, as demonstrated in Fig. 7, which shows the fraction of the total BCtr and BCdp mass suspended in the atmosphere which is contained between 9 km and the top of our sampling domain (30 km). For BCdp, this fraction increases throughout the 48 days considered in our study and for some cities it approaches the value of 1 (Fig. 7b and d).

The SH megacity plumes have a similar behavior compared to their NH counterparts. Midlatitude megacity plumes show initially a faster horizontal transport while ultimately the tropical and subtropical latitude megacity plume have a larger transport distance. However, the transport for mid latitude cities is here faster than in the NH and the plumes reach faster their hemispheric asymptotic distance. In fact Fig. 4c shows that for BCtr all the plumes but the Lima and Jakarta plumes are transported more rapidly compared to the NH plumes. This behavior is likely related to the stronger westerlies in the SH. Jakarta shows a typical behavior of a tropical megacity as previously discussed

for the NH, while Lima shows the slowest horizontal transport speed of all the megacity plumes investigated.

4 Statistical measures of regional pollution export

L07 defined several metrics to characterize the dispersion of the megacity plumes and their potential impact on a regional scale. Here we will calculate within our modeling framework two of their metrics of pollutant Export over Long-Range (ELRcol and ELR1km), and compare our results with theirs. ELRcol is defined as the fraction of the total BC mass in the atmosphere which is more than 1000 km away from the source at any altitude. ELR1km is the same as ELRcol but for the mass contained in the first 1 km a.t.l. Contrary to L07 we considered only whole grid cells and did not split grid cell according to their fractions within or beyond the 1000 km radius from the source, however, our grid has finer resolution (1° versus 1.9°), which compensates for this less accurate treatment. Also, the simulations of L07 were for the year 1995 while our simulations are for the years 2003–2005. Also, as before, we will investigate the effect of the deposition thus extending the results of L07.

Our age class treatment generalizes these metrics to any decay time within the 48-day range considered. The ELR measure has been calculated as cumulated mass exported up to a specific age normalized with the total mass of the substance suspended in the atmosphere cumulated up to the same age. This means that the value of the last age class corresponds to the total exported mass fraction.

L07 report in detail the ELR1km values for a scalar with a ten days decaying time scale. Figure 8a shows that the value of ELR1km for BCtr becomes, after about one week, almost constant for most cities. This means that all tracers with a decay time scale larger than about one week will behave similarly to a purely passive tracer without decay. This simply reflects the partitioning of tracer between the lowest 1 km and the rest of the atmosphere, as basically all of the tracer will be exported beyond 1000 km after a week. However, using the discretized time dependence information present in

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacitiesM. Cassiani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the age class partitioning we can explicitly apply an exponential decay process with 10 day time scale to allow a more precise comparison of our results to those of L07. Table 2 compares the ELR1km results for the 48 days age class for a conserved tracer with those for a tracer with ten days lifetime (BC10). The results of L07 for BC10 are also reported. It is possible to see that the difference between BC10 and the conserved tracer, BCtr, is minor in most of the cases. However, in general and as expected, the results obtained for the decaying scalar agree better with the results reported by L07. Considering that the results of L07 were for a different year, used different meteorological input data (NCEP reanalysis) and a different transport model (MATCH), the agreement between the two studies is very good. The values of ELR1km obtained in our study range from 6.3 % to 30.1 % compared with 3.2 % to 34 % in L07.

A scatter plot (Fig. 9) shows that the ELR1km values for European, North American and mid latitude Asian cities agree reasonably well in the two studies. Southern Hemisphere megacities have generally higher values in the present calculation based on FLEXPART. The worst agreement is achieved for Indian subcontinent, Asian tropical, and South American megacities, although the differences are not systematic. Jakarta stands out with ELR1km values of 7 % and 3.2 % in FLEXPART and MATCH, respectively. This difference must be related to the different convection schemes used in FLEXPART (Emanuel and Zivcovic – Rothman scheme, implemented by Forster et al., 2008) and MATCH (plume ensemble scheme, proposed by Lawrence and Rasch, 2005). The low fraction in MATCH means that in this model the BC10 tracer must have been vented out of the boundary layer very efficiently, with only small compensating downward fluxes. This result is obtained, in somewhat less extreme form, also for other tropical cities such as Manila, Bangkok, San Paulo and Rio de Janeiro. On the other hand, FLEXPART has lower ELR1km values than MATCH for the cities on the Indian subcontinent. This is probably due to different representation of the winter monsoon in the two models, where FLEXPART shows a relatively rapid low-level southward transport to the intertropical convergence zone, followed by rapid lifting (Stohl et al., 2002).

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Figure 8b reports the metric ELR1km computed for BCdp. The export efficiency for the BCdp tracer depends not only on the combined action of horizontal wind and vertical transport, but also on the removal rates. In general the effect of deposition is to reduce the value of ELR1km and the reduction is stronger for higher local deposition rates, i.e. low value of T_{ji} . However, the megacities in the Indian subcontinent plus the city of Cairo have a noticeable increase of ELR1km for BCdp with respect to that of BCtr. A possible mechanism for this happening is if precipitations act mainly on rising air masses outside the 1000 km radius from the source. This is consistent with the winter monsoon behavior for the Indian subcontinent, as described above. The city of Cairo is instead characterized by an arid local climate (it is the city with the highest T_{ji}) and therefore, since there are very low precipitations close to the city, it is plausible that precipitations act mainly on rising air masses outside the 1000 km radius. In general this additional factor adds variability to the possible range of ELR1km for BCdp, with values spanning about a factor of ten, compared to only a factor of four for BCtr.

The two lowest total values of ELR1km for BCtr and BCdp are found for Jakarta and Lagos (see Table 2) and it is interesting to examine why. Lagos is the city with the slowest horizontal transport of the plume during the first 3 days after tracer release (Fig. 4a, b, left). Jakarta on the other hand is the second slowest in the range $3 < t < 14$ days (Fig. 4a, b, right). This difference explain also why Jakarta has an initial peak in the value of ELR1km for BCtr (Fig. 8a). When deposition is accounted for, Jakarta is the city with the shortest T_{ji} and therefore the deposition generates a stronger decrease of ELR1km compared to that of any other city (including Lagos). As a consequence the difference between the two lowest values of ELR1km becomes more marked for BCdp. It is also interesting to discuss why Mexico City has initially the lowest value of ELR1km and why it has a quite low overall ELR1km, despite having a quite fast overall horizontal transport (Fig. 4a, b, right). This city is characterized by slow initial transport ($t < 1$ day, see Fig. 4a, b) and it has the highest initial plume elevation among all cities (Fig. 5), as a consequence, the plume is initially transported slowly and subsequently, when the transport becomes faster, it is mostly transported at an elevation above 1 km a.t.l. The

two top low level exporters irrespective of deposition are the cities of Moscow and Saint Petersburg and these are cities with a quite fast initial transport (Fig. 4) but also and especially very low average elevation (Fig. 5).

As previously reported by L07, results for the total mass exported to a distance of more than 1000 km (ELRcol) are considerably less variable. In particular for the 48 days age class we find a very narrow range of values from 85 % to 95 % (Fig. 10). This is expected since in this case the values should converge towards the same asymptotic limit of 98.8 % for a scalar well mixed in the hemisphere, where an area bounded by the 1000 km radius covers 1.2 % of the hemisphere. The values of ELRcol for BCtr in the two weeks age class goes from about 60 % to 85 %. This is indicative of the value of a scalar subject to a superimposed decay of a similar duration, and indeed this range is quite similar to the cumulated value of ELRcol for BC10 which ranges from about 59 % to 82 %. The range of values of BC10 is again in good agreement with the results of L07 for the ten days decaying scalar for which they reported a range from 62 % to 84 %.

It is interesting to note that Lima, which has clearly the lowest value of ELRcol for any age, was not among the cities having a very low ELR1km. This city has an extremely slow horizontal transport (Fig. 4) which is perfectly reflected in the low ELRcol value. However, contrary to Jakarta and Lagos that have a very low ELR1km, Lima is not characterized by a very fast uplifting of the plume (Fig. 5) and this increases the value of ELR1km. The deposition modifies again the picture by widening the range of values. However, in this case the interpretation is simpler with respect to ELR1km since deposition systematically decreases the value of ELRcol. The cities with the highest values of ELRcol are those characterized by a high value of T_{ij} and Cairo (which is the city with the highest T_{ij}) becomes the top exporter. The fact that Cairo was not the top low level exporter for BCdp underlines again that for ELR1km it is very important that the plume is transported at low elevation, which is not the case for Cairo.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Air pollution and BC deposition in the polar regions

Analysis of BC concentration in the polar regions is important due to the positive radiative forcing of this compound both in the atmosphere and when deposited on snow or ice and the high efficacy of this forcing for climate warming (e.g. Hansen and Nazarenko, 2004; Flanner et al., 2009). Our simulations allow us to quantify the efficiency with which BC originating from megacities in different parts of the world can reach the polar atmosphere and deposit on the ice/snow cover. This study is somewhat complementary to the works of Stohl (2006), Stohl and Sodemann (2010) and Hirdman et al. (2010) who studied the possible sources of BC in the Arctic and Antarctic atmosphere from a receptor-oriented perspective. Here, we take a source-oriented view by following the emissions from the world's megacities.

5.1 Air pollution

Figure 11a, b shows the total BC_{tr} and BC_{dp} mass in the lowest 1 km a.t.l. of the NH polar atmosphere (north of 66° N), scaled for the mass emitted in the respective age class interval, while Fig. 11c, d provides the same information for the SH (south of 66° S). As discussed extensively in Stohl (2006), the contribution to mass in this layer in the NH depends mainly on the latitude of the source, with largest contributions from northern Eurasia. This is confirmed here for the BC_{tr} results. Specifically, we find that releases from Saint Petersburg give the largest fractional contributions of all megacities and for all age classes. As much as 10 % of the emissions can be located in the Arctic already as early as about 2 days after the emission. In fact, the contribution decreases after about 2 weeks, as the Saint Petersburg plume is initially preferably transported to the north and subsequently disperses into the entire NH. Similar behavior can be expected for pollution export from other Baltic and Scandinavian cities that are relatively close to Saint Petersburg and jointly contribute up to 36 % of the total BC emissions north of 60° N. This group of cities is consequently likely to be the largest “localized” contributor of anthropogenic pollution in the lower layer of the Arctic atmosphere for substances

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with lifetimes on the order of days to a few weeks. Furthermore, while Moscow makes a smaller BC_{tr} fractional contribution to the pollution in the Arctic atmosphere during the first 3 days after emission, the contributions at later times are almost identical to those of Saint Petersburg. The age class sampling allows appreciating that, for any substance with a decaying time scale of less than about 10 days, the contributions from non European cities are very small. Only after this elapsed time, contributions from North American megacities start to be more relevant and soon after also from East Asian mid latitude cities. Contributions from tropical locations are even for the last age class considered (48 days) an order of magnitude smaller than contributions from Saint Petersburg. Table 3 (column 1 and 3) reports the cumulative amount of mass irrespective of the age classes and expressed both as mass and as a percentage of total emitted mass. The table confirms that cities in Europe have significantly larger fractional contributions but in terms of absolute numbers (i.e. taking into account also the different emission strengths), New York is the largest contributor and Tokyo, Seoul and Beijing have numbers comparable to those of most European cities. However, we remind the reader that, as shown by the age class plots, these cities have smaller contributions relative to European cities when the lifetime is significantly shorter than 48 days.

When taking into account deposition processes (BC_{dp}), the fractional contributions from all cities become much smaller (Fig. 11b). However, the reductions are particularly large for non-European cities and, thus, significant contributions to the BC_{dp} loading in the lowest 1 km of the Arctic atmosphere are confined to Northern European sources with small contributions from North American cities. Saint Petersburg and Moscow give similarly high contributions in the age classes going from 2 to 20 days, although both London and Ruhr valley are also relevant. The cumulative numbers reported in Table 3 (column 2 and 4) show that, for BC_{dp}, all the megacities in Asia contribute less than the sole city of Saint Petersburg and that New York City has a similar contribution to Saint Petersburg despite having almost 20 times the population.

possible to see how the Po-valley stands out as the largest contributor to BCdp in the Arctic stratosphere.

The current results reporting a contribution mainly from European megacities to the mass of BC in the lowest kilometer of the Arctic atmosphere and a progressively increasing contribution of North American and East Asian megacities when moving up to the free troposphere and the stratosphere is in agreement with the findings of Hirdman et al. (2010), which reported a substantial contribution to BC measurements at lower altitude stations only from northern Eurasian sources, while significant contributions from more remote sources (e.g. East Asian) was observed for measurements on top of the Greenland ice sheet.

5.2 Deposition

For quantifying BC deposition in the Polar Regions we consider the accumulated mass over an area of interest during a period (i.e. year, season). When normalized quantities are shown this is done considering the total mass available for deposition during the period considered, given by the initial mass suspended in the atmosphere at the beginning of the period plus the mass emitted during the period. Figure 13 shows the normalized (left panels) and un-normalized (right panels) deposition north/south of 66° N/66° S within 48 days from the release. The values cumulated over all age classes are reported in Table 3. Saint Petersburg has the largest fraction of its BCdp emissions deposited in the Arctic (9.4 %), followed by Moscow (5.6 %) and the Ruhr valley and London (both about 4 %). However, the absolute deposition in the Arctic is comparable for these cities and also the Po-valley and Paris cause significant deposition values. The North American megacities of New York and Los Angeles give normalized deposition values that are more than an order of magnitude smaller, but because of the larger BC emissions of these American cities, the actual deposition values are only about 2 to 3 times lower than the contributions of European megacities. Absolute and relative contributions from megacities on other continents are much lower. This means, as shown in Table 3, that the sole city of Saint Petersburg (population 2.6 million) produces a

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BCdp deposition in the Arctic that is marginally larger, than the BCdp deposition from all the megacities in Asia (population roughly 500 million). This shows again that most of the Arctic BC deposition from megacities must be attributed to northern Eurasian sources.

5 The BCdp deposition over land during the winter season for latitudes north of 40° is also reported in Table 3 (columns 7 and 8). During winter, a significant fraction of land above this latitude may be covered by snow and the BC deposited may have an influence in accelerating the melting of the snow in spring (e.g. Quinn et al., 2008; Flanner et al., 2009). For this region, the fractional contributions from European and North
10 American megacities are comparable, and the Asian megacities of Beijing, Tianjing, and Teheran give significant fractional contribution followed by Cairo in North Africa. Similar considerations are valid for the absolute mass deposition and in this case New York City emerges as the largest contributor to deposited BCdp.

5.3 The Southern Hemisphere

15 Although there are just eight megacities in the SH, they represent about 12 % of the total emission (see Fig. 2) in the SH from the south pole to a latitude of 25° S. The export of BCtr and BCdp from SH megacities to the Antarctic region is much smaller than from NH megacities to the Arctic, and even export efficiencies relative to the emission strength are lower. In fact, due to the generally longer distances of the SH megacities
20 to the Antarctic region, the fraction of BCtr mass transported to the lower layer of the Antarctic atmosphere (0–1 km) reaches maximum values of 2×10^{-2} (Fig. 11c) in the last age class. Values for ages of less than about 20 days are lower and there is no peak in the early age class, again due to the considerable distance of the sources from the polar region. Moreover, these values are reached only for Buenos Aires and Sidney while the other megacities and especially Lima and Jakarta, give substantially
25 lower contributions. For the depositing tracer BCdp, no more than a fraction of 10^{-4} of the emitted mass in any age class, reaches the Antarctic atmosphere (Fig. 11d). Summary numbers, cumulating the contribution from all age classes, are reported in

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3 for both BCtr and BCdp. They show that for BCtr the total megacities emissions in the SH generate a suspended mass in the Antarctic atmosphere comparable to that produced by the single city of St. Petersburg in the Arctic. If deposition is furthermore taken into account, the suspended mass is entirely negligible with less than 100 kg of suspended mass overall.

The fraction of BCdp mass depositing south of 66° S is as well very low. Our simulations estimate a total annual deposition of BCdp in the Antarctic by all the megacities of about 45 tons (see Table 3), and most of it is coming from just four megacities, Sidney, Johannesburg, Santiago and Buenos Aires, with a marginal contribution from San Paulo and Rio de Janeiro. The comparison with the total deposition of 2330 tons of BCdp from NH megacities in the Arctic shows that this deposition is very small, also considering that the total emission from NH megacities is just about 5.5 times the SH megacity emissions. However, this 45 tons of deposition is comparable to the local emissions, below 60° south, of BC from shipping (100 tons) as reported by Stohl and Sodemann (2010) based on the emission inventory of Bond et al. (2007), with tourist ships contributing about 20 tons (Stohl and Sodemann, 2010). Also, if we consider the 12% megacity emissions representative of all SH emissions south of 25° S, we can estimate that less than 400 tons of BCdp are deposited in the South Pole region from all continental sources.

6 Exposure of human population

In this section, population (P) and annual mean ground level concentration (GLC) maps are combined to derive simple but quantitative measures of the exposure of the human population to air pollution from megacities. Particular focus will be on the relative importance of population exposure inside and outside the megacity. We use the GLC multiplied by population (CmP) as a measure of total population exposure to air pollution and define this metric as a function of time (i.e. for any age class),

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$CmP(t) = CmP_{it} = \sum_{\text{all } ij} P_{ij} \sum_{l=1}^{it} GLC_{ij,l}.$$

Here i , j are grid indices, indicating that the metric is integrated over the entire globe, and it is the index of the age class considered. Therefore, CmP is a function of time since emission and the concentration used is given by the summation of the concentration associated with the current and the previous age classes. The values computed for the 48 days age class are reported in Table 1 for both BCtr and BCdp. Notice that since the lifetime of BCdp is substantially shorter than 48 days for all cities, the metric measures the total impact of BCdp. The table shows that the largest values of CmP are reached for the megacities in Asia, with Dhaka setting the records of 149 and 104 ($10^{10} \text{ ng m}^{-3} \text{ people}$) respectively for BCtr and BCdp. The first non Asian megacity is the extended New York area which ranks seventh, with values of 40 and 22 for BCtr and BCdp, while the first-ranking European city is Moscow with CmP values of 19 and 14.

Figure 14 shows the values for all the age classes, and it can be seen that when deposition is considered the CmP exposure metric for BCdp (Fig. 14, right) is quite flat and determined mainly by the exposure on the very first day. This is due to a combination of three factors: (i) the highest population density at the location of the release (i.e. within the megacity), (ii) all BCdp is initially located near the surface, and (iii) the quick loss of BCdp mass with time. This suggests that most of the population exposure to BCdp occurs within or very close to the megacity. Also for the conserved tracer BCtr, CmP values do not grow substantially after the first day (Fig. 14, left), suggesting that even for long-lived pollutants most of the population exposure occurs in the city or its vicinity. This is also the reason why the CmP values for BCtr and BCdp are quite similar for most cities, suggesting that population exposure is not very sensitive to a substance's lifetime, if it is longer than a day.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In Table 1 we also report the population weighted concentration for the city (PWC_{city}) for BCtr and BCdp:

$$PWC_{\text{city}} = \frac{1}{P_{\text{city}}} \sum_{ij \in \text{city}} P_{ij} \sum_{l=1}^{Na} GLC_{ij,l},$$

i.e. the same as the CmP metric but computed only for the grid cells defining each megacity, with GLC integrated over all the age classes ($Na = 10$), and normalized with the megacity's total population (P_{city}). This metric measures the concentration to which the population inside the city is exposed to, on average. Some interesting features arise when comparing the megacities. The highest value of PWC can be found in Asian megacities with Seoul having value of 1621 and 1535 ng m^{-3} respectively for BCtr and BCdp. The lowest values are registered for Istanbul, respectively 64 and 61 ng m^{-3} for BCtr and BCdp. It should be noted that for cities covering only one cell this value is the same as the maximum concentration in the domain, while it can be significantly lower for larger megacities.

A further parameter of interest is CmP normalized by CmP obtained when integrating only over the megacity grid cells, called here CmP_c . This metric (CmP/CmP_c) measures the total population exposure relative to the exposure inside the city. These values are plotted as a function of the age in Fig. 15. First, we note that for the non depositing tracer BCtr, several cities have an impact on surrounding regions which is at least half as large as on the population of the city itself (i.e. $\text{CmP}/\text{CmP}_c > 1.5$), but few cities have an impact outside the city's boundaries larger than the impact inside the city boundary (i.e. $\text{CmP}/\text{CmP}_c > 2$) and only three cities reach this thresholds if deposition is taken into account (Fig. 15, right panel). For BCtr, Saint Petersburg shows the largest CmP/CmP_c values but only for ages greater than 10 days. This is due both to the fact that it is surrounded but not closely surrounded, by relatively densely populated areas and that it has the smallest population among the cities considered (thus, reducing the impact inside the city boundaries). Southern Hemisphere megacities show generally somewhat lower CmP/CmP_c values, with minima obtained for Buenos Aires and San

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Paulo, since these are densely populated areas surrounded by relatively sparsely populated areas and are close to the ocean. In the NH, Tokyo stands out with the lowest value of the ratio, since most of its pollution is exported to the ocean. If deposition is accounted for (BCdp) Sydney also has a very low CmP/CmP_c ratio while the cities showing highest values of CmP/CmP_c are the densely populated Chinese megacities of Tianjin and Beijing, which are actually surrounded by regions with extremely high population density outside the city boundaries, and the city of Karachi. This last city shows high impact only after about 5 days, thus this should be due to the Karachi's plume impacting on densely populated Indian urban areas. It is interesting to note that the cities with high CmP/CmP_c ratio also have a quite high index of low level BCdp export (ELR1km, Fig. 8b).

In general and especially for the BCdp tracer, it seems that the impact of megacities on population outside their boundaries relative to their impact inside city boundaries is low, with only three of the megacities (Tianjin, Beijing and Karachi) showing an impact comparable or greater than the impact inside the cities, while most of the cities have $CmP/CmP_c < 1.4$. It is also of some interest to note that based on the time related behavior of this index one could speculate whether the impact on population is transnational. For example when deposition is involved the Chinese megacities of Tianjin and Beijing have an index $CmP/CmP_c > 1.6$ already in the first age class (less than a day) and this value does not grow substantially with age. Therefore, most of their influence is on national population. On the contrary, Karachi reaches a similar index only in the second age class (3 days) so it is probably related to transboundary pollution transport.

Based on the age class spectrum it is also easily possible to extrapolate the behavior for decaying scalars but in case of BCdp any superimposed decay with characteristic time greater than about ten days would basically leave the results unchanged.

7 Summary and conclusions

A study of the characteristics of air pollution plumes dispersing from a group of thirty-six of world's megacities has been performed using the Lagrangian particle model FLEX-PART. The study focuses on plumes generated by BC emissions. BC was chosen since:

(i) this is a compound well suited for a Lagrangian investigation because it can be represented in a linear modeling framework, (ii) it is representative of combustion-related emissions and (iii) it is an important air pollutant and short-lived climate forcer. Based on BC emissions two artificial tracers were modeled, a purely conserved tracer with a cut-off lifetime at 48 days – BC_{tr} – and one resembling more closely the properties of true BC and subject to wet and dry deposition – BC_{dp}. Four aspects of megacities' plumes have been investigated: (1) physical dispersion characteristics, e.g. transport distance and average altitude, (2) pollution export in the lower troposphere over more than one thousand kilometers (ELR1km), (3) concentrations and deposition of BC in the polar regions and (4) exposure of human population with a focus on comparing the inside-the-city versus outside-the-city exposure values.

Although local conditions can be very important the main factor influencing the physical dispersion characteristics was found to be the latitude of the emitting cities; initially, plumes from higher-latitude cities travel faster but, after about three days, the higher elevation reached by the plumes in tropical/subtropical latitude generated a complementary faster horizontal advection. The proximity with the ocean was found to be an important factor in determining a fast early phase of plume transport. Among the European megacities the Po-valley shows a unique behavior since the tracer is trapped there by mountains on three sides, which strongly limits the early phase of transport but, overall, in the NH the megacity with the slowest horizontal transport up to 10 days is the tropical city of Lagos. However, Lima is the city showing the slowest horizontal transport.

It was found that the cities showing high level of regional pollution export in the lower troposphere (ELR1km) are generally high latitude cities since these are characterized

ACPD

12, 26351–26400, 2012

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

by an initially fast transport and by a low plume elevation. The influence of deposition was mainly to reduce the value of ELR1km and this reduction was found to be stronger for cities with low values of the local characteristic deposition time scale T_{li} . However, few exceptions (Indian subcontinent megacities and Cairo) were observed where the value of ELR1km increased due to the effect of deposition. This behavior could be explained with the possibility of low level transport at distance larger than 1000 km followed by a preferential deposition in uplifting air masses. Noteworthy a good agreement between our simulations and previous results by Lawrence et al. (2007) for ELR1km was observed for a non depositing scalar with a ten days decaying time scale.

The contribution to lower tropospheric concentrations and deposition in the Arctic must be mainly attributed to northern European megacities (St. Petersburg, Moscow, London and Ruhr Valley). The sole city of St. Petersburg (population about 2 million) contributes up to 10 % of the BC emission north of 60° N during winter and when deposition is considered (BCdp) its contribution to suspended mass of BCdp in the Arctic atmosphere and deposition on the Arctic surface is larger than that of the whole group of Asian megacities (population about 500 millions). When considering the upper layer of the arctic troposphere (2–9 km) and stratosphere (9–30 km) the contribution of remote sources become progressively more important. In the stratosphere the contribution from Asian megacities is the largest, followed by those from North American megacities and European megacities with the Po-Valley standing out among this latter group. However, all the megacities can be considered significant for pollution in this layer of arctic atmosphere.

The export of BCtr and BCdp from Southern Hemisphere megacities to the Antarctic region is much smaller than from Northern Hemisphere megacities to the Arctic, and even the export efficiency relative to the emission strength is lower. Our simulations estimate a total annual deposition of BCdp in the Antarctic by all the Southern Hemisphere megacities of about 45 tons, compared to the total deposition of 2330 tons of BCdp from Northern Hemisphere megacities in the Arctic. Note that the contribution by

all megacities to the deposition in the Antarctic is, therefore, estimated to be smaller than the emissions coming from the local shipping activities.

Regarding the exposure of the human population, we found that for the passive tracer Bctr ten megacities have an impact outside their boundaries greater than their impact inside the cities' boundaries. If Bcdp is considered, only three megacities have larger impacts outside than inside their boundaries. These three cities are Beijing, Tianjin and Karachi and they were shown to have quite high values of low level pollution export - ELR1km. Therefore the combined conditions of high ELR1km and high population densities outside cities' boundaries were necessary to have a higher impact on population outside the city than that on population inside the city. In general, however, it was found that most of the impact on population is related to pollution inside the city boundary.

Acknowledgements. This study was funded in the European Union's Seventh Framework Programme FP/2007–2013 within the projects MEGAPOLI (grant agreement no. 212520) and ECLIPSE (grant agreement no. 282688) and by the Norwegian Research Council in the framework of the project CLIMSLIP.

References

- Baumbach, G., Vogt, U., Hein, K. R. G., Oluwole, A. F., Ogunsola, O. J., Olaniyi, H. B., and Akeredolu, F. A.: Air pollution in a large tropical city with a high traffic density – results of measurements in Lagos, Nigeria. *Science of the total environment*, 169, 25–32, doi:10.1016/0048-9697(95)04629-F, 1995.
- Bigi, A., Ghermandi, G., and Harrison, R. M.: Analysis of the air pollution climate at a background site in the Po valley, *J. Environ. Monit.*, 14, 552–563, 2012.
- Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G., Fernandes, S., and Trautmann, N.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850–2000, *Global Biogeochem. Cy.*, 21, GB2018, doi:10.1029/2006GB002840, 2009.
- Emanuel, K. A. and Zivkovic-Rothman, M.: Development and evaluation of a convection scheme for use in climate models, *J. Atmos. Sci.*, 56, 1766–1782, 1999.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Gurjar, B. R. and Lelieveld, J.: New Directions: Megacities and global change, *Atmos. Environ.*, 39 391–393, 2005.
- Gurjar, B. R., Butler, T. M., Lawrence, M. G., and Lelieveld, J.: Evaluation of emissions and air quality in megacities, *Atmos. Environ.*, 42, 1593–1606, 2008.
- 5 Guttikunda, S. K., Carmichael, G. R., Calori, G., Eck, C., and Woo, J. H.: The contribution of megacities to regional sulfur pollution in Asia, *Atmos. Environ.*, 37, 11–22, 2003.
- Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan, V., and Rasch, P. J.: Springtime warming and reduced snow cover from carbonaceous particles, *Atmos. Chem. Phys.*, 9, 2481–2497, doi:10.5194/acp-9-2481-2009, 2009.
- 10 Forster, C., Stohl, A., and Seibert, P.: Parameterization of convective transport in a Lagrangian particle dispersion model and its evaluation, *Jo. Appl. Meteorol. Climatol.*, 46, 403–422, doi:10.1175/JAM2470.1, 2007.
- Hansen, J. and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *P. Natl. Acad. Sci.*, 101, 423–428, 2004.
- 15 Hertel, O., Christensen, J., Runge, E. H., Asman, W. A. H., Berkowicz, R., Hovmand, M. F., and Hov, O.: Development and testing of a new variable scale air pollution model – ACDEP. *Atmos. Environ.*, 29, 1267–1290, 1995.
- Hirdman, D., Sodemann, H., Eckhardt, S., Burkhardt, J. F., Jefferson, A., Mefford, T., Quinn, P. K., Sharma, S., Ström, J., and Stohl, A.: Source identification of short-lived air pollutants in the Arctic using statistical analysis of measurement data and particle dispersion model output, *Atmos. Chem. Phys.*, 10, 669–693, doi:10.5194/acp-10-669-2010, 2010.
- 20 Lawrence, M. G. and Rasch, P. J.: Tracer transport in deep convective updrafts: plume ensemble versus bulk formulations, *J. Atmos. Sci.*, 62, 2880–2894, 2005.
- Lawrence, M. G., Butler, T. M., Steinkamp, J., Gurjar, B. R., and Lelieveld, J.: Regional pollution potentials of megacities and other major population centers, *Atmos. Chem. Phys.*, 7, 3969–3987, doi:10.5194/acp-7-3969-2007, 2007.
- 25 McMahan, T. A. and Denison, P. J.: Empirical atmospheric deposition parameters – a survey. *Atmos. Environ.*, 13, 571–585, 1979.
- Mena-Carrasco, M., Carmichael, G. R., Campbell, J. E., Zimmerman, D., Tang, Y., Adhikary, B., D'allura, A., Molina, L. T., Zavala, M., García, A., Flocke, F., Campos, T., Weinheimer, A. J., Shetter, R., Apel, E., Montzka, D. D., Knapp, D. J., and Zheng, W.: Assessing the regional impacts of Mexico City emissions on air quality and chemistry, *Atmos. Chem. Phys.*, 9, 3731–3743, doi:10.5194/acp-9-3731-2009, 2009.
- 30

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Quinn, P. K., Bates, T. S., Baum, E., Doubleday, N., Fiore, A. M., Flanner, M., Fridlind, A., Garrett, T. J., Koch, D., Menon, S., Shindell, D., Stohl, A., and Warren, S. G.: Short-lived pollutants in the Arctic: their climate impact and possible mitigation strategies, *Atmos. Chem. Phys.*, 8, 1723–1735, doi:10.5194/acp-8-1723-2008, 2008.
- 5 Real, E., Law, K. S., Schlager, H., Roiger, A., Huntrieser, H., Methven, J., Cain, M., Holloway, J., Neuman, J. A., Ryerson, T., Flocke, F., de Gouw, J., Atlas, E., Donnelly, S., and Parrish, D.: Lagrangian analysis of low altitude anthropogenic plume processing across the North Atlantic, *Atmos. Chem. Phys.*, 8, 7737–7754, doi:10.5194/acp-8-7737-2008, 2008.
- 10 Riahi, K., Grubler, A., and Nakicenovic, N.: Scenarios of long-term socio-economic and environmental development under climate stabilization, *Technological Forecasting and Social Change*, 74, 887–935, 2007.
- Schneider, A., Friedl, M. A., and Potere, D.: A new map of global urban extent from MODIS satellite data, *Environ. Res. Lett.*, 4, 044003, doi:10.1088/1748-9326/4/4/044003, 2009.
- Slinn, W. G. N.: Predictions for particle deposition to vegetative canopies, *Atmos. Environ.*, 16, 1785–1794, 1982.
- 15 Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys. Res.*, 111, D11306, doi:10.1029/2005JD006888, 2006.
- Stohl, A. and Sodemann, H.: characteristics of atmospheric transport into the Antarctic troposphere, *J. Geophys. Res.*, 112, D02305, doi:10.1029/2009JD012536, 2010.
- 20 Stohl, A. and Thomson, D. J.: A density correction for Lagrangian particle dispersion models, *Bound. Lay. Meteorol.*, 90, 155–167, doi:10.1023/A:1001741110696, 1999.
- Stohl, A., Hittenberger, M., and Wotawa, G.: Validation of the Lagrangian particle dispersion model FLEXPART against large-scale tracer experiment data, *Atmos. Environ.* 32, 4245–4264 doi:10.1016/S1352-2310(98)00184-8, 1998.
- 25 Stohl, A., Eckhardt, S., Forster, C., James, P., and Spichtinger, N.: On the pathways and timescales of intercontinental air pollution transport, *J. Geophys. Res.*, 107, 4684, doi:10.1029/2001JD001396, 2002.
- Stohl, A., Forster, C., Eckhardt, S., Spichtinger, N., Huntrieser, H., Heland, J., Schlager, H., Wilhelm, S., Arnold, F., and Cooper, O.: A backward modeling study of intercontinental pollution transport using aircraft measurements, *J. Geophys. Res.*, 108, 4370, doi:10.1029/2002JD002862, 2003.
- 30

Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, doi:10.5194/acp-5-2461-2005, 2005.

Thomson, D. J.: Criteria for the selection of the stochastic models of particle trajectories in turbulent flows, *J. Fluid Mech.*, 180, 529–556, 1987.

Vignati, E., Karl, M., Krol, M., Wilson, J., Stier, P., and Cavalli, F.: Sources of uncertainties in modelling black carbon at the global scale, *Atmos. Chem. Phys.*, 10, 2595–2611, doi:10.5194/acp-10-2595-2010, 2010.

Voss, P. B., Zaveri, R. A., Flocke, F. M., Mao, H., Hartley, T. P., DeAmicis, P., Deonandan, I., Contreras-Jiménez, G., Martínez-Antonio, O., Figueroa Estrada, M., Greenberg, D., Campos, T. L., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Crouse, J. D., Wennberg, P. O., Apel, E., Madronich, S., and de Foy, B.: Long-range pollution transport during the MILAGRO-2006 campaign: a case study of a major Mexico City outflow event using free-floating altitude-controlled balloons, *Atmos. Chem. Phys.*, 10, 7137–7159, doi:10.5194/acp-10-7137-2010, 2010.

Wesely, M. L. and Hicks, B. B.: Some factors that affect the deposition rates of sulfur dioxide and similar gases on vegetation. *J. Air Poll. Contr. Assoc.*, 27, 1110–1116, 1977.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Several quantities for all the megacities being part of this study are reported. T_{li} and T_{lo} are two decaying time scales obtained for the deposition process (see text for details). PWC is the population-weighted ground level concentration and measures the concentration to which the population inside the city is exposed to, on average. $GLC \times Pop$, is the product of population and ground level concentration. Other quantities are self explanatory.

Properties	N. cells	City population	Emission	T_{lo}	T_{li}	PWC	PWC	GLC. \times Pop.	
		10^6 people	10^6 kg yr $^{-1}$	days	days	ng m $^{-3}$	ng m $^{-3}$	10^{10} ng m $^{-3}$ people	
Cities name (short name)								BCtr	BCdp
Europe									
Istanbul (ISTANB)	4	13.9	2.17	7.8	5.0	61	64	2.4	1.2
London(LONDON)	2	14.2	5.85	5.7	1.5	379	398	10	6.8
Moscow(MOSCOW)	1	14.6	6.74	5.7	2.4	884	931	19	14
Paris(PARIS)	1	9.9	5.06	6.3	1.8	543	563	10	6.7
Po valley(POVALL)	11	22.7	14.8	7.6	3.7	282	335	17	8.8
Ruhr valley(RUHRVA)	3	13.6	7.98	6.0	1.7	362	388	14	8.1
St Petersburg(STPBRG)	2	2.6	3.42	5.5	1.4	235	234	3	0.9
North America									
Chicago(CHICAG)	1	5.5	2.42	5.9	2.1	227	236	2.9	1.7
Los Angeles(LOSANG)	4	17.2	12.7	7.9	6.0	583	606	15	11
New York(NEWYOR)	15	45.5	38.0	5.7	1.7	454	518	40	22
Central /South America									
Buenos Aires(BAIRES)	1	13.0	5.07	5.9	2.4	556	582	8.4	7.6
Lima(LIMA)	4	8.1	5.22	7.4	6.0	164	181	2.4	1.6
Mexico City(MEXICO)	2	25.2	8.03	7.8	5.9	340	360	12	9.4
Rio de Janeiro(RIODEJ)	1	10.7	5.00	6.3	1.7	368	429	6	4.4
San Paulo(SPAULO)	1	21.2	8.55	6.1	1.9	548	623	15	12
Santiago(SNTAGO)	1	6.0	5.53	6.9	5.3	512	520	3.9	3.3
Asia									
Bangkok(BANGKO)	1	11.4	7.01	6.7	2.2	593	633	11	7.8
Beijing(BEIJIN)	1	9.7	12.1	6.2	3.5	1273	1315	37	27
Delhi(DELHI)	1	25.5	8.41	8.5	6.1	1184	1228	61	53
Dhaka(DHAKA)	11	138	32.5	7.1	2.4	637	753	149	104
Hong Kong(HONKON)	4	51.7	44.2	6.0	1.7	1109	1281	105	69
Jakarta(JAKART)	1	25.7	10.4	6.6	1.0	802	968	31	23
Karachi(KARACH)	1	9.3	1.91	8.9	8.2	137	138	3.5	2.5
Kolkata(KOLKAT)	1	28.1	7.66	7.3	2.8	901	949	42	32
Manila(MANILA)	2	22.5	7.65	6.8	1.2	220	265	8.6	5.3
Mumbai(MUMBAL)	4	30.0	12.9	8.6	5.2	312	333	21	13
Osaka(OSAKA)	1	16.0	8.32	5.9	1.8	534	572	13	9.8
Seoul(SEOUL)	2	23.8	25.9	5.9	2.6	1535	1621	60	44
Shanghai(SHANGH)	1	18.1	7.87	6.1	2.2	626	657	26	18
Teheran(TEHERA)	1	11.5	3.51	8.4	8.2	349	358	6.7	5
Tianjin(TIAAJ)	1	8.9	7.67	6.2	3.4	855	882	23	16
Tokyo(TOKYO)	1	28.7	15.3	5.8	1.6	1206	1306	46	37
Africa									
Cairo(CAIRO)	1	22.3	5.59	9.6	10.5	472	473	15	13
Johannesburg(JOBUrg)	2	7.6	11.4	7.4	5.3	759	778	8.3	7
Lagos(LAGOS)	1	9.7	2.84	7.5	3.4	220	261	4.9	2.9
Australia									
Sidney(SYDNEY)	5	5.1	7.44	6.8	3.0	145	158	1.2	0.8
Averaged value		20.8	10.5	6.86	3.52	565	608	23.7	17.4

Table 2. Comparison of ELR1km for a conserved tracer with no decay (BCtr) and one subject to deposition (BCdp), for the BCtr tracer with a superimposed exponential decay process with a time scale of 10 days (BC10, see text for clarification) and a tracer with 10 days decay time scale computed by L07 using the model MATCH for the year 1995.

ELR1km (%)	BCtr	BCdp	BC10 FLEX	BC10 MATCH
Cities				
Europe				
ISTANB	19.1	21.8	20.4	21.7
LONDON	25.5	20.7	25.9	26.0
MOSCOW	28.6	25.4	29.6	34.2
PARIS	23.4	20.6	23.8	24.8
POVALL	19.5	17.3	17.6	18.5
RUHRVA	24.9	20.3	25.1	–
STPBRG	29.3	24.9	30.1	–
North America				
CHICAG	20.3	19.2	19.6	19.3
LOSANG	15.3	15.9	15.8	16.7
NEWYOR	20.5	18.7	19.9	16.0
Central & South America				
BAIRES	19.8	16.8	19.1	14.7
LIMA	13.1	12.7	15.7	10.9
MEXICO	10.1	6.2	6.6	5.5
RIODEJ	13.2	5.8	9.0	6.0
SPAULO	13.3	6.2	9.2	5.4
SNTAGO	16.5	8.1	12.7	–
Asia				
BANGKO	9.1	6.3	9.1	6.6
BEIJIN	18.3	17.9	18.1	19.8
DELHII	11.3	21.8	12.9	18.9
DHAKA	9.2	13.3	9.0	13.3
HONKON	10.9	10.4	10.1	12.9
JAKARTA	7.3	2.6	7.0	3.2
KARACH	11.5	18.7	14.4	17.1
KOLKAT	9.3	15.2	9.8	13.5
MANILA	8.7	7.3	9.8	5.8
MUMBAI	9.7	12.5	11.9	14.5
OSAKA	16.6	13.3	15.0	13.2
SEOUL	18.2	16	18.0	16.7
SHANGH	14.8	15.1	16.3	16.8
TEHERA	15.7	14	13.1	17.2
TIAAJI	18.4	17.8	18.6	18.5
TOKYO	18	10.7	15	11.5
Africa				
CAIRO	15.3	20.4	17.8	21.6
JOBURG	13.6	8	9.3	6.5
LAGOS	8.8	4.6	6.3	6.0
Australia				
SYDNEY	18.2	14	15.8	11.4

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. The Table reports: (1) cumulative value of mass of BCdp deposited (Dep66) in the polar regions ($|\text{Lat}| > 66^\circ$); (2) mass of BCtr and Bcdp suspended (Sus66) in the polar atmosphere up to an elevation $h < 1$ km above terrain; (3) the mass of BCdp deposited over land in the Northern Hemisphere during winter for ($\text{Lat} > 40^\circ$), Dep40l. The mass is reported in tons, kg and as a fraction (% , ‰) of the total averaged emitted mass within the 48 days age class limit; (4) the mass in the polar stratosphere (Strat66).

Cities	Mass										
	BCtr Sus66 ton	Bcdp Sus66 kg	Bctr Sus66 %	BCdp Sus66 ‰	BCdp Dep66 ton	BCdp Dep66 %	Bcdp Dep40l ton	BCdp Dep40l %	BCtr Strat66 ton	BCdp Strat66 kg	
Europe											
ISTANB	4.01	40	1.43	0.14	19.4	0.89	305	45.6	1.07	38	
LONDON	31.4	569	4.09	0.74	256	4.35	795	44.4	2.58	36	
MOSCOW	54.8	1423	6.2	1.61	382	5.61	1544	75	3.42	71	
PARIS	19.9	231	3.02	0.35	126	2.49	879	57.2	2.59	81	
POVALL	31.3	276	1.61	0.14	249	1.67	2084	46	9.96	626	
RUHRVA	40.4	723	3.86	0.69	355	4.41	1421	58.2	3.87	98	
STPBRG	31.1	886	6.94	1.98	326	9.43	679	67	1.53	34	
North America											
CHICAG	6.06	56	1.9	0.17	12.4	0.51	456	61.7	1.98	35	
LOSANG	9.24	211	0.56	0.13	104	0.81	388	9.91	9.88	276	
NEWYOR	97.2	764	1.95	0.15	180	0.47	3169	27.3	29.5	582	
Central & South America											
BAIRES	7.02	18	1.06	0.03	6.08	0.12	–	–	1.96	29	
LIMAPE	0.08	0	0.01	0	0.27	0.01	–	–	0.43	3	
MEXICO	1.14	11	0.11	0.01	12.4	0.15	68.1	2.72	3.99	156	
RIODEJ	1.87	3	0.29	0	1.2	0.02	–	–	2.05	21	
SPAULO	3.21	2	0.29	0	1.73	0.02	–	–	3.54	27	
SNTAGO	4.03	11	0.55	0.01	9.2	0.16	–	–	2.17	44	
Asia											
BANGKO	0.57	1	0.06	0	0.87	0.01	11.6	0.74	3.04	51	
BEIJIN	20	168	1.26	0.11	49.2	0.4	780	20.9	9.2	164	
DELHI	0.74	1	0.07	0	2.66	0.03	14.8	0.56	4.12	169	
DHAKA	2.48	5	0.06	0	10.9	0.03	64.3	0.61	16.8	363	
HONKON	10.7	31	0.18	0.01	25.1	0.06	87.4	0.64	27.9	345	
JAKART	0.16	0	0.01	0	0.28	0.00	–	–	1.33	14	
KARACH	0.17	1	0.07	0	0.84	0.04	5	0.82	0.77	51	
KOLKAT	0.43	1	0.04	0	1.52	0.02	10.5	0.44	3.5	103	
MANILA	0.61	1	0.06	0	1.07	0.01	9.24	0.39	3.11	63	
MUMBAI	0.45	2	0.03	0	2.66	0.02	19.7	0.48	4.28	271	
OSAKA	11.9	36	1.09	0.03	18.2	0.22	46.5	1.82	7.27	94	
SEOUL	45.6	195	1.34	0.06	58.1	0.22	245.6	3.09	19.6	194	
SHANGH	5.92	18	0.57	0.02	8.85	0.11	60.2	2.48	5.52	71	
TEHERA	4.42	96	0.96	0.21	35.	1.01	312.8	28.7	2.31	60	
TIAAJI	12.9	87	1.28	0.09	28.6	0.37	473.5	20	5.77	98	
TOKYO	29.2	117	1.45	0.06	42.0	0.27	104.2	2.2	13	188	
Africa											
CAIRO	3.39	66	0.46	0.09	22.1	0.39	191	10.8	1.95	97	
JOBURG	5.11	4	0.34	0	7.06	0.06	–	–	4.3	77	
LAGOS	0.06	0	0.02	0	0.36	0.01	2.84	0.32	0.53	25	
Australia											
SYDNEY	8.98	46	0.92	0.05	19.39	0.26	–	–	3.21	82	
Total NH	476.09	6016			2330.2		14227.28		199.04	4440	
Total SH	30.46	84			45.2				18.99	297	




Fig. 1. Location and extension of the megacities investigated.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[⏪](#) | [⏩](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 2. Relative contribution of the megacity emissions for BC, carbon monoxide (CO), nitrogen oxides (NO_x) and sulfur dioxide (SO₂) accumulated from the poles towards the equator, relative to the accumulated global emission as a function of the latitude south of the North Pole (**b, d**) and north of the South Pole (**a, c**) and divided as winter (**a, b**) and summer (**c, d**).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 3. Concentration of BCdp normalized with the concentration of BCtr (BCdp/BCtr) as function of the age for the Northern Hemisphere (NH, **a**) and the Southern Hemisphere (SH, **b**). Lower panels (**c**, **d**) as above but using a linear scale.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Transport distance ($\bar{d}(t)$) from the source megacity location for the vertically integrated concentration of BCtr and BCdp for the NH (a, b) and SH (c, d). The smaller panels on the left are a zoom for 0–3 days using a logarithmic scale.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 5. Altitude (above sea level) of the center of mass of horizontally integrated concentration ($\bar{h}(t)$) for BCtr (a, c) and BCdp (b, d) in the NH (a, b) and in the SH (c, d).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 6. Ratio of the averaged (over all cities) horizontal distance of centers of mass, $D(t)$, and vertical position of center of mass, $H(t)$, for BCdp over BCtr.




Fig. 7. Mass in the upper troposphere and stratosphere (9–30 km) normalized by the total mass suspended in the atmosphere as a function of tracer age for BCtr and BCdp, for the NH (**a, b**) and SH (**c, d**).

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 8. Fraction of BCtr (a) and BCdp (b) mass exported over 1000 km but with elevation below 1 km above ground level: ELR1km. The fraction is calculated as cumulated mass exported up to a specific age normalized with the total suspended mass in the atmosphere up to this age.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 9. Scatter plot of ELR1km for the values of BC10 reported in Table 2, produced by FLEXPART for the years 2003–2005 and MATCH for the year 1995.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 10. Fraction of mass transported over 1000 km away from the source over the whole air column (ELRcol) for BCtr and for BCdp. The normalization is the same used before in Fig. 8.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 11. BCtr (left panels) and BCdp (right panels) mass content in a layer 0–1000 m a.t.l. in the Arctic (a, b) and Antarctic (c, d) regions, scaled for the total mass emitted in any of its respective age class.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 12. BCtr (left panels) and BCdp (right panels) mass content in the Arctic region scaled for the total mass emitted in the respective age class for layers of the atmosphere extending from 2 to 9 km (**a**, **b**) and from 9 to 30 km (**c**, **d**) above sea level. The insert in panel (**c**) shows the same information but on a linear scale.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 13. Deposition of BC at latitudes north of 66° N (a, b) and south of 66° S (c, d), relative to the mass available for deposition (left) and absolute value (right).




Fig. 14. The CmP metric for BCtr (left) and BCdp (right) as a function of tracer age.

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The dispersion characteristics of air pollution from world's megacities

M. Cassiani et al.




Fig. 15. CmP/CmP_c as a function of tracer age, for BCtr (left) and BCdp (right). In the left panel, CmP/CmP_c values for the city of St. Petersburg exceed the scale and reach about 4.5 at 48 days.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

