

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Comparing two years of Saharan dust source activation obtained by regional modeling and satellite observations

I. Tegen¹, K. Schepanski², and B. Heinold²

¹Leibniz Institute for Tropospheric Research, Leipzig, Germany

²School of Earth and Environment, University of Leeds, Leeds, UK

Received: 3 August 2012 – Accepted: 1 October 2012 – Published: 22 October 2012

Correspondence to: I. Tegen (itegen@tropos.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

12, 27667–27691, 2012

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

A regional-scale dust model is used to simulate Saharan dust emissions and atmospheric distributions in the years 2007 and 2008. The model results are compared to dust source activation events compiled from infrared dust index imagery from the geostationary Meteosat Second Generation (MSG) satellite. The observed morning maximum in dust source activation frequencies indicates that the breakdown of nocturnal low-level jets is responsible for a considerable number of dust source activation events in the Sahara. The comparison shows that the time of the day of the onset of dust emission is delayed in the model compared to the observations. Also, the simulated number of dust emission events associated with nocturnal low level jets in mountainous regions is underestimated in the model. The MSG dust index observations indicate a strong increase in dust source activation frequencies in the year 2008 compared to 2007, the difference between the two years is less pronounced in the model. The quantitative comparison of simulated dust optical thicknesses with observations at stations of the sunphotometer network AERONET shows, however, good agreement for both years, indicating that the number of observed dust activation events is only of limited use for estimating actual dust emission fluxes in the Sahara.

1 Introduction

Soil dust aerosol is considered to be an important factor in the climate system (Solomon et al., 2007). Dust aerosol can impact upon climate directly by modifying the atmospheric radiation balance or indirectly by influencing cloud properties and biogeochemical cycles (e.g. Sokolik et al., 2001; DeMott et al., 2003; Jickells et al., 2005). An adequate description of dust source regions as well as the meteorological processes leading to dust emission are prerequisites for evaluating the influence of dust aerosol on the climate, as well as for understanding the response of dust distribution to changing climate conditions.

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Sahara desert is the largest dust source worldwide. Saharan dust distributions are strongly variable at different timescales (e.g., Mahowald et al., 2006). Information on the sub-daily and seasonal occurrence of Saharan dust emission events is available from the geostationary Meteosat Second Generation (MSG) satellite infrared (IR) dust index since the year 2004 (Schepanski et al., 2007). The time of day when dust emissions are initiated can be an indicator for the meteorological processes causing dust emissions. The MSG observations suggest that approximately 60 % of the Saharan dust events are initiated in the morning hours before 9UTC (Schepanski et al., 2009), indicating the importance of turbulent downward mixing of momentum from nocturnal low level jets (LLJs) for activating dust sources (Washington and Todd, 2005). Seasonal changes in emissions and atmospheric transport of dust are characterized by variations in near-surface meteorology in connection with the seasonal displacement of the intertropical convergence zone and the African monsoon system (Schepanski et al., 2009; BouKaram et al., 2009). The causes of Saharan dust variability at multi-annual timescales are not well known. While observed variations in transported dust have been linked to drought conditions in the Sahel and North Atlantic Oscillation patterns (Prospero and Lamb, 2003; Chiapello et al., 2005), the study of variability of dust emissions is problematic due to the lack of available observations.

While the MSG dust source activation (DSA) dataset provides useful information about the occurrence of dust emissions events, models of dust emission and transport are useful to understand the relationships between the frequency of dust emission events, emission fluxes, and atmospheric dust loads. In an intercomparison of 14 different global dust models Huneeus et al. (2011) show that the seasonal cycles and largescale transport patterns of dust aerosol can be well reproduced by such models. However, large discrepancies in the results of the different models for results at individual locations indicate that dust processes may not be adequately resolved in such models. Regional-scale models that are commonly used for dust forecasts e.g. in the Mediterranean region (see <http://sds-was.aemet.es/>) are expected to better predict the meteorological features leading to dust emissions. Such processes include syn-

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



optic scale features like mesoscale convective systems in the Southern Sahara, LLJs, and microscale processes related to dry convection, which have different seasonal and diurnal characteristics (Knippertz and Todd, 2012). Synoptic scale meteorological patterns leading to dust emissions like the Sharav cyclone in late winter/early spring in the northern part of the Sahara (Alpert and Ziv, 1989) are well reproduced by regional scale simulations. Problematic at the regional (as well as global) scale are simulations of dust emission processes related to wet convection (Knippertz et al., 2009; Reinfried et al., 2009) or dry convective events at microscale (e.g., dust devils, Koch and Renno, 2005). In this work we use the regional-scale Saharan dust model COSMO-MUSCAT (Heinold et al., 2007) to simulate the dust emission events and dust optical thicknesses for the years 2007 and 2008 using observed DSA from MSG dust index data for evaluation of the spatiotemporal distribution of modeled dust emission events.

2 Methods

2.1 Model description

The regional dust model system developed in the framework of Saharan Mineral Dust experiment SAMUM is described in detail by Heinold et al. (2007). It is based on the COnsortium for Small scale MOdelling (COSMO) meteorological model (Steppler et al., 2003), the operational weather model of the German Weather Service (Deutscher Wetterdienst, DWD), the online-coupled Chemistry-Transport-Model Multi-Scale Chemistry Aerosol Transport (MUSCAT) model (Wolke et al., 2004), and a dust emission scheme (Tegen et al., 2002). In MUSCAT, advection of chemical species and aerosol particles is computed by a third-order upstream scheme. Temporal integration is carried out by an implicit-explicit method. Dust emissions are computed by the model in non-vegetated areas. Emission fluxes depend on surface wind friction velocities, surface roughness, soil particle size distribution, and soil moisture (Heinold et al., 2007). Wind fields are assimilated from the COSMO model. Threshold friction

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



velocities that must be reached to initiate dust emission depend on soil particle size distribution following Marticorena and Bergametti (1995). Within each model grid cell constant surface roughness lengths are assumed. Soil particle size distributions were derived from soil texture data (Zobler, 1986), assigning a composite of four particle size modes with mode diameters at 2 μm , 15 μm , 158 μm and 720 μm to each texture class (Tegen et al., 2002). Model grid cells where dust emissions are not observed in the MSG satellite observations during the years 2006 to 2009 (Schepanski et al., 2007) are excluded as dust source regions in the model. Thereby, model constraints due to uncertainties in soil data are fixed, while the actual placement and the predicted amount of dust emissions are controlled by COSMO surface winds. In the transport model MUSCAT dust is transported in five independent size classes with radius limits between 0.1 μm and 24 μm as dynamic tracer. Dust removal is computed considering dry and wet deposition processes. Dry deposition is computed following Seinfeld and Pandis (1998) and wet deposition (rain-out and wash-out) is parameterized according to Berge (1997). Wavelength-dependent dust optical thicknesses are computed from simulated dust concentrations, particle size distribution and extinction efficiencies as:

$$\tau = \sum_i \frac{3 Q_{\text{ext}}(\lambda, r_i) M_i}{4 r_i \rho} \quad (1)$$

where Q_{ext} is the dimensionless specific particle extinction efficiency (depending on wavelength λ , Lacis and Mishchenko, 1995), ρ is the particle density, and M_i is the aerosol mass load of size fraction i characterized by the effective radius r_i .

The computation of radiative transfer of short- and longwave radiation in COSMO takes into account effects of scattering, absorption, and emission by aerosols, cloud droplets, and gases. It additionally considers the size-resolved dust optical thickness computed in the dust transport model. Thus the computation of radiation fluxes accounts for the spatially and temporally varying atmospheric dust load (Helmert et al., 2007).

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A horizontal grid resolution of 28 km is used for the area between 0.2° N, 32.3° W and 41.1° N, 32.9° E, including relevant Saharan dust sources, parts of Europe and the Eastern Tropical Atlantic. This domain ensures that synoptic and regional scale weather systems and local dust emissions are reproduced. The model is operated with 40 vertical layers of a pressure-based, terrain-following vertical coordinate; the lowest layer extends 68 m. Simulations were performed for the full two years 2007 and 2008. COSMO simulations of the regional meteorology were initialized and the lateral boundary conditions are updated 6-hly with analysis fields from the global model GME (Global Modell of the DWD). The simulations were re-initialized every 48 h to keep the meteorology of the regional model close to the analysis fields. After a spin-up of 24 h the COSMO model is coupled with MUSCAT to compute dust emission and transport, using modeled dust concentration from the previous cycle as initialization. The model has been extensively tested with observations from the SAMUM field study and available station and remote sensing data (Heinold et al., 2009, 2011).

2.2 Meteosat second generation infrared dust index

Brightness temperature differences converted from radiances of infrared wavelength channels measured by the SEVIRI (Spinning Enhanced Visible and InfraRed Imager) instrument on board the geostationary MSG satellite were used for Saharan dust detection at day- and nighttime. The qualitative index indicating the presence of mineral dust is computed from brightness temperatures of three different IR channels (8.7 μm , 10.8 μm , and 12.0 μm). The spatiotemporal resolution of the dust signal is determined by the sampling rate of 15 min and the spatial resolution of 3 km at nadir. No dust emission can be detected if thick clouds cover dust sources or the column water vapour is too high (Brindley et al., 2012). This signal was used to localize dust source activation events by visually tracing dust plume patterns back to their origin by inspecting consecutive images during dust events. Dust emitting areas were marked as DSA event on a 1° × 1° gridded map for the Saharan region between 5° N, 20° W and 40° N, 40° E. The monthly DSA frequencies (DSAF) were calculated for each grid cell in this re-

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gion (Schepanski et al., 2007). High DSAF were identified in the Bodélé depression in Chad, but also near mountain foothill areas where fine sediment is available from fluvial sources, and high surface wind speeds frequently initiate dust emission events. The observed seasonal and diurnal patterns of dust emission help to understand the meteorological processes controlling the spatio-temporal patterns of dust emission in the different regions of in the Saharan desert (Schepanski et al., 2009). A good agreement of the spatio-temporal variability of dust sources derived from the MSG dust index data and ground-based dust observations at meteorological stations in the Sahara has been found for a case study in spring/summer 2006 (Laurent et al., 2010). The dust source regions inferred from the backtracking of dust plumes differ from satellite-inferred dust source regions that are based on a high frequency of high daily dust aerosol loads, e.g. from the Ozone Monitoring Instrument Aerosol Index (OMI AI) or aerosol optical thickness derived from the Moderate Resolution Imaging Spectroradiometer (MODIS) (both flying on the NASA A-Train constellation). The differences were attributed to the temporal resolution of the different satellite retrievals (Schepanski et al., 2012).

3 Results

3.1 Dust source activation frequencies

Dust emissions in the Sahara desert are computed for the 2-yr model period. Simulated total annual dust emissions reach values of approximately 1900 Mtyr^{-1} for 2007 and 2300 Mtyr^{-1} for 2008 (Table 1). That the simulated emissions are higher compared to earlier publications (e.g. Laurent et al., 2008) can be due to the considered particle radius up to $24 \mu\text{m}$, while in most publications only particles with radii less than $10 \mu\text{m}$ are considered.

For direct comparison with the MSG data, the model results were re-binned to a resolution of $1^\circ \times 1^\circ$. To facilitate comparisons with the observed DSA from the MSG observations the occurrence of dust emissions were counted when the emission flux in

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a model grid cell during the day exceeded $0.6 \times 10^{-4} \text{ kg m}^{-2} \text{ s}^{-1}$ in a 3-h time interval. Only higher dust emission fluxes were counted as dust events to account for the fact that very minor dust events would not be detectable in the MSG data. The emission flux limit was chosen such that the 2-yr sum of counts of Saharan dust emission events for all Sahara gridboxes from both model and observations agreed with each other. To account for the fact that dust emissions cannot be detected from the satellite data when cloud cover is present, model DSA events were not counted when the model-simulated cloud cover in the grid cell during the respective day was higher than 50 %. The two-year sum of daily modeled dust emission counts for the Sahara were 25 582 compared to 26 249 counts for the MSG DSA (Table 1).

Results for the monthly sums of DSA for the Saharan domain including information on the time-of-day of the beginning of dust emissions are shown in Fig. 1. The modeled monthly total numbers of Saharan dust emission events show a pronounced seasonal cycle with a clear spring maximum, while the seasonality is less pronounced in the observations. The number of DSA in the MSG dataset increases by more than 100 % in 2008 compared to 2007 (see also Table 1). The model results show also an increase in the number of DSA in 2008, but only by 27 %. The higher number of DSA observed in 2008 is mainly due to increased emissions in the morning hours between 06:00 and 12:00 UTC (Fig. 1). The higher dust source activation at morning hours in 2008 points towards an increase in the occurrence of nocturnal LLJs that were identified to be major contributor to dust emission events in the Sahara (Washington and Todd, 2005; Schepanski et al., 2009). LLJs develop as consequence of the nocturnal stabilization of the boundary layer; their momentum is mixed to the surface by turbulence setting in after sunrise when the surface heats up. In average 65 % of the simulated dust emission events occur before noontime, which indicates that the breakdown of the nocturnal LLJ is also a dominant dust mobilization mechanism in the model.

While it cannot be ruled out that the DSA observations are biased towards the morning hours, the less pronounced diurnal cycle of DSA activation in the model compared to the observations is likely to be related to shortcomings in the turbulence parameteri-

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



zation and the lower boundary conditions like model topography and surface roughness (Todd et al., 2008; Heinold et al., 2011). The results suggest that the downward mixing of momentum during the morning degradation of LLJs is delayed in the model. Thus, dust emission events related to the LLJ breakdown can occur later in the day compared to the observations. The boundary layer scheme may not cope with the nocturnal stable stratification resulting in too weak jets. In addition, the characteristic sharp morning wind peak is not always reproduced, as the turbulent mixing takes place too gradually.

The spatial distribution of dust source activation frequencies are compared for the MSG observations and numbers of model dust emission events are shown in Fig. 2. Model results from the COSMO-MUSCAT and the MSG infrared dust source activation (DSA) events for the years 2007 and 2008 were averaged for months, in which similar meteorological conditions were dominant causes for dust emissions (Fig. 2, left and middle panels). In addition, the occurrence of LLJs in the model is shown (Fig. 2, right panels). The occurrence of nocturnal LLJs in the individual grid cells was computed from simulated wind speed differences above the nocturnal boundary layer. A nocturnal LLJ was assumed to have developed in a model grid cell if the wind speed difference between the layer directly above the boundary layer at 06:00 UTC and the layer located 200 hPa above this layer exceeds 5 m s^{-1} (this value was chosen in agreement with Schepanski et al., 2009). Approximately 40 % of the simulated DSA events in the model are associated with the occurrence of LLJs on the same day (Table 1). In October to January (Fig. 2, lower panels) Saharan dust emissions are dominant in the Bodélé depression, here the occurrence of nocturnal LLJs are coincident with days of dust emissions. In the months from February to May the passages of Sharav cyclones are the cause of major dust emission events in Northern Sahara (e.g. Alpert and Ziv, 1989; Horvath et al., 2006) (Fig. 2, top panels). In the summer months (June to September, Fig. 2, middle panels) mesoscale convective systems are major causes for dust emissions in the Southern Sahara (Knippertz and Todd, 2012). Those events should occur less frequently than dust source activation by LLJs, but may be more important in terms of dust emission fluxes.

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The figures illustrate the dominance of the Bodélé dust source in the winter months, the good agreement of the model and observations in this region, and the important role of LLJs for dust mobilization in the Bodélé, where nearly all dust emission events are occurring at the same days when LLJs develop. An overestimate of DSAs that are unrelated to the development of LLJs is evident in the Northern Sahara. This may suggest that here the MSG observations underestimate DSA, as during these large scale events dust sources may become activated during the passage of Sharav cyclones below the passing dust cloud. These events would not be detected by the MSG imagery as new DSA, leading to an underestimation of DSA in the observations.

The MSG DSA data indicate a strongly increased dust activity in the year 2008 compared to 2007. However, this increase is less evident in the model simulations (Figs. 1, 3). In both years the agreement between modeled and observed dust emission events is very good in the Bodélé region in Chad, which is the most active dust source in the Sahara desert. While the over-prediction of DSA in the annual averages occurs mostly in Northern Saharan grid cells in both years, missed emissions are most evident for the annual average in the Sudan and the mountainous region along the border between Algeria, Mali and Niger (Fig. 3). In these regions dust emissions that are observed in the MSG record may be partly caused by high wind speeds that occur with the degradation of nocturnal LLJs when turbulent mixing sets in during the morning hours. As mentioned above, in the Bodélé dust hot spot almost all modeled dust emission events occur on days when a LLJ developed. In addition to the total number of Saharan dust emissions in the model compared to the MSG record the number of grid cells is given (summed over one year for the area between 15° W and 32° E, 10° N and 40° N), for which the modeled monthly dust emission frequencies either exceed (“overpredictions” in Table 1) or are below the observed frequencies (“underpredictions” in Table 1). In 29 % and 23 % of the gridcells modeled DSA are overpredicted compared to the observations in 2007 and 2008, respectively. An overprediction can be caused by insufficient decoupling of the nocturnal boundary layer and too high surface winds at night and early morning hours in the model. As an overprediction of

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



morning wind peaks due to the LLJ breakdown is unlikely, the misrepresentation of other peak wind generating processes acting at the same time or a wrong placement of the LLJ are more likely reasons. Unrelated to the occurrence of LLJs, dust emissions that are caused by wet convective events, i.e. meso-scale convective systems in the Southern Sahara and Sahel in summer and density currents in mountainous areas, may be misrepresented in the model (e.g. Reinfried et al., 2009; Marsham et al., 2011). On the other hand, potential deficits in the model capability to reproduce moist convection may be also obscured in favor of the model by the restriction of the MSG satellite product to cloud-free conditions. The percentage of grid cells where modeled dust emission events are underestimated compared to the observations is higher in 2008 (16 %) compared to 2007 (10 %) (Table 1). LLJs can develop but do not always lead to dust emissions in the model (Figs. 2 and 3). Nevertheless, increased LLJ occurrence in the summer months of the year 2008 compared to 2007 (Fig. 3) that corroborates the higher MSG DSA observations in 2008 points towards the importance of this phenomenon for dust mobilization even in the summer season. The reason for the lack of modeled dust mobilization following the development of nocturnal LLJs in mountainous regions may be due to too weakly developed LLJs or insufficient turbulent mixing of momentum towards the surface during the breakdown of the LLJs in the morning hours in the COSMO model. Also, the assumption of too high roughness lengths in the dust emission model that suppresses dust emission may not be representative for fluvial dust sources that may in fact be highly productive dust sources. If the number of days with LLJ occurrences but without dust emissions are added to the dust emission events in the respective model grid cells (i.e. implicitly assuming that all LLJ occurrences are followed by dust emission events), a close agreement is achieved between the model and observations with only 3 % or less of the model grid cells remaining to under-predict dust emission events (Table 1).

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Dust optical thickness

While the spatiotemporal distribution of dust source activation events provides necessary information for relating dust particle properties to soil properties in the source regions and estimate the effects of changing climate conditions on dust emissions, the actual effects of atmospheric dust aerosol on the radiation balance, precipitation distribution and fertilization of ecosystems depends on the atmospheric dust loads, that is not necessarily related to dust source activation frequencies in particular regions. In the annual average, the modeled dust aerosol optical thickness (AOT) reaches values up to 0.3 that occur over the central and western parts of the Sahara, with highest values in the Southern Sahara and the Bodélé region in Chad (Fig. 4a, b). As for the simulated DSAFs, AOTs are slightly higher in 2008 compared to 2007. Maximum values of annual average optical AOTs are 0.29 in 2007 compared to 0.32 in 2008. A qualitative comparison of model AOT averaged at 12:00 UTC with the annual averaged absorbing aerosol index (AI) measured by the Ozone Monitoring Instrument (OMI) (Torres et al., 2007) for the two years (Fig. 4c, d) shows similarities to the spatial distribution of the simulated dust aerosol. However it should be noted that the OMI AI is only a qualitative indicator of the presence of dust aerosol, and is also sensitive to biomass burning smoke aerosol in the Sahel region, which is not considered in the model. Similar to the model results the OMI AI shows slightly increased values in the year 2008 compared to 2007, particularly in the northern part of the Sahara.

Quantitative evaluation of the model results at individual location is done by comparison with sunphotometer data from the Aerosol Robotic Network AERONET (Holben et al., 1998). We compare optical thickness observations at 440 nm with model results of dust optical thickness at 500 nm wavelength. Daily averaged AOT retrievals from the AERONET stations are compared to modelled daily averaged dust optical thickness results at the respective location. Here the daily model results are averaged for the time period between 09:00 and 15:00 UTC to account for the fact that the measurements can only be performed during daytime. Since anthropogenic aerosols are not simu-

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lated in this version of the regional model, only days are compared when the Ångström parameter is lower than 0.7, indicating a dominance of large dust particles on that day and location (Dubovik et al., 2002). Only few AERONET stations continuously operate in the dust-dominated regions in the Saharan desert. Here we compare measurements and model results for the years 2007 and 2008 for the stations Tamanrasset (22.8° N, 5.5° E), Saada (31.6° N, 8.2° W) Dakar (14.4° N, 17.0° W) and Agoufou (15.3° N, 1.5° W) representing dust conditions in the northern, western, central and southern part of the Sahara, respectively (Fig. 5).

At these stations there is no clear difference in AOT magnitude for the two years, an increase in dust AOT in 2008 compared to 2007 is not observed or simulated. The seasonal cycle and the large day-to-day variability in the Saharan dust load is however evident in the observations and model results. The magnitudes and timing of most individual AOT peaks indicating strong dust events are matched well by the model results. Ehile there is good agreement at the station Saada in the Northern Sahara, notable discrepancies between the model results and the measurements occur at the station Agoufou in Northern Hemisphere spring and even more evident at Tamanrasset in the summer months (Fig. 5), where the model does not reproduce a number of dust events in both years. At Agoufou the underestimate may be due to influences from other aerosols at the location, which is close to the Sahel and may be influenced by biomass burning and anthropogenic aerosols. At the station Tamanrasset (located in Algeria in the vicinity of the Ahaggar mountains) the modeled dust AOT in the summer does not match several observed dust events and does not reach more than approximately 30 % of the observed AOT magnitudes in both years This strong underestimate is likely due to failures in the model to reproduce the meteorological conditions correctly that lead to dust emission upwind of this location at this time of the year. It could be either due to dust emitted by wet convective events which the model does not reproduce, or due to insufficiently resolved topography resulting in incorrect wind fields in this mountainous area. It may also be possible that the emission scheme used in the model is unsuitable for this terrain and should be modified accordingly. Dedicated sensitivity studies of

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



selected dust storm cases should be carried out to clarify the likely causes of this model shortcoming. It is notable that both regions influencing the AERONET stations – Agoufou in Northern Hemisphere spring and Tamanrasset in the summer months – are affected by the occurrence of LLJs (compare Fig. 2). Thus here it could be argued that the underestimate in modeled dust events and AOTs could be at least partly due to the underestimate in dust emission fluxes by the morning break-down of nocturnal LLJs.

4 Conclusions

Comparing the results of a two-year regional model simulation of Saharan dust distributions with the DSA record compiled from the MSG IR dust index highlighted some strengths and weaknesses of both the model and the observational dataset. The observations from the MSG IR dust index derived with a backtracking method point toward the foothills of mountains as notable source regions in addition to hot spots like the Bodélé depression (Schepanski et al., 2012). The data also highlight the important role of the development of nocturnal LLJs for the frequency of dust emission events for large parts of the Sahara. LLJs develop above the stable nocturnal boundary layer; after sunrise when the surface heats up their momentum is mixed to the surface by turbulence. In the model this process is also reproduced, but LLJs lead less frequently (and later in the day) to dust emission compared to the observations. While it is possible that LLJs develop too weakly and their degradation is delayed in the model due to an inappropriate boundary layer parameterization, it is also possible that the spatial model resolution is still too coarse to resolve strong gradients in the topography. This would mean that a model gridcell with a horizontal gridsize of 28 km is represented by a high roughness length that leads to suppression of dust emissions in these areas. In contrast, fluvial sediments at mountainsides or foothills of mountains are frequently observed to be active dust sources. The dust emission scheme used in the regional model will need to be adapted to represent such sources appropriately.

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



While the model may be deficient in reproducing dust emission events in mountainous regions, the satellite observations in turn may miss dust emissions that occur below an existing dust cloud. This may be the case for dust emissions related to the passage of cyclonic fronts in the Northern Sahara in the late winter and spring months and moist convective systems in the Sahel and Southern Sahara in summertime. These issues need to be considered when conducting model-data comparisons of dust source activation events.

The model results agree generally well with the AERONET optical thickness data while underpredicting the year-to-year changes in DSA numbers compared to the MSG retrievals. This may be an indication that the frequency of dust emission events is not necessarily a good indicator for dust emission fluxes in the Sahara desert. In addition, the MSG infrared dust index data need to be interpreted carefully, since the magnitude of the IR signal depends on factors like the specific humidity and temperature profiles. Ideally the model should be modified such that it directly simulates the satellite observables to facilitate a quantitative comparison.

Acknowledgement. The authors thank the Deutscher Wetterdienst (DWD) for good cooperation and support. We thank NASA GES DISC for development and maintenance of Giovanni online system providing OMI A1 data. We thank P. Goloub, D. Tanré, B. Mougenot and B. Duchemin, as well as E. Cuevas-Agullo for their efforts in establishing and maintaining the AERONET sites Agoufou, Dakar, Saada and Tamanrasset. A part of this work has been funded by the the BMBF (Bundesministerium für Bildung und Forschung) as part of the SOPRAN project, which is a German national contribution to the International SOLAS project. KS and BH also acknowledge funding from European Research Council Grant No. 257543 “Desert Storms”.

References

- Alpert, P. and Ziv, B.: The Sharav Cyclone: observations and some theoretical considerations, J. Geophys. Res., 94, 18495–18514, doi:10.1029/JD094iD15p18495, 1989.
- Berge, E.: Transboundary air pollution in Europe, in: MSC-W Status Report 1997, Part 1 and 2, EMEP/MSC-W Report 1/97, The Norwegian Meteorological Institute, Oslo, 1997.

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bou Karam, D., Flamant, C., Tulet, P., Chaboureaud, J.-P., Dabas, A., and Todd, M. C.: Estimate of Sahelian dust emissions in the intertropical discontinuity region of the West African Monsoon, *J. Geophys. Res.*, 114, D13106, doi:10.1029/2008JD011444, 2009.

Brindley, H., Knippertz, P., Ryder, C., and Ashpole, I.: A critical evaluation of the ability of the Spinning Enhanced Visible and Infrared Imager (SEVIRI) thermal infrared red-green-blue rendering to identify dust events: theoretical analysis, *J. Geophys. Res.*, 117, D07201, doi:10.1029/2011JD017326, 2012.

Chiappello, I., Moulin, C., and Prospero, J. M.: Understanding the long-term variability of African dust transport across the Atlantic as recorded in both Barbados surface concentrations and large-scale Total Ozone Mapping Spectrometer (TOMS) optical thickness, *J. Geophys. Res.*, 110, D18S10, doi:10.1029/2004JD005132, 2005.

DeMott, P. J., Sassen, K., Poellot, M. R., Baumgardner, D., Rogers, D. C., Brooks, S. C., Prenni, A. J., and Kreidenweis, S. M.: African dust aerosols as atmospheric ice nuclei, *Geophys. Res. Lett.*, 30, 1732, doi:10.1029/2003GL017410, 2003.

Dubovik, O., Holben, B. N., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D., and Slutsker, I.: Climatology of absorption and optical properties of key aerosol types, *J. Atmos. Sci.*, 59, 590–608, 2002.

Heinold, B., Helmert, J., Hellmuth, O., Wolke, R., Ansmann, A., Marticorena, B., Laurent, B., and Tegen, I.: Regional modeling of Saharan dust events using LM-MUSCAT: model description and case studies, *J. Geophys. Res.*, 112, D11204, doi:10.1029/2006JD007443, 2007.

Heinold, B., Tegen, I., Esselborn, M., Kandler, K., Knippertz, P., Müller, D., Schladitz, A., Tesche, M., Weinzierl, B., Ansmann, A., Althausen, D., Laurent, B., Massling, A., Müller, T., Petzold, A., Schepanski, K., and Wiedensohler, A.: Regional Saharan dust modelling during the SAMUM 2006 campaign, *Tellus B*, 61, 307–324, 2009.

Heinold, B., Tegen, I., Schepanski, K., Tesche, M., Esselborn, M., Freudenthaler, V., Gross, S., Kandler, K., Knippertz, P., Müller, D., Schladitz, A., Toledano, C., Weinzierl, B., Ansmann, A., Althausen, D., Müller, T., Petzold, A., and Wiedensohler, A.: Regional modelling of Saharan dust and biomass-burning smoke, *Tellus B*, 63, 781–799, doi:10.1111/j.1600-0889.2011.00570.x, 2011.

Helmert, J., Heinold, B., Tegen, I., Hellmuth, O., and Wendisch, M.: On the direct and semi-direct effect of Saharan dust over Europe: a case study, *J. Geophys. Res.*, 112, D13208, doi:10.1029/2006JD007444, 2007.

Holben B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y., Nakajima, T., Lavenu, F., Jankowiak I., and Smirnov, A.: AERONET – a federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16, 1998.

5 Horvath, K., Fita, L., Romero, R., and Ivancan-Picek, B.: A numerical study of the first phase of a deep Mediterranean cyclone: cyclogenesis in the lee of the Atlas Mountains, *Meteor. Z.*, 15, 133–146, doi:10.1127/0941-2948/2006/0113, 2006.

Huneus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A., Horowitz, L., Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J.-J., Myhre, G., Penner, J., Perlwitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model intercomparison in AeroCom phase I, *Atmos. Chem. Phys.*, 11, 7781–7816, doi:10.5194/acp-11-7781-2011, 2011.

15 Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, N. N., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., LaRoche, J., Liss, P. J., Mahowald, N. M., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry and climate, *Science*, 308, 67–71, 2005.

20 Koch, J. and Renno, N. O.: The role of convective plumes and vortices on the global aerosol budget, *Geophys. Res. Lett.*, 32, L18806, doi:10.1029/2005GL023420, 2005.

Knippertz, P. and Todd, M. C.: Mineral dust aerosols over the Sahara: meteorological controls on emission and transport and implications for modeling, *Rev. Geophys.*, 50, RG1007, doi:10.1029/2011RG000362, 2012.

25 Knippertz, P., Trentmann, J., and Seifert, A.: Highresolution simulations of convective cold pools over the Northwestern Sahara, *J. Geophys. Res.*, 114, D08110, doi:10.1029/2008JD011271, 2009.

Lacis, A. A. and Mishchenko, M. I.: Climate forcing, climate sensitivity and climate response: a radiative modeling perspective on atmospheric aerosols, in: *Aerosol Forcing of Climate*, edited by: Charlson, R. J. and Heintzenberg, J., John Wiley & Sons, Berlin, 11–42, 1995.

30 Laurent, B., Marticorena, B., Bergametti, G., Léon, J. F., and Mahowald, N. M.: Modeling mineral dust emissions from the Sahara desert using new surface and soil developments, *J. Geophys. Res.*, 113, D14218, doi:10.1029/2007JD009484, 2008.

ACPD

12, 27667–27691, 2012

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Laurent, B., Tegen, I., Heinold, B., Schepanski, K., Weinzierl, B., and Esselborn, M.: A model study of Saharan dust emissions and distributions during the SAMUM-1 campaign, *J. Geophys. Res.*, 115, D21210, doi:10.1029/2009JD012995, 2010.
- Mahowald, N. M., Muhs, D. R., Levis, S., Rasch, P. J., Yoshioka, M., Zender, C. S., and Luo, C.: Change in atmospheric mineral aerosols in response to climate: last glacial period, preindustrial, modern, and doubled carbon dioxide climates, *J. Geophys. Res.*, 111, D10202, doi:10.1029/2005JD006653, 2006.
- Marticorena, B. and Bergametti, G.: Modeling the atmospheric dust cycle. 1. Design of a soil-derived dust production scheme, *J. Geophys. Res.*, 100, 16415–16430, 1995.
- Marshall, J. H., Knippertz, P., Dixon, N. S., Parker, D. J., and Lister, G. M. S.: The importance of the representation of deep convection for modeled dust-generating winds over West Africa during summer, *Geophys. Res. Lett.*, 38, L16803, doi:10.1029/2011GL048368, 2011.
- Prospero, J. M. and Lamb, P. J.: African droughts and dust transport to the Caribbean: climate change implications, *Science*, 302, 1024–1027, 2003.
- Reinfried, F., Tegen, I., Heinold, B., Hellmuth, O., Schepanski, K., Cubasch, U., Hübener, H., and Knippertz, P.: Density currents in the Atlas Region leading to dust mobilization: a model sensitivity study, *J. Geophys. Res.*, 114, D08127, doi:10.1029/2008JD010844, 2009.
- Schepanski, K., Tegen, I., Laurent, B., Heinold, B., and Macke, A.: A new Saharan dust source activation frequency map derived from MSG-SEVIRI IR-channels, *Geophys. Res. Lett.*, 34, 18803, doi:10.1029/2007GL030168, 2007.
- Schepanski, K., Tegen, I., Todd, M. C., Heinold, B., Bönisch, G., Laurent, B., and Macke, A.: Meteorological processes forcing Saharan dust emission inferred from MSG-SEVIRI observations of sub-daily source activation and numerical models, *J. Geophys. Res.*, 114, D10201, doi:10.1029/2008JD010325, 2009.
- Schepanski, K., Tegen, I., and Macke, A.: Comparison of satellite based observations of Saharan dust source areas, *Remote Sens. Environ.*, 123, 90–97, doi:10.1016/j.rse.2012.03.019, 2012.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics*, John Wiley & Sons, New York, 1326 pp., 1998.
- Sokolik, I. N., Winker, D. M., Bergametti, G., Gillette, D. A., Carmichael, G., Kaufman, Y. J., Gomes, L. Schuetz, L., and Penner, J. E.: Introduction to special section: outstanding problems in quantifying the radiative impacts of mineral dust, *J. Geophys. Res.*, 106, 18015–18027, doi:10.1029/2000JD900498, 2001.

Comparing two years of Saharan dust source activation

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Comparing two years
of Saharan dust
source activation**

I. Tegen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solomon, S., Qin, D., Manning, M., Marquis, M., Averyt, K., Tignor, M. M. B., and Miller, H. L. (Eds.): Climate Change 2007: The Physical Science Basis, Cambridge University Press, UK, 989 pp., 2007.

Steppeler, J., Doms, G., Schättler, U., Bitzer, H. W., Gassmann, A., Damrath, U., and Gregoric, G.: Meso-gamma scale forecasts using the nonhydrostatic model LM, Meteorol. Atmos. Phys., 82, 75–96, 2003.

Tegen, I., Harrison, S. P., Kohfeld, K. E., and Prentice, I. C.: Impact of vegetation and preferential source areas on global dust aerosol: results from a model study, J. Geophys. Res., 107, 4576, doi:10.1029/2001JD000963, 2002.

Todd, M. C., Bou Karam, D., Cavazos, C., Bouet, C., Heinold, B., Baldasano, J. M., Cautenet, G., Koren, I., Perez, C., Solmon, F., Tegen, I., Tulet, P., Washington, R., and Zakey, A.: Quantifying uncertainty in estimates of mineral dust flux: An intercomparison of model performance over the Bodélé Depression, Northern Chad, J. Geophys. Res., 113, D24107, doi:10.1029/2008JD010476, 2008.

Torres, O., Tanskanen, A., Veihelmann, B., Ahn, C., Braak, R., Bhartia, P. K., Veefkind, P., and Levelt, P.: Aerosols and surface UV products from ozone monitoring instrument: an overview, J. Geophys. Res., 112, D24S47, doi:10.1029/2007JD008809, 2007.

Washington, R. and Todd, M. C.: Atmospheric controls on mineral dust emission from the Bodele depression, Chad: the role of the low level jet, Geophys. Res. Lett., 32, L17701, doi:10.1029/2005GL023597, 2005.

Wolke, R., Hellmuth, O., Knoth, O., Schröder, W., Heinrich, B., and Renner, E.: The chemistry-transport modeling system LM-MUSCAT: description and CITYDELTA applications, in: Air Pollution Modeling and its Application XVI, edited by: Borrego, C. and Incecik, S., Kluwer Academic/Plenum Publishers, New York, USA, 427–439, 2004.

Zobler, L.: A world soil file for global climate modeling, Tech. Rep. NASA TM-87802, NASA, Washington, D. C., 32 pp., 1996.

Comparing two years of Saharan dust source activation

I. Tegen et al.

Table 1. Sahara dust results from the regional model COSMO-MUSCAT for the years 2007 and 2008. DSA: Dust source activation. Over- and underpredictions refer to the percentage of grid cells between 15° W, 10° N and 32° E, 40° N that are activated compared to MSG IR dust index.

Date	2007	2008
Model annual dust emission (Mt)	1870	2330
Emission count (Model)	11 140	14 442
Emission events with LLJ	40 %	39 %
Emission count (MSG)	7139	19 110
Percent grid cells overpredicting DSA	29 %	23 %
Overpredictions with LLJ	12 %	9 %
Overpredictions without LLJ	17 %	14 %
Percent grid cells underpredicting DSA	10 %	16 %
Underpredictions with added LLJs	1 %	3 %

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 1. Number of dust emission events in the Sahara for the years 2007 and 2008 derived from the MSG infrared dust index (top) and computed with the COSMO-MUSCAT model (bottom). The colors indicate the time of day when dust emission starts. Grey: 00:00–06:00 UTC, blue: 06:00–12:00 UTC, orange: 12:00–18:00 UTC, purple: 18:00–00:00 UTC.




Fig. 2. Distribution of the percent of days when dust source activation occurs in the COSMO-MUSCAT model simulations (left panels) and derived from the MSG infrared dust index (middle panels). Right panels: occurrence of nocturnal low level jets in the COSMO model simulations. Results are average values for the years 2007 and 2008, averaged for the months February to May (top panels), June to September (middle panels) and October to January (bottom panels). The model results are binned into $1^\circ \times 1^\circ$ gridcells to facilitate comparison with the MSG data.

Comparing two years of Saharan dust source activation

I. Tegen et al.




Fig. 3. Number of days per month with dust emission derived from the MSG SEVIRI IR dust index (a, b) and simulated with the COSMO-MUSCAT model (c, d); for year 2007 (left) and 2008 (right). The red and orange contours indicate areas where the occurrence of low level jets in the model is higher than 5 and 10 days per months, respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Fig. 4. Annual averaged dust optical thickness for the years 2007 (left panels) and 2008 (right panels) computed with COSMO-MUSCAT (**a, b**); annual averaged absorbing aerosol index from the OMI satellite instrument (**c, d**).

Comparing two years of Saharan dust source activation

I. Tegen et al.




Fig. 5. Time series of daily modeled dust aerosol optical thickness (red line) and AOT measurements at 440 nm from AERONET at the stations Tamanrasset (22.8° N, 5.5° E), Saada (31.6° N, 8.2° W) Dakar (14.4° N, 17.0° W) and Agoufou (15.3° N; 1.5° W) (black plus signs) for 2007 and 2008. The black dots represent AOT measurements for Ångström parameters larger than 0.7 where dust AOT is not dominant.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)