

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A one-year comprehensive chemical characterisation of fine aerosol (PM_{2.5}) at urban, suburban and rural background sites in the region of Paris (France)

M. Bressi^{1,2}, J. Sciare¹, V. Ghersi³, N. Bonnaire¹, J. B. Nicolas^{1,2}, J.-E. Petit¹, S. Moukhtar³, A. Rosso³, N. Mihalopoulos⁴, and A. Féron¹

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE, UMR8212, CNRS-CEA-UVSQ, Gif-sur-Yvette, 91191, France

²French Environment and Energy Management Agency, ADEME, 20 avenue du Grésillé, BP90406 49004, Angers Cedex 01, France

³AIRPARIF, Surveillance de la Qualité de l'Air en Ile-de-France, Paris, 75004, France

⁴Environmental Chemical Processes Laboratory, ECPL, Heraklion, Voutes, Greece

Received: 26 March 2012 – Accepted: 25 October 2012 – Published: 15 November 2012

Correspondence to: M. Bressi (michael.bressi@ensiacet.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

Studies describing the chemical composition of fine aerosol (PM_{2.5}) in urban areas are often conducted during few weeks only, and at one sole site, giving thus a narrow view of their temporal and spatial characteristics. This paper presents a one-year (11 September 2009–10 September 2010) survey of the daily chemical composition of PM_{2.5} in the region of Paris, which is the second most populated “Larger Urban Zone” in Europe. Five sampling sites representative of suburban (SUB), urban (URB), north-east (NER), northwest (NWR) and south (SOR) rural backgrounds were implemented. The major chemical components of PM_{2.5} were determined including elemental carbon (EC), organic carbon (OC), and the major ions. OC was converted to organic matter (OM) using the chemical mass closure methodology, which leads to conversion factors of 1.95 for the SUB and URB sites, and 2.05 for the three rural ones. On average, gravimetrically determined PM_{2.5} annual mass concentrations are 15.2, 14.8, 12.6, 11.7 and 10.8 µg m⁻³ for SUB, URB, NER, NWR and SOR sites, respectively. The chemical composition of fine aerosol is very homogeneous at the five sites and is composed of OM (38–47 %), nitrate (17–22 %), non-sea-salt sulfate (13–16 %), ammonium (10–12 %), EC (4–10 %), mineral dust (2–5 %) and sea salt (3–4 %). This chemical composition is in agreement with those reported in the literature for most European environments. On the annual scale, Paris (URB and SUB sites) exhibits its highest PM_{2.5} concentrations during late autumn, winter and early spring (higher than 15 µg m⁻³ on average, from December to April), intermediates during late spring and early autumn (between 10 and 15 µg m⁻³ during May, June, September, October, and November) and the lowest during summer (below 10 µg m⁻³ during July and August). PM levels are mostly homogeneous at the regional scale, on the whole duration of the project (e.g. for URB plotted against NER sites: slope = 1.06, $r^2 = 0.84$, $n = 330$), suggesting the importance of mid-or long-range transport, and regional instead of local scale phenomena. During this one-year project, two third of the days exceeding the PM_{2.5} 2015 EU annual limit value of 25 µg m⁻³ were due to continental import from countries located northeast, east of

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



France. This result questions the efficiency of local, regional and even national abatement strategies during pollution episodes, pointing the need for a wider collaborative work with the neighbourhood countries on these topics. Nevertheless, emissions of local anthropogenic sources lead to higher levels at the URB and SUB sites compared to the others (e.g. 26% higher on average at the URB than at the NWR site for PM_{2.5}, during the whole campaign), which can even be emphasised by specific meteorological conditions such as low boundary layer heights. OM and secondary inorganic species (nitrate, non-sea-salt sulfate and ammonium, noted SIA) are mainly imported by mid- or long-range transport (e.g. for NWR plotted against URB sites: slope = 0.79, $r^2 = 0.72$, $n = 335$ for OM, and slope = 0.91, $r^2 = 0.89$, $n = 335$ for SIA) whereas EC is primarily locally emitted (e.g. for SOR plotted against URB sites: slope = 0.27; $r^2 = 0.03$; $n = 335$). This database will serve deepest investigations of carbonaceous aerosols, metals as well as the main sources and geographical origins of PM in the region of Paris.

1 Introduction

Adverse health effects of aerosols and especially of fine particles (PM_{2.5} i.e. particulate matter with aerodynamic diameter, AD, below 2.5 μm) have been widely demonstrated (Bernstein et al., 2004; Pope et al., 2004), especially in urban areas (Lawrence et al., 2007). Paris is highly concerned by these impacts as it is the second "Larger Urban Zone" in Europe with 11 million inhabitants, i.e. 18% of the French population (Eurostat, 2011). A recent study (Aphekom, 2011) reported that a reduction of PM_{2.5} concentrations in Paris (average 2004–2006: 16.4 $\mu\text{g m}^{-3}$; Airparif, 2012) towards the World Health Organization recommendation value (10 $\mu\text{g m}^{-3}$) would lead to a gain in life expectancy of 5.8 months for 30 yr old and older persons. In addition, epidemiologists and toxicologists suggest investigating chemical and physical characteristics of particles in order to better assess their toxicity (Schlesinger, 2007; Ramgolam et al., 2008).

Besides, climate effects of aerosols have been a subject of concern for more than 20 yr (WCP, 1983; Ramanathan et al., 1987). Whereas long-lived greenhouse gases and ozone contribute a positive radiative forcing (RF) of $+2.9 (\pm 0.3) \text{ W m}^{-2}$, the combined aerosol direct and cloud albedo effect have a median RF of -1.3 W m^{-2} and a -2.2 to -0.5 W m^{-2} 90 % confidence range (Forster et al., 2007). To better estimate climate effects of aerosols, their chemical composition has to be exhaustively documented as each chemical component will play a specific role in the direct (i.e. the scattering and absorbance of solar and infrared radiation in the atmosphere) and indirect effects (i.e. the modification of the formation and precipitation efficiency of liquid water, ice and mixed-phase clouds) on climate (Forster et al., 2007; Isaksen et al., 2009).

Because of health and climate impacts of particles, limit values of $\text{PM}_{2.5}$ and PM_{10} (particles with an AD below $10 \mu\text{m}$) determined by the European Union (EU) became more stringent in recent years. As $\text{PM}_{2.5}$ represents 50–90 % of PM_{10} mass in most European environments (Putaud et al., 2010), the conclusions drawn in this paper will also help for the understanding of PM_{10} . Concerning PM_{10} , the actual EU daily limit value is $50 \mu\text{g m}^{-3}$ and has not to be exceeded more than 35 days per year (European Directive 2008/50/EC). During 2010, this limit value has been exceeded from 42 to 176 days at seven traffic sites in the region of Paris, thus affecting 1.8 million inhabitants i.e. 16 % of the regional population (Airparif, 2012). In May 2011, France has even been summoned to the Court of Justice of the EU because of these PM_{10} exceedances. Concerning $\text{PM}_{2.5}$, since 2007 annual levels have constantly been around $20 \mu\text{g m}^{-3}$ at an urban background (Airparif, 2012), which is under the 2015 EU annual limit value of $25 \mu\text{g m}^{-3}$, but equal to the one planned for 2020. Therefore, there is a clear need to better understand the origin and chemical composition of PM over the region of Paris to tackle health, climate and legislative issues.

Studies describing the chemical composition of aerosols in the region of Paris are however scarce and were mostly conducted over short time periods (typically few weeks), at only one sampling site (e.g. Hodzic et al., 2006; Favez et al., 2007; Gros et al., 2007; Sciare et al., 2010, 2011). Although they bring valuable information on

Chemical characterisation of $\text{PM}_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the physical and chemical characteristics of PM in this region, they do not address their temporal and spatial evolution on a large scale. In particular, the seasonality of the chemical processes governing PM mass and chemical composition, as well as the annual evolution of the major emission sources are still poorly known.

To fill these gaps, a research project involving the Climate and Environmental Sciences Laboratory (LSCE) and the regional air quality network of Paris (AIRPARIF) has been implemented. This LSCE-AIRPARIF “Particles” project goes beyond the scope of this paper and its full description can be found in Ghersi et al. (2010) and Airparif and LSCE (2012). Briefly, its general strategy aims at documenting the daily chemical composition of PM₁₀ and PM_{2.5} in different types of environments representative of urban, suburban, rural and traffic sites, during a one-year period (11 September 2009–10 September 2010). It should allow the identification of the major sources of PM as well as their geographical origins in order to implement effective abatement policies (Ghersi et al., 2012). In addition, a detailed study of carbonaceous aerosols has been performed and will help in better documenting its main sources (including traffic and domestic wood burning) and its atmospheric processing. Finally, a focus on metal concentrations will be made in order to describe their main primary sources, size distribution and temporality in the region of Paris. This “Particles” project will be helpful to put into perspective the recent intensive field campaigns performed within the European project MEGAPOLI (Megacities: emissions, urban, regional and Global Atmospheric POLLution and climate effects, and Integrated tools for assessment and mitigation), by providing a spatially and temporally extended view of the PM chemical composition in the entire region of Paris.

A focus will be made here on the daily chemical composition of fine aerosol determined during one-year at one urban, one suburban and three rural sites of the region of Paris. This paper aims at presenting this chemical dataset including a description and evaluation of measurement methods, and a first analysis of its spatio-temporal variability. More precisely, Sect. 2 will describe (i) sampling sites, (ii) analytical techniques, (iii) chemical mass closure methodology and (iv) organic carbon to organic matter

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



conversion factor's methodology. Section 3 will present (i) the representativeness of the studied period, (ii) first results regarding the chemical composition of PM_{2.5}, (iii) an evaluation of measurement methods and (iv) the adequate OC to OM conversion factor to use. Finally, Sect. 4 will discuss (i) the temporal variability of PM_{2.5} with an emphasis on meteorological parameters that can explain daily and seasonal variations of specific chemical compounds; and (ii) the spatial variability of PM_{2.5} and major chemical compounds hence giving an insight into their geographical origins (local versus regional or transboundary).

2 Material and methods

2.1 Sampling sites description

Six sampling sites were implemented, covering the region of Paris (see Fig. 1). These sites are part of the AIRPARIF Air Quality monitoring network, and are regarded as being representative of background conditions (Gherzi et al., 2010). They were categorised according to criteria proposed by the European Environment Agency (Larssen et al., 1999). The first site is an urban (URB) station located in the city centre of Paris (4th district, 48°50'56" N, 02°21'55" E, 20 m above ground level, a.g.l.). The second site is a near-city or suburban (SUB) station located at 10 km northeast of the UR station (48°52'54" N, 02°30'23" E, 5 m a.g.l.). The third, fourth and fifth sites are rural stations located respectively at ca. 65 km northeast (NER), 50 km northwest (NWR) and 60 km south (SOR) of the URB station (49°05'15" N, 03°04'35" E, 5 m a.g.l.; 49°03'48" N, 01°51'59" E, 5 m a.g.l. and 48°21'49" N, 02°14'07" E, 5 m a.g.l., respectively). A traffic (TR) station located at the ring road of Paris, at 9 km west of the UR station was also implemented (48°51'02" N, 2°15'09" E, 5 m a.g.l.). This last station will not be described here due to its specificity (traffic sources) whereas this paper aims at describing representative urban, suburban and rural backgrounds.

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Aerosol sampling

Fine aerosol particles ($PM_{2.5}$) were collected at each site every day during 24 h (from 00:00 to 23:59 LT) during one year (from 11 September 2009 to 10 September 2010). Filter sampling was performed using two collocated Leckel low volume samplers (SEQ47/50) at each station running at $2.3\text{ m}^3\text{ h}^{-1}$. One Leckel sampler was equipped with quartz filters (QMA, Whatman, 47 mm diameter) for carbon analyses, the second with Teflon filters (PTFE, Pall, 47 mm diameter, $2.0\text{ }\mu\text{m}$ porosity) for gravimetric and ion measurements. Before being sampled, QMA filters were baked at $480\text{ }^\circ\text{C}$ for 48 h and PTFE filters pre-weighed as reported in Sciare et al. (2003). Field blanks were taken every two weeks for PTFE filters, and every week for QMA filters. A total of 4040 filters has been collected including 2085 QMA and 1955 PTFE filters. Few samples have been discarded because of power failures, chemical analysis problems, etc. (Supplement Table S1) and represent ca. 5 % of QMA and 5 % of PTFE filters. Once sampled, filters were stored at $-20\text{ }^\circ\text{C}$ in a freezer prior to chemical analyses.

2.3 Chemical analyses

2.3.1 Gravimetry

In order to minimise the influence of water adsorption, loaded and unloaded PTFE filters were equilibrated for 48 h at ambient temperature and below 30 % relative humidity (RH) prior to being weighed (MacMurry, 2000; Sciare et al., 2005). PTFE filters were then weighed with a microbalance (Sartorius, MC21S) with $1\text{ }\mu\text{g}$ sensitivity. Filter weighing was repeated until having a difference between two weighs below $5\text{ }\mu\text{g}$. The overall error is estimated to be $10\text{ }\mu\text{g}$ which represents $3 \pm 2\%$ of loaded filters ($n = 1723$; all sites, one-year measurements, blank filters excluded). Aerosol masses (PM_{grav}) were deduced from the gravimetric measurements done before and after sampling. Field blanks taken on the field show an averaged PM_{grav} of $7 \pm 11\text{ }\mu\text{g}$ ($n = 130$),

Chemical characterisation of $PM_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which represents 2 ± 1 % of loaded filters ($n = 1723$), being thus below the overall microbalance error.

2.3.2 Ions

The following water-soluble major ions were analysed by Ion Chromatographs (IC): chloride, nitrate, sulfate, sodium, ammonium, potassium, magnesium and calcium. The analytical protocol followed is thoroughly described by Sciare et al. (2008) and Guinot et al. (2007). Briefly, filter samples were extracted in 15 ml of Milli-Q water during 45 min in a sonic bath. To prevent bacteria activity, 50 μ l of chloroform was added. Samples were then filtered using Acrodisc filters (Pall Gelmann) with a porosity of 0.4 μ m. Cations were analysed on a 2-mm diameter CS12 pre-column and column with an IC (Dionex, Model DX-600, USA), anions on a 2-mm diameter AS11 pre-column and column with an IC (Dionex, Model DX-600, USA). Both IC apparatus were equipped with a reagent free system (automated eluent generation and self-regenerating suppression). Semi-annual laboratory IC inter-comparison studies were performed (accessible at <http://qasac-america.org/lis/summary/44>) and showed errors of less than 5 % for every cited ion. Field blank measurement medians ($n = 130$) are 26, 29, 4, 9, 9, 4, 2 and 22 ppb for Cl^- , NO_3^- , SO_4^{2-} , Na^+ , NH_4^+ , K^+ , Mg^{2+} , and Ca^{2+} , respectively, which represent 16, 2, 0, 5, 1, 3, 6 and 24 % of the medians of loaded filters ($n = 1723$), respectively. Blank corrections have only been performed for chloride and calcium ions by subtracting blank medians to the loaded filter values.

2.3.3 Carbon

Elemental carbon (EC) and organic carbon (OC) were determined by a thermal-optical method using a Sunset Laboratory Carbonaceous Analyser (Sunset Lab., OR, USA) and the EUSAAR_2 protocol defined by Cavalli et al. (2010). A detailed description concerning thermal-optical methods and the relevance of the protocol chosen can be found elsewhere (Chow et al., 1993; Schmid et al., 2001; Cavalli et al., 2010). The

Chemical characterisation of $\text{PM}_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



detection limit and the uncertainty given by the Sunset Company are estimated to be 0.2 µgC and 5 %, respectively, for EC and OC measurements. EC was not detected in field blanks ($0.0 \pm 0.1 \mu\text{gC cm}^{-1}$, $n = 252$), and OC was found with an average value of $1.1 \pm 0.5 \mu\text{gC cm}^{-2}$ ($n = 252$). On average, this field blank value represents $15 \pm 7 \%$ of sampled filter's OC concentrations ($n = 1723$). Blank corrections have been performed for OC concentrations by subtracting the blank average to the sampled filter values.

2.4 Chemical Mass Closure

Aerosol Chemical Mass Closure (CMC) consists in comparing the sum of the major aerosol chemical species (PM_{chem}) with gravimetric measurements (PM_{grav}). When achieved (i.e. when $\text{PM}_{\text{chem}} = \text{PM}_{\text{grav}}$), CMC attests the consistency of chemical analyses, and confirms that all the major aerosol chemical species are taken into account. PM_{chem} calculation is here expressed as:

$$[\text{PM}_{\text{chem}}] = [\text{Sea Salt}] + [\text{Dust}] + [\text{Secondary Inorganic Aerosols}] + [\text{Carbonaceous Matter}] \quad (1)$$

Sea salt (ss-) concentrations are calculated from the six major ions accounting for more than 99 % of the mass of salts dissolved in seawater:

$$[\text{Sea salt}] = [\text{Na}^+] + [\text{Cl}^-] + [\text{Mg}^{2+}] + [\text{ss-K}^+] + [\text{ss-Ca}^{2+}] + [\text{ss-SO}_4^{2-}] \quad (2)$$

with $[\text{ss-K}^+] = 0.036 \times [\text{Na}^+]$; $[\text{ss-Ca}^{2+}] = 0.038 \times [\text{Na}^+]$ and $[\text{ss-SO}_4^{2-}] = 0.252 \times [\text{Na}^+]$

Typical seawater ion ratios based on the average seawater composition are taken from Seinfeld and Pandis (1998).

Different methods are used to calculate the mineral dust fraction in $\text{PM}_{2.5}$, and are based on its average elemental composition from specific sites, or on specific tracers (Pettijohn, 1975; Malm et al., 1994; Guieu et al., 2002). Recently, nss-calcium has been used to estimate mineral dust in aerosols because of its abundance (Putaud et al., 2004a; Sciare et al., 2005). We used the 15 % contribution of nss-calcium in mineral

Chemical characterisation of $\text{PM}_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dust determined by Guinot et al. (2007) in Paris:

$$[\text{Dust}] = [\text{nss-Ca}^{2+}] / 0.15 \quad (3)$$

This is in agreement with the ratio of 18 % reported by Putaud et al. (2004b) in Monte Cimone (Italy) during non-Saharan dust periods. The resulting proportion of dust in PM_{2.5} during the whole campaign ranges from 2 to 5 % on average at the five sites (see Sect. 4.2.2). By changing this ratio by ± 3 %, the resulting proportion of dust in PM_{2.5} remains very low at the five sites (3 to 6 % and 2 to 4 % of PM_{2.5} with nss-Ca²⁺ to dust ratios of 12 and 18 %, respectively). In the frame of chemical mass closure, further investigation allowing the estimation of dust will thus not be conducted.

Secondary inorganic aerosols (SIA) are calculated as:

$$[\text{Secondary Inorganic Aerosols}] = [\text{nss-SO}_4^{2-}] + [\text{NO}_3^-] + [\text{NH}_4^+] \quad (4)$$

where $[\text{nss-SO}_4^{2-}] = [\text{SO}_4^{2-}] - [\text{ss-SO}_4^{2-}]$, “nss-” standing for “non-sea-salt”.

Finally, carbonaceous matter can be expressed as:

$$[\text{Carbonaceous Matter}] = [\text{EC}] + [\text{OM}] \quad (5)$$

with $[\text{OM}] = f_{\text{OC-OM}} \times [\text{OC}]$.

The estimation of organic matter and more specifically of $f_{\text{OC-OM}}$ will now be discussed.

2.5 Organic carbon to organic matter conversion factor

Organic matter (OM) is here inferred from filter OC measurements determined by the Sunset Laboratory analyser. The estimation of OM is of high complexity due to its varied chemical composition which changes according to location, season and time of the day (Turpin et al., 2000; Andrews et al., 2000). Turpin and Lim (2001) recommend the measurement of the average molecular weight per carbon weight in the location of interest; such measurements were not conducted in this study. In that case, they suggest the use of an OC to OM conversion factor ($f_{\text{OC-OM}}$) of 1.6 ± 0.2 for urban aerosols

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and 2.1 ± 0.2 for nonurban aerosols. These factors were widely used in recent peer reviewed publications (Terzi et al., 2010; Cheung et al., 2011; Rengarajan et al., 2011) despite their spatial and temporal dependences. We decided to estimate $f_{\text{OC-OM}}$ from our dataset by using a method adapted from Guinot et al. (2007) who used the chemical mass closure technique as a tool to assess $f_{\text{OC-OM}}$ from OC measurements.

To infer $f_{\text{OC-OM}}$, we assumed that CMC is achieved ($\text{PM}_{\text{grav}} = \text{PM}_{\text{chem}}$), and used Eq. (1) to write the following equation:

$$f_{\text{OC-OM}} = 1/[\text{OC}] \times ([\text{PM}_{\text{grav}}] - ([\text{Sea Salt}] + [\text{Dust}] + [\text{SIA}] + [\text{EC}])) \quad (6)$$

This will allow us to find the “best guess” $f_{\text{OC-OM}}$ value from our dataset. To simplify Eq. (6), a chemical fraction named “Remaining mass” (RM) has been defined in Eq. (7):

$$[\text{Remaining Mass}] = [\text{PM}_{\text{grav}}] - ([\text{Sea Salt}] + [\text{Dust}] + [\text{S/A}] + [\text{EC}]) \quad (7a)$$

thus leading to:

$$[\text{Remaining Mass}] = f_{\text{OC-OM}} \times [\text{OC}] \quad (7b)$$

Two different methods were then used to estimate $f_{\text{OC-OM}}$. In the first method, RM concentrations are plotted against OC concentrations and $f_{\text{OC-OM}}$ is the slope of the linear regression (Eq. 7). We decided to use a linear function without y-intercept because of the analytical form of this equation. In the second method, we numerically calculated $f_{\text{OC-OM}}$ day-by-day from Eq. (6). Days with $f_{\text{OC-OM}}$ higher than 3 and lower than 1 were considered as physically meaningless and were therefore excluded from the dataset. These days represent 6 to 14 % of the samples according to sites. Results will be discussed in Sect. 3.5.

2.6 Additional measurements

Meteorological parameters such as temperature, pressure, precipitation, wind speed and wind direction were provided by the French national meteorological service

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



“Meteo-France” from measurements recorded at Montsouris (14th district, 48°49′20″ N, 02°20′18″ E) located at about 5 km south of the URB station. Boundary layer heights (BLH) were determined by radiosondes by Meteo-France at Trappes (48°46′26″ N; 02°00′31″ E) located at ca. 30 km southwest of the URB station. Air mass back trajectories were calculated using the Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPPLIT) model (Draxler and Rolph, 2011).

3 Results

3.1 Representativeness of the campaign

The meteorological conditions and the PM level representativeness of the studied period were investigated and reported in Supplement Fig. S1. Air temperatures measured during our campaign were comparable to standard values determined by Meteo-France (calculated following Arguez and Vose, 2011), showing however lower values during winter (DJF, Fig. S1a). Levels of precipitation showed discrepancies in comparison with standard values (Fig. S1b). More specifically, September 2009 and April 2010 were particularly dry, unusual snow events occurred during January 2010 and heavy rains during July and August 2010. Paris is typically characterised by a dominance of south to southwest winds (ca. 35 %) and to a lesser extent north to northeast winds (ca. 20 %) (Fig. S1c). The studied period showed similar trends although a stronger contribution of the northeast sector was observed.

Finally, our yearly average PM_{2.5} mass concentration of 18.4 µg m⁻³ was characteristic of a usual year (R&P Tapered Element Oscillating Microbalance – Filter Dynamic Measurement System, TEOM-FDMS, (Patashnik and Rupprecht, 1991) data corrected for semi-volatile materials at the urban site). It was lower than in 2007 (ca. -11 %), but higher than in 2008 (ca. +16 %, values calculated from TEOM-FDMS data measured at a similar urban site of the Airparif network).

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Chemical composition of PM_{2.5}

Table 1 reports statistics on the chemical composition of PM_{2.5} at the five sites, on the whole sample set. PM_{2.5} levels range from 10.8 to 15.2 µg m⁻³ on average according to sites. Fine aerosols are primarily made of OC (2.1–3.2 µg m⁻³), nitrate (2.2–2.9 µg m⁻³), sulfate (1.8–2.1 µg m⁻³) and ammonium (1.2–1.5 µg m⁻³); and to a lesser extent of EC (0.4–1.4 µg m⁻³) and minor ions (less than 0.2 µg m⁻³ for Cl⁻, Na⁺, K⁺, Ca²⁺ and Mg²⁺). Further discussion will be conducted in Sect. 4.

3.3 Comparison between filter and on-line determined masses

We investigated the atmospheric consistency of our PM_{2.5} measurements, and attempted to estimate artefacts associated with filter sampling (Zhang and McMurry, 1987; Mc Dow and Huntzicker, 1990; Turpin et al., 2000). A conventional on-line automatic system (TEOM-FDMS) was running during the campaign at the urban station only and is used by Airparif since 2007 (Airparif, 2012). Gravimetric and TEOM-FDMS determined masses are compared in Figs. 2 and 3. TEOM-FDMS was running at 30 °C and was not corrected for semi-volatile materials in order to be as close as possible to our laboratory conditions. Very similar temporal variations are observed in Fig. 2 for both datasets for the whole duration of the campaign. TEOM-FDMS data plotted against gravimetric mass concentrations show a very good correlation ($r^2 = 0.94$, $n = 318$, Fig. 3). However, filter measurements exhibit mass concentrations about 6% higher than the on-line method (slope ± 1 standard error = 0.938 ± 0.007), which can be related to temperature differences during filter sampling (ambient temperature) and TEOM-FDMS measurements (30 °C). Absorption of volatile organic compounds (VOC) (Turpin et al., 1994, 2000) and/or water onto filters (Quinn and Coffman, 1998; Speer et al., 2003; Hueglin et al., 2005) could also enhance filter masses. In addition, taking into account semi-volatile materials in TEOM-FDMS measurements leads to higher concentrations compared with the gravimetric method (18.4 versus 14.8 µg m⁻³, respectively, on average at the URB site). The former method is regarded as an equivalent

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



method to the EU reference method (EN 14907), whereas the latter does not fulfil EU requirements (by operating below 30 % RH instead of at 50 % RH). It should thus be borne in mind that our gravimetric method will underestimate PM_{2.5} mass compared to EU reference methods by ca. 20 % on average.

3.4 Comparison between filters and on-line determined chemical components

The chemical results obtained at the urban site were compared with real-time chemical analysers that were set in the Paris urban area (ca. 2 km south of the URB site) as part of the wintertime intensive field experiment of the European program MEGAPOLI (15 January–15 February 2010). Such comparison may provide insights on the importance of positive and negative artefacts associated with our undenuded filter sampling of semi-volatile species (typically ammonium nitrate and organic aerosols).

Concerning ion concentrations, filter measurements were compared for a period of 40 days (6 January–15 February 2010) with a Particle-Into-Liquid-Sampler (PILS; Orsini et al., 2003) coupled with two IC results. The PILS-IC instrument measures selected anions and cations (Cl⁻, NO₃⁻, SO₄²⁻, Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺) every 10 min in PM_{2.5}. Settings used here for the PILS-IC measurements are similar to those reported in Sciare et al. (2011) and will be presented elsewhere (Crippa et al., 2012). The liquid-based aerosol collection principle used in the PILS-IC avoids the positive and negative filter sampling artefacts usually associated with the collection of semi-volatile ammonium nitrate in PM_{2.5}. Uncertainties of the PILS-IC measurements are typically within 20 % (Weber et al., 2003; Orsini et al., 2003; Hogrefe et al., 2004; Takegawa et al., 2005). They are compared with the filter sampling results in Fig. 4a for the three major ions (NH₄⁺; NO₃⁻; SO₄²⁻). Very satisfactory results are obtained (r^2 ranging from 0.88 to 0.94) with slopes close to 1 (ranging from 0.91 to 1.15) and y-intercepts close to zero (ranging from -0.19 to +0.56 $\mu\text{g m}^{-3}$), i.e. in the range of uncertainties given by the two techniques. One of the most important points is the absence of significant discrepancies concerning nitrate, which is a semi-volatile species exhibiting high concentrations during winter in Paris (Favez et al., 2007). Note that the comparison results obtained

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for the other ions (Cl^- , Na^+ , Mg^{2+}) are also very satisfactory (r^2 ranging from 0.85 to 0.90) with slopes close to 1 (ranging from 1.11 to 1.19).

Filter sampling EC and OC concentrations were compared for a period of 70 days (6 January–15 March 2010) with semi-continuous hourly measurements of VOC denuded EC and OC concentrations in $\text{PM}_{2.5}$, obtained using an OCEC Sunset field instrument (Sunset Laboratory, Forest Grove, OR, USA; Bae et al., 2004). The default thermal program (National Institute for Occupational Safety and Health, NIOSH; Birch and Cary, 1996) was used in this instrument. Measurement uncertainty given by the OCEC Sunset field instrument is poorly described in the literature and an estimate of 20% was considered following Peltier et al. (2007). Comparisons between filter sampling and semi-continuous EC and OC measurements are performed in Fig. 4b and show a relatively good agreement (r^2 of 0.69 and 0.84) with slopes of 0.75 and 1.20, and y-intercepts of $+0.32$ and $+0.17 \mu\text{g m}^{-3}$ for EC and OC, respectively. Slope differences may partly originate from the different thermal programs used, having a nearly 200°C difference for the last temperature plateau under Helium (Cavalli et al., 2010). In order to test this assumption, a comparison was performed for Total Carbon (TC) measurements and showed a very good correlation ($r^2 = 0.87$) with a slope close to one (1.09) and a y-intercept close to zero ($+0.4 \mu\text{g m}^{-3}$).

3.5 OC to OM conversion factor

Results concerning the OC to OM conversion factor are reported in Table 2. Concerning the first method (linear regression), very good correlations between RM and OC concentrations were observed for the whole duration of the project, at every site (r^2 higher than 0.90, Fig. S2). This suggests that the unidentified chemical fraction RM is directly related to OC. In other terms, this confirms that all the major chemical components are taken into account in Eq. (1), and that the defined “Remaining Mass” fraction can be regarded as organic matter like. It should be noted that no significant amount of water is adsorbed onto filters because weighing is performed under dry conditions

Chemical characterisation of $\text{PM}_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(RH below 30 %, Sect. 2.3.1). Conversion factors are slightly lower for the urban and suburban stations than for the rural ones (1.95, 1.98, 2.03, 2.08 and 2.12 for SUB, URB, NER, NWR and SOR, respectively). Organic matter is generally more oxidised in rural areas, leading to higher conversion factors (Turpin and Lim, 2001). The small discrepancy observed between the different types of environments in the region of Paris may be explained by the homogeneity of organic carbon concentrations, suggesting the importance of imported sources (see Sect. 4.2). Seasonal variations of f_{OC-OM} are illustrated in Fig. S3. This factor is relatively stable all along the year at the five sites, ranging between 1.8 and 2.2; different patterns are observed from one season to another according to sites, thus suggesting the absence of clear seasonal variations.

Very similar conclusions can be drawn concerning the second method (day-by-day calculation), which leads to slightly lower ($p < 0.005$ using the Welch's t-test; Welch, 1947) conversion factors at SUB and URB than at the rural sites (1.96, 1.92, 2.08, 2.05 and 2.09 for SUB, URB, NER, NWR and SOR, respectively). High relative standard deviations are observed at every site, ranging from 15 to 20 %. This suggests that the organic matter chemical composition is strongly daily dependant, which can be related to the daily changing meteorological conditions (as air mass origins and air temperatures; see Sect. 4.1). No clear seasonal pattern (Fig. S3) is observed with this second method as well. Comparable f_{OC-OM} values are found with both methods in every environment, showing relative differences below 3 % for each site. The first method is however preferred as no days are excluded from the dataset and because of its more advanced mathematical approach.

We chose to apply an OC to OM conversion factor equal to 1.95 for the suburban and urban sites, and of 2.05 for the three rural sites, for the whole duration of the project. By choosing these factors, we wanted to give an insight of the general chemical properties of OM in the region, and to allow its comparison between sites. The chosen conversion factors set among the values suggested by Turpin and Lim (2001) of 1.6 ± 0.2 for urban areas and 2.1 ± 0.2 for non-urban areas, and Kiss et al. (2002) of 1.9–2.0 in a rural area. It is higher than the factors used in previous studies in the region of Paris by

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Guinot et al. (2007) of 1.8 and Favez et al. (2009) of 1.7 (this last factor was recalculated from water-soluble organic carbon and water-insoluble organic carbon contents), which can be related to differences in OC/EC separation methods (e.g. differences in thermo-optical protocols).

Gravimetric, chemically reconstructed and on-line determined $PM_{2.5}$ mass concentrations are compared in Fig. 2. Very good correlations are found between the three datasets, with r^2 of 0.98 and 0.94, and slopes of 1.00 and 1.05 for chemically reconstructed against gravimetric, and against TEOM-FDMS determined mass concentrations, respectively. This confirms the consistency of our measurements and the conversion factors chosen to estimate organic matter.

4 Discussion

Section 4.1 will describe the temporal variability of $PM_{2.5}$ mass and major chemical constituents, whereas Sect. 4.2 will focus on their spatial variability.

4.1 Temporal variability of $PM_{2.5}$

4.1.1 Daily temporal variability of fine aerosols

Daily temporal variability of fine aerosol chemical composition at the suburban site is reported in Fig. 5. Similar patterns are observed at the five stations (Fig. S4 and Sect. 4.2) making the conclusions drawn for this site relevant for the four others. Strong variability can be observed from one day to another, for PM mass and chemical composition. On the whole duration of the project PM_{grav} is on average $15.2 \pm 10.5 \mu g m^{-3}$, and ranges from 3.7 to $62.6 \mu g m^{-3}$ (Table 1; unless otherwise stated all the figures mentioned in this Sect. 4.1 refer to the SUB site). Most pollution events occur during late autumn, winter and early spring (from December to April) and are associated with marked increases of secondary inorganic species (e.g. on 26 January 2010: PM_{grav}

Chemical characterisation of $PM_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and SIA concentrations are 58.7 and 32.1 $\mu\text{g m}^{-3}$, respectively). Organic matter also significantly contributes to the enhancement of PM mass during pollution events (e.g. more than 60 % of PM_{grav} on 7 January 2010, with $[\text{PM}_{\text{grav}}] = 62.6 \mu\text{g m}^{-3}$), contrarily to sea salt, dust and EC. The strong daily variability of PM concentration and composition can be explained by the variations of source emission intensities, atmospheric processes (e.g. Healy et al., 2012) and meteorological parameters (e.g. Galindo et al., 2011; Martin et al., 2011; Georgoulias and Kourtidis, 2011). A focus will be made on the latter variable and especially on temperature, boundary layer height (BLH), precipitation and air mass origins. Although those meteorological parameters can be highly correlated, their individual influence will be highlighted during specific polluted and clean conditions.

4.1.2 Influence of meteorological parameters

Temperature modifies the emission of secondary PM precursors such as biogenic VOCs during summer (Fowler et al., 2009, and references therein), the formation of temperature inversion during winter (Stull, 1988) or the condensation of high saturation vapour pressure compounds such as nitric acid (Monks et al., 2009; Hueglin et al., 2005). In the city of Paris, low temperatures (daily average below 0 °C) often lead to high pollution events ($[\text{PM}_{\text{grav}}] > 40 \mu\text{g m}^{-3}$) mainly due to the increased contribution of SIA (Fig. 6). During winter (DJF) SIA concentrations exceed 10 $\mu\text{g m}^{-3}$ during 35 days, contributing $50 \pm 9\%$ of PM_{grav} under temperatures of $-0.7 \pm 2.9^\circ\text{C}$ on average. SIA are thus a major cause of the high $\text{PM}_{2.5}$ concentrations observed in the region of Paris during cold events, hence asking for particular attention for the implementation of efficient abatement strategies. Whereas nitrate concentrations are enhanced during cold periods because of thermodynamic processes (Clegg et al., 1998), high nss-sulfate concentrations are certainly due to mid- or long-range transport episodes that are related to anticyclonic conditions, eastern air masses and low temperatures (see below and Sect. 4.2). (In the following, mid- or long-range transport

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



will refer to transport from outside the region of Paris, the exact origin not being quantitatively assessed at the current state of analysis.) In fact, nss-sulfate is mostly produced from cloud processing over large scales rather than from local gas phase oxidation of SO₂ (Putaud et al., 2004a). Finally, nitrate and nss-sulfate are fully neutralized by ammonium ($[\text{NO}_3^-] + 2[\text{SO}_4^{2-}]$ versus $[\text{NH}_4^+]$ in mol m⁻³: slope = 0.96; $r^2 = 0.96$, $n = 348$), which therefore follows the same pattern. Similar correlations between high atmospheric concentrations and low temperatures are also found for organic matter, which can be explained by strong biomass burning sources – related to domestic heating – in the region of Paris (Favez et al., 2009; Sciare et al., 2011; Healy et al., 2012) and probably to the condensation of semi-volatile organic species as observed in many European areas (Putaud et al., 2004a).

However, days exhibiting high SIA and PM concentrations cannot solely be explained by low temperatures (e.g. 14 April 2010, 10 May 2010, etc.), meteorological parameters such as precipitation and boundary layer height should be regarded as well (Fig. S5). BLH plays an important role in determining the transport, storage and dispersion of atmospheric pollutants (Salmond and McKendry, 2005) and is responsible for high PM daily variability in Paris during all the campaign. As an illustration, during two following days (31 October 2009 and 1 November 2009), PM_{2.5} concentrations were reduced by a factor of 3.5 (37.7 and 10.6 μg m⁻³, respectively) mainly due to the increase by a factor of 2.8 of the BLH (288 and 820 m, respectively). In addition, the wet removal by precipitation is known to be the most efficient atmospheric aerosol sink (Radke et al., 1980) and contribute to low concentration days ($< 10 \mu\text{g m}^{-3}$) in Paris, especially from November to April because of heavy rains (typically higher than 5 mm per day).

Finally, air mass origins can be a strong cause of PM daily variability because of the flat topography of the region, and the contrasted surrounding environments (marine versus continental areas). This is illustrated in Fig. S5 where two-days back trajectories were calculated every four hours for two typical types of air masses in the region, using HYSPLIT (Draxler and Rolph, 2011). Air masses originating from continental Europe (northeast to east of France) lead to high PM loadings and high contents of

SIA, especially during winter, whereas air masses coming from marine sectors (west and north of France) bring much lower aerosol content. In fact, one of the highest PM pollution peaks of the campaign (26 January 2010, $[PM_{\text{grav}}] = 58.7 \mu\text{g m}^{-3}$) is mainly due to mid- or long-range transport from continental Europe (concentrations ranging from 47.2 to 61.4 $\mu\text{g m}^{-3}$ at the other sites) and is predominantly made of SIA (50 % of $[PM_{\text{grav}}]$). (Mid- or long-range transport from continental Europe will be named continental transport or continental import later on.) Contrarily, under marine air masses that are poorly influenced by anthropogenic pollutions, $PM_{2.5}$ concentrations are typically below 10 $\mu\text{g m}^{-3}$ (e.g. 6.5 and 8.6 $\mu\text{g m}^{-3}$ for 13 September 2009 and 25 October 2009, respectively). These conclusions, regarding the significant contribution of eastern mid- or long-range pollution in the region of Paris, are in agreement with what Bessagnet et al. (2005), Sciare et al. (2010, 2011) and Healy et al. (2012) reported in short-term studies in this city in winter and spring. We attempted to quantify this phenomenon by focusing on days showing PM_{grav} higher than the $PM_{2.5}$ 2015 EU annual limit value of 25 $\mu\text{g m}^{-3}$, which represents 50 days on the whole duration of the campaign. Three categories were defined: days with air masses originating from the northeast to the east of France, days with BLH below 400 m (representing the 10th percentile of BLH values) and days that do not fit either of both above mentioned criteria. We found that 66 % of the polluted days can be attributed to continental import (among which 12 % show low BLH), 20 % exhibit low BLH with other air mass origins, and 14 % cannot be explained by the previous factors. Therefore, in Paris, two third of the days exceeding the 2015 EU annual $PM_{2.5}$ limit value are due to continental import from the northeast to the east of France, questioning the efficiency of local, regional and even national abatement strategies during pollution episodes, suggesting instead collaborative works with neighbourhood countries on these topics. It should be added that French emissions can also impact surrounding areas as reported in Bessagnet et al. (2005), being significantly influential on Great Britain, Belgium, the Netherlands and Eastern Europe.

**Chemical
characterisation of
 $PM_{2.5}$ in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1.3 Monthly and seasonal variability of PM_{2.5}

Keeping in mind the high daily variability of PM mass and chemical composition, their monthly and seasonal trends have been studied (Figs. 7 and S6). On the annual scale, Paris (URB and SUB sites) exhibits its highest PM_{2.5} concentrations during late autumn, winter and early spring (higher than $15 \mu\text{g m}^{-3}$ on average, from December to April), intermediates during late spring and early autumn (between 10 and $15 \mu\text{g m}^{-3}$ during May, June, September, October and November) and the lowest during summer (below $10 \mu\text{g m}^{-3}$ during July and August). This pattern is mainly driven by OM and SIA concentrations that are the main components of fine aerosol (Fig. 7). Figure S6 shows that OM monthly mean concentrations significantly increase from autumn to winter (e.g. 2.5 times higher from November to January), slightly decrease from winter to early spring (e.g. -28% from February to April), and remain fairly constant until the end of the campaign ($4.0 \pm 0.7 \mu\text{g m}^{-3}$ from June to September). During autumn and winter this pattern can be explained by stronger emissions of wood burning sources (Favez et al., 2009; Sciare et al., 2011, Sect. 4.2.3), whereas during spring and summer OM concentrations are likely related to biogenic emissions and secondary organic aerosol formations (Jacobson et al., 2000, and references therein). Nitrate concentrations are, on the other hand, on average significantly higher during winter and early spring months (JFMAM) than the rest of the year ($4.6 \pm 1.2 \mu\text{g m}^{-3}$ and $1.1 \pm 0.8 \mu\text{g m}^{-3}$, respectively), partly because of thermodynamic conditions favouring the partitioning of this molecule into the particulate phase (Clegg et al., 1998). Non-sea-salt sulfate exhibits lower concentrations ($< 1.5 \mu\text{g m}^{-3}$) during autumn (OND) and mid-summer (JA), because Paris is less exposed to continental advection during these months (Airparif and LSCE, 2012). Ammonium follows seasonal variations halfway between nss-sulfate and nitrate as it fully neutralizes both compounds. EC and mineral dust do not display any seasonal pattern with stable concentrations all along the year that are 1.3 ± 0.3 and $0.5 \pm 0.1 \mu\text{g m}^{-3}$, respectively, on average and calculated from monthly means. Finally sea salt monthly variations are clearly related to wind

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



directions, and show slightly higher concentrations during autumn (OND) as air masses coming from marine regions were prevalent (Airparif and LSCE, 2012).

4.2 Spatial variability of PM_{2.5}

4.2.1 PM_{2.5} mass concentrations

Spatial variability of fine aerosol in the region of Paris will first be discussed by comparing atmospheric mass concentrations determined gravimetrically at the five sites (Fig. 8). PM concentrations are surprisingly very similar at the regional scale, during most of the one-year project. Very good correlations are found between urban and suburban sites ($r^2 = 0.94$, slope = 0.99, $n = 335$), good correlations are found between rural sites (e.g. for NWR plotted against SOR sites, $r^2 = 0.83$, slope = 1.08, $n = 351$) and, more surprisingly, between urban and rural sites (e.g. for URB plotted against NER sites, $r^2 = 0.84$, slope = 1.06, $n = 330$). Interestingly, when focusing on individual days a very high temporal variability is generally observed at the same time at the five sites; this is for instance clearly illustrated at the end of January where PM_{grav} is reduced by a factor of 10 to 14 according to sites after only two days (from 27 to 29 January 2010), with concentrations shifting from a range of 49.4–62.0 $\mu\text{g m}^{-3}$ to 3.9–5.9 $\mu\text{g m}^{-3}$, respectively. This suggests that, in the region of Paris, background PM_{2.5} levels are mainly controlled by regional instead of local scale phenomena, which is in agreement with the above mentioned influences of mesoscale meteorological parameters and mid- or long-range transport (Sect. 4.1; Sciare et al., 2010).

Discrepancies between the different sites can however be noticed, with decreased concentrations when shifting from urban and suburban to rural sites, which is in agreement with most European environments (Querol et al., 2004; Van Dingenen et al., 2004; Putaud et al., 2004a). In fact, annual PM_{grav} mean concentrations are 15.2, 14.8, 12.6, 11.7 and 10.8 $\mu\text{g m}^{-3}$ for SUB, URB, NER, NWR and SOR sites, respectively (Table 1). Discrepancies between URB and SUB mean concentrations are not statistically significant ($p > 0.25$) and are due to differences between the sampling days discarded in

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



each station (Table S1), whereas discrepancies between NER, NWR and SOR sites cannot be explained by the former argument. The same conclusions can be drawn when comparing datasets reconstructed by excluding at every site each day missing at one site minimum. In a general way, differences observed between PM_{2.5} site's concentrations can mainly be attributed to local source emissions and chemical processes as it will be discussed later (Sects. 4.3.2 and 4.3.3). It is noteworthy that most pollution episodes that are related to variations of BLH show a very high concentration gradient between urban and rural sites (e.g. about 40 % higher at URB and SUB than at rural sites on 23 January 2010, with a BLH of 273 m) which can be related to enhanced effects of local emissions on atmospheric PM concentrations. To summarise, background PM_{2.5} mass concentrations are most of the time homogeneous at the regional scale on the whole duration of the project; however urban and suburban sites show higher PM levels than the rural ones because of emissions of local anthropogenic sources that can be emphasised by meteorological conditions.

4.2.2 Annual average chemical composition of PM_{2.5}

The annual average chemical composition of fine aerosol is depicted in Fig. 9 and shows a very similar pattern at the five sites, confirming the homogeneous feature of fine aerosol at the regional scale. The major chemical component is organic matter, accounting for 38 to 47 % of PM_{grav} according to sites, and showing a slightly higher contribution at the NER station for reasons given later on (Sect. 4.2.3). It is followed by nitrate (17–22 %), nss-sulfate (13–16 %) and ammonium (10–12 %) i.e. secondary inorganic aerosols. The highest SIA contributions are found at two rural sites (NWR and SOR) and are due to lower influences of local anthropogenic sources compared to the other locations. This leads to higher proportions of SIA even though mass concentrations remain approximately the same at the regional scale (5.2 to 6.4 μg m⁻³). EC has a PM_{2.5} mass contribution of 4 to 10 %, and is about 3 times higher (in absolute concentrations) at URB and SUB than at rural sites because of its mainly local traffic source origin (Healy et al., 2012). Finally, dust and sea salts are minor components of fine

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



aerosol in the region of Paris, representing 2 to 5 % and 3 to 4 % of its mass, respectively. The methodology developed to estimate the OC-OM conversion factor allows us to have a very small proportion of unaccounted mass (0 to 2 %). This overall chemical composition is consistent with what is found in other European environments (Putaud et al., 2010, and references therein), exhibiting very high proportions of carbonaceous and secondary inorganic aerosols.

4.2.3 Major chemical compounds of PM_{2.5}

A detailed discussion of the spatial variability of each major chemical compound will now be given from the highest to the lowest contributor to fine aerosol mass concentrations (see Figs. 10 and 11). To begin with, OM shows fairly homogeneous concentrations at the regional scale, on the whole duration of the project with good correlations between for example SUB and URB ($r^2 = 0.82$, slope = 1.08, $n = 335$), NWR and URB (0.72, 0.79, 335) or NWR and SOR sites (0.73, 1.05, 351) and slopes fairly close to 1. As for PM, comparable OM temporal variations are observed between sites, and very high levels can be followed by very low ones after only few days (e.g. on 27 and 29 January 2010, OM concentrations were 21.7–24.8 $\mu\text{g m}^{-3}$ and 1.0–3.2 $\mu\text{g m}^{-3}$ according to sites, respectively). This implies that a significant part of OM is either imported from outer regions of Paris, or spatially uniformly distributed over the region because of similar primary emission sources. The latter argument is very unlikely because primary emission sources of OM, such as wood burning (WB) (Puxbaum et al., 2007, and references therein) and traffic (Thorpe and Harrison, 2008; Pio et al., 2011) are thought to be rather local and site-dependent. In fact concerning WB, good correlations between OM and levoglucosan (not shown here) at the NER site during cold days assert the presence of a local source, explaining higher OM concentrations at this site, and less satisfactory correlations with the other rural sites (e.g. for NER plotted against SOR sites, $r^2 = 0.55$, slope = 1.46, $n = 330$). Moreover, during low BLH days, clearly higher OM concentrations are found at URB, SUB and NER sites compared to NWR and SOR ones (e.g. 31 October 2009) that can be explained by the enhancement of local

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



WB sources. It is however likely that a part of WB aerosols is also due to mid-range transport as suggested by the comparison of levoglucosan levels between sites and as reported for biomass burning (including WB) in other regions of the world (Niemi et al., 2005, 2009; Stohl et al., 2007; Mochida et al., 2010). To conclude, in the region of Paris, OM is therefore mainly imported by mid- or long-range transport, although local sources such as traffic and wood burning also contribute to its atmospheric concentrations.

SIA including nitrate, nss-sulfate and ammonium also shows very good site-to-site covariations during this study (Figs. 10 and 11). As already mentioned in Sect. 4.1.2, SIA concentrations highly depend on air mass origins and temperatures, and are mainly due to transboundary mid- or long-range transport from countries located east of France (Bessagnet et al., 2005; Sciare et al., 2010, 2011). However, local emissions of precursor gases such as NO_x , NH_3 and SO_2 can partly explain the discrepancies observed between sites for nitrate and ammonium concentrations, and to a lower extent for nss-sulfate (Figs. 9, 10 and 11). In fact, when focusing again on the end of October (that was under low BLH conditions), nitrate shows a clear gradient exhibiting its highest to lowest concentrations at URB, SUB, NER, NWR and SOR sites (11.6, 8.5, 8.0, 6.9, and $5.2 \mu\text{g m}^{-3}$, respectively, the 31 October 2009). On the whole duration of the campaign, nitrate concentrations are 4, 11, 13 and 20 % higher at URB than at SUB, NWR, SOR and NER sites, respectively (values determined from the slope of the correlations between sites). This suggests a local production of ammonium nitrate that could be related to higher local traffic emissions of NO_x at the URB and SUB sites (Alary et al., 1995; Frangi et al., 1996; Airparif, 2012), knowing that ammonia is in excess in the region of Paris (www.emep.int). In addition, local chemical processes have been observed in this region by Healy et al. (2012) and could explain the differences observed between sites for specific days. For instance on 18 January 2010, a local thick fog was observed at the URB site and led to higher concentrations of SIA compared to the other sites (3 to 7 times higher according to sites) due to heterogeneous processing (Reid et al., 1998; Schaap et al., 2004). To summarise, SIA are mainly imported from

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



countries located east of France, even if local sources and chemical processes could lead to discrepancies between sites on very specific days.

EC shows a completely different pattern, characteristic of a local source. A detailed discussion of EC sources and mixing state in Paris can be found in Healy et al. (2012).

In this study, 88 % and 12 % of EC particle mass was apportioned to fossil fuel and biomass burning, respectively, and 79 % and 21 % to local emissions and continental transport, respectively, during a one-month winter period. In our study, no or poor correlations are found between rural sites (e.g. $r^2 = 0.01$, slope = 0.76, $n = 351$, for SOR plotted against NWR sites), URB and SUB sites ($r^2 = 0.45$, slope = 0.84, $n = 335$) or urban and rural sites (e.g. $r^2 = 0.03$, slope = 0.27, $n = 335$, for SOR plotted against URB sites) confirming the importance of local emissions. In addition, as clearly illustrated in Fig. 11, EC concentrations are significantly higher at URB and SUB sites (1.4 ± 0.7 and $1.3 \pm 0.7 \mu\text{g m}^{-3}$ on average on the whole year, respectively) than at the rural ones (0.6 ± 0.3 , 0.5 ± 0.3 , $0.4 \pm 0.3 \mu\text{g m}^{-3}$ on average at the NER, NWR and SOR sites, respectively) which is in agreement with its traffic origin. This can also be illustrated on specific days such as on 28 October 2009 which exhibits a very low BLH (182 m), the highest EC concentrations of the year at URB and SUB sites (5.3 and $4.5 \mu\text{g m}^{-3}$, respectively) and comparatively low ones at the rural sites (1.6 , 1.9 and $0.9 \mu\text{g m}^{-3}$ at NER, NWR and SOR sites, respectively).

The interpretation of the mineral dust dataset is complex because of the substantial uncertainties associated with the methodology chosen for its estimation, and because of the low contribution of this source in fine aerosol (Maenhaut and Cafmeyer, 1998; Gehrig et al., 2001; Putaud et al., 2004a, 2010). For these reasons, its spatial variability will not be discussed. Note however that the very high levels of dust estimated at the URB site during specific days (e.g. $7.2 \mu\text{g m}^{-3}$ on 17 May 2010) are due to renovation of building façades located nearby.

Sea salt concentrations are, as expected, similar at the five sites most of the time (Figs. 10 and 11). The highest concentration is reached on 23 November 2009 (e.g. $3.6 \mu\text{g m}^{-3}$ at the URB site) and is associated with air masses coming from the Atlantic

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ocean. At the exception of about 10 days exceeding $1.5 \mu\text{g m}^{-3}$ at the five sites, sea salt concentrations remained very low ($< 1 \mu\text{g m}^{-3}$) during the entire project.

5 Conclusion and perspectives

This study allowed the construction of a one-year (11 September 2009–10 September 2010) database describing the daily chemical composition of fine aerosol ($\text{PM}_{2.5}$) in the region of Paris. It comprises EC, OC, Cl^- , NO_3^- , SO_4^{2-} , Na^+ , NH_4^+ , K^+ , Mg^{2+} , and Ca^{2+} aerosol compositions at five sites being characteristic of urban, suburban and rural backgrounds. The chemical mass closure methodology was applied for the determination of the OC to OM conversion factor ($f_{\text{OC-OM}}$). Comparable $f_{\text{OC-OM}}$ values were determined for the URB and SUB sites (1.95) and for the rural sites (2.05), which is consistent with the strong homogeneity of OC concentrations observed in the region and its aged (i.e. oxidised) nature. Although clear daily variations are observed, this factor does not show a significant seasonal dependence despite the expected temporality of organic aerosol sources (biogenic, wood burning, etc.) and atmospheric processes (e.g. photochemical ageing).

The temporal variability of $\text{PM}_{2.5}$ was investigated, showing a very strong daily variability at the five sites for PM mass and chemical compositions. The influence of specific meteorological parameters was highlighted: anticyclonic conditions, low temperatures, low BLH and mid- or long-range transport of air masses originating from the continental European sector lead to the highest pollution events of the year, whereas precipitations or western (oceanic) air masses lead to the lowest PM levels. On the annual scale, $\text{PM}_{2.5}$ concentrations in the city of Paris are the highest during late autumn, winter and early spring (higher than $15 \mu\text{g m}^{-3}$ on average from December to April), intermediates during late spring and early autumn (between 10 and $15 \mu\text{g m}^{-3}$ during May, June, September, October, and November) and the lowest during summer (below $10 \mu\text{g m}^{-3}$ during July and August). A seasonal dependence is observed for specific

Chemical characterisation of $\text{PM}_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



chemical species only (e.g. nitrate and OM) whereas others were rather stable along the year (e.g. EC and mineral dust).

The spatial variability of fine aerosol is very similar at the regional scale during most of the year, suggesting the substantial influence of mesoscale meteorological parameters and mid- or long-range transport. Nevertheless, emissions of local anthropogenic sources lead to higher levels at the URB and SUB sites. Gravimetrically determined $PM_{2.5}$ annual mean concentrations are 15.2, 14.8, 12.6, 11.7 and $10.8 \mu\text{g m}^{-3}$ for SUB, URB, NER, NWR and SOR sites, respectively. The annual average chemical composition is also comparable at every site, showing high proportions of OM (38–47 % of $PM_{2.5}$ mass on average at the five sites), nitrate (17–22 %), non-sea-salt sulfate (13–16 %), ammonium (10–12 %), EC (4–10 %), mineral dust (2–5 %) and sea salt (3–4 %), which is in agreement with fine aerosol chemical composition commonly reported in Europe. Organic aerosols are mainly imported by mid- or long-range transport (e.g. for NWR plotted against URB sites: slope = 0.79, $r^2 = 0.72$, $n = 335$), although local sources such as traffic and wood burning may also contribute to its atmospheric concentrations. Secondary inorganic aerosols are also mainly imported (e.g. for NWR plotted against URB sites: slope = 0.91, $r^2 = 0.89$, $n = 335$) from continental European countries (east to northeast sectors), although local sources and chemical processes (e.g. fog) can lead to discrepancies between sites on very specific days. EC is primarily locally emitted (e.g. for SOR plotted against URB sites: slope = 0.27; $r^2 = 0.03$; $n = 335$) and associated with fossil fuel sources (predominantly traffic).

The very large quality controlled aerosol database presented here within the LSCE-AIRPARIF “Particles” project will allow the identification of the major sources of PM as well as their geographical origins, which are prerequisites to implement effective abatement policies. In conjunction with the use of specific tracers, this aerosol database will also help in better documenting primary and secondary sources as well as atmospheric processing of carbonaceous aerosols. Furthermore, this chemical database should help modellers to better understand the transport and transformation of aerosols in the

Chemical characterisation of $PM_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



region, or toxicologists to better assess the toxicity of individual chemical compounds, to mention only a few.

Supplementary material related to this article is available online at:

<http://www.atmos-chem-phys-discuss.net/12/29391/2012/>

[acpd-12-29391-2012-supplement.pdf](#).

Acknowledgements. The author would like to acknowledge the Agence de l'Environnement et de la Maîtrise de l'Energie (ADEME) and the Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA) for funding his Ph.D. Région Ile-de-France, Ville de Paris, Centre National de la Recherche Scientifique and Université de Versailles Saint-Quentin-en-Yvelines are also thanked for their support. The French institute of meteorology "Meteo-France" is acknowledged for providing meteorological data. M. Reynaud and M. Artufel are gratefully acknowledged for their technical and analytical help. Finally, the Scientific Committee associated with this project is thanked for his fruitful comments.

Disclaimer. This document has been subjected to Airparif review and approved for publication. Approval does not signify that the contents reflect the policies or views of the association.



The publication of this article is financed by CNRS-INSU.

ACPD

12, 29391–29442, 2012

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

Alary, R., Donati, J., and Viellard, H.: Automobile pollution in Paris: influence of traffic and meteorological conditions, *Sci. Total Environ.*, 169, 53–61, 1995.

Airparif: <http://www.airparif.asso.fr>, last access: 7 February 2012.

5 Airparif and LSCE: http://www.airparif.asso.fr/_pdf/publications/rapport-particules-anglais-120829.pdf, last access: 10 October 2012.

Andrews, E., Saxena, P., Musarra, S. I., Hildemann, L. M., Koutrakis, P., McMurry, P. H., Olmez, I., and White, W. H.: Concentration and composition of atmospheric aerosols from the 1995 SEAVS experiment and a review of the closure between chemical and gravimetric measurements, *J. Air Waste Manage.*, 50, 648–664, 2000.

10 Aphekom: <http://www.aphekom.org/web/aphekom.org/home>, last access: 31 August 2011.

Arguez, A. and Vose, R. S.: The Definition of the Standard WMO Climate Normal: The Key to Deriving Alternative Climate Normals, *B. Am. Meteorol. Soc.*, 92, 699–704, doi:10.1175/2010BAMS2955.1, 2011.

15 Bae, M. S., Schauer, J. J., DeMinter, J. T., Turner, J. R., Smith, D., and Cary, R. A.: Validation of a semi-continuous instrument for elemental, carbon and organic carbon using a thermal-optical method, *Atmos. Environ.*, 38, 2885–2893, 2004.

Bernstein, J. A., Alexis, N., Barnes, C., Bernstein, I. L., Bernstein, J. A., Nel, A., Peden, D., Diaz-Sanchez, D., Tarlo, S. M., and Williams, P. B.: Health effects of air pollution, *J. Allergy Clin. Immunol.*, 114, 1116–1123, 2004.

20 Bessagnet, B., Hodzic, A., Blanchard, O., Lattuati, M., Le Bihan, O., Marfaing, H., and Rouil, L.: Origin of particulate matter pollution episodes in wintertime over the Paris Basin, *Atmos. Environ.*, 39, 6159–6174, 2005.

Birch, M. E. and Cary, R. A.: Elemental carbon-based method for monitoring occupational exposures to particulate diesel exhaust, *Aerosol Sci. Technol.*, 25, 221–241, 1996.

25 Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3, 79–89, doi:10.5194/amt-3-79-2010, 2010.

Cheung, K., Daher, N., Kam, W., Shafer, M. M., Ning, Z., Schauer, J., and Sioutas, C.: Spatial and temporal variation of chemical composition and mass closure of ambient coarse particulate matter (PM_{10-2.5}) in the Los Angeles area, *Atmos. Environ.*, 45, 2651–2662, 2011.

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chow, J. C., Watson, J. G., Chen, L. W. A., Arnott, W. P., and Moosmuller, H.: The DRI Thermal/Optical Reflectance carbon analysis system: Description, evaluation, and applications in U.S. air quality studies, *Atmos. Environ.*, 27A, 1185–1201, 1993.

Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system H^+ - NH_4^+ - SO_4^{2-} - NO_3^- - H_2O at Tropospheric Temperatures, *J. Phys. Chem. A*, 102, 2137–2154, 1998.

Crippa, M., DeCarlo, P. F., Slowik, J. G., Mohr, C., Heringa, M. F., Chirico, R., Poulain, L., Freutel, F., Sciare, J., Cozic, J., Di Marco, C. F., Elsasser, M., José, N., Marchand, N., Abidi, E., Wiedensohler, A., Drewnick, F., Schneider, J., Borrmann, S., Nemitz, E., Zimmermann, R., Jaffrezo, J.-L., Prévôt, A. S. H., and Baltensperger, U.: Wintertime aerosol chemical composition and source apportionment of the organic fraction in the metropolitan area of Paris, *Atmos. Chem. Phys. Discuss.*, 12, 22535–22586, doi:10.5194/acpd-12-22535-2012, 2012.

Draxler, R. R. and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY Website (<http://ready.arl.noaa.gov/HYSPLIT.php>), NOAA Air Resources Laboratory, Silver Spring, MD, 2011.

Eurostat: <http://epp.eurostat.ec.europa.eu>, last access: 9 January 2011.

Favez, O., Cachier, H., Sciare, J., and Le Moulllec, Y.: Characterization and contribution to $\text{PM}_{2.5}$ of semi-volatile aerosols in Paris (France), *Atmos. Environ.*, 41, 7969–7976, 2007.

Favez, O., Cachier, H., Sciare, J., Sarda-Estève, R., and Martinon, L.: Evidence for a significant contribution of wood burning aerosols to $\text{PM}_{2.5}$ during the winter season in Paris, France, *Atmos. Environ.*, 43, 3640–3644, 2009.

Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., van Dorland, R., Bodeker, G., Boucher, O., Collins, W. D., Conway, T. J., Dlugokencky, E., Elkins, J. W., Etheridge, D., Foukal, P., Fraser, P., Geller, M., Joos, F., Keeling, C. D., Kinne, S., Lassey, K., Lohmann, U., Manning, A. C., Montzka, S., Oram, D., O’Shaughnessy, K., Piper, S., Plattner, G.-K., Ponater Michael, Ramankutty, N., Reid, G., Rind, D., Rosenlof, K., Sausen, R., Schwarzkopf, D., Solanki, S. K., Stenchikov, G., Stuber, N., Takemura, T., Textor, C., Wang, R., Weiss, R., and Whorf, T.: Changes in atmospheric constituents and in radiative forcing, in: *Climate Change 2007: the Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, United Kingdom and New York, NY, USA, 2007.

ACPD

12, 29391–29442, 2012

Chemical characterisation of $\text{PM}_{2.5}$ in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fowler, D., Pilegaard, K., Sutton, M. A., Ambus, P., Raivonen, M., Duyzer, J., Simpson, D., Fagerli, H., Schjoerring, J. K., Neftel, A., Burkhardt, J., Daemmgen, U., Neiryink, J., Perronne, E., Wichink-Kruit, R., Butterbach-Bahl, K., Flechard, C., Tuovinen, J.P., Coyle, M., Gerosa, G., Loubet, B., Altimir, N., Gruenhage, L., Ammann, C., Cieslik, S., Paoletti, E., Mikkelsen, T. N., Ro-Poulsen, H., Cellier, P., Cape, J. N., Horvath, L., Loreto, F., Niinemets, U., Palmer, P. I., Rinne, J., Misztal, P., Nemitz, E., Nilsson, D., Pryor, S., Gallagher, M. W., Vesala, T., Skiba, U., Brüeggemann, N., Zechmeister-Boltenstern, S., Williams, J., O'Dowd, C., Facchini, M. C., de Leeuw, G., Flossman, A., Chaumerliac, N., and Erisman, J. W.: Atmospheric composition change: ecosystems and atmosphere exchange, *Atmos. Environ.*, 43, 5192–5263, 2009.

Frangi, J. P., Jacquemoud S., Puybonnieux-Texier, V., and Lazard, H.: Suivi spatio-temporel de la concentration en NO_x en Ile de France, *C. R. Acad. Sci. II A*, 323, 373–379, 1996.

Galindo, N., Varea, M., Gil-Molto, J., Yubero, E., and Nicolas, J.: The Influence of Meteorology on Particulate Matter Concentrations at an Urban Mediterranean Location, *Water Air Soil Pollut.*, 215, 365–372, 2011.

Gehrig, R., Hüglin, C., and Hofer, P.: Contributions of Road Traffic to Ambient PM₁₀ and PM_{2.5} concentrations, 1st Swiss Transport Research Conference, Monet Verità/Ascona, 1–3, 2001.

Georgoulas, A. K. and Kourtidis, K. A.: On the aerosol weekly cycle spatiotemporal variability over Europe, *Atmos. Chem. Phys.*, 11, 4611–4632, doi:10.5194/acp-11-4611-2011, 2011.

Gherzi, V., Rosso, A., Moukhtar, S., Lameloise, P., Sciare, J., Bressi, M., Nicolas, J., Féron, A., Bonnaire, N.: A comprehensive source apportionment study of fine aerosol (PM_{2.5}) in the region of Paris, France, *Pollution Atmosphérique, Numéro Spécial*, 63–72, 2010.

Gherzi, V., Rosso, A., Moukhtar, S., Léger, K., Sciare, J., Bressi, M., Nicolas, J., Féron, A., and Bonnaire, N.: Sources of fine aerosol (PM_{2.5}) in the region of Paris, *Pollution Atmosphérique*, submitted, 2012.

Gros, V., Sciare, J., and Yu, T.: Air-quality measurements in megacities: Focus on gaseous organic and particulate pollutants and comparison between two contrasted cities, Paris and Beijing, *C. R. Geosci.*, 339, 764–774, 2007.

Guieu, C., Loye-Pilot, M. D., Ridame, C., and Thomas, C.: Chemical characterization of the Saharan dust end-member: Some biogeochemical implications for the western Mediterranean Sea, *J. Geophys. Res.-Atmos.*, 107, 4258, doi:10.1029/2001JD000582, 2002.

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Guinot, B., Cachier, H., and Oikonomou, K.: Geochemical perspectives from a new aerosol chemical mass closure, *Atmos. Chem. Phys.*, 7, 1657–1670, doi:10.5194/acp-7-1657-2007, 2007.

5 Healy, R. M., Sciare, J., Poulain, L., Kamili, K., Merkel, M., Müller, T., Wiedensohler, A., Eckhardt, S., Stohl, A., Sarda-Estève, R., McGillicuddy, E., O'Connor, I. P., Sodeau, J. R., and Wenger, J. C.: Sources and mixing state of size-resolved elemental carbon particles in a European megacity: Paris, *Atmos. Chem. Phys.*, 12, 1681–1700, doi:10.5194/acp-12-1681-2012, 2012.

10 Hodzic, A., Vautard, R., Chazette, P., Menut, L., and Bessagnet, B.: Aerosol chemical and optical properties over the Paris area within ESQUIF project, *Atmos. Chem. Phys.*, 6, 3257–3280, doi:10.5194/acp-6-3257-2006, 2006.

Hogrefe, O., Schwab, J. J., Drewnick, F., Lala, G. G., Peters, S., Demerjian, K. L., Rhoads, K., Felton, H. D., Rattigan, O. V., Husain, L., and Dutkiewicz, V. A.: Semicontinuous PM_{2.5} sulfate and nitrate measurements at an urban and a rural location in New York: PMTACS-NY summer 2001 and 2002 campaigns, *J. Air Waste Manage.*, 54, 1040–1060, 2004.

15 Hueglin, C., Gherig, R., Baltensperger, U., Gysel, M., Monn, C., and Vonmont, H.: Chemical characterisation of PM_{2.5}, PM₁₀ and coarse particles at urban, near-city and rural sites in Switzerland, *Atmos. Environ.*, 39, 637–651, 2005.

20 Isaksen, I. S. A., Benestad, R., Berntsen, T. K., Bosquet, P., Collins, W., Cox, T., Dalsoren, S. B., Eyring, V., Fowler, D., Fuzzi, S., Gauss, M., Granier, C., Joeckel, P., Klimont, Z., Laj, P., Lohmann, U., Maione, M., Monks, P. S., Myhre, G., Prevot, A. S. H., Raes, F., Richter, A., Rognerud, B., Schulz, M., Shindell, D., Stevenson, D., Storelvmo, T., Wang, W.-C., van Weele, M., Wild, M., and Wuebbles, D.: Atmospheric Composition Change: Climate-Chemistry Interaction, *Atmos. Environ.*, 43, 5138–5191, 2009.

25 Jacobson, M. C., Hansson, H. C., Noone, K. J., and Charlson, R. J.: Organic atmospheric aerosols: review and state of the science, *Rev. Geophys.*, 38, 267–294, 2000.

Kiss, G., Varga, B., Galambos, I., and Ganszky, I.: Characterization of water-soluble organic matter isolated from atmospheric fine aerosol, *J. Geophys. Res.-Atmos.*, 107, 8339, doi:10.1029/2001JD000603, 2002.

30 Larssen, S., Sluyster, R., and Helmis, C.: Criteria for EUROAIRNET, the EEA Air Quality Monitoring and Information Network, <http://www.eea.europa.eu/publications/TEC12>, last access: 23 September 2011, 1999.

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lawrence, M. G., Butler, T. M., Steinkamp, J., Gurjar, B. R., and Lelieveld, J.: Regional pollution potentials of megacities and other major population centers, *Atmos. Chem. Phys.*, 7, 3969–3987, doi:10.5194/acp-7-3969-2007, 2007.
- MacMurry, P. H.: A review of atmospheric aerosol measurements, *Atmos. Environ.*, 34, 1959–1999, 2000.
- Maenhaut, W. and Cafmeyer, J.: Long-term atmospheric aerosol study at urban and rural sites in Belgium using multi-elemental analysis by particle-induced X-ray emission spectrometry and short-irradiation instrumental neutron activation analysis, *X-ray Spectrom.*, 27, 236–246, 1998.
- Malm, W. C., Sisler, J. F., Huffman, D., Eldred, R. A., and Cahill, T. A.: Spatial and seasonal trends in particle concentration and optical extinction in the United States, *J. Geophys. Res.*, 99, 1347–1370, 1994.
- Martin, C. L., Allan, J. D., Crosier, J., Choularton, T. W., Coe, H., and Gallagher, M. W.: Seasonal variation of fine particulate composition in the centre of a UK city, *Atmos. Environ.*, 45, 4379–4389, 2011.
- McDow, S. R. and Huntzicker, J. J.: Vapor adsorption artifact in the sampling of organic aerosol: face velocity effects, *Atmos. Environ.*, 24A, 2563–2571, 1990.
- Mochida, M., Kawamura, K., Fu, P. Q., and Takemura, T.: Seasonal variation of levoglucosan in aerosols over the western North Pacific and its assessment as a biomass-burning tracer, *Atmos. Environ.*, 44, 3511–3518, 2010.
- Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M., Baklanov, A., Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K., Cooper, O. R., Dentener, F., Fowler, D., Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V., Guenther, A., Hansson, H. C., Henne, S., Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen, I. S., Jenkin, M. E., Kaiser, J., Kanakidou, M., Klimont, Z., Kulmala, M., Laj, P., Lawrence, M. G., Lee, J. D., Liousse, C., Maione, M., McFiggans, G., Metzger, A., Mieville, A., Moussiopoulos, N., Orlando, J. J., O'Dowd, C. D., Palmer, P. I., Parrish, D. D., Petzold, A., Platt, U., Pöschl, U., Prevot, A. S., Reeves, C. E., Reimann, S., Rudich, Y., Sellegri, K., Steinbrecher, R., Simpson, D., ten Brink, H., Theloke, J., van der Werf, G. R., Vautard, R., Vestreng, V., Vlachokostas, C., and von Glasow, R.: Atmospheric composition change – Global and regional air quality, *Atmos. Environ.*, 43, 5268–5350, 2009.
- Niemi, J. V., Tervahattu, H., Vehkamäki, H., Martikainen, J., Laakso, L., Kulmala, M., Aarnio, P., Koskentalo, T., Sillanpää, M., and Makkonen, U.: Characterization of aerosol particle

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



episodes in Finland caused by wildfires in Eastern Europe, *Atmos. Chem. Phys.*, 5, 2299–2310, doi:10.5194/acp-5-2299-2005, 2005.

Niemi, J. V., Saarikoski, S., Aurela, M., Tervahattu, H., Hillamo, R., Westphal, D. L., Aarnio, P., Koskentalo, T., Makkonen, U., Vehkamäki, H., and Kulmala, M.: Long-range transport episodes of fine particles in southern Finland during 1999–2007, *Atmos. Environ.*, 43, 1255–1264, 2009.

Orsini, D., Ma, Y., Sullivan, A., Sierau, B., Baumann, K., and Weber, R.: Refinements to the Particle-Into-Liquid Sampler (PILS) for Ground and Airborne Measurements of Water Soluble Aerosol Composition, *Atmos. Environ.*, 37, 1243–1259, 2003.

Patashnik, H. and Rupprecht, E. G.: Continuous PM₁₀ measurements using a tapered element oscillating microbalance, *J. Air Waste Manage.*, 41, 1079–1083, 1991.

Peltier, R. E., Weber, R. J., and Sullivan, A. P.: Investigating a liquid-based method for online organic carbon detection in atmospheric particles, *Aerosol Sci. Tech.*, 41, 1117–1127, 2007.

Pettijohn, F. J.: *Sedimentary Rocks*, Earth Data, NY, Harper and Row, 1975.

Pio, C., Cerqueira, M., Harrison, R. M., Nunes, T., Mirante, F., Alves, C., Oliveira, C., Sanchez de la Campa, A., Artinano, B., and Matos, M.: OC/EC ratio observations in Europe: Rethinking the approach for apportionment between primary and secondary organic carbon, *Atmos. Environ.*, 45, 6121–6132, 2011.

Pope, C. A., Burnett, R. T., Thurston, G. D., Thun, M. J., Calle, E. E., Krewski, D., and Godleski, J. J.: Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease, *Circulation*, 109, 71–77, 2004.

Putaud, J.-P., Raes, F., Van Dingenen, R., Baltensperger, J. P. U., Brüggemann, E., Facchini, M. C., Decesari, S., Fuzzi, S., Gehrig, R., Hansson, H. C., Hüglin, C., Laj, P., Lorbeer, G., Maenhaut, W., Mihalopoulos, N., Müller, K., Querol, X., Rodriguez, S., Schneider, J., Spindler, G., ten Brink, H., Tørseth, K., Wehner, B., and Wiedensohler, A.: A European Aerosol Phenomenology. 2: chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2579–2595, 2004a.

Putaud, J.-P., Van Dingenen, R., Dell'Acqua, A., Raes, F., Matta, E., Decesari, S., Facchini, M. C., and Fuzzi, S.: Size-segregated aerosol mass closure and chemical composition in Monte Cimone (I) during MINATROC, *Atmos. Chem. Phys.*, 4, 889–902, doi:10.5194/acp-4-889-2004, 2004b.

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Putaud, J.-P., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Herrmann, H., Hitztenberger, R., Huglin, C., Jones, A. M., Kasper-Giebl, A., Kiss, G., Kousa, A., Kuhlbusch, T. A. J., Loschau, G., Maenhaut, W., Molnar, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A., and Raes, F.: A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, *Atmos. Environ.*, 44, 1308–1320, 2010.
- Puxbaum, H., Caseiro, A., Sanchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencser, A., Preunkert, S., Legrand, M., and Pio, C. A.: Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background, *J. Geophys. Res.*, 112, D23S05, doi:10.1029/2006JD008114, 2007.
- Quinn, P. K. and Coffman, D. J.: Local closure during the First Aerosol Characterization Experiment (ACE 1): Aerosol mass concentration and scattering and backscattering coefficients, *J. Geophys. Res.*, 103, 16575–16596, 1998.
- Radke, L. F., Hobbs, P. V., and Eltgroth, M. W.: Scavenging of aerosol particles by precipitation, *J. Atmos. Sci.*, 19, 715–722, 1980.
- Ramanathan, V., Callis, L., Cess, R., Hansen, J., Isaksen, I. S. A., Kuhn, W., Lacic, A., Luther, F., Mahlman, J., Reck, R., and Schlesinger, M.: Climate-chemical interactions and effects of changing atmospheric trace gases, *Review of Geophysics and Space Physics*, 25, 1441–1482, 1987.
- Ramgolam, K., Chevaillier, S., Marano, F., Baeza-Squiban, A., and Martinon, L.: Proinflammatory effect of fine and ultrafine particulate matter using size-resolved urban aerosols from Paris, *Chemosphere*, 72, 1340–1346, 2008.
- Reid, J. S., Hobbs, P. V., Ferek, R. J., Blake, D. R., Martins, J. V., Dunlap, M. R., and Liousse, C.: Physical, chemical and optical properties of regional hazes dominated by smoke in Brazil, *J. Geophys. Res.*, 103, 32059–32080, 1998.
- Rengarajan, R., Sudheer, A. K., and Sarin, M. M.: Wintertime PM_{2.5} and PM₁₀ carbonaceous and inorganic constituents from urban site in western India, *Atmos. Res.*, 102, 420–431, 2011.
- Salmond, J. A. and McKendry, I. G.: A review of turbulence in the very stable nocturnal boundary layer and its implications for air quality, *Prog. Phys. Geog.*, 29, 171–188, 2005.

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schaap, M., van Loon, M., ten Brink, H. M., Dentener, F. J., and Bultjes, P. J. H.: Secondary inorganic aerosol simulations for Europe with special attention to nitrate, *Atmos. Chem. Phys.*, 4, 857–874, doi:10.5194/acp-4-857-2004, 2004.

Schlesinger, R. B.: The health impact of common inorganic components of fine particulate matter (PM_{2.5}) in ambient air – a critical review, *Inhal. Toxicol.*, 19, 811–832, 2007.

Schmid, H., Laskus, L., Abraham, H. J., Baltensperger, U., Lavanchy, V., Bizjak, M., Burba, P., Cachier, H., Crow, D., Chow, J., Gnauk, T., Even, A., ten Brink, H. M., Giesen, K.-P., Hitznerberger, R., Hueglin, C., Maenhaut, W., Pio, C., Carvalho, A., Putaud, J.-P., Toom-Saunty, D., and Puxbaum, H.: Results of the “carbon conference” international aerosol carbon round robin test stage 1, *Atmos. Environ.*, 35, 2111–2121, 2001.

Sciare, J., Cachier, H., Oikonomou, K., Ausset, P., Sarda-Estève, R., and Mihalopoulos, N.: Characterization of carbonaceous aerosols during the MINOS campaign in Crete, July–August 2001: a multi-analytical approach, *Atmos. Chem. Phys.*, 3, 1743–1757, doi:10.5194/acp-3-1743-2003, 2003.

Sciare, J., Oikonomou, K., Cachier, H., Mihalopoulos, N., Andreae, M. O., Maenhaut, W., and Sarda-Estève, R.: Aerosol mass closure and reconstruction of the light scattering coefficient over the Eastern Mediterranean Sea during the MINOS campaign, *Atmos. Chem. Phys.*, 5, 2253–2265, doi:10.5194/acp-5-2253-2005, 2005.

Sciare, J., Sarda-Estève, R., Favez, O., Cachier, H., Aymoz, G., and Laj, P.: Nighttime residential wood burning evidenced from an indirect method for calculating real-time concentration of particulate organic matter (POM), *Atmos. Environ.*, 42, 2158–2172, 2008.

Sciare, J., d’Argouges, O., Zhang, Q. J., Sarda-Estève, R., Gaimoz, C., Gros, V., Beekmann, M., and Sanchez, O.: Comparison between simulated and observed chemical composition of fine aerosols in Paris (France) during springtime: contribution of regional versus continental emissions, *Atmos. Chem. Phys.*, 10, 11987–12004, doi:10.5194/acp-10-11987-2010, 2010.

Sciare, J., d’Argouges, O., Sarda-Estève, R., Gaimoz, C., Dolgorouky, C., Bonnaire, N., Favez, O., Bonsang, B., and Gros, V.: Large contribution of water-insoluble secondary organic aerosols in the region of Paris (France) during wintertime, *J. Geophys. Res.*, 116, D22203, doi:10.1029/2011JD015756, 2011.

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: From air pollution to climate change*, John Wiley, Hoboken, N. J., 1360 pp., 1998.

Speer, R. E., Edney, E. O., and Kleindienst, T. E.: Impact of organic compounds on the concentrations of liquid water in ambient PM₁₀, *J. Aerosol Sci.*, 34, 63–77, 2003.

- Stohl, A., Berg, T., Burkhardt, J. F., Fjærevaa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth, K., Treffeisen, R., Virkkunen, K., and Yttri, K. E.: Arctic smoke – record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmos. Chem. Phys.*, 7, 511–534, doi:10.5194/acp-7-511-2007, 2007.
- Stull, R. B.: An introduction to boundary layer meteorology, 1st Edn., Dordrecht, Kluwer Academic Publishers, 666 pp., 1988.
- Takegawa, N., Miyazaki, Y., Kondo, Y., Komazaki, Y., Miyakawa, T., Jimenez, J. L., Jayne, J. T., Worsnop, D. R., Allan, J. D., and Weber, R. J.: Characterization of an aerodyne aerosol mass spectrometer (AMS): intercomparison with other aerosol instruments, *Aerosol Sci. Technol.*, 39, 760–770, 2005.
- Terzi, E., Argyropoulos, G., Bougatioti, A., Mihalopoulos, N., Nikolaou, K., and Samara, C.: Chemical composition and mass closure of ambient PM₁₀ at urban sites, *Atmos. Environ.*, 44, 2231–2239, 2010.
- Thorpe, A. and Harrison, R. M.: Sources and properties of non-exhaust particulate matter from road traffic: A review, *Sci. Total Environ.*, 400, 270–282, 2008.
- Turpin, B. J. and Lim, H. J.: Species contribution to PM_{2.5} mass concentrations: revisiting common assumptions for estimating organic mass, *Aerosol Sci. Technol.*, 35, 602–610, 2001.
- Turpin, B. J., Hering, S. V., and Huntzicker, J. J.: Investigation of organic aerosol sampling artifacts in the Los Angeles Basin, *Atmos. Environ.*, 28, 3061–3071, 1994.
- Turpin, B. J., Saxena, P., and Andrews, E.: Measuring and simulating particulate organics in the atmosphere: problems and prospects, *Atmos. Environ.*, 34, 2983–3013, 2000.
- Van Dingenen, R., Raes, F., Putaud, J.-P., Baltensperger, U., Charron, A., Facchini, M.-C., Decesari, S., Fuzzi, S., Gehrig, R., Hansson, H.-C., Harrison, R. M., Hüglin, C., Jones, A. M., Laj, P., Lorbeer, G., Maenhaut, W., Palmgren, F., Querol, X., Rodriguez, S., Schneider, J., ten Brink, H., Tunved, P., Tørseth, K., Wehner, B., Weingartner, E., Wiedensohler, A., and Wahlin, P.: A European aerosol phenomenology 1: physical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2561–2577, 2004.
- Weber, R., Orsini, D., Duan, Y., Baumann, K., Kiang, C. S., Chameides, W., Lee, Y. N., Brechtel, F., Klotz, P., Jongejan, P., ten Brink, H., Slanina, J., Boring, C. B., Genfa, Z., Dasgupta, P., Hering, S., Stolzenburg, M., Dutcher, D. D., Edgerton, E., Hartsell, B., Solomon, P., and Tanner, R.: Intercomparison of near real time monitors of PM_{2.5} nitrate and sulfate at the U.S.

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Environmental Protection Agency Atlanta Supersite, J. Geophys. Res.-Atmos., 108, 8421, doi:10.1029/2001JD001220, 2003.

Welch, B. L.: The generalization of “student’s” problem when several different population variances are involved, Biometrika, 34, 28–35, 1947.

- 5 World Climate Research Programme (WCP): Aerosols and their climatic effects, report of experts meeting, Williamsburg, Va., 28–30 March 1983, Rep. WCP-55, World Meteorological Organization, Geneva, 1983.

Zhang, X. Q. and McMurry, P. H.: Theoretical analysis of evaporative losses from impactor and filter deposits, Atmos. Environ., 21, 1779–1789, 1987.

ACPD

12, 29391–29442, 2012

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Table 1. Chemical composition ($\mu\text{g m}^{-3}$) of PM_{2.5} at the five sites during the one-year period.

	SUBURBAN					URBAN					NORTH EAST RURAL					NORTH WEST RURAL					SOUTH RURAL				
	<i>n</i> = 348					<i>n</i> = 335					<i>n</i> = 330					<i>n</i> = 359					<i>n</i> = 351				
	min	max	med	av	std	min	max	med	av	std	min	max	med	av	std	min	max	med	av	std	min	max	med	av	std
Mass	3.7	62.6	11.9	15.2	10.5	3.9	62.8	11.6	14.8	9.6	2.8	48.6	9.6	12.6	8.6	2.1	61.4	8.7	11.7	9.5	0.7	49.4	7.9	10.8	8.4
OC	0.5	19.8	2.5	3.2	2.5	0.6	12.7	2.6	3.0	1.7	0.4	13.1	2.2	2.9	2.2	0.2	12.8	1.6	2.2	1.9	0.3	10.6	1.7	2.1	1.5
NO ₃ ⁻	<LOQ	17.5	1.1	2.9	3.6	<LOQ	17.1	1.1	2.9	3.7	<LOQ	16.6	0.9	2.2	2.9	<LOQ	18.3	0.9	2.6	3.5	0.0	18.2	0.8	2.2	3.1
SO ₄ ²⁻	0.2	10.1	1.7	2.1	1.7	0.2	11.2	1.6	2.0	1.6	0.2	9.6	1.5	1.9	1.4	0.2	10.9	1.5	1.9	1.6	0.1	9.6	1.4	1.8	1.6
NH ₄ ⁺	0.1	8.1	0.8	1.5	1.6	0.0	9.0	0.7	1.4	1.6	0.1	7.2	0.7	1.2	1.3	<LOQ	8.0	0.7	1.3	1.5	0.1	7.9	0.7	1.2	1.4
EC	0.3	5.3	1.2	1.3	0.7	0.1	4.5	1.3	1.4	0.7	0.1	1.6	0.5	0.5	0.3	0.1	2.3	0.4	0.5	0.3	0.0	1.8	0.4	0.4	0.3
Cl ⁻	<LOQ	1.65	0.11	0.18	0.21	<LOQ	1.88	0.11	0.19	0.23	<LOQ	1.73	0.08	0.16	0.21	<LOQ	1.42	0.09	0.18	0.22	<LOQ	1.72	0.07	0.14	0.19
Na ⁺	<LOQ	1.08	0.11	0.16	0.15	<LOQ	1.19	0.11	0.18	0.17	<LOQ	1.22	0.09	0.14	0.14	<LOQ	0.94	0.11	0.16	0.15	<LOQ	1.01	0.08	0.13	0.13
K ⁺	<LOQ	0.74	0.08	0.13	0.12	<LOQ	0.73	0.07	0.12	0.12	<LOQ	1.59	0.07	0.12	0.13	<LOQ	0.72	0.06	0.11	0.12	<LOQ	0.61	0.06	0.10	0.10
Ca ²⁺	<LOQ	1.29	0.06	0.08	0.09	<LOQ	1.09	0.09	0.12	0.12	<LOQ	0.34	0.04	0.05	0.04	<LOQ	0.31	0.05	0.06	0.05	<LOQ	0.30	0.03	0.04	0.04
Mg ²⁺	<LOQ	0.18	0.02	0.03	0.02	<LOQ	0.16	0.02	0.03	0.02	<LOQ	0.16	0.02	0.02	0.02	<LOQ	0.13	0.02	0.03	0.02	<LOQ	0.15	0.01	0.02	0.02

LOQ: Limit Of Quantification.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Table 2. Determination of the OC to OM conversion factor ($f_{\text{OC-OM}}$) at the five sites, from 11 September 2009 to 10 September 2010, using two analytical methods.

Methods	SUBURBAN		URBAN		NORTH EAST RURAL		NORTH WEST RURAL		SOUTH RURAL	
	1	2	1	2	1	2	1	2	1	2
n	348	326	335	307	330	297	359	309	351	306
$f_{\text{OC-OM}}$	1.95 ± 0.02	1.96 ± 0.33	1.98 ± 0.02	1.92 ± 0.33	2.03 ± 0.02	2.08 ± 0.34	2.08 ± 0.02	2.05 ± 0.38	2.12 ± 0.02	2.09 ± 0.38
r^2	0.94	–	0.90	–	0.94	–	0.92	–	0.90	–

Method 1: linear regression “ $y = a \cdot x$ ” from Eq. (7). $f_{\text{OC-OM}}$ is the slope of the linear regression. Standard deviation is calculated from the standard error of the slope.

Method 2: day-by-day calculation from Eq. (6). Days with $f_{\text{OC-OM}}$ higher than 3 and lower than 1 were excluded.

$f_{\text{OC-OM}}$ is the arithmetic mean.

n : number of samples.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 1. Spatial distribution of the sampling sites in the region of Paris (Ile-de-France region). Source: Google Earth. Legend: SUB: SUBurban, URB: URBan, NER: North-East Rural, NWR: North-West Rural, SOR: SOuth Rural.

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.




Fig. 2. Comparison between gravimetric (PM_{grav}), chemically reconstructed (PM_{chem}) and TEOM-FDMS ($PM_{TEOM-FDMS}$) daily mass concentrations ($\mu\text{g m}^{-3}$) at the urban site from 11 September 2009 to 10 September 2010. Note: TEOM-FDMS measurements are conducted at 30°C and do not take into account semi-volatile materials.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.




Fig. 3. Comparison between gravimetric (PM_{grav}) and TEOM-FDMS ($PM_{TEOM-FDMS}$) daily mass concentrations ($\mu\text{g}/\text{m}^3$) at the urban site from 11 September 2009 to 10 September 2010 ($n = 318$). Note: TEOM-FDMS measurements are conducted at 30°C and do not take into account semi-volatile materials. The slope is given ± 1 standard error.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.




Fig. 4. Comparison between filter and on-line measurements for a 40-day period for the major ions (a), and for a 70-day period for carbonaceous matter (b).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.




Fig. 5. Daily variation of fine aerosol chemical composition at the suburban site from 11 September 2009 to 10 September 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.




Fig. 6. Comparison between daily nitrate, nss-sulfate and ammonium concentrations ($\mu\text{g m}^{-3}$) and temperatures ($^{\circ}\text{C}$) at the suburban site from 11 September 2009 to 10 September 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.




Fig. 7. Monthly mean concentrations ($\mu\text{g m}^{-3}$) of fine aerosol chemical composition at the sub-urban site from 11 September 2009 to 10 September 2010. Error bars represent the standard deviation ($\pm 1\sigma$) of fine aerosol mass concentrations.

**Chemical
characterisation of
PM_{2.5} in Paris**

M. Bressi et al.




Fig. 8. Comparison between PM_{2.5} daily mass concentrations (gravimetric measurements) at the five sites during the one-year period.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.




Fig. 9. Annual average chemical composition of PM_{2.5} ($\mu\text{g m}^{-3}$; %) at the five sites from 11 September 2009 to 10 September 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.




Fig. 10. Daily mass concentrations ($\mu\text{g m}^{-3}$) of the major chemical compounds of fine aerosol at the five sites from 11 September 2009 to 10 September 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chemical characterisation of PM_{2.5} in Paris

M. Bressi et al.




Fig. 11. Monthly mass concentrations ($\mu\text{g}/\text{m}^3$) of the major chemical compounds of fine aerosol at the five sites from 11 September 2009 to 10 September 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion