

RECONCILE

M. von Hobe et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Reconciliation of essential process parameters for an enhanced predictability of Arctic stratospheric ozone loss and its climate interactions

M. von Hobe¹, S. Bekki², S. Borrmann³, F. Cairo⁴, F. D'Amato⁵, G. Di Donfrancesco³², A. Dörnbrack⁶, A. Ebersoldt⁷, M. Ebert⁸, C. Emde⁹, I. Engel¹⁰, M. Ern¹, W. Frey³, S. Griessbach¹¹, J.-U. Groß¹, T. Gulde¹², G. Günther¹, E. Hösen¹³, L. Hoffmann¹¹, V. Homonnai¹⁴, C. R. Hoyle¹⁰, I. S. A. Isaksen¹⁵, D. R. Jackson¹⁶, I. M. Jánosi¹⁴, K. Kandler⁸, C. Kalicinsky¹³, A. Keil¹⁷, S. M. Khaykin¹⁸, F. Khosrawi¹⁹, R. Kivi²⁰, J. Kuttippurath², J. C. Laube²¹, F. Lefèvre², R. Lehmann²², S. Ludmann²³, B. P. Luo¹⁰, M. Marchand², J. Meyer¹, V. Mitev²⁴, S. Molleker³, R. Müller¹, H. Oelhaf¹², F. Olschewski¹³, Y. Orsolini²⁵, T. Peter¹⁰, K. Pfeilsticker²³, C. Piesch¹², M. C. Pitts²⁶, L. R. Poole²⁷, F. D. Pope^{28,*}, F. Ravegnani⁴, M. Rex²², M. Riese¹, T. Röckmann²⁹, B. Rognerud¹⁵, A. Roiger⁶, C. Rolf¹, M. L. Santee³⁰, M. Scheibe⁶, C. Schiller¹, H. Schlager⁶, M. Siciliani de Cumis⁵, N. Sitnikov¹⁸, O. A. Søvde¹⁵, R. Spang¹, N. Spelten¹, F. Stordal¹⁵, O. Sumińska-Ebersoldt^{1,12}, S. Viciani⁵, C. M. Volk¹³, M. vom Scheidt¹³, A. Ulanovski¹⁸, P. von der Gathen²², K. Walker³¹, T. Wegner¹, R. Weigel³,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**S. Weinbuch⁸, G. Wetzel¹², F. G. Wienhold¹⁰, J. Wintel¹³, I. Wohltmann²²,
W. Woiwode¹¹, I. A. K. Young^{28,**}, V. Yushkov¹⁸, B. Zobrist¹⁰, and F. Strohl¹**

¹Forschungszentrum Jülich, Institute for Energy and Climate Research (IEK-7), Jülich, Germany

²LATMOS-IPSL, UPMC Univ. Paris 06, Université Versailles St.-Quenitn; CNRS/INSU, Paris, France

³Max Planck Institute for Chemistry, Particle Chemistry Department, Mainz, Germany

⁴Institute of Atmospheric Science and Climate, ISAC-CNR, Rome, Italy

⁵CNR-INO (Istituto Nazionale di Ottica), Largo E. Fermi, 6, 50125 Firenze, Italy

⁶Institut für Physik der Atmosphäre, DLR Oberpfaffenhofen, Germany

⁷Institute for Data Processing and Electronics, Karlsruhe Institute of Technology, Karlsruhe, Germany

⁸Technische Universität Darmstadt, Institut für Angewandte Geowissenschaften, Umweltmineralogie, Darmstadt, Germany

⁹Meteorologisches Institut, Ludwig-Maximilians-Universität, München, Germany

¹⁰ETH Zurich, Institute for Atmospheric and Climate Science, Zurich, Switzerland

¹¹Jülich Supercomputing Centre (JSC), Forschungszentrum Jülich GmbH, Germany

¹²Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

¹³Department of Physics, University of Wuppertal, Germany

¹⁴Department of Physics of Complex Systems, Eötvös Loránd University, Pázmány P. s. 1/A, 1117 Budapest, Hungary

¹⁵Department of Geosciences, University of Oslo, Oslo, Norway

¹⁶Met Office, Exeter, UK

¹⁷Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Frankfurt, Germany

¹⁸Central Aerological Observatory, Dolgoprudny, Moscow Region, Russia

¹⁹MISU, Stockholm University, Stockholm, Sweden

²⁰Finnish Meteorological Institute, Arctic Research, Sodankylä, Finland

²¹University of East Anglia, School of Environmental Sciences, Norwich, UK

ACPD

12, 30661–30754, 2012

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- ²² Alfred Wegener Institute for Polar and Marine Research, Potsdam, Germany
²³ Institut für Umweltphysik, University of Heidelberg, Germany
²⁴ CSEM Centre Suisse d'Electronique et de Microtechnique SA, Neuchâtel, Switzerland
²⁵ Norwegian Institute for Air Research, Kjeller, Norway
²⁶ NASA Langley Research Center, Hampton, VA 23681, USA
²⁷ Science Systems and Applications, Inc. Hampton, VA 23666, USA
²⁸ University of Cambridge, Department of Chemistry, Cambridge, UK
²⁹ Institute for Marine and Atmospheric Research Utrecht (IMAU), Utrecht University, Utrecht, The Netherlands
³⁰ JPL/NASA, California Institute of Technology, Pasadena, California, USA
³¹ Department of Physics, University of Toronto, Toronto, Canada
³² Ente Nazionale per le Nuove tecnologie, l'Energia e l'Ambiente, Roma, Italy
* now at: School of Geography, Earth and Environmental Sciences, University of Birmingham, UK
** now at: The British Museum, London, UK

Received: 11 November 2012 – Accepted: 14 November 2012 – Published: 27 November 2012

Correspondence to: M. von Hobe (m.von.hobe@fz-juelich.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Significant reductions in stratospheric ozone occur inside the polar vortices each spring when chlorine radicals produced by heterogeneous reactions on cold particle surfaces in winter destroy ozone mainly in two catalytic cycles, the ClO dimer cycle and the ClO/BrO cycle. Chlorofluorocarbons (CFCs), which are responsible for most of the chlorine currently present in the stratosphere, have been banned by the Montreal Protocol and its amendments, and the ozone layer is predicted to recover to 1980 levels within the next few decades. During the same period, however, climate change is expected to alter the temperature, circulation patterns and chemical composition in the stratosphere, and possible geo-engineering ventures to mitigate climate change may lead to additional changes. To realistically predict the response of the ozone layer to such influences requires the correct representation of all relevant processes. The European project RECONCILE has comprehensively addressed remaining questions in the context of polar ozone depletion, with the objective to quantify the rates of some of the most relevant, yet still uncertain physical and chemical processes. To this end RECONCILE used a broad approach of laboratory experiments, two field missions in the Arctic winter 2009/10 employing the high altitude research aircraft M55-Geophysica and an extensive match ozone sonde campaign, as well as microphysical and chemical transport modelling and data assimilation. Some of the main outcomes of RECONCILE are as follows: (1) vortex meteorology: the 2009/10 Arctic winter was unusually cold at stratospheric levels during the six-week period from mid-December 2009 until the end of January 2010, with reduced transport and mixing across the polar vortex edge; polar vortex stability and how it is influenced by dynamic processes in the troposphere has led to unprecedented, synoptic-scale stratospheric regions with temperatures below the frost point; in these regions stratospheric ice clouds have been observed, extending over $> 10^6 \text{ km}^2$ during more than 3 weeks. (2) Particle microphysics: heterogeneous nucleation of nitric acid trihydrate (NAT) particles in the absence of ice has been unambiguously demonstrated; conversely, the synoptic scale ice clouds also

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



change induced changes in stratospheric conditions and their impact on ozone are still not easy to predict. The record-breaking ozone loss that occurred over the Arctic in spring 2011 (Manney et al., 2011; Kuttippurath et al., 2012a) has shown that stratospheric ozone is still a critical issue, and that climate change could bring about ozone hole like conditions in the Arctic as long as stratospheric chlorine levels are still high enough.

With its focus primarily on the Arctic stratosphere, the EU funded project RECONCILE, short for “Reconciliation of essential process parameters for an enhanced predictability of Arctic stratospheric ozone loss and its climate interactions”, runs from 2009 to 2013 and aims at understanding how the physical and chemical processes affecting stratospheric ozone will be influenced by and feedback on climate change. In Sect. 2, a brief overview of past research on polar ozone depletion is given, and the motivation and objectives of RECONCILE are stated. The overall strategy as well as the tools and activities adopted are described in Sect. 3, and a summary of the main scientific findings of the RECONCILE project is presented in Sect. 4. Section 5 explains how these results are used to advance our ability to predict the future state of the ozone layer using chemistry climate models (CCMs), i.e. global circulation models (GCMs) that include chemical reactions. Section 6 gives a brief overview on the record Arctic ozone depletion 2011 and its implications.

2 Historical background and objectives

Below, the history of research and policies on stratospheric ozone is briefly reviewed and the state of knowledge, including open questions, at the beginning of the RECONCILE project is described. Some key events – from the early warnings of possible dangers to the ozone layer expressed in the 1970s to recent international research projects with a focus on polar stratospheric ozone – are illustrated in Fig. 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 The discovery of the ozone hole and the Montreal Protocol

In 1985 Joe Farman, Brian Gardiner and Jonathan Shanklin from the British Antarctic survey reported “that the spring values of total O₃ in Antarctica have now fallen considerably” (Farman et al., 1985), and others soon confirmed this observation (Chubachi and Kajiwara, 1986; Gernandt, 1987; Stolarski et al., 1986). Even though a number of scientists had expressed concerns that anthropogenic emissions of CFCs (Cicerone et al., 1974; Crutzen, 1974; Molina and Rowland, 1974), nitrous oxide (N₂O) from fertilizer use (Crutzen, 1976), and reactive nitrogen compounds from high-flying aircraft (Johnston, 1971; Crutzen, 1971) would potentially lead to faster catalytic ozone removal rates and hence lower stratospheric ozone concentrations, the “Ozone Hole” came as a surprise to atmospheric scientists. But with the research that had already been carried out in the 1970s as a solid foundation, a theory was soon developed that the dramatic ozone destruction over the Antarctic in spring is the result of a complex interplay of atmospheric dynamics, heterogeneous reactions, and catalytic reactions involving chlorine compounds (Fig. 2).

A link was made between these stratospheric chlorine compounds and CFCs, which had been introduced as refrigerating agents and as propellants in spray cans in the middle of the 20th century. Being inert in the lower atmosphere, these compounds are transported to the stratosphere, where they release chlorine upon UV irradiation. We undoubtedly know today that the attribution of CFCs as the ultimate cause of the ozone hole has been correct: Morgenstern et al. (2008) and Newman et al. (2009) recently demonstrated that this first widely visible global environmental threat caused by human activities had the potential for a global catastrophe.

Fortunately, the 1987 Montreal Protocol and its subsequent amendments and adjustments essentially banned CFCs and certain other chlorinated and brominated compounds from being produced. As a result, stratospheric chlorine and bromine levels have started to decline since the late 1990s and will continue to do so over the next decades, although this will be a slow process due to their long atmospheric lifetimes

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(WMO, 2010). In most regions, column ozone is expected to return to 1980 levels (1980 is usually taken as the reference year for the so-called ozone recovery) around the middle of the century (Eyring et al., 2010; WMO, 2010). However, the state-of-the-art CCM simulations used to estimate the recovery dates, clearly show that the decline in ozone depleting substances (ODS) is only part of the story. The alterations to atmospheric dynamics and composition induced by global climate change (see Figs. 3–22 in WMO, 2010) become increasingly important and have a significant impact on stratospheric ozone and consequently on recovery date projections (Eyring et al., 2010; Li et al., 2009; Waugh et al., 2009; WMO, 2010).

2.2 Arctic variability and interactions with climate change

Ozone is a greenhouse gas and radiatively heats the stratosphere. The reduced stratospheric heating related to the substantially reduced ozone amounts in Antarctic spring has led to notable changes in atmospheric dynamics in the Southern Hemisphere (Polvani et al., 2010; Gillett and Son, 2012 and references therein) and may be the cause for observed increases in Antarctic sea ice (Turner et al., 2009). Increases in UV radiation caused by ozone depletion may affect ecosystems with possible feedbacks on the carbon cycle.

On the other hand, climate change affects biogeochemical processes as well as atmospheric dynamics, leading to alterations in emissions and transport times of many trace gases that play a role in stratospheric chemistry and affect ozone (Engel et al., 2009 and references therein; Waugh et al., 2009). A major factor in Arctic ozone depletion, the stability of the polar vortex, directly depends on planetary wave forcing, which is also sensitive to climate change (e.g. Shindell et al., 1999; Sigmond et al., 2008; Dameris and Baldwin, 2012 and references therein; McLandress and Shepherd, 2009). A particular susceptibility of the Arctic stratosphere to climate change may manifest in the trend of cold stratospheric winters in the Arctic getting colder, amplifying heterogeneous chlorine activation and ozone depletion (Rex et al., 2004, 2006).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Over longer periods, climate change and possible geoengineering ventures to mitigate climate change (e.g. Crutzen, 2006) may radically alter the temperature, circulation patterns and chemical composition in the stratosphere. It has been argued (e.g. Tilmes et al., 2008) that the artificial injection of sulphur into the stratosphere considered as a geoengineering venture (e.g. Crutzen, 2006) would increase the extent of Arctic ozone depletion in cold winters and would considerably delay the recovery of the Antarctic ozone hole. To realistically predict the response of the ozone layer to these changes and the future evolution of Arctic ozone, a complete and correct representation of all relevant processes is necessary.

2.3 Major breakthroughs in understanding polar ozone depletion through previous projects and field campaigns

A large number of international research projects (included on the timeline shown in Fig. 1) carried out during the past 25 yr have continuously advanced our understanding of the processes involved in ozone depletion. Virtually all major breakthroughs (e.g. role of chlorine radicals in ozone destruction by Anderson et al., 1989; detection of large nitric acid trihydrate (NAT) particles by Fahey et al., 2001; heterogeneous nucleation and growth of NAT particles by Voigt et al., 2005) have been based on outstanding experimental results gathered during large scale field campaigns. Some of these results were somewhat unexpected or even surprising and have triggered new research.

Since the Airborne Antarctic Ozone Experiment (AAOE) in 1987 (Tuck et al., 1989), numerous campaigns for the exploration of polar as well as mid-latitude and tropical UTLS (upper troposphere and lower stratosphere) processes have been carried out (e.g. Turco et al., 1990; Anderson and Toon, 1993; Newman et al., 2002; Vaughan et al., 2008; Cairo et al., 2010), employing state-of-the-art instrumentation and elaborate flight strategies that exploit the full horizontal and vertical manoeuvrability of high-altitude research aircraft like the US ER-2 and Russian M55 Geophysica and complemented by parallel balloon-borne and lower flying aircraft measurements and extensive modelling

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



activities. The RECONCILE project and the Arctic field campaign described in this paper stand in and continue this fruitful tradition.

2.4 Objectives of RECONCILE

At the beginning of the RECONCILE project in 2009, the lack of understanding in stratospheric dynamics and chemistry was most palpable for issues including the catalytic $\text{ClO}_x/\text{BrO}_x$ chemistry, chlorine activation on cold stratospheric aerosol, NAT nucleation mechanisms, and mixing and transport of processed air to lower latitudes. Therefore, the following specific objectives were defined for RECONCILE:

- Fully understand and correctly represent in global circulation models the transport and mixing across the edge of the polar vortex.

The edge of the polar vortex can be defined in several ways (e.g. Dameris et al., 1995; Harvey et al., 2002; Nash et al., 1996; Waugh and Polvani, 2010). Dynamically, it is roughly centred in the polar night jet that develops in the stratosphere over the winter poles in response to seasonal cooling over the respective polar region. The strong circumpolar westerly winds act as a transport barrier and affect both the extent of ozone depletion and the meridional exchange of ozone depleted air with lower latitudes. Due to the distinct land-sea distribution in the Northern Hemisphere, tropospheric weather regimes and associated planetary wave activity modify the strength of the polar night jet and lead to instabilities. Therefore, the persistence of the polar vortex and, eventually, the strength of transport barrier are rather variable during each winter and vary interannually. A misrepresentation of the transport and mixing across the vortex edge in global circulation models affects our ability to reliably predict both quantities in future climate scenarios.

- Elucidate in detail the processes of NAT nucleation leading to polar stratospheric cloud (PSC) formation, in particular the role of meteoritic dust and other refractory material as condensation nuclei, and the mechanism by which NAT particles grow to rather large sizes forming so-called NAT rocks.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The conventional understanding of PSC formation based on NAT nucleation depending on prior ice formation has been questioned based on in-situ and ground-based lidar observations (Drdla et al., 2002; Pagan et al., 2004; Voigt et al., 2005). However, these in-situ and ground-based lidar observations relied on spatially and temporally limited data and left room for uncertainties. One issue were temperature histories along air parcel trajectories, which were calculated using gridded data of numerical weather prediction (NWP) models. Due to the limited spatial and temporal resolution, these temperature histories were not reliable because, for example, effects of stratospheric mountain waves were not taken into account (Dörnbrack et al., 1998, 1999).

The possible role of meteoritic dust and other refractory material as condensation nuclei has been pointed out (Curtius et al., 2005; Voigt et al., 2005). NAT particles with diameters of tens of microns have been observed and termed “NAT rocks” (Brooks et al., 2003; Fahey et al., 2001; Fueglistaler et al., 2002). When gravitationally sedimenting, these particles efficiently transport and, eventually, redistribute nitric acid, HNO₃. Later in the polar winter, this redistribution impacts the potential for further PSC formation and the rate of chlorine deactivation into chlorine nitrate, ClONO₂.

- Investigate the importance of chlorine activation on background aerosol.

Katja Drdla (Drdla, 2005; Drdla and Müller, 2012) pointed out that at sufficiently low temperatures, chlorine is activated on the background binary aerosol, and that the actual formation of PSCs is not a prerequisite for chlorine activation. This is of particular importance in the context of enhanced stratospheric water vapour concentrations fostering an enhanced total reactive surface area (Anderson et al., 2012), or of geo-engineering concepts to cool the Earth’s surface by deliberately enhancing the stratospheric loading of sulphate aerosol (Crutzen, 2006) or other aerosol compositions (Pope et al., 2012). Clearly, the process of chlorine activation on binary aerosol could have consequences for stratospheric ozone on a global scale.

- Reduce uncertainties and reach an adequate level of confidence concerning the parameters governing the rate of catalytic ozone loss

Uncertainties in various kinetic parameters and the stratospheric bromine budget led to significant uncertainties in model simulations of ozone loss (e.g. Frieler et al., 2006). The single most critical parameter governing the rate of catalytic ozone loss in polar spring is the ClOOCl photolysis rate J_{ClOOCl} , i.e. the product of the actinic flux and the ClOOCl photolysis cross sections σ_{ClOOCl} . In 2007, values of σ_{ClOOCl} determined in numerous laboratory experiments differed by up to a factor of 10, and an experiment by Pope et al. (2007) that suggested the lowest absorption cross sections ever published even put into question our qualitative understanding of the catalytic ozone loss (von Hobe et al., 2007; Schiermeier, 2007; von Hobe, 2007) leading to speculations on previously unknown additional ozone loss mechanisms (e.g. Salawitch et al., 2009; Vogel et al., 2006).

Besides investigating these dynamic, microphysical and chemical processes, and thus completing our knowledge and understanding of the processes leading to polar ozone depletion, the central aim of RECONCILE is to develop reliable process parameterisations and implement them in a global CCM. It is worth pointing out that the model spread in stratospheric ozone simulations and projections is particularly wide for the Arctic region (SPARC, 2010). This can be attributed to high natural variability but also to very significant model differences in the representations of polar processes.

3 RECONCILE strategy and activities

Figure 3 gives a graphical representation of the RECONCILE overall approach. To address the uncertainties and open questions as named in Sect. 1.4, a comprehensive strategy of laboratory experiments, aircraft and match campaigns, microphysical and chemical transport modelling, and data assimilation is employed. The results are used to produce reliable parameterisations of key processes in Arctic stratospheric ozone depletion, which are implemented in a global CCM. The quality of the simulations is assessed against observations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1 Laboratory experiments

3.1.1 The ClOOCl absorption spectrum

A dual wavelength range spectrometer system that can simultaneously perform single pass UV absorption spectroscopy (wavelength range 200–350 nm) and broadband cavity enhanced absorption spectroscopy (BBCEAS, wavelength range 510–560 nm) has been used to investigate the UV and visible absorption cross sections of molecular chlorine (Cl_2) and ClOOCl at different temperatures (Young et al., 2011). With this spectrometer, the experiment carried out by Pope et al. (2007) – i.e. purifying ClOOCl by cold trapping prior to spectral analysis – was repeated, but now with the possibility to simultaneously monitor and unambiguously subtract the Cl_2 contribution to the spectrum, i.e. the impurity that proved to cause major uncertainties in the previous experiment.

3.1.2 Heterogeneous NAT nucleation

The heterogeneous NAT nucleation rate on meteoritic nuclei was investigated by revisiting earlier experiments of Biermann et al. (1996). Back then, an upper bound had been reported for freezing rates of 6 different ternary $\text{H}_2\text{SO}_4/\text{HNO}_3/\text{H}_2\text{O}$ -solutions with immersed micrometeorites under stratospheric conditions. Using a Differential Scanning Calorimeter (DSC) (e.g. Koop and Zobrist, 2009; Marcolli et al., 2007; Zobrist et al., 2008), new bulk experiments were carried out on fragments of the meteorites Orgueil and Murchison with radii varying between 100 and 500 nm, provided in a collaboration with CSNSM (Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, Paris, France).

3.2 Field observations

Extensive field measurements to study processes related to transport and mixing within and at the edge of the polar vortex, PSCs, chlorine chemistry and ozone loss were

30673

ACPD

12, 30661–30754, 2012

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



carried out in the Arctic winter 2009/10. This winter was marked by a sequence of tropospheric disturbances and associated planetary wave activity impacting the formation and evolution of the polar vortex in late November and early December 2009. Afterwards, a period of exceptionally cold temperatures, leading to extensive PSC formation, followed in January 2010 due to the development of a positive phase of the Western Pacific (WP) teleconnection pattern in the troposphere, as described by Nishii et al. (2011) and Orsolini et al. (2009). A major sudden stratospheric warming (SSW) near the end of January 2010 concluded the anomalously cold vortex period and prevented an extensive ozone loss in the RECONCILE winter (cf. Sect. 4.1). A detailed overview of the meteorological situation and evolution of the polar vortex has been given in Dörnbrack et al. (2012). An additional match campaign as well as investigations based on satellite observations were carried out in the subsequent Arctic winter 2010/11 that saw record breaking ozone loss (Manney et al., 2011, cf. Sect. 6).

3.2.1 The Geophysica aircraft campaign

During two measurement periods (17 January–2 February and 27 February–10 March 2010) to cover early and late winter conditions, 12 RECONCILE flights of the high-altitude aircraft M55-Geophysica totalling 57 flight hours were carried out from Kiruna, Sweden, complemented by one flight of the PremierEX project funded by ESA (Spang et al., 2011). An overview of all flights with the respective scientific missions is given in Table 1 and the flight tracks are shown in Fig. 4. An overview of the instruments deployed onboard the M55-Geophysica during the RECONCILE campaign is given in Table 2. The data will be made available online at <https://www.fp7-reconcile.eu/reconciledata.html>.

Flight planning was based on results of the Integrated Forecast System (IFS) of the European Centre of Medium-Range Weather Forecasts (ECMWF). Different thermodynamical fields and derived quantities from the deterministic run were visualized on a web-site as part of the mission support system being developed at DLR (Deutsches Zentrum für Luft und Raumfahrt, Rautenhaus et al., 2012). Chemical forecasts were

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



made using a CLaMS (Chemical Lagrangian Model of the Stratosphere, cf. Sect. 3.5) simulation that started at 1 December 2009 and was run using up-to-date ECMWF analyses and ECMWF forecasts 3 to 4 days into the future.

Air mass trajectories for Lagrangian match experiments were also based on ECMWF forecasts. The idea of the match method consists in probing an individual air parcel twice in order to obtain information on the temporal evolution of the concentration of chemical species (cf. Sect. 3.2.2). During the RECONCILE aircraft campaign it was applied to three flights. During two self-match flights air parcels probed during the outbound flight leg were sampled again during the return flight leg to investigate the temporal evolution of the concentration of chlorine species during sunrise. During one match flight air parcels probed during one flight were sampled again three days later to investigate the transport and mixing processes in the stratosphere (Fig. 5).

3.2.2 Match sonde and balloon activities

A match campaign in the winter 2009/10 consisted of two elements, ozone sonde matches and COBALD backscatter sonde activities, including water vapour soundings.

The match technique to estimate winter ozone loss has been first described by Rex et al. (1999). 31 ozone sounding stations (Fig. 6) participated in the 2010 match campaign and 573 ozone sondes were launched from 29 of these stations. The ozone sonde match campaign started on 5 January and was ended on 28 February, after a major warming had occurred and the meteorological conditions for continuing the campaign deteriorated. The factors that led to the decision to terminate the campaign included large shear and deformation in the flow, expected small scale filaments in the ozone field and the position of the remnants of the polar vortex which were over Eastern Siberia at that time, out of reach of the network of match stations. A vortex remnant returned to Canada and later to the European sector of the Arctic during mid-March. Coordinated sondes were launched into this remnant to assess whether ozone loss had continued after the end of the match campaign.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

An additional ozone sonde match campaign was carried out in the winter 2010/11 with about 270 ozone sonde launches from 13 stations actively participating in the campaign. The campaign started on 12 January 2011 and was stopped on 27 April 2011.

The recently developed light-weight, high-power-LED-based aerosol backscatter sonde COBALD (Compact Optical Backscatter and Aerosol Detector, Wienhold, 2012) was successfully employed during the RECONCILE campaign under Arctic winter conditions (Engel et al., 2012a; Khaykin et al., 2012). During the PSC phase in January 2010, 8 COBALD launches with PSC observations were carried out from Ny-Ålesund, with the advantage of reaching altitudes higher than the M55-Geophysica ceiling altitude. Matches (planned and confirmed using trajectory calculations) with the Geophysica aircraft and with the Cloud-Aerosol Lidar (CALIPSO) satellite (see below) were carried out to allow for a comparison of different particle backscatter measurements and for the observation of cloud changes.

A series of balloon-borne soundings has been carried out in the framework of LAPBIAT-II (Lapland Atmosphere-Biosphere Facility) campaign held at Sodankylä, Finland (360 km East of Kiruna, Sweden) under the premises of Finnish Meteorological Institute's Arctic Research Center (FMI-ARC). A total number of 172 individual balloon instruments including radio and water vapour sondes, aerosol and ozone sensors were flown during the LAPBIAT-II campaign. The balloon flights were coordinated with the aircraft flights and the sounding programme at Sodankylä was carried out in two phases, with 18 payloads flown between 17 January and 6 February 2010 and 15 payloads flown between 10 and 24 March 2010. The stratospheric water vapour measurements were conducted using two types of hygrometers, the Cryogenic Frostpoint Hygrometer (CFH, Vömel et al., 2007) and the Fluorescence Lyman-alpha Stratospheric Hygrometer (FLASH-B, Yushkov et al., 1998). The intercomparison of the data obtained by these two instruments during LAPBIAT-II shows good agreement of $0.73 \pm 5\%$ and consistency in reproducing vertical structures, allowing for the use of FLASH-B and CFH water vapour measurements as interchangeable data series

(Khaykin et al., 2012). During the first phase of LAPBIAT-II the CFH instrument was flown on board of 15 payloads, 11 of which included COBALD backscatter sondes, 5 payloads included FLASH-B sondes and 2 payloads included the backscatter sondes of the University of Wyoming (Rosen and Kjome, 1991).

In coordination with the RECONCILE Geophysica flights, a balloon flight with a remote sensing payload consisting of the instruments MIPAS-B (Michelson Interferometer for Passive Atmospheric Sounding – Balloonborne), TELIS (TERahertz and submillimeter LimbSounder) and mini-DOAS was carried out from Esrange, Kiruna, Sweden on 24 January 2010 (de Lange et al., 2012; Wetzel et al., 2012) to study the diurnal variation of reactive chlorine and nitrogen oxides inside the Arctic vortex from nighttime photochemical equilibrium until local noon covering the full vertical extent of the activated region. Along with O₃, H₂O and tracers the complete nitrogen and chlorine families were measured, allowing to study the partitioning and the budgets of these ozone-controlling substances from the tropopause up to about 34 km.

3.2.3 Satellite observations

Observations made by a number of satellite instruments have been used extensively to support the flight planning during the field campaigns and in scientific process studies. In the following, the satellite instruments and data products that have been of particular importance for the RECONCILE project are briefly described.

PSCs have been observed since June 2006 by the lidar system on the CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation) satellite, which provides extensive daily measurement coverage up to 82 degrees latitude in both hemispheres. CALIPSO PSCs are detected at 180-m vertical resolution and at horizontal resolutions ranging from 5 km to 135 km, and are classified by composition (super-saturated ternary solution STS, liquid/NAT mixtures, ice, and wave ice) based on the measured 532-nm scattering ratio and aerosol depolarization ratio (Pitts et al., 2009, 2011).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The Microwave Limb Sounder (MLS) is an instrument on the EOS (Earth Observing System) Aura satellite, which is in a polar orbit of 705 km at an inclination of 98°, and has provided continuous measurements since August 2004 (Waters et al., 2006). MLS provides about 3500 profiles each day from Earth's surface to ~ 90 km altitude between 82° N and 82° S. We use observations of gas-phase HNO₃ (Santee et al., 2007a), H₂O (Lambert et al., 2007), HCl (Froidevaux et al., 2008), ClO (Santee et al., 2007b), O₃ (Froidevaux et al., 2008) and temperature from retrieval version 3.3 (Livesey et al., 2012). In the stratosphere, the vertical resolution and precision of the observations are 3.5–4.5 km and 0.7 ppb for HNO₃, 3 km and 15 % for H₂O, 3 km and 0.2–0.4 ppb for HCl, 3–4.5 km and 0.1 ppb for ClO, 3 km and 0.04–0.1 ppm for O₃, and 4 km and 1 K for temperature

The Atmospheric Chemistry Experiment–Fourier Transform Spectrometer (ACE-FTS) is a solar occultation instrument on SCISAT-1, flying in a circular orbit at 650 km at an inclination of 74° and has made measurements since 2004 (Bernath et al., 2005). ACE-FTS provides daily vertical profiles for up to 15 sunrises and 15 sunsets with latitudinal coverage exhibiting an annual cycle between 85° S to 85° N. We use observations of ClONO₂, N₂O₅ (Wolff et al., 2008) and HCl (Mahieu et al., 2005) from retrieval version 3 with a vertical resolution of 3–4 km.

The Sub-Millimetre Radiometer (SMR) onboard the Odin satellite observes the thermal emission from the Earth limb and provides vertical trace gas profiles including O₃, N₂O, and HNO₃. Stratospheric mode measurements are generally performed in the altitude range from 7 to 70 km with a resolution of 1.5 km below 50 km tangent altitude and 5.5 km above, normally covering the latitude range between 82.5° S and 82.5° N (Urban et al., 2009; Murtagh et al., 2002).

To assess the impact of the 2011 Arctic ozone hole on surface UV radiation and compare the results to previous Arctic and Antarctic winters (Sect. 6), local-noon clear-sky UV index data products from a combined MSR/SCIAMACHY (Scanning Imaging Absorption Spectrometer for Atmospheric CHartographY) dataset (Eskes et al., 2005; van der A et al., 2010; Allaart et al., 2004) were averaged as a function of day-of-

year and latitude (von Hobe et al., 2012). The comparison of clear-sky data isolates the ozone effect on UV-radiation, i.e. interannual variability due to changes in albedo (snow cover), clouds and aerosol loading is excluded in the analysis.

3.3 Microphysical modelling

To understand STS, NAT, and ice formation in the Arctic stratosphere, the Zurich Optical/Microphysical box Model (ZOMM) is used. ZOMM has been described previously by Luo et al. (2003) and Meilinger et al. (1995) and was applied very recently in detailed cirrus modelling (Brabec et al., 2012). Along the lines of previous understanding of PSC formation (Peter, 1997) – kinetic growth of STS by simultaneous uptake of H_2O and HNO_3 , homogeneous nucleation of ice in these solution droplets (Koop et al., 1995) and then heterogeneous nucleation of NAT on pre-existing ice particles (Carslaw et al., 1998) along air parcel trajectories – the pre-RECONCILE version of the model focussed on simulating exactly these processes. RECONCILE observations triggered the implementation of additional new heterogeneous formation pathways for NAT and ice. To this end, solid nuclei were implemented by assuming the presence of non-volatile particles with a radius of 20 nm and a uniform number density of 7.5 cm^{-3} throughout the Arctic stratosphere. These assumptions on size and number density reflect the in-situ measurements by the heated condensation nuclei counter COPAS on board of Geophysica during RECONCILE and EUPLEX (cf. Sect. 4.2.2).

These heterogeneous nuclei for NAT and ice formation (Hoyle et al., 2012; Engel et al., 2012b) are then allowed to compete with the accepted pathway of NAT formation via homogeneous nucleation of ice, which typically requires cold localized pools in mountain-wave-driven temperature fluctuations (Carslaw et al., 1998; Peter, 1997). In the absence of better knowledge of the nucleation properties of these nuclei, immersion freezing of NAT and ice was assumed to occur on active sites much like heterogeneous ice nucleation on Arizona test dust demonstrated in previous laboratory experiments (Marcolli et al., 2007). This approach still needs to be reconciled with the laboratory experiments of Biermann et al. (1996), who established upper limits of NAT nucleation

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rates and classified direct heterogeneous NAT nucleation on pre-existing solid particles as relatively inefficient.

3.4 Chemistry transport modelling

The Chemical Lagrangian Model of the Stratosphere (CLaMS, Konopka et al., 2004; McKenna et al., 2002a,b; Grooß et al., 2005) is a hierarchy of models ranging from a box model to a 3-D Chemistry Transport Model (CTM) based on a Lagrangian transport concept, in which the chemical evolution within individual air parcels is simulated along trajectories determined from ECMWF temperature data and wind fields. Stratospheric chemistry in CLaMS is an update of that described in McKenna et al. (2002a) and now includes reactions of importance in the upper stratosphere. 143 reactions of 45 variable chemical species now explicitly contain reactions involving H radicals as well as N₂O and CFCs. Chemical reaction rates and absorption cross sections are based on JPL 2011 (Sander et al., 2011) except when explicitly noted otherwise. Heterogeneous reaction rates are parameterised as described in Wegner et al. (2012a), and a particle based denitrification scheme (Grooß et al., 2005) is implemented.

For RECONCILE, CLaMS has been used extensively on all spatial, temporal and complexity scales to investigate individual physical and chemical processes using field observations and to develop mathematical parameterisations for use in global prognostic models (cf. below). Hemispheric simulations have been carried out with horizontal resolutions up to 70 km. For the mixing parameterisation (McKenna et al., 2002b; Konopka et al., 2004), a time step of 24 h and a critical Lyapunov coefficient of 1.5 day⁻¹ were used (see also Riese et al., 2012). To study transport and mixing processes, and to distinguish between dynamic and chemical processes acting on the concentrations of certain trace gases, dynamical tracers that are not influenced by chemistry such as passive ozone, passive NO_y, or artificial air mass origin tracers are used. The evolution of the polar vortex composition during the winter 2009/10 has been investigated utilizing a suite of 9 inert tracers defined by PV thresholds based on an algorithm by Nash et al. (1996). The tracer fields were initialized on 1 December 2009 according

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the initial position of the air masses in the modified potential vorticity (PV) space (see Günther et al., 2008 and references therein). To further quantify the in-mixing of air masses from lower latitudes into the polar vortex during the vortex split in mid December 2009, an additional simulation starting at the beginning of January 2010 with re-initialized tracers was performed.

ATLAS is another global Chemistry Transport Model with a focus on the stratosphere (Wohltmann et al., 2010; Wohltmann and Rex, 2009), based on the same Lagrangian approach as CLaMS. Transport and chemistry are driven by ECMWF ERA Interim data for RECONCILE. The model includes a stratospheric chemistry module that comprises 47 species and more than 180 reactions, detailed heterogeneous chemistry on Polar Stratospheric Clouds and a particle-based Lagrangian denitrification module. For RECONCILE, ATLAS has been used for global modelling studies of the 2009/10 winter (Wohltmann et al., 2012).

3.5 Data assimilation

A data assimilation scheme to estimate chemical ozone loss in the Arctic winter stratosphere based on a 3D-Var version of the operational Met Office assimilation system (Jackson, 2007) has been developed by Jackson and Orsolini (2008) to address some potential shortcomings of vortex-averaged and other methods used to estimate ozone loss, namely, the definition of the vortex edge, and estimates of mixing across the vortex edge. Jackson and Orsolini's results showed that the data assimilation method is very promising and can lead to potentially more accurate ozone-loss estimates than other, more established methods.

EOS MLS and SBUV/2 ozone observations are assimilated, together with dynamical observations from satellites, aircraft, radiosondes and surface stations. A realistic ozone field is produced by running the ozone assimilation (referred to as ASSIM) for one week. The ASSIM run is continued for several more weeks, while a reference run (referred to as REF) is also started using the same initial conditions, but continued without further ozone observations being assimilated. Hence in REF ozone is initialized

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with EOS MLS and SBUV/2 observations and thereafter transported passively by assimilated winds. The one week initialization step ensures that the reference ozone is realistic and shares the same potential observational biases as the assimilated ozone in the ASSIM run. To estimate chemical ozone loss, REF ozone is subtracted from the ASSIM ozone field. The dynamical fields in both runs are near-identical and differences between the assimilated and reference ozone fields can be attributed to chemistry. In the middle and high-latitude winter lower stratosphere, it is likely that these chemical changes will be overwhelmingly due to heterogeneous PSC-related chemistry. Applying this method for the Arctic winter 2005 between early February and mid March at 450 K, Jackson and Orsolini (2008) reported chemical ozone loss of 0.6 ppm, at the lower edge of the 0.6–1.2 ppm range for ozone loss estimates for this period based on other methods.

3.6 Global modelling

Global models play a crucial role in understanding the chemical and dynamical processes in the atmosphere. CCMs provide the only tool for predictions and projections, therefore their validation and the evaluation and improvement of the performance of parameterizations are important parts of the continuous improvement.

CCM activities within RECONCILE are carried out with the LMDz-Reprobus model (Marchand et al., 2012; Jourdain et al., 2008) that couples the LMDZ general circulation model (Hourdin et al., 2006, 2012; Le Treut et al., 1998; Sadourny and Laval, 1984) and the Reprobus stratospheric chemistry package (Lefèvre et al., 1994, 1998). The performances of LMDz-Reprobus simulations of stratospheric ozone and climate have been evaluated against a wide range of observations and against other chemistry-climate models (Austin, 2010; Jourdain et al., 2008; Gettelman et al., 2010; Hegglin, 2010). In RECONCILE, the configuration of LMDz (LMDZ5) is identical to that employed for the Coupled Model Intercomparison Project Phase 5 simulations (CMIP5, Hourdin et al., 2012): the model extends from the ground up to 65 km on 39 hybrids pressure vertical levels, and the horizontal resolution is 3.75° in longitude and 1.875° in

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

latitude. The chemical package provides a comprehensive description of 55 chemical species of stratospheric interest, by means of the usual set of gas-phase and heterogeneous reaction rates recommended by the JPL evaluation (Sander et al., 2011). In its pre-RECONCILE version, the CCM scheme employed for the treatment of polar stratospheric clouds is a simple equilibrium scheme that predicts the existence of liquid (binary $\text{H}_2\text{O}/\text{H}_2\text{SO}_4$ or ternary $\text{H}_2\text{O}/\text{H}_2\text{SO}_4/\text{HNO}_3$ solutions) and solid (NAT or H_2O ice) aerosols from the local thermodynamic conditions in each model grid box. When saturation is reached, solid particles are assumed to form instantaneously at a fixed radius (two modes of 0.5 and 6.5 μm for NAT, 10 μm for water-ice) and only their number density (particles cm^{-3}) is variable. As for all equilibrium schemes, the scheme has no memory of the temperature history of the air mass and the particles are not transported separately from one time step to the other. This usual treatment of PSCs in CCMs is computationally cheap but cannot represent properly the slow growth of large solid particles (such as “NAT rocks”) and their consequences on denitrification and ozone loss.

The main objective of the global modelling activity in RECONCILE is to assess the impact of the results from the process studies related to the field and laboratory measurements on predictions of polar stratospheric ozone loss. This involves an ensemble of long-term transient simulations covering the period 1960–2050 with different parameterisations of polar stratospheric clouds. The reference simulations of LMDz-Reprobus for the past and future evolution of the ozone layer use the standard reaction rates recommended by JPL 2011 and the pre-RECONCILE PSC equilibrium scheme described above. These simulations also include the same scenarios for the emissions of greenhouse and ozone-depleting gases as those published in the latest WMO assessment (WMO, 2010). On the other hand, the improved version of the CCM takes full benefit of the results of RECONCILE obtained in the laboratory, in the field, and by CTM modelling. In this version, a coherent set of kinetic parameters for ClO_x chemistry that is consistent with observations (Table 3 and Sect. 4.3) is adopted, and a more

realistic parameterisation of PSCs has been developed based on the results described in Sect. 4.2.

The improved version of LMDz-Reprobus will be integrated over the same timescales as the reference simulation and the impact of the new parameterisations developed from RECONCILE will be evaluated against the observed ozone trends since the 1960s and for the future period. A full analysis of the CCM runs carried out within the RECONCILE framework will be presented in a future paper.

While the last big cross-validation project CCMVal-2 (Eyring et al., 2010; Gillett et al., 2011; SPARC, 2010) provided comprehensive bias estimates for several models, we extended the methodology by comparing dynamical features of empirical data and time series obtained by the LMDZrepro simulations. The main focus was on a global analysis of spectral properties, two-point correlations determined by detrended fluctuation analysis (DFA), and detecting higher order correlations by the spectral surrogate method (see Sect. 5.2).

4 Processes in the Arctic vortex

In this section, we summarize the key results from RECONCILE in terms of improved understanding of dynamic, microphysical and chemical processes relevant to polar ozone loss. The field measurements in 2010 provided good opportunities for process studies, even though 2009/10 was a moderate winter in terms of ozone loss. From the match campaign, maximum loss rates on the order of 60 ppb day^{-1} in mid-February and an overall column ozone loss of about 66 DU inside the polar vortex were determined.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1 Vortex dynamics and mixing

4.1.1 The evolution of the 2009/10 Arctic vortex and factors influencing vortex stability

The cooling of the air inside the polar vortex generally followed the 21-yr ECMWF climatological mean from summer until mid-November 2009, when the minimum temperature (T_{MIN}) at the 50 hPa pressure surface had dropped to about 200 K (Fig. 7). From this time until mid-December 2009, the minimum temperatures inside the vortex were well above the climatological mean. The period was characterized by significant planetary wave activity (see Fig. 8 in Dörnbrack et al., 2012). As a consequence of these disturbances, the polar vortex split into two unequally strong lobes during the first ten days of December 2009 (Fig. 8, left hand column). The two lobes rejoined, including some mid-latitude air in the process, and the new vortex cooled gradually through mid-January 2010 (second column in Fig. 8). There was a significant drop in T_{MIN} to values as much as 9 K below the climatological mean (Fig. 7). These coldest conditions in the entire winter occurred during a brief period between two sudden stratospheric warmings (SSWs). Indeed, the polar stratospheric cooling and vortex strengthening during this period were a response of the weakened planetary wave activity (Dörnbrack et al., 2012). The formation of such an exceptionally cold vortex corresponds to the development of a positive phase of the Western Pacific (WP) teleconnection pattern in the troposphere, as described by Nishii et al. (2011) and Orsolini et al. (2009). The anomalously cold vortex period was concluded near the end of January 2010 by the onset of a major SSW that was also related to tropospheric features (Ayarzaguena et al., 2011). In early February, the vortex again broke apart into two lobes (third column in Fig. 8) which rejoined again in early March with patches of mid-latitude air that were included in the process. The new single vortex remained intact throughout March (Fig. 8, right hand column). A climatological analysis revealed the surprising result that the 2009/10 winter was the third warmest winter in the 21 yr period from 1989 to 2009 measured by the polar cap temperatures at 50 hPa (see Tables in Dörnbrack et al., 2012).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A comprehensive analysis of SSWs in past Arctic winters also suggests profound tropospheric forcing for the observed major SSWs (Kuttippurath and Nikulin, 2012). The authors also present a statistical analysis for the past 17 Arctic winters showing that ozone loss closely relates to the intensity and timing of major warmings in each winter and argue that a particularly high frequency of major warmings observed in recent Arctic winters could have significant implications for stratospheric ozone trends in the Northern Hemisphere.

4.1.2 Mixing within the polar vortex and across its edge

According to the CLaMS artificial tracer simulations, 60 % of vortex air were affected by mixing as a result of the vortex split in December 2009 (Fig. 8, left hand column). After a re-organization before January 2010 the vortex stayed coherent with significant mixing only below 500 K (Fig. 8, second column). Following the second vortex split enhanced transport of air from lower latitudes into the reformed vortex occurred (Fig. 8, third column), leading to a heterogeneous origin distribution inside the vortex interior during late winter (Fig. 8, right hand column). The results from the CLaMS simulations were validated with observations obtained by the CRISTA (CRyogenic Infrared Spectrometers and Telescopes for the Atmosphere) (Kalicinski et al., 2012) and HAGAR (Hösen et al., 2012) instruments. Analysis of tracer-tracer correlations observed by HAGAR also indicate ongoing irreversible mixing of the intruded air masses inside the vortex in March 2010 (Hösen et al., 2012), which is also captured by the CLaMS simulations.

A detailed case study of model uncertainties in reproducing observed tracer concentrations in a region of strong gradient was carried out for the RECONCILE match flight (Wegner et al., 2010). The comparison of mixing ratios of the tracer N_2O at the start and end points of trajectories connecting the two match flights on 30 January and 2 February revealed that the matches were not always as successful as planned (Fig. 5), and that the determination of matching air masses on the timescale of several days solely by trajectory calculations is insufficient. The mean uncertainty of the trajectory

calculations over three days was 65 km, even when they were based on high resolution ECMWF wind fields. This could potentially impair ozone loss estimates by the match technique in an inhomogeneous vortex.

4.1.3 Data assimilation

5 Issues related to mixing in and at the edge of the polar vortex were also investigated for the assimilation-based method. This was part of a broader study where ozone loss estimates made using the assimilation-based method were compared to ozone loss estimates obtained by running the University of Oslo CTM (Sovde et al., 2011). The focus was on the winter of 2006/07, and the assimilation used 3D-Var, while the CTM
10 constituents were advected using ECMWF analyses produced by 4D-Var. The results indicated that the assimilation-system vertical transport is found to be too fast, which affects the ozone reference field used for the loss estimation, and thus points to an underestimation in the ozone loss estimate. However, horizontal transport errors in the assimilation system were found to be small compared to the vertical transport errors.
15 Improving the ozone reference field, by advecting the Met Office ozone field by the ECMWF 4D-Var winds instead of the Met Office 3D-Var assimilated winds, produced results in better agreement with those produced by the CTM. These results are in agreement with other studies that indicate that stratospheric transport is improved when meteorological analyses are produced using 4D-Var instead of 3D-Var.

4.2 PSC evolutions and microphysical processes

4.2.1 PSC observations

A large and complementary dataset on PSC properties has been obtained during the RECONCILE winter by ground-based, aircraft and space-borne measurements. The evolution of PSCs during the 2009–2010 Arctic winter is detailed in Pitts et al. (2011).
25 During this unusual winter, more PSCs were observed by CALIPSO than in the

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

previous three Arctic seasons combined. Examples of CALIPSO PSC observations during the season are shown in Fig. 9. The early period from 15–30 December 2009 was characterized primarily by patchy, tenuous low number density liquid/NAT mixtures (Fig. 9a). No ice clouds were observed by CALIPSO during this early phase, suggesting that the NAT clouds were formed through a non-ice nucleation mechanism. From 31 December 2009–14 January 2010, there were frequent mountain wave ice clouds that nucleated widespread NAT particles throughout the vortex (Fig. 9b). Synoptic-scale temperatures in the Arctic fell below the frost point during the period from 15–21 January 2010, which led to a rare outbreak of widespread ice PSCs (Fig. 9c). The final period of the PSC season (22–28 January) was characterized by a major stratospheric warming that distorted the vortex, displacing the cold pool from the vortex centre. CALIPSO observed primarily STS PSCs during this final period, although NAT particles may have been present in low number densities, but masked from detection by the more abundant STS droplets at colder temperatures (Fig. 9d). The evolution of PSCs in January 2010 observed by CALIPSO was also seen by ground based lidar observations near Kiruna (Khosrawi et al., 2011). PSCs have also been observed by COBALD in every flight within the period from 17 to 25 January 2010. Closely matched measurements between CALIPSO and COBALD agree well in their backscatter profiles at PSC levels (Fig. 10) and also the agreement between COBALD and the Ny Ålesund lidar is convincing.

The in situ optical particle spectrometers onboard the Geophysica FSSP (Forward Scattering Spectrometer Probe) and CIP (Cloud Imaging Probe) observed PSCs during the first five RECONCILE flights. Measurements with these instruments show the existence of so-called NAT-rocks up to sizes of roughly 25 μm in diameter, confirming earlier observations (Brooks et al., 2003; Fahey et al., 2001; Northway et al., 2002). The probably first images of NAT-rocks recorded by the greyscale CIP (optical array probe) provide an additional proof of their existence with an optical detection technique other than forward scattering. Those images were observed exclusively when the penetration of NAT fields during a RECONCILE flight was confirmed also by the MAL

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Miniature Aerosol Lidar), MAS (Multiwavelength Aerosol Scattersonde), and SIOUX (Stratospheric Observation Unit for nitrogen oxides) instruments (cf. Table 2). The CIP greyscale image data showed mainly 2–4 pixel features which correspond to particles of roughly 15 μm to 30 μm in diameter. Larger particles were not detected. The images recorded during a PSC penetration on 25 January 2010 over a time period of 30 min amounts to a number concentration of about 10^{-3} cm^{-3} , which is in the same range as measured by the FSSP. The NO_y instrument (SIOUX) capable of resolving large NO_y -containing particles supports the optical data, including the assumed composition of the particles and consequently the assumption of the refractive index of NAT for the processing of FSSP data. Overall, the high number concentration of large NAT particles amounts to condensed phase HNO_3 concentrations approaching the theoretical limit of available NO_y , raising the question whether the composition of these particles is exclusively NAT. Alternatively, non-spherical shapes of the NAT particles might contribute to a certain overestimation in the particle size and therefore in the total volume of the particle phase. Figure 11 shows a size distribution averaged over a period of 22 min that was measured by the FSSP-100 instrument in one of the dense NAT fields. This size distribution is compared to the theoretical size distribution that Fahey et al. (2001) retrieved from their NO_y measurements made in 2000.

Beside the large NAT particles, the FSSP-300 detected a particle mode around 0.5–1 μm which can be attributed to STS particles. It shows volume to temperature behaviour close to the one measured (Dye et al., 1992) and modelled for STS (Peter, 1997). Throughout all proven PSC penetration events, the particles larger than about 1 μm in diameter were detected only at ambient temperatures below the NAT equilibrium temperature (T_{NAT}), but mostly well above the frost point (T_{ICE}) so that pure ice particles can most likely be excluded. Only at a few short parts along some of the flight tracks ambient air temperature just fell down to T_{ICE} . There was no indication for the presence of ice from the observations of gas phase and total water by FLASH (FLuorescent Airborne Stratospheric Hygrometer) and FISH (Fast In situ Stratospheric Hygrometer). Consistent with the in situ observations, the presence of ice-particles at

or below the flight altitude is not supported by the spectral signatures seen by CRISTANF. Spectra simulated for in situ measured particle size distributions (similar to the one shown in Fig. 11) compare reasonably well with the observed IR limb spectra under the assumption that the large particles consist of NAT (Griessbach et al., 2012; Kalicinsky et al., 2012).

The series of balloon-borne measurements of water vapour and aerosol acquired within the LAPBIAT-II campaign in Sodankylä during the second half of January revealed an unprecedented evidence of water redistribution in the Arctic stratosphere as a consequence of ice PSC formation and concurrent irreversible dehydration by up to 1.5 ppm within 20–24 km altitude range followed by a rehydration in a 2 km thick layer below. The source region and the spatiotemporal evolution of the dehydrated air masses were well established using AURA MLS observations (Khaykin et al., 2012). For the first time, an active phase of ice PSC formation was captured by the simultaneous balloon measurements of water vapour and aerosol on 17 January, providing a unique high-resolution snapshot of repartitioning of water vapour into ice particles. This observation, bearing important implications for the PSCs formation thresholds, is analyzed in detail by Engel et al. (2012a) using microphysical box modelling.

4.2.2 Characterisation of background aerosol

The measurements with the COndensation PArTicle counting System (COPAS, Weigel et al., 2009) during the RECONCILE winter agree well with the data from Curtius et al. (2005). While outside the polar vortex, the fraction of non-volatile (at 250 °C) particles remains nearly constant with altitude, it increases noticeably with altitude inside the vortex, reaching up to 80 % of detected sub-micron refractory particles deeply inside the Arctic vortex ($\Theta \approx 470$ K).

Nevertheless, the hypothesis by Curtius et al. (2005) that the enhanced non-volatile aerosol in the polar vortex mainly consists of meteoritic smoke particles needs further refinements. Weigel et al. (2012) present a detailed chemical analysis of the refractory aerosol material from 13 (one for each flight) in-vortex particle samples that

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

were collected using a miniaturized impactor system placed downstream of the COPAS aerosol inlet, generally at $\Theta > 410$ K. Subsequently, the samples are processed by using offline individual particle analysis with Environmental Scanning Electron Microscopy (ESEM) and Energy Dispersive X-ray (EDX) analysis. Many of the samples contained spheres and aggregates of globules (like “a bunch of grapes”, Bigg et al., 1970) in the size range 500 nm–2 μ m, chemically dominated by metallic species such as Al, Cr, Mn, Fe or Ni. The chemical composition and morphology indicates that these particles most likely originate to a large extent from space debris. Apart from these metallic particles, also fractured alumo-silicate particles, showing the character of crustal material, were found. In many samples for smaller particles, i.e. $d_p < 500$ nm, also a high number of soot and lead-bearing particles were found, indicating anthropogenic origin. Based on these results, metallic particles (from meteoritic or space debris ablations) are not the major compound of the vortex refractory aerosol material. In particular the fact that surface aerosol material and even anthropogenic particles were found in the subsiding air masses over the Arctic in winter is an essential new finding. It indicates efficient transport pathways for aerosol particles to reach elevated atmospheric altitude, i.e. the stratosphere or even in the mesosphere, to re-enter the lowermost stratosphere due to the developing polar vortex. It is conceivable, that the abundance of non-volatile particles (up to 80 %) in the Arctic winter stratosphere significantly aids the formation of PSCs, and furthermore influences the tropospheric cloud formation after the polar vortex has dissolved in the Arctic spring season.

4.2.3 New insights into particle nucleation and PSC formation

The CALIPSO observations during the RECONCILE winter were analyzed in detail to investigate the heterogeneous nucleation pathway of NAT and ice on pre-existing solid particles. The presence of an ample supply of such nuclei in the polar vortex is justifiable as described in Sect. 4.2.2.

For every distinct PSC period defined by Pitts et al. (2011) and described in Sect. 4.2.1, optical parameters simulated by ZOMM (cf. Sect. 3.3) were compared to

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

selected CALIPSO PSC observations. The best agreement between model results and observations is achieved by (1) allowing NAT and ice to nucleate heterogeneously, and (2) superimposing small scale temperature fluctuations onto the trajectories (which otherwise remain unresolved along the ECMWF trajectories). Only the combination of both effects can provide good agreement in particle types and captures the backscatter and depolarization ratios measured by CALIPSO (Hoyle et al., 2012; Engel et al., 2012c). Figure 12 shows the new scheme of PSC formation and phase transitions that will be presented and discussed in detail in two upcoming papers (Hoyle et al., 2012; Engel et al., 2012b).

At first sight these results appear to be at odds with the previous laboratory-based understanding of NAT nucleation, namely that ice would be the first solid, nucleating homogeneously within liquid aerosol particles (Koop et al., 1995), and that NAT nucleated only subsequently on ice (Carslaw et al., 1998). In these days direct heterogeneous NAT nucleation on pre-existing solid particles was thought to be unlikely (Biermann et al., 1996). Within RECONCILE we presently undertake additional laboratory experiments on meteoritic material. Preliminary experiments corroborate the early results of Biermann et al. (1996), but also seem to suggest that the heterogeneous NAT nucleation rate may indeed reach the upper limits specified by Biermann and colleagues. With today's new insights into low number density NAT clouds, as observed by CALIPSO in December 2009, and NAT-rocks, described by Fueglistaler et al. (2002), nuclei consisting of coagulated meteoritic smoke particles or micrometeorites become favourable candidates to explain those observations.

4.2.4 Vertical redistribution of NO_y

One of the RECONCILE objectives is to understand the processes causing the vertical NO_y -redistribution, i.e. denitrification above about 19 km altitude and a renitrification below.

In 2009/10, Odin/SMR observations showed the strongest denitrification since the start of the Odin measurement period in 2001. However, this was exceeded in the

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2010/11 Arctic winter (cf. Sect. 6). Box model simulations together with Odin/SMR, Aura-MLS and CALIPSO observations of HNO_3 , water vapour and PSC particles respectively show that the strong denitrification observed in mid January 2010 is linked to an unusually strong synoptic cooling event in mid January with ice particle formation on NAT as a possible formation mechanism. The denitrification in the beginning of January was more likely caused by the sedimentation of NAT particles that formed on mountain wave ice clouds (Khosrawi et al., 2011).

During the aircraft campaign, NO_y compounds were observed in situ by the SIOUX instrument and also by remote sensing using MIPAS-STR and CRISTA-NF (Woiwode et al., 2012b; Ungermann et al., 2012). HNO_3 observations from the ACE-FTS satellite instrument are also available for the entire winter.

One critical parameter determining the rate and extent of vertical HNO_3 redistribution in the polar vortex is the nucleation rate for NAT particle formation. A CLaMS simulation of denitrification was performed in a Lagrangian way using particle trajectories that follow individual NAT particles. Previous simulations (Groß et al., 2005) that assumed globally constant nucleation rates for all locations with $T < T_{\text{NAT}}$ produced reasonable results for the winter 2002/3 using a nucleation rate of $7.8 \times 10^{-6} \text{ cm}^{-3} \text{ h}^{-1}$. Using the same nucleation rate, the observed vertical NO_y redistribution in the winter 2009/10 was underestimated, suggesting faster NAT nucleation. In CLaMS simulations with the new nucleation rate parameterisation described in Sect. 3.3, the vertical redistribution of NO_y compares better to the observations, but still underestimates the observed denitrification in early March.

Significant HNO_3 enhancements attributed to renitrification processes were observed by MIPAS-STR and CRISTA-NF in the polar vortex after the end of the PSC phase in the Arctic winter 2009/10. Standard retrieval results from MIPAS-STR showing vertical distributions of HNO_3 versus potential temperature are presented in Fig. 13 for the flight on 30 January and for two flights in March (vortex air identified by measured tracer distributions). A significant HNO_3 maximum peaking around 405 K is identified for the flight in January directly after the PSC phase of this winter. The redistribution of

5 HNO_3 as a source for NO_2 is closely linked to the deactivation of active chlorine species into ClONO_2 . A dedicated study is carried out involving refined MIPAS-STR retrievals and CLaMS simulations considering the new temperature-dependent NAT nucleation parameterisation to verify whether the representation of HNO_3 redistribution in CLaMS is improved and whether ClONO_2 formation is simulated adequately (Woiwode et al., 2012a).

4.3 Stratospheric halogen chemistry and catalytic ozone depletion

4.3.1 Heterogeneous chlorine activation

10 As expected, the cold vortex temperatures and extensive PSC coverage in December 2009 and January 2010 led to substantial chlorine activation. A broad activated layer of enhanced ClO between 16 and 26 km altitude was detected by MIPAS-B on 24 January (Wetzel et al., 2012), corresponding nicely to the vertical extent of PSCs/cold temperatures observed by MIPAS-B in the same data set, as well as CALIPSO (Pitts et al., 2011) and ground based lidars (Khosrawi et al., 2011) in the time period preceding the flight. ClONO_2 observations indicate no or very little activation in the air masses below and above this layer.

15 The RECONCILE flights on 20, 22 and 24 January showed temperatures where chlorine activation would be expected and were analysed by CLaMS simulations along seven-day backward trajectories (chemistry was initialised from a hemispheric CLaMS simulation, cf. Sect. 3.4). Up to 1 ppb of additional chlorine activation was produced, and the simulated ClO_x values are consistent with HALOX (HALogen Oxide monitor) observations along the corresponding flight tracks (Fig. 14). Virtually no difference in chlorine activation exists between simulations employing the full PSC scheme with the aerosol surface area density (SAD) increasing with STS, NAT and ice formation and simulations keeping the SAD at background levels, demonstrating that at sufficiently low temperatures heterogeneous chemistry on the cold binary background aerosol is sufficient to produce the same amount of active chlorine as the full PSC scheme for the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



case considered (Wegner et al., 2012a). The same paper describes another CLaMS investigation for an episode in the Arctic winter 2004/5 constrained by aircraft and satellite data. Significant chlorine activation was produced by CLaMS along 96 h trajectories leading up to a flight on 7 March 2005 (von Hobe et al., 2006) with the resulting ClO_x levels comparing well to the observations. Temperatures were low enough for chlorine activation to occur but significant denitrification probably prevented the formation of NAT and STS, so that chlorine activation occurred to a very large extent on cold background binary aerosol.

Wegner et al. (2012a) corroborate the conclusions of their CLaMS investigations by an analysis of Aura-MLS observations for seven Arctic winters (2004/5–2010/11), focusing on HCl as an indicator for chlorine activation and HNO_3 as an indicator for the formation of PSC particles. Below 550 K potential temperature, the vortex average removal rates of gas-phase HCl and HNO_3 are not correlated. By 1 January, the vortex average shows that for all considered winters HCl has been removed from the gas-phase to a similar extent while HNO_3 shows great interannual variability. This indicates that at these altitudes the additional surface area provided by PSCs does not directly influence the rate of chlorine activation. However, at higher altitudes, where the background aerosol begins to thin out, ice PSCs may play an important role. Between 600 and 650 K, the maximum HCl depletion is observed in the winter 2009/10, which was unique in terms of occurrence of ice PSCs. Other than in the Arctic, HCl depletion cannot be used as an indicator for chlorine activation in the cold vortex core of the Antarctic, where condensation of HCl into STS presents a viable pathway to sequester HCl from the gas-phase. This has been demonstrated by Wegner et al. (2012c) who argue that during the polar night a major part of HCl is dissolved in STS particles, because heterogeneous reactions, especially with ClONO_2 , cannot explain the observed loss of gas-phase HCl due to shortage of NO_x even if the photolysis of condensed phase HNO_3 is taken into account as a potential additional NO_x source.

A comprehensive set of sensitivity runs to examine the impact of known uncertainties in heterogeneous chemistry and PSC microphysics was carried out using the ATLAS

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

model (Wohltmann et al., 2012). The simulations were compared to MLS and ACE-FTS observations and in-situ measurements onboard the Geophysica during the aircraft campaign. Results show that ozone loss is a surprisingly robust quantity with regard to changes in the microphysical assumptions with column ozone loss never varying more than 10% in all sensitivity runs (Fig. 15, right panel). In agreement with Wegner et al. (2012a), chlorine activation on liquid ternary aerosols alone and even on the binary background aerosol was sufficient to explain the observed order of magnitude and morphology of the abundances of chlorine, reservoir gases and ozone (Fig. 15). Assuming default values for NAT number density and supersaturation, runs with different rates for the heterogeneous reactions $\text{ClONO}_2 + \text{HCl}$ and $\text{ClONO}_2 + \text{H}_2\text{O}$ on liquid aerosols (Hanson and Ravishankara, 1994; Shi et al., 2001) and on NAT (Carslaw and Peter, 1997; Carslaw et al., 1997; based on laboratory studies by Abbatt and Molina, 1992; Hanson and Ravishankara, 1993) produce very similar results. Only if an appreciable NAT surface area density is assumed and the reactions are not dominated by liquid aerosols, the rate and degree of chlorine activation becomes significantly smaller with the rates based on Abbatt and Molina (1992).

Figure 16 summarizes the rate of the most important heterogeneous chlorine activation reaction, $\text{HCl} + \text{ClONO}_2 \rightarrow \text{Cl}_2 + \text{HNO}_3$ on different particle types as a function of temperature. Above about 192 K, binary aerosol can activate chlorine just as fast as NAT. Below 192 K, the activation rate increases along a smooth transition from binary to ternary aerosol. Below the frost point, ice PSCs can increase the activation rate again by about a factor of 10.

4.3.2 ClO_x catalysed ozone destruction

The UV/Vis spectrum (Young et al., 2012) obtained from the laboratory experiment described in Sect. 3.1 agrees well with the one reported by Papanastasiou et al. (2009) that has been adopted for the most recent JPL recommendation (Sander et al., 2011). Furthermore it is in close agreement with all the ClOOCl spectra which appeared subsequent to Pope et al. (2007), namely: Chen et al. (2009), Lien et al. (2009), Wilmouth

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al. (2009), Jin et al. (2010) and the spectrum reported by von Hobe et al. (2009) which is scaled to the Lien et al. (2009) measurement. It is clearly not in agreement with the Pope et al. (2007) results which likely suffered from subtracting too much Cl_2 from the measured composite $\text{Cl}_2/\text{ClOOCl}$ spectrum. The measurement of the much weaker visible spectrum of ClOOCl is the first time this spectrum has been measured. It provides important information to constrain the long wavelength absorption of ClOOCl and hence the impact of this absorption on ClO_x catalysed polar O_3 cycles.

Results from the laboratory experiment are consistent with atmospheric observations made during the self-match flight (cf. Table 2) published by Sumińska-Ebersoldt et al. (2012). The observed increase of ClO after sunrise is consistent with the Papanastasiou et al. (2009) cross sections, and essentially rules out ClOOCl absorption cross sections lower than the spectrum published by von Hobe et al. (2009) scaled to absolute cross sections near the peak of the spectrum (246 nm) observed by Lien et al. (2009). The field measurements do not support an additional ClOOCl absorption band of atmospheric significance in the visible region of the spectrum. Remaining uncertainties in the ClOOCl cross sections influence J_{ClOOCl} approximately on the same order of magnitude as parameters influencing the actinic flux (Fig. 17). The presence of PSCs can have a particularly strong influence on the actinic flux and photochemistry. Wetzel et al. (2012) showed that, during the balloon flight on 24 January, the onset of ClO production from ClOOCl photolysis was delayed by about 45 min after sunrise in the stratosphere due to the occurrence of stratospheric clouds.

The thermal equilibrium constant K_{EQ} reported by Plenge et al. (2005) is supported by HALOX ClO and preliminary ClOOCl observations made before sunrise during two RECONCILE Geophysica flights (Stroh et al., 2012). The higher K_{EQ} currently recommended by JPL (Sander et al., 2011) would imply unrealistically high total stratospheric chlorine (Cl_y) around 5 ppb to be consistent with the ClO observations (Suminska-Ebersoldt et al., 2012).

Table 3 provides parameterisations for ClO_x kinetics that consistently and reasonably accurately simulate atmospheric observations of ClO_x partitioning and ozone loss. No

indication was found for alternative ClO_x chemistry that was suggested during the time of the RECONCILE proposal (e.g. Salawitch et al., 2009; Vogel et al., 2006).

4.3.3 Bromine chemistry

The bromine budget and photochemistry – important for the rate of the $\text{ClO}-\text{BrO}$ -cycle – was also investigated by aircraft and balloon observations during RECONCILE. First on 7 and 8 September 2009 the stratospheric photochemistry of bromine was probed for the high latitude fall circulation turn-over, where in the stratosphere major reactive bromine species are at daytime BrO and at night time BrONO_2 . By observing the transition of BrO into its night-time reservoir gas BrONO_2 at dusk and dawn with direct sun solar occultation and Limb spectroscopy and by radiative transfer and photochemical modelling, Kreycky et al. (2012) found that the ratio of the photolysis rate of BrONO_2 J_{BrONO_2} to the three body formation reaction ($\text{BrO} + \text{NO}_2 + \text{M}$) rate coefficient of BrONO_2 $k_{\text{BrO}+\text{NO}_2}$, i.e. $J_{\text{BrONO}_2}/k_{\text{BrO}+\text{NO}_2}$, is by a factor 1.69 ± 0.04 larger than recommended by the most recent JLP-2011 compilation (Sander et al., 2011). Major consequences of the findings for the stratosphere are that (1) recent assessments of total stratospheric bromine (Br_y) using the inorganic method for high stratospheric NO_x loadings may have overestimated the necessary correction for the BrO to Br_y ratio (for this flight, Br_y should be 1.4 ppt or 6.8% of the total lower using the revised $J_{\text{BrONO}_2}/k_{\text{BrO}+\text{NO}_2}$ ratio as compared to the JPL-2011 recommendations), and (2) a larger $J_{\text{BrONO}_2}/k_{\text{BrO}+\text{NO}_2}$ ratio may cause a small increase in the bromine-mediated ozone loss in the stratosphere (maximum -0.8% , see Fig. 18).

The trend of total stratospheric bromine (Fig. 1.21 in WMO, 2010) was also continued with data collected during RECONCILE, auxiliary field campaigns performed for example in 2008 in the tropics, at high latitudes in fall 2009, in winter 2010 and spring 2011, and lately during the EU project SHIVA (Stratospheric Ozone: Halogen Impacts in a Varying Atmosphere) in the Western Pacific in fall 2011. The collected data indicate that from its peak in the early 2000s, Br_y continues to decrease by about 10% per

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



decade and may now have reached about 19 ppt in 4 yr old air. Together with the finding of Kreygy et al. (2012) the collected field data, in particular during SHIVA, indicate that the contribution of the so-called very short-lived species (VSLS) inferred using the inorganic stratospheric method may be lower than stated in WMO (2010), i.e. 4 ± 2.5 ppt instead of 6 (range 3–8) ppt. The former estimate is also in better agreement with past and recent estimates of VSLS based on the organic bromine method, which indicate [VSLS] = 2.25 ± 0.25 ppt, where 90 % is due to contributions from CH_2Br_2 and CHBr_3 (e.g. Brinckmann et al., 2012).

4.3.4 Halogen source gases

A number of important long-lived brominated and chlorinated source gases (i.e. the three major CFCs and HCFCs, the two major halons, CCl_4 and CH_3CCl_3) were sampled by the Whole Air Sampler (WAS) onboard the Geophysica and subsequently analysed in the laboratory. Together with age-of-air measurements inferred from SF_6 , the results have been used for a re-evaluation of the stratospheric lifetimes of these compounds as well as their fractional release (i.e. the fraction that has released its halogens) and Ozone Depletion Potentials (ODPs), all of which are policy-relevant quantities. As outlined in Laube et al. (2012b) there are indications for considerably smaller fractional release for some of these gases. Alongside with possibly longer lifetimes this would not only lead to smaller ODPs in many cases, but also has the potential to significantly delay the recovery of the ozone layer. WAS measurements of other climate-relevant trace gases have improved the knowledge of their atmospheric distributions (e.g. Laube et al., 2010b, 2012a; Oram et al., 2012).

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.4 Studies of the Antarctic ozone hole

4.4.1 Minimum ozone mixing ratios

Although the main focus of RECONCILE is on Arctic ozone loss, the same processes are at play in the Antarctic, and there is still a lot to be learned from studies looking at the much more severe and annually reproducible Antarctic ozone hole. An investigation was carried out with CLaMS to explain the extremely low ozone mixing ratios below about 10 ppb observed by ozone sondes in Antarctic spring (Solomon et al., 2005). While in principle the mechanisms of ozone depletion are largely known, it was not clear what controls the ozone and chlorine chemistry causing such low ozone concentrations and what determines the lowest ozone values that can be reached. The ozone observations since the 1990s show a very similar behaviour from year to year, that is a minimum value changing with the time of year. This lower envelope of the observations reaches a minimum in late September and increases again in October and November. The complex interaction between gas-phase and heterogeneous chemistry under these conditions is described in detail by Grooß et al. (2011). To correctly simulate the behaviour of the ozone observations, it is necessary to include the oxidation chains of CO and CH₄, especially the reaction of the methyl peroxy radical CH₃O₂ with ClO (Crutzen et al., 1992) is important here.

4.4.2 Ozone trends

A long-term assessment of ozone loss and ozone trends in Antarctica has recently been published by Kuttippurath et al. (2012b). It reveals a slow but significant increase of the September–November average column ozone since the late 1990s, indicating that ozone recovery may have indeed started in the Antarctic.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Progress in global chemistry climate modelling

5.1 Amendments to the model

Based on the results described in Sect. 4, the following updates were implemented in the LMDZ-Reprobus CCM:

- A new PSC scheme is used with an improved parameterization of the growth of NAT particles. This new scheme includes a tracer keeping the memory of the time spent under the theoretical temperature of NAT formation, allowing a more realistic slow growth of NAT particles, rather than the instantaneous formation of large particles previously used in the so-called equilibrium scheme of the CCM. Maximum radius and number density of liquid and solid PSC particles are tuneable parameters.
- Based on the results described in Sect. 4.3.3., no new reactions driving catalytic ozone loss were included. ClO_x parameterisations from Table 5 were adopted as opposed to the reference model that uses parameterisations from Sander et al. (2011).

In addition to LMDZ, scientific findings from the RECONCILE project have been incorporated in the Whole Atmosphere Community Climate Model (WACCM). PSC processes have been improved in WACCM to give a better representation of mixed PSCs and the formation threshold for ice PSCs has been modified to agree with observations (Wegner et al., 2012b).

5.2 Model evaluation

Before carrying out long-term CCM simulations, the new PSC scheme developed during the project was tested with the CTM version of REPROBUS driven by ECMWF analysis for the Arctic winter 2009/10. The results obtained with this new microphysical parameterisation show a clearly improved HNO_3 distribution when compared to Aura

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MLS measurements, with less denitrification than with the reference model. The impact of this effect on ozone and climate over long timescales now needs to be investigated with the CCM version.

The scanning algorithms necessary for data analysis on global scales and the de-trended fluctuation analysis (DFA) intended to reveal the correlation properties of total column ozone (TO) time series were implemented successfully and the performance was cross-checked by reproducing the earlier results on satellite measurements (Kiss et al., 2007). While the original analysis used mostly the records of two instruments (Nimbus-7 and Earth Probe), the longer and continuous database (daily TO time series between 1978 and 2011) compiled by Bodeker Scientific (<http://www.bodekerscientific.com/>) was exploited in the present validation (Struthers et al., 2009). DFA runs on the CCMVal-2 outputs from the LMDZrepro reference simulation (1960–2006) indicate that the two-point correlation properties of model data are almost fully consistent with the measurements (Fig. 19a, b).

As for the spectral analysis, various regions exhibit different behaviours. Three significant peaks show up characterizing a half-year, annual and quasi-biennial oscillations (QBO) with different amplitudes. In order to estimate the importance of these modes, the spectral weight of the peaks were determined by integrating the area under the given peak and normalizing by the total power (total area of the frequency spectrum). As an example, a comparison of the spectral weight distribution of the semi-annual peak for the empirical and simulated data is shown in Fig. 19c, d. Interesting dynamical features in the simulations are the apparent shift and increased amplitude at the equatorial Pacific region, and a semi-annual oscillation above the Tibetan Plateau missing from the observations.

The QBO along the Equator is apparent in empirical TO data. However it is well known that numerical models can hardly reproduce this dynamical mode at the usual spatial and temporal resolutions. In an extended analysis we compared simulated time series of a CCM model, which is able to reproduce QBO (the Japanese Meteorological

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Research Institute model, MRI, in the CCMVal-2 data bank), but we could easily detect other anomalies in the correlation and spectral properties in the MRI time series.

6 Arctic ozone depletion in 2011

In terms of geographical extent and persistence of PSC conditions the winter 2010/11 was one of the coldest Arctic winters on record (Fig. 7), and measureable ozone loss rates persisted longer into spring than in any previous winter and exceeded the maximum rates previously measured in the Arctic. Based on Aura Satellite observations (OMI and MLS) as well as results from the 2010/11 match campaign, Manney et al. (2011) have shown that the vertical loss profile is mostly within the range of Antarctic ozone losses and far outside the range of previous ozone loss in the Arctic (Fig. 20). More than 80% of ozone was lost at the surface subsiding to 460 K by late March. Manney et al. (2011) also provided an in depth investigation of the processes leading to the record ozone loss, highlighting the remarkable degrees of chlorine activation and denitrification. In addition, they presented a comprehensive comparison to the winter 1996/97, which was among the coldest previous Arctic winters. Kuttippurath et al. (2012a) also discussed the dynamical situation in both winters and extended the analysis of Manney et al. (2011) by describing the contribution of NO_x cycle at higher altitudes.

Balis et al. (2011) show that dynamics and in particular weak transport from lower latitudes were important in setting the scene for the record low Arctic ozone in 2011. The respective roles of transport and chemical loss during 2010 and 2011 in simulations with the University of Oslo CTM (Sovde et al., 2011) are compared in Fig. 21. The upper panel shows that in 2010, ozone is replenished more efficiently at high latitudes than in 2011, which results from both isentropic mixing and from transport by the Brewer–Dobson circulation. This indicates that the lack of transport into the high latitudes played an important role in the ozone deficit in 2011. The second panel compares the large relative difference in transport to high latitudes in 2010 vs. 2011 (dotted

line) to the absolute contribution of heterogeneous chemistry to ozone loss in 2011 (full line, signed reversed). While the chemical loss in 2011 was much higher than in 2010, the relative weakness of transport into the very stable and narrow vortex was a stronger contributor to the column ozone deficit poleward of 60° N.

5 Von Hobe et al. (2012a) carried out an analysis of a 30-yr record of surface UV data products from satellite measurements and could show that for Arctic winters with strong ozone depletion including 2011, increases in surface UV radiation did not reach proportions like in the Antarctic. The enhancements were most significant at high latitudes (> 60° N) in March and early April, when absolute UV levels are still moderate. At
10 lower latitudes, where UV increases pose a more serious threat to human health and ecosystems, low column ozone due to transport from the subtropics has a stronger effect on surface UV than spring Arctic ozone loss.

7 Conclusions

RECONCILE has substantially advanced our understanding of the processes destroying ozone in the polar vortex. Three particular achievements are (1) the much improved elucidation of the NAT nucleation process and its impact on denitrification and thus ozone loss in the polar vortex, (2) the gathering of substantial evidence for the importance of heterogeneous chlorine activation on cold binary aerosol, and (3) the reconciliation of kinetic parameters governing ClO_x catalysed polar ozone loss. It should
15 be mentioned that many other substantial contributions besides RECONCILE have been made in the past few years, e.g. a recent SPARC initiative (Kurylo and Sinnhuber, 2009) and related laboratory studies to investigate ClOOCl photolysis (Lien et al., 2009; Papanastasiou et al., 2009; von Hobe et al., 2009). The representation of polar ozone depletion and related processes in global models has also been improved
20 (WMO, 2010, and references therein).

25 But besides helping to reach closure with respect to some existing uncertainties, some of the RECONCILE results also raised new questions. The observation of the

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

coldest Arctic winter on record in 2010/11 prompting an unprecedented degree of Arctic ozone loss (Manney et al., 2011) and related increases in UV radiation (Bernhard et al., 2012; von Hobe et al., 2012a), and the lack of prognostic tools that could have predicted such an event, made it evident that efforts to understand exactly what drives the stability and interannual variability of the Arctic polar vortex and the development of predictive capabilities are imperative. The finding of a large number of non-volatile particles with sometimes surprising chemical composition inside the Arctic vortex challenges some of our presumptions on how material, and especially aerosols, are transported to and distributed through the stratosphere. This is particularly important if we want to understand and model how climate change might affect the composition of the stratosphere in the future.

These questions concern the preconditions for Arctic ozone depletion rather than the actual ozone destruction process. The answers lie, to a large extent, in the tropics. The tropical tropopause is the main “gate to the stratosphere” (e.g. Fueglistaler et al., 2009) for trace gases and particles, which are then transported to higher latitudes via the Brewer–Dobson (BD) circulation. Changes in the chemical composition of the stratosphere related to climate change are likely, given the impact of climate on tropospheric sources and sinks as well as chemical processing of gases and particles, on convective transport into the tropical tropopause region, and on transport through the stratosphere via the BD. For example, it has been shown by Zhou et al. (2012) that the upwelling branch of the BD circulation is strongly influenced by tropical and subtropical wave activity. Also the turnaround latitudes of the BD circulation are likely governed by an interplay of wave activity at midlatitudes and the tropics/subtropics. Since tropical and subtropical wave activity is expected to increase in a changing climate, this is expected to have an important strengthening influence on the BD circulation (Butchart and Scaife, 2001; Fomichev et al., 2007) with possible influences on ozone depletion at high latitudes. Possibly even more important for ozone loss in the Arctic than potential changes in chemical composition, Naoe and Shibata (2012) pointed out that climate change may lead to changes in the Holton–Tan-mechanism that links the stability of

the polar vortex to the QBO (Garfinkel et al., 2012; Holton and Tan, 1980). Future investigations that address these questions will be needed to complement the work done in RECONCILE and further advance our understanding of Arctic ozone depletion.

5 The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

Acknowledgements. RECONCILE is funded by the European Commission under the grant number RECONCILE-226365-FP7-ENV-2008-1. This funding made the science described in this paper possible and is greatly appreciated. We are also grateful to the EC scientific officers Claus Brüning and Franz Immler for their support. We also acknowledge important contributions from institutional funding and thank the involved scientists and administrative and technical staff at the partner institutions who supported the project work in many ways. We thank MDB and in particular the M55-Geophysica pilots Oleg Shepetkov and Oleg Kononenko and the ground crew led by G. Belyaev for their fantastic work during the aircraft campaign. For invaluable support with logistics, flight permissions, and many other aspects of the aircraft campaign, we thank Rolf Maser and Harald Franke from ENVISCOPE, Heinz Finkenzeller from FINKCAS, Kurt Mäki and the Kiruna airport crew, and the Longyearbyen airport crew. Support for M. C. Pitts is provided under the auspices of the NASA CALIPSO/CloudSat Science Team. Support for L. R. Poole is provided under NASA contract NNL11AA10D. Work at the Jet Propulsion Laboratory, California Institute of Technology, was done under contract with the National Aeronautics and Space Administration. The Atmospheric Chemistry Experiment (ACE), also known as SCISAT, is a Canadian-led mission mainly supported by the Canadian Space Agency. Work at the University of East Anglia was made possible through funding from the UK Natural Environment Research Council (Research fellowships NE/F015585/1 & NE/I021918/1). The atmospheric sounding campaign at FMI Sodankylä was supported by the EU through the Lapland Atmosphere-Biosphere Facility (LAPBIAT2) project. The Geophysica flight on 10 March 2010 was funded by ESA under the PremierEx project. The PremierEx project also funded two Geophysica flights in autumn 2009 from Oberpfaffenhofen, Germany, that provided an invaluable opportunity for instrument testing prior to the Kiruna activities. We thank Bodeker Scientific for the total column ozone data and ECMWF for providing forecast, analysis and reanalysis data.

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Some of these data were available through the special project “Effect of non-hydrostatic gravity waves on the stratosphere above Scandinavia” by one of the authors (A.D.).

References

- 5 Abbatt, J. P. D. and Molina, M. J.: Heterogeneous interactions of ClONO₂ and HCl on nitric acid trihydrate at 202 K, *J. Phys. Chem.*, 96, 7674–7679, 1992.
- Allaart, M., van Weele, M., Fortuin, P., and Kelder, H.: An empirical model to predict the UV-index based on solar zenith angles and total ozone, *Meteorol. Appl.*, 11, 59–65, 2004.
- Anderson, J. G. and Toon, O. B.: Airborne Arctic Stratospheric Expedition .2. an overview, *Geophys. Res. Lett.*, 20, 2499–2502, 1993.
- 10 Anderson, J. G., Brune, W. H., and Proffitt, M. H.: Ozone destruction by chlorine radicals within the Antarctic vortex – the spatial and temporal evolution of ClO–O₃ anticorrelation based on in situ ER-2 data, *J. Geophys. Res.*, 94, 11465–11479, 1989.
- Anderson, J. G., Wilmouth, D. M., Smith, J. B., and Sayres, D. S.: UV dosage levels in summer: increased risk of ozone loss from convectively injected water vapor, *Science*, 337, 835–839, 2012.
- 15 Austin, J., Struthers, H., Scinocca, J., Plummer, D. A., Akiyoshi, H., Baumgaertner, A. J. G., Bekki, S., Bodeker, G. E., Braesicke, P., Brühl, C., Butchart, N., Chipperfield, M. P., Cugnet, D., Dameris, M., Dhomse, S., Frith, S., Garny, H., Gettelman, A., Hardiman, S. C., Jöckel, P., Kinnison, D., Kubin, A., Lamarque, J. F., Langematz, U., Mancini, E., Marchand, M., Michou, M., Morgenstern, O., Nakamura, T., Nielsen, J. E., Pitari, G., Pyle, J., Rozanov, E., Shepherd, T. G., Shibata, K., Smale, D., Teyssède, H., and Yamashita, Y.: The decline and recovery of total column ozone using a multi-model time series analysis, *J. Geophys. Res.*, 115, D00M10, doi:10.1029/2010JD013857, 2010.
- 20 Ayarzagüena, B., Langematz, U., and Serrano, E.: Tropospheric forcing of the stratosphere: a comparative study of the two different major stratospheric warmings in 2009 and 2010, *J. Geophys. Res.*, 116, D18114, doi:10.1029/2010JD015023, 2011.
- 25 Balis, D., Isaksen, I. S. A., Zerefos, C., Zyrichidou, I., Eleftheratos, K., Tourpali, K., Bojkov, R., Rognerud, B., Stordal, F., Søvde, O. A., and Orsolini, Y.: Observed and modelled record ozone decline over the Arctic during winter/spring 2011, *Geophys. Res. Lett.*, 38, L23801, doi:10.1029/2011gl049259, 2011.
- 30

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Baumgardner, D., Jonsson, H., Dawson, W., O'Connor, D., and Newton, R.: The cloud, aerosol and precipitation spectrometer: a new instrument for cloud investigations, *Atmos. Res.*, 59, 251–264, 2001.
- Bernath, P. F., McElroy, C. T., Abrams, M. C., Boone, C. D., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P. F., Colin, R., DeCola, P., DeMazière, M., Drummond, J. R., Dufour, D., Evans, W. F. J., Fast, H., Fussen, D., Gilbert, K., Jennings, D. E., Llewellyn, E. J., Lowe, R. P., Mahieu, E., McConnell, J. C., McHugh, M., McLeod, S. D., Michaud, R., Midwinter, C., Nassar, R., Nichitui, F., Nowlan, C., Rinsland, C. P., Rochon, Y. J., Rowlands, N., Semeniuk, K., Simon, P., Skelto, R., Sloan, J. J., Soucy, M. A., Strong, K., Tremblay, P., Turnbull, D., Walker, K. A., Walkty, I., Wardle, D. A., Wehrle, V., Zander, R., and Zou, J.: Atmospheric Chemistry Experiment (ACE): mission overview, *Geophys. Res. Lett.*, 32, 5 pp., 2005.
- Bernhard, G., Manney, G., Fioletov, V., Grooß, J.-U., Heikkilä, A., Johnsen, B., Koskela, T., Lakkala, K., Müller, R., Myhre, C. L., and Rex, M.: Arctic ozone and UV radiation, *B. Am. Meteorol. Soc.*, 93, 129–132, doi:10.1175/2012BAMSStateoftheClimate.1, 2012.
- Biele, J., Tsias, A., Luo, B. P., Carslaw, K. S., Neuber, R., Beyerle, G., and Peter, T.: Nonequilibrium coexistence of solid and liquid particles in Arctic stratospheric clouds, *J. Geophys. Res.*, 106, 22991–23007, 2001.
- Biermann, U. M., Presper, T., Koop, T., Mossinger, J., Crutzen, P. J., and Peter, T.: The unsuitability of meteoritic and other nuclei for polar stratospheric cloud freezing, *Geophys. Res. Lett.*, 23, 1693–1696, 1996.
- Bigg, E. K., Ono, A., and Thompson, W. J.: Aerosols at altitudes between 20 and 37 km, *Tellus*, 22, 550–563, 1970.
- Brabec, M., Wienhold, F. G., Luo, B. P., Vömel, H., Immler, F., Steiner, P., Hausammann, E., Weers, U., and Peter, T.: Particle backscatter and relative humidity measured across cirrus clouds and comparison with microphysical cirrus modelling, *Atmos. Chem. Phys.*, 12, 9135–9148, doi:10.5194/acp-12-9135-2012, 2012.
- Brasseur, G. P.: Creating knowledge from the confrontation of observations and models: the case of stratospheric ozone, in: *Climate Variability and Extremes During the Past 100 Years*, edited by: Brännimann, S., et al., Springer, 303–316, 2008.
- Brinckmann, S., Engel, A., Bönisch, H., Quack, B., and Atlas, E.: Short-lived brominated hydrocarbons – observations in the source regions and the tropical tropopause layer, *Atmos. Chem. Phys.*, 12, 1213–1228, doi:10.5194/acp-12-1213-2012, 2012.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Brooks, S. D., Baumgardner, D., Gandrud, B., Dye, J. E., Northway, M. J., Fahey, D. W., Bui, T. P., Toon, O. B., and Tolbert, M. A.: Measurements of large stratospheric particles in the Arctic polar vortex, *J. Geophys. Res.*, 108, 4652–4662, 2003.
- 5 Buontempo, C., Cairo, F., Di Donfrancesco, G., Morbidini, R., Viterbini, M., and Adriani, A.: Optical measurements of atmospheric particles from airborne platforms: in situ and remote sensing instruments for balloons and aircrafts, *Ann. Geophys.*, 49, 57–64, 2006, <http://www.ann-geophys.net/49/57/2006/>.
- 10 Butchart, N. and Scaife, A. A.: Removal of chlorofluorocarbons by increased mass exchange between the stratosphere and troposphere in a changing climate, *Nature*, 410, 799–802, 2001.
- Cairo, F., Pommereau, J. P., Law, K. S., Schlager, H., Garnier, A., Fierli, F., Ern, M., Streibel, M., Arabas, S., Borrmann, S., Berthelie, J. J., Blom, C., Christensen, T., D'Amato, F., Di Donfrancesco, G., Deshler, T., Diedhiou, A., Durry, G., Engelsen, O., Goutail, F., Harris, N. R. P., Kerstel, E. R. T., Khaykin, S., Konopka, P., Kylling, A., Larsen, N., Lebel, T., Liu, X., MacKenzie, A. R., Nielsen, J., Oulanowski, A., Parker, D. J., Pelon, J., Polcher, J., Pyle, J. A., Ravegnani, F., Rivière, E. D., Robinson, A. D., Röckmann, T., Schiller, C., Simões, F., Stefanutti, L., Stroh, F., Some, L., Siegmund, P., Sitnikov, N., Vernier, J. P., Volk, C. M., Voigt, C., von Hobe, M., Viciani, S., and Yushkov, V.: An introduction to the SCOUT-AMMA stratospheric aircraft, balloons and sondes campaign in West Africa, August 2006: rationale and roadmap, *Atmos. Chem. Phys.*, 10, 2237–2256, doi:10.5194/acp-10-2237-2010, 2010.
- 20 Carlsaw, K. S. and Peter, T.: Uncertainties in reactive uptake coefficients for solid stratospheric particles, 1. Surface chemistry, *Geophys. Res. Lett.*, 24, 1743–1746, 1997.
- Carlsaw, K. S., Peter, T., and Müller, R.: Uncertainties in reactive uptake coefficients for solid stratospheric particles, 2. Effect on ozone depletion, *Geophys. Res. Lett.*, 24, 1747–1750, 1997.
- 25 Carlsaw, K. S., Wirth, M., Tsias, A., Luo, B. P., Dörnbrack, A., Leutbecher, M., Volkert, H., Renger, W., Bacmeister, J. T., Reimer, E., and Peter, T.: Increased stratospheric ozone depletion due to mountain-induced atmospheric waves, *Nature*, 391, 675–678, 1998.
- Chen, H. Y., Lien, C. Y., Lin, W. Y., Lee, Y. T., and Lin, J. J.: UV absorption cross sections of ClOOCl are consistent with ozone degradation models, *Science*, 324, 781–784, 2009.
- 30 Chubachi, S. and Kajiwara, R.: Total ozone variations at Syowa, Antarctica, *Geophys. Res. Lett.*, 13, 1197–1198, 1986.

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Cicerone, R. J., Stolarski, R. S., and Walters, S.: Stratospheric ozone destruction by man-made chlorofluoromethanes, *Science*, 185, 1165–1167, 1974.
- Crutzen, P. J.: Ozone production rates in an oxygen-hydrogen-nitrogen oxide atmosphere, *J. Geophys. Res.*, 76, 7311–7327, 1971.
- 5 Crutzen, P. J.: Estimates of possible future ozone reductions from continued use of fluorochloro-methanes (CF_2Cl_2 , CFCl_3), *Geophys. Res. Lett.*, 1, 205–208, 1974.
- Crutzen, P. J.: Upper limits on atmospheric ozone reductions following increased application of fixed nitrogen to soil, *Geophys. Res. Lett.*, 3, 169–172, 1976.
- Crutzen, P. J.: Albedo enhancement by stratospheric sulfur injections: a contribution to resolve a policy dilemma?, *Climatic Change*, 77, 211–219, 2006.
- 10 Crutzen, P. J., Müller, R., Brühl, C., and Peter, T.: On the potential importance of the gas-phase reaction $\text{CH}_3\text{O}_2 + \text{ClO} \rightarrow \text{ClOO} + \text{CH}_3\text{O}$ and the heterogeneous reaction $\text{HOCl} + \text{HCl} \rightarrow \text{H}_2\text{O} + \text{Cl}_2$ in ozone hole chemistry, *Geophys. Res. Lett.*, 19, 1113–1116, 1992.
- 15 Curtius, J., Weigel, R., Vössing, H.-J., Wernli, H., Werner, A., Volk, C.-M., Konopka, P., Krebsbach, M., Schiller, C., Roiger, A., Schlager, H., Dreiling, V., and Borrmann, S.: Observations of meteoric material and implications for aerosol nucleation in the winter Arctic lower stratosphere derived from in situ particle measurements, *Atmos. Chem. Phys.*, 5, 3053–3069, doi:10.5194/acp-5-3053-2005, 2005.
- 20 Dameris, M. and Baldwin, M. P.: Impact of climate change on the stratospheric ozone layer, in: *Stratospheric Ozone Depletion and Climate Change*, edited by: Müller, R., RSC Publishing, Cambridge, 214–252, 2012.
- Dameris, M., Wirth, M., Renger, W., and Grewe, V.: Definition of the polar vortex edge by LIDAR data of the stratospheric aerosol: a comparison with values of potential vorticity, *Beitr. Phys. Atmos.*, 68, 113–119, 1995.
- 25 Dameris, M., Peter, T., Schmidt, U., and Zellner, R.: The ozone hole and its causes, *Chem. Unserer Zeit*, 41, 152–168, 2007.
- de Lange, A., Birk, M., de Lange, G., Friedl-Vallon, F., Kiselev, O., Koshelets, V., Maucher, G., Oelhaf, H., Selig, A., Vogt, P., Wagner, G., and Landgraf, J.: HCl and ClO in activated Arctic air; first retrieved vertical profiles from TELIS submillimetre limb spectra, *Atmos. Meas. Tech.*, 5, 487–500, doi:10.5194/amt-5-487-2012, 2012.
- 30 de Reus, M., Borrmann, S., Bansemmer, A., Heymsfield, A. J., Weigel, R., Schiller, C., Mitev, V., Frey, W., Kunkel, D., Kürten, A., Curtius, J., Sitnikov, N. M., Ulanovsky, A., and Ravegnani, F.:

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Evidence for ice particles in the tropical stratosphere from in-situ measurements, *Atmos. Chem. Phys.*, 9, 6775–6792, doi:10.5194/acp-9-6775-2009, 2009.
- Dörnbrack, A., Leutbecher, M., Volkert, H., and Wirth, M.: Mesoscale forecasts of stratospheric mountain waves, *Meteorol. Appl.*, 5, 117–126, 1998.
- 5 Dörnbrack, A., Leutbecher, M., Kivi, R., and Kyrö, E.: Mountain wave induced record low stratospheric temperatures above Northern Scandinavia, *Tellus A*, 51, 951–963, 1999.
- Dörnbrack, A., Pitts, M. C., Poole, L. R., Orsolini, Y. J., Nishii, K., and Nakamura, H.: The 2009–2010 Arctic stratospheric winter – general evolution, mountain waves and predictability of an operational weather forecast model, *Atmos. Chem. Phys.*, 12, 3659–3675, doi:10.5194/acp-12-3659-2012, 2012.
- 10 Drdla, K.: Temperature thresholds for polar stratospheric ozone loss, AGU Fall 2005 Meeting, San Francisco, 5–9 December 2005, A31D-03, 2005.
- Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the polar stratosphere, *Ann. Geophys.*, 30, 1055–1073, doi:10.5194/angeo-30-1055-2012, 2012.
- 15 Drdla, K., Schoeberl, M. R., and Browell, E. V.: Microphysical modeling of the 1999–2000 Arctic winter: 1. Polar stratospheric clouds, denitrification, and dehydration, *J. Geophys. Res.*, 108, 8312–8332, 2003.
- Dye, J. E., Baumgardner, D., Gandrud, B. W., Kawa, S. R., Kelly, K. K., Loewenstein, M., Ferry, G. V., Chan, K. R., and Gary, B. L.: Particle size distributions in Arctic polar stratospheric clouds, growth and freezing of sulfuric acid droplets, and implications for cloud formation, *J. Geophys. Res.*, 97, 8015–8034, 1992.
- 20 Emde, C. and Mayer, B.: Simulation of solar radiation during a total eclipse: a challenge for radiative transfer, *Atmos. Chem. Phys.*, 7, 2259–2270, doi:10.5194/acp-7-2259-2007, 2007.
- Emde, C., Buras, R., and Mayer, B.: ALIS: an efficient method to compute high spectral resolution polarized solar radiances using the Monte Carlo approach, *J. Quant. Spectrosc. Ra.*, 25 112, 1622–1631, doi:10.1016/j.jqsrt.2011.03.018, 2011.
- Engel, A., Moebius, T., Boenisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nat. Geosci.*, 2, 28–31, 2009.
- 30 Engel, I., Luo, B., Khaykin, S., Wienhold, F. G., Kivi, R., Vömel, H., Lykov, A., and Peter, T.: Arctic stratospheric dehydration, Part II: a microphysical modeling study, in preparation, 2012a.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Engel, I., Luo, B. P., Hoyle, C. R., Pitts, M. C., Poole, L. R., Groöb, J.-U., and Peter, T.: Heterogeneous formation of polar stratospheric clouds, Part II: Nucleation of synoptic scale ice clouds, in preparation, 2012b.

Engel, I., Pitts, M. C., Luo, B., Hoyle, C., Zobrist, B., Jacot, L., Poole, L. R., Groöb, J. U., Weigel, R., Borrmann, S., Ebert, M., Duprat, J., and Peter, T.: Unprecedented evidence for large scale heterogeneous nucleation of polar stratospheric clouds, likely by nanometer-sized meteoritic particles, AGU Fall Meeting, San Francisco, USA, San Francisco, 3–7 December 2012, A53K-0309, 2012c.

Eskes, H. J., van der A, R. J., Brinksma, E. J., Veefkind, J. P., de Haan, J. F., and Valks, P. J. M.: Retrieval and validation of ozone columns derived from measurements of SCIAMACHY on Envisat, *Atmos. Chem. Phys. Discuss.*, 5, 4429–4475, doi:10.5194/acpd-5-4429-2005, 2005.

Eyring, V., Cionni, I., Bodeker, G. E., Charlton-Perez, A. J., Kinnison, D. E., Scinocca, J. F., Waugh, D. W., Akiyoshi, H., Bekki, S., Chipperfield, M. P., Dameris, M., Dhomse, S., Frith, S. M., Garny, H., Gettelman, A., Kubin, A., Langematz, U., Mancini, E., Marchand, M., Nakamura, T., Oman, L. D., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Shepherd, T. G., Shibata, K., Tian, W., Braesicke, P., Hardiman, S. C., Lamarque, J. F., Morgenstern, O., Pyle, J. A., Smale, D., and Yamashita, Y.: Multi-model assessment of stratospheric ozone return dates and ozone recovery in CCMVal-2 models, *Atmos. Chem. Phys.*, 10, 9451–9472, doi:10.5194/acp-10-9451-2010, 2010.

Fahey, D. W., Gao, R. S., Carslaw, K. S., Kettleborough, J., Popp, P. J., Northway, M. J., Holecek, J. C., Ciciora, S. C., McLaughlin, R. J., Thompson, T. L., Winkler, R. H., Baumgardner, D. G., Gandrud, B., Wennberg, P. O., Dhaniyala, S., McKinney, K., Peter, T., Salawitch, R. J., Bui, T. P., Elkins, J. W., Webster, C. R., Atlas, E. L., Jost, H., Wilson, J. C., Herman, R. L., Kleinbohl, A., and von Konig, M.: The detection of large HNO₃-containing particles in the winter arctic stratosphere, *Science*, 291, 1026–1031, 2001.

Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica reveal seasonal ClO_x/NO_x interaction, *Nature*, 315, 207–210, 1985.

Fomichev, V. I., Jonsson, A. I., De Grandpre, J., Beagley, S. R., McLandress, C., Semeniuk, K., and Shepherd, T. G.: Response of the middle atmosphere to CO₂ doubling: results from the Canadian Middle Atmosphere Model, *J. Climate*, 20, 1121–1144, 2007.

Frieler, K., Rex, M., Salawitch, R. J., Canty, T., Streibel, M., Stimpfle, R. M., Pfeilsticker, K., Dorf, M., Weisenstein, D. K., and Godin-Beekmann, S.: Toward a better

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



quantitative understanding of polar stratospheric ozone loss, *Geophys. Res. Lett.*, 33, doi:10.1029/2005GL025466, L10812, 2006.

Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Fuller, R. A., Marcy, T. P., Popp, P. J., Gao, R. S., Fahey, D. W., Jucks, K. W., Stachnik, R. A.,
 5 Toon, G. C., Christensen, L. E., Webster, C. R., Bernath, P. F., Boone, C. D., Walker, K. A., Pumphrey, H. C., Harwood, R. S., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder HCl measurements, *J. Geophys. Res.*, 113, D15S25, doi:10.1029/2007JD009025, 2008.

10 Fueglistaler, S., Luo, B. P., Voigt, C., Carslaw, K. S., and Peter, Th.: NAT-rock formation by mother clouds: a microphysical model study, *Atmos. Chem. Phys.*, 2, 93–98, doi:10.5194/acp-2-93-2002, 2002.

Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008rg000267, 2009.

15 Garfinkel, C. I., Shaw, T. A., Hartmann, D. L., and Waugh, D. W.: Does the Holton–Tan mechanism explain how the quasi-biennial oscillation modulates the Arctic polar vortex?, *J. Atmos. Sci.*, 69, 1713–1733, doi:10.1175/jas-d-11-0209.1, 2012.

Gernandt, H.: The vertical ozone distribution above the GDR research base, Antarctica in 1985, *Geophys. Res. Lett.*, 14, 84–86, 1987.

20 Gettelman, A., Hegglin, M. I., Son, S.-W., Kim, J., Fujiwara, M., Birner, T., Kremser, S., Rex, M., Añel, J. A., Akiyoshi, H., Austin, J., Bekki, S., Braesicke, P., Brühl, C., Butchart, N., Chipperfield, M., Dameris, M., Dhomse, S., Garny, H., Hardiman, S. C., Jöckel, P., Kinnison, D. E., Lamarque, J. F., Mancini, E., Marchand, M., Michou, M., Morgenstern, O., Pawson, S., Pitari, G., Plummer, D. A., Pyle, J. A., Rozanov, E., Scinocca, J., Shepherd, T. G., Shibata, K., Smale, D., Teyssère, H., and Tian, W.: Multimodel assessment of the upper troposphere and lower stratosphere: tropics and global trends, *J. Geophys. Res.*, 115, D00M08, doi:10.1029/2009JD013638, 2010.

25 Gillett, N. P. and Son, S.-W.: Impact of polar ozone loss on the troposphere, in: *Stratospheric Ozone Depletion and Climate Change*, edited by: Müller, R., RSC Publishing, Cambridge, 190–213, 2012.

30 Gillett, N. P., Akiyoshi, H., Bekki, S., Braesicke, P., Eyring, V., Garcia, R., Karpechko, A. Yu., McLinden, C. A., Morgenstern, O., Plummer, D. A., Pyle, J. A., Rozanov, E., Scinocca, J.,

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and Shibata, K.: Attribution of observed changes in stratospheric ozone and temperature, *Atmos. Chem. Phys.*, 11, 599–609, doi:10.5194/acp-11-599-2011, 2011.

Grooß, J.-U., Günther, G., Müller, R., Konopka, P., Bausch, S., Schlager, H., Voigt, C., Volk, C.M., and Toon, G. C.: Simulation of denitrification and ozone loss for the Arctic winter 2002/2003, *Atmos. Chem. Phys.*, 5, 1437–1448, doi:10.5194/acp-5-1437-2005, 2005.

Grooß, J.-U., Brauttsch, K., Pommrich, R., Solomon, S., and Müller, R.: Stratospheric ozone chemistry in the Antarctic: what determines the lowest ozone values reached and their recovery?, *Atmos. Chem. Phys.*, 11, 12217–12226, doi:10.5194/acp-11-12217-2011, 2011.

Günther, G., Müller, R., von Hobe, M., Stroh, F., Konopka, P., and Volk, C. M.: Quantification of transport across the boundary of the lower stratospheric vortex during Arctic winter 2002/2003, *Atmos. Chem. Phys.*, 8, 3655–3670, doi:10.5194/acp-8-3655-2008, 2008.

Hanson, D. R. and Ravishankara, A. R.: Reaction of ClONO₂ with HCl on NAT, NAD and frozen sulfuric acid and hydrolysis of N₂O₅ and ClONO₂ on frozen sulfuric acid, *J. Geophys. Res.*, 98, 22931–22936, 1993.

Hanson, D. R. and Ravishankara, A. R.: Reactive uptake of ClONO₂ onto sulfuric acid due to reaction with HCl and H₂O, *J. Phys. Chem.*, 98, 5728–5735, 1994.

Harvey, V. L., Pierce, R. B., and Hitchman, M. H.: A climatology of stratospheric polar vortices and anticyclones, *J. Geophys. Res.*, 107, 4442, doi:10.1029/2001JD001471, 2002.

Hegglin, M. I., Gettelman, A., Hoor, P., Krichevsky, R., Manney, G. L., Pan, L. L., Son, S. W., Stiller, G., Tilmes, S., Walker, K. A., Eyring, V., Shepherd, T. G., Waugh, D., Akiyoshi, H., Añel, J. A., Austin, J., Baumgaertner, A., Bekki, S., Braesicke, P., Brühl, C., Butchart, N., Chipperfield, M., Dameris, M., Dhomse, S., Frith, S., Garny, H., Hardiman, S. C., Jöckel, P., Kinnison, D. E., Lamarque, J. F., Mancini, E., Michou, M., Morgenstern, O., Nakamura, T., Olivié, D., Pawson, S., Pitari, G., Plummer, D. A., Pyle, J. A., Rozanov, E., Scinocca, J. F., Shibata, K., Smale, D., Teyssere, H., Tian, W., and Yamashita, Y.: Multimodel assessment of the upper troposphere and lower stratosphere: extratropics, *J. Geophys. Res.*, 115, D00M09, doi:10.1029/2010JD013884, 2010.

Holton, J. R. and Tan, H.-C.: The influence of the equatorial quasi-biennial oscillation on the global circulation at 50 mb, *J. Atmos. Sci.*, 37, 2200–2208, 1980.

Homan, C. D., Volk, C. M., Kuhn, A. C., Werner, A., Baehr, J., Viciani, S., Ulanovski, A., and Ravegnani, F.: Tracer measurements in the tropical tropopause layer during the AMMA/SCOUT-O3 aircraft campaign, *Atmos. Chem. Phys.*, 10, 3615–3627, doi:10.5194/acp-10-3615-2010, 2010.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Hösen, E., Volk, C. M., vom Scheidt, M., Wintel, J., Ulanovsky, A., Ravegnani, F., Grooß, J. U., Günther, G., and Walker, K. A.: Transport, mixing and ozone loss in the 2010 Arctic vortex region from in-situ tracer observations during RECONCILE, EGU General Assembly, Vienna, 2012.
- 5 Hourdin, F., Musat, I., Bony, S., Braconnot, P., Codron, F., Dufresne, J.-L., Fairhead, L., Filiberti, M.-A., Friedlingstein, P., Grandpeix, J.-Y., Krinner, G., LeVan, P., Li, Z.-X., and Lott, F.: The LMDZ4 general circulation model: climate performance and sensitivity to parametrized physics with emphasis on tropical convection, *Clim. Dynam.*, 27, 787–813, 2006.
- 10 Hourdin, F., Foujols, M.-A., Codron, F., Guemas, V., Dufresne, J.-L., Bony, S., Denvil, S., Guez, L., Lott, F., Ghattas, J., Braconnot, P., Marti, O., Meurdesoif, Y., and Bopp, L.: Impact of the LMDZ atmospheric grid configuration on the climate and sensitivity of the IPSL-CM5A coupled model, *Clim. Dynam.*, doi:10.1007/s00382-012-1411-3, 2012.
- 15 Hoyle, C. R., Engel, I., Luo, B. P., Pitts, M. C., Poole, L. R., Grooß, J.-U., and Peter, T.: Heterogeneous formation of polar stratospheric clouds, Part I: Nucleation of NAT clouds, 2012, in preparation.
- Isaksen, I. S. A., Zerefos, C., Wang, W.-C., Balis, D., Eleftheratos, K., Rognerud, B., Stordal, F., Berntsen, T. K., LaCasce, J. H., Søvde, O. A., Olivié, D., Orsolini, Y. J., Zyrichidou, I., Prather, M., and Tuinder, O.: Attribution of Arctic ozone loss in March 2011, *Geophys. Res. Lett.*, doi:10.1029/2012GL053876, in press, 2012.
- 20 Jackson, D. R.: Assimilation of EOSMLS ozone observations in the Met Office data assimilation system, *Q. J. Roy. Meteor. Soc.*, 133, 1771–1788, 2007.
- Jackson, D. R. and Orsolini, Y. J.: Estimation of Arctic ozone loss in winter 2004/05 based on assimilation of EOS MLS and SBUV/2 observations, *Q. J. Roy. Meteor. Soc.*, 134, 1833–1841, 2008.
- 25 Jin, B., Chen, I. C., Huang, W. T., Lien, C. Y., Guchhait, N., and Lin, J. J.: Photodissociation cross section of ClOOCl at 330 nm, *J. Phys. Chem. A*, 114, 4791–4797, doi:10.1021/jp909374k, 2010.
- Johnston, H.: Reduction of stratospheric ozone by nitrogen oxide catalysts from supersonic transport exhaust, *Science*, 173, 517–522, 1971.
- 30 Jourdain, L., Bekki, S., Lott, F., and Lefèvre, F.: The coupled chemistry–climate model LMDZ-REPROBUS: description and evaluation of a transient simulation of the period 1980–1999, *Ann. Geophys.*, 26, 1391–1413, doi:10.5194/angeo-26-1391-2008, 2008.

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kaiser, J., Engel, A., Borchers, R., and Röckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N₂O isotope distribution, *Atmos. Chem. Phys.*, 6, 3535–3556, doi:10.5194/acp-6-3535-2006, 2006.
- Khaykin, S., Engel, I., Vömel, H., Luo, B., Wienhold, F., Peter, T., Kivi, R., Lykov, A., Yushkov, V., Santee, M., Spelten, N., Schiller, C., and Shur, G.: Arctic stratospheric dehydration, Part I: Unprecedented observations of vertical redistribution of water, in preparation, 2012.
- Khosrawi, F., Urban, J., Pitts, M. C., Voelger, P., Achtert, P., Kaphlanov, M., Santee, M. L., Manney, G. L., Murtagh, D., and Fricke, K.-H.: Denitrification and polar stratospheric cloud formation during the Arctic winter 2009/2010, *Atmos. Chem. Phys.*, 11, 8471–8487, doi:10.5194/acp-11-8471-2011, 2011.
- Kiss, P., Müller, R., and Janosi, I. M.: Long-range correlations of extrapolar total ozone are determined by the global atmospheric circulation, *Nonlin. Proc. Geophys.*, 14, 435–442, 2007.
- Konopka, P., Steinhorst, H. M., Groß, J.-U., Günther, G., Müller, R., Elkins, J. W., Jost, H. J., Richard, E., Schmidt, U., Toon, G., and McKenna, D. S.: Mixing and ozone loss in the 1999–2000 Arctic vortex: simulations with the 3-dimensional Chemical Lagrangian Model of the Stratosphere (CLaMS), *J. Geophys. Res.*, 109, D02315, doi:10.1029/2003JD003792, 2004.
- Koop, T. and Zobrist, B.: Parameterizations for ice nucleation in biological and atmospheric systems, *Phys. Chem. Chem. Phys.*, 11, 10839–10850, 2009.
- Koop, T., Biermann, U. M., Raber, W., Luo, B. P., Crutzen, P. J., and Peter, T.: Do stratospheric aerosol droplets freeze above the ice frost point?, *Geophys. Res. Lett.*, 22, 917–920, 1995.
- Kreycy, S., Camy-Peyret, C., Chipperfield, M. P., Dorf, M., Feng, W., Hossaini, R., Kritten, L., Werner, B., and Pfeilsticker, K.: Atmospheric test of the $J(\text{BrONO}_2)/k_{\text{BrO}+\text{NO}_2}$ ratio: implications for total stratospheric Br_y and bromine-mediated ozone loss, *Atmos. Chem. Phys. Discuss.*, 12, 27821–27845, doi:10.5194/acpd-12-27821-2012, 2012.
- Kurylo, M. J. and Sinnhuber, B.-M.: The role of halogen chemistry in polar stratospheric ozone depletion, SPARC, 2009.
- Kuttippurath, J. and Nikulin, G.: A comparative study of the major sudden stratospheric warmings in the Arctic winters 2003/2004–2009/2010, *Atmos. Chem. Phys.*, 12, 8115–8129, doi:10.5194/acp-12-8115-2012, 2012.
- Kuttippurath, J., Godin-Beekmann, S., Lefèvre, F., Nikulin, G., Santee, M. L., and Froidevaux, L.: Record-breaking ozone loss in the Arctic winter 2010/2011: comparison with 1996/1997, *Atmos. Chem. Phys.*, 12, 7073–7085, doi:10.5194/acp-12-7073-2012, 2012a.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Kuttippurath, J., Lefèvre, F., Pommereau, J.-P., Roscoe, H. K., Goutail, F., Pazmiño, A., and Shanklin, J. D.: Antarctic ozone loss in 1989–2010: evidence for ozone recovery?, *Atmos. Chem. Phys. Discuss.*, 12, 10775–10814, doi:10.5194/acpd-12-10775-2012, 2012b.
- 5 Lambert, A., Read, W. G., Livesey, N. J., Santee, M. L., Manney, G. L., Froidevaux, L., Wu, D. L., Schwartz, M. J., Pumphrey, H. C., Jimenez, C., Nedoluha, G. E., Cofield, R. E., Cuddy, D. T., Daffer, W. H., Drouin, B. J., Fuller, R. A., Jarnot, R. F., Knosp, B. W., Pickett, H. M., Perrun, C. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Waters, J. W., Jucks, K. W., Toon, G. C., Stachnik, R. A., Bernath, P. F., Boone, C. D., Walker, K. A., Urban, J., Murtagh, D., Elkins, J. W., and Atlas, E.: Validation of the Aura Microwave Limb
- 10 Sounder middle atmosphere water vapor and nitrous oxide measurements, *J. Geophys. Res.*, 112, D24S36, doi:10.1029/2007JD008724, 2007.
- Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Sturges, W. T., Braß, M., and Röckmann, T.: Fractional release factors of long-lived halogenated organic compounds in the tropical stratosphere, *Atmos. Chem. Phys.*, 10, 1093–1103, doi:10.5194/acp-10-1093-2010, 2010a.
- 15 Laube, J. C., Martinerie, P., Witrant, E., Blunier, T., Schwander, J., Brenninkmeijer, C. A. M., Schuck, T. J., Bolder, M., Röckmann, T., van der Veen, C., Bönisch, H., Engel, A., Mills, G. P., Newland, M. J., Oram, D. E., Reeves, C. E., and Sturges, W. T.: Accelerating growth of HFC-227ea (1,1,1,2,3,3,3-heptafluoropropane) in the atmosphere, *Atmos. Chem. Phys.*, 10, 5903–5910, doi:10.5194/acp-10-5903-2010, 2010b.
- 20 Laube, J. C., Hogan, C., Newland, M. J., Mani, F. S., Fraser, P. J., Brenninkmeijer, C. A. M., Martinerie, P., Oram, D. E., Röckmann, T., Schwander, J., Witrant, E., Mills, G. P., Reeves, C. E., and Sturges, W. T.: Distributions, long term trends and emissions of four perfluorocarbons in remote parts of the atmosphere and firn air, *Atmos. Chem. Phys.*, 12, 4081–4090, doi:10.5194/acp-12-4081-2012, 2012a.
- 25 Laube, J. C., Keil, A., Bönisch, H., Engel, A., Röckmann, T., Volk, C. M., and Sturges, W. T.: Observation-based assessment of stratospheric fractional release, lifetimes, and Ozone Depletion Potentials of ten important source gases, *Atmos. Chem. Phys. Discuss.*, 12, 28525–28557, doi:10.5194/acpd-12-28525-2012, 2012b.
- 30 Le Treut, H., Forichon, M., Boucher, O., and Li, Z.: Sulfate aerosol indirect effect and CO₂ greenhouse forcing: equilibrium response of the LMD GCM and associated cloud feedbacks, *J. Climate*, 11, 1673–1694, 1998.

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lefèvre, F., Brasseur, G. P., Folkins, I., Smith, A. K., and Simon, P.: Chemistry of the 1991–1992 stratospheric winter: three-dimensional model simulations, *J. Geophys. Res.*, 199, 8183–8195, 1994.
- Lefèvre, F., Figarol, F., Carslaw, K. S., and Peter, T.: The 1997 Arctic ozone depletion quantified from three-dimensional model simulations, *Geophys. Res. Lett.*, 25, 2425–2428, 1998.
- Li, F., Stolarski, R. S., and Newman, P. A.: Stratospheric ozone in the post-CFC era, *Atmos. Chem. Phys.*, 9, 2207–2213, doi:10.5194/acp-9-2207-2009, 2009.
- Lien, C. Y., Lin, W. Y., Chen, H. Y., Huang, W. T., Jin, B., Chen, I. C., and Lin, J. J.: Photodissociation cross sections of ClOOCl at 248.4 and 266 nm, *J. Chem. Phys.*, 131, 174301, doi:10.1063/1.3257682, 2009.
- Livesey, N. J., Read, W. G., Froidevaux, L., Lambert, A., Manney, G. L., Pumphrey, H. C., Santee, M. L., Schwartz, M. J., Wang, S., Cofeld, R. E., Cuddy, D. T., Fuller, R. A., Jarnot, R. F., Jiang, J. H., Knosp, B. W., Stek, P. C., Wagner, P. A., and Wu, D. L.: Version 3.3 Level 2 data quality and description document, Jet Propulsion Laboratory, Pasadena, California, USA, 2012.
- Luo, B. P., Voigt, C., Fueglistaler, S., and Peter, T.: Extreme NAT supersaturations in mountain wave ice PSCs: a clue to NAT formation, *J. Geophys. Res.*, 108, 4441, doi:10.1029/2002JD003104, 2003.
- Mahieu, E., Zander, R., Duchatelet, P., Hannigan, J. W., Coffey, M. T., Mikuteit, S., Hase, F., Blumenstock, T., Wiacek, A., Strong, K., Taylor, J. R., Mittermeier, R. L., Fast, H., Boone, C. D., McLeod, S. D., Walker, K. A., Bernath, P. F., and Rinsland, C. P.: Comparison between ACE-FTS and ground-based measurements of stratospheric HCl and ClONO₂ loadings at northern latitudes, *Geophys. Res. Lett.*, 32, L15S08, doi:10.1029/2005GL022396, 2005.
- Manney, G. L., Santee, M. L., Rex, M., Livesey, N. J., Pitts, M. C., Veeckind, P., Nash, E. R., Wohltmann, I., Lehmann, R., Froidevaux, L., Poole, L. R., Schoeberl, M. R., Haffner, D. P., Davies, J., Dorokhov, V., Gernandt, H., Johnson, B., Kivi, R., Kyro, E., Larsen, N., Levelt, P. F., Makshtas, A., McElroy, C. T., Nakajima, H., Concepcion Parrondo, M., Tarasick, D. W., von der Gathen, P., Walker, K. A., and Zinoviev, N. S.: Unprecedented Arctic ozone loss in 2011, *Nature*, 478, 469–475, doi:10.1038/nature10556 2011.
- Marchand, M., Keckhut, P., Lefebvre, S., Claud, C. C. D., Hauchecorne, A., Lefèvre, F., Lefebvre, M.-P., Jumelet, J., Lott, F., Hourdin, F., Thuillier, G., Poulain, V., Bossay, S., Lemennais, P., David, C., and Bekki, S.: Dynamical amplification of the stratospheric solar response simu-

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

lated with the chemistry–climate model LMDz-Reprobus, *J. Atmos. Sol.-Terr. Phys.*, 75–76, 147–160, doi:10.1016/j.jastp.2011.11.008, 2012.

Marculli, C., Gedamke, S., Peter, T., and Zobrist, B.: Efficiency of immersion mode ice nucleation on surrogates of mineral dust, *Atmos. Chem. Phys.*, 7, 5081–5091, doi:10.5194/acp-7-5081-2007, 2007.

Matthey, R., Cacciani, M., Fiocco, G., Martinez, A. A., Martucci, G., Mitev, V., Pace, G., and Stefanutti, L.: Observations of aerosol and clouds with the ABLE and MAL lidars during the mid-latitude and arctic ENVISAT validation campaigns, in: 16th ESA Symposium on European Rocket and Balloon Programmes and Related Research, Proceedings, ESA Special Publications, 579–584, St. Gallen, Switzerland, 2–5 June 2003, 2003.

Mayer, B.: Radiative transfer in the cloudy atmosphere, *Eur. Phys. J. Conf.*, 1, 75–99, doi:10.1140/epjconf/e2009-00912-1, 2009.

Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations – description and examples of use, *Atmos. Chem. Phys.*, 5, 1855–1877, doi:10.5194/acp-5-1855-2005, 2005.

McElroy, M. B., Salawitch, R. J., Wofsy, S. C., and Logan, J. A.: Reductions of Antarctic ozone due to synergistic interactions of chlorine and bromine, *Nature*, 321, 759–762, 1986.

McIntyre, M. E.: On the Antarctic ozone hole, *J. Atmos. Terr. Phys.*, 51, 29–43, 1989.

McKenna, D. S., Grooß, J. U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS) – 2. Formulation of chemistry scheme and initialization, *J. Geophys. Res.*, 107, 4256–4269, doi:10.1029/2000JD000113, 2002a.

McKenna, D. S., Konopka, P., Grooß, J. U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS) – 1. Formulation of advection and mixing, *J. Geophys. Res.*, 107, 4309–4323, 2002b.

McLandress, C. and Shepherd, T. G.: Impact of climate change on stratospheric sudden warmings as simulated by the Canadian Middle Atmosphere Model, *J. Climate*, 22, 5449–5463, doi:10.1175/2009JCLI3069.1, 2009.

Meilinger, S. K., Koop, T., Luo, B. P., Huthwelker, T., Carslaw, K. S., Krieger, U., Crutzen, P. J., and Peter, T.: Size-dependent stratospheric droplet composition in Lee wave temperature fluctuations and their potential role in PSC freezing, *Geophys. Res. Lett.*, 22, 3031–3034, 1995.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Molina, L. T. and Molina, M. J.: Production of Cl_2O_2 from the self-reaction of the ClO radical, *J. Phys. Chem.*, 91, 433–436, 1987.
- Molina, M. J. and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes – chlorine atomic-catalysed destruction of ozone, *Nature*, 249, 810–812, 1974.
- 5 Morgenstern, O., Braesicke, P., Hurwitz, M. M., O'Connor, F. M., Bushell, A. C., Johnson, C. E., and Pyle, J. A.: The world avoided by the Montreal Protocol, *Geophys. Res. Lett.*, 35, L16811, doi:10.1029/2008gl034590, 2008.
- Moyna, B. P., Oldfield, M. L., Goizel, A. S., Gerber, D., Siddans, R., Reburn, W. J., Matheson, D. N., Kerridge, B. J., De Maagt, P. J. I., Langen, J., and Klein, U.: MARSCHALS: air-borne simulator of a future space instrument to observe millimetre-wave limb emission from the upper troposphere and lower stratosphere, in: *Sensors, Systems, and Next-Generation Satellites X*, Proceedings of the Society of Photo-Optical Instrumentation Engineers (Spie), 75–86, 2006.
- 10 Müller, R.: A brief history of stratospheric ozone research, *Meteorol. Z.*, 18, 3–24, 2009.
- 15 Murtagh, D. P., Frisk, U., Merino, F., Ridal, M., Jonsson, A., Stegman, J., Witt, G., Eriksson, P., Jimenez, C., Megie, G., de la Noe, J., Ricaud, P., Baron, P., Pardo, J. R., Hauchcorne, A., Llewellyn, E. J., Degenstein, D. A., Gattinger, R. L., Lloyd, N. D., Evans, W. F. J., McDade, I. C., Haley, C. S., Sioris, C., von Savigny, C., Solheim, B. H., McConnell, J. C., Strong, K., Richardson, E. H., Leppelmeier, G. W., Kyrola, E., Auvinen, H., and Oikarinen, L.: An overview of the Odin atmospheric mission, *Can. J. Phys.*, 80, 309–319, 2002.
- 20 Naoe, H. and Shibata, K.: Future changes in the influence of the quasi-biennial oscillation on the northern polar vortex simulated with an MRI chemistry climate model, *J. Geophys. Res.*, 117, D03102, doi:10.1029/2011JD016255, 2012.
- Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective determination of the polar vortex using Ertel's potential vorticity, *J. Geophys. Res.*, 101, 9471–9478, 1996.
- 25 Newman, P. A., Harris, N. R. P., Adriani, A., Amanatidis, G. T., Anderson, J. G., Braathen, G. O., Brune, W. H., Carslaw, K. S., Craig, M. S., DeCola, P. L., Guirlet, M., Hipskind, R. S., Kurylo, M. J., Kullmann, H., Larsen, N., Megie, G. J., Pommereau, J. P., Poole, L. R., Schoeberl, M. R., Stroh, F., Toon, O. B., Trepte, C. R., and Van Roozendaal, M.: An overview of the SOLVE/THESEO 2000 campaign, *J. Geophys. Res.-Atmos.*, 107, 8259–8284, 2002.
- 30 Newman, P. A., Oman, L. D., Douglass, A. R., Fleming, E. L., Frith, S. M., Hurwitz, M. M., Kawa, S. R., Jackman, C. H., Krotkov, N. A., Nash, E. R., Nielsen, J. E., Pawson, S., Stolarski, R. S., and Velders, G. J. M.: What would have happened to the ozone layer if

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

chlorofluorocarbons (CFCs) had not been regulated?, *Atmos. Chem. Phys.*, 9, 2113–2128, doi:10.5194/acp-9-2113-2009, 2009.

Nickolaisen, S. L., Friedl, R. R., and Sander, S. P.: Kinetics and mechanism of the ClO + ClO reaction – pressure and temperature dependences of the bimolecular and termolecular channels and thermal-decomposition of chlorine peroxide, *J. Phys. Chem.*, 98, 155–169, 1994.

Nishii, K., Nakamura, H., and Orsolini, Y. J.: Geographical dependence observed in blocking high influence on the stratospheric variability through enhancement and suppression of upward planetary-wave propagation, *J. Climate*, 24, 6408–6423, doi:10.1175/JCLI-D-10-05021.1, 2011.

Northway, M. J., Gao, R. S., Popp, P. J., Holecek, J. C., Fahey, D. W., Carslaw, K. S., Tolbert, M. A., Lait, L. R., Dhaniyala, S., Flagan, R. C., Wennberg, P. O., Mahoney, M. J., Herman, R. L., Toon, G. C., and Bui, T. P.: An analysis of large HNO₃-containing particles sampled in the Arctic stratosphere during the winter of 1999/2000, *J. Geophys. Res.*, 107, 8298–8319, 2002.

Oram, D. E., Mani, F. S., Laube, J. C., Newland, M. J., Reeves, C. E., Sturges, W. T., Penkett, S. A., Brenninkmeijer, C. A. M., Röckmann, T., and Fraser, P. J.: Long-term tropospheric trend of octafluorocyclobutane (c-C₄F₈ or PFC-318), *Atmos. Chem. Phys.*, 12, 261–269, doi:10.5194/acp-12-261-2012, 2012.

Orsolini, Y. J., Karpechko, A. Y., and Nikulin, G.: Variability of the Northern Hemisphere polar stratospheric cloud potential: the role of North Pacific disturbances, *Q. J. Roy. Meteor. Soc.*, 135, 1020–1029, doi:10.1002/qj.409, 2009.

Pagan, K. L., Tabazadeh, A., Drdla, K., Hervig, M. E., Eckermann, S. D., Browell, E. V., Legg, M. J., and Foschi, P. G.: Observational evidence against mountain-wave generation of ice nuclei as a prerequisite for the formation of three solid nitric acid polar stratospheric clouds observed in the Arctic in early December 1999, *J. Geophys. Res.*, 109, D04312, doi:10.1029/2003JD003846, 2004.

Papanastasiou, D. K., Papadimitriou, V. C., Fahey, D. W., and Burkholder, J. B.: UV absorption spectrum of the ClO dimer (Cl₂O₂) between 200 and 420 nm, *J. Phys. Chem.*, 113, 13711–13726, doi:10.1021/jp9065345, 2009.

Peter, T.: Microphysics and heterogeneous chemistry of polar stratospheric clouds, *Annu. Rev. Phys. Chem.*, 48, 785–822, 1997.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Pitts, M. C., Poole, L. R., and Thomason, L. W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, doi:10.5194/acp-9-7577-2009, 2009.
- Pitts, M. C., Poole, L. R., Dörnbrack, A., and Thomason, L. W.: The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, doi:10.5194/acp-11-2161-2011, 2011.
- Plenge, J., Köhl, S., Vogel, B., Müller, R., Strohm, F., von Hobe, M., Flesch, R., and Rühl, E.: Bond strength of chlorine peroxide, *J. Phys. Chem.*, 109, 6730–6734, 2005.
- Polvani, L. M., Waugh, D. W., Correa, G. J. P., and Son, S.-W.: Stratospheric ozone depletion: the main driver of twentieth-century atmospheric circulation changes in the Southern Hemisphere, *J. Climate*, 24, 795–812, doi:10.1175/2010jcli3772.1, 2010.
- Pope, F. D., Hansen, J. C., Bayes, K. D., Friedl, R. R., and Sander, S. P.: Ultraviolet absorption spectrum of chlorine peroxide, ClOOCl, *J. Phys. Chem.*, 111, 4322–4332, 2007.
- Pope, F. D., Braesicke, P., Grainger, R. G., Kalberer, M., Watson, I. M., Davidson, P. J., and Cox, R. A.: Stratospheric aerosol particles and solar-radiation management, *Nature Climate Change*, 2, 713–719, 2012.
- Proffitt, M. H., Powell, J. A., Tuck, A. F., Fahey, D. W., Kelly, K. K., Krueger, A. J., Schoeberl, M. R., Gary, B. L., Margitan, J. J., Chan, K. R., Loewenstein, M., and Podolske, J. R.: A chemical definition of the boundary of the Antarctic ozone hole, *J. Geophys. Res.*, 94, 11437–11448, 1989.
- Rautenhaus, M., Bauer, G., and Dörnbrack, A.: A web service based tool to plan atmospheric research flights, *Geosci. Model Dev.*, 5, 55–71, doi:10.5194/gmd-5-55-2012, 2012.
- Rex, M., Von Der Gathen, P., Braathen, G. O., Harris, N. R. P., Reimer, E., Beck, A., Alfier, R., Krüger-Carstensen, R., Chipperfield, M., De Backer, H., Balis, D., O'Connor, F., Dier, H., Dorokhov, V., Fast, H., Gamma, A., Gil, M., Kyrö, E., Litynska, Z., Mikkelsen, I. S., Molyneux, M., Murphy, G., Reid, S. J., Rummukainen, M., and Zerefos, C.: Chemical ozone loss in the Arctic winter 1994/95 as determined by the match technique, *J. Atmos. Chem.*, 32, 35–59, doi:10.1023/a:1006093826861, 1999.
- Rex, M., Salawitch, R. J., von der Gathen, P., Harris, N. R. P., Chipperfield, M. P., and Naujokat, B.: Arctic ozone loss and climate change, *Geophys. Res. Lett.*, 31, L04116, doi:10.1029/2003GL018844, 2004.
- Rex, M., Salawitch, R. J., Deckelmann, H., von der Gathen, P., Harris, N. R. P., Chipperfield, M. P., Naujokat, B., Reimer, E., Allaart, M., Andersen, S. B., Bevilacqua, R., Braa-

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



then, G. O., Claude, H., Davies, J., De Backer, H., Dier, H., Dorokhov, V., Fast, H., Gerd-
 ing, M., Godin-Beekmann, S., Hoppel, K., Johnson, B., Kyroe, E., Litynska, Z., Moore, D.,
 Nakane, H., Parrondo, M. C., Risley Jr., A. D., Skrivankova, P., Stuebi, R., Viatte, P.,
 Yushkov, V., and Zerefos, C.: Arctic winter 2005: implications for stratospheric ozone loss
 and climate change, *Geophys. Res. Lett.*, 33, L23808, doi:10.1029/2006GL026731, 2006.

Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact
 of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative
 effects, *J. Geophys. Res.*, 117, D16305, doi:10.1029/2012JD017751, 2012.

Rosen, J. M. and Kjome, N. T.: Backscattersonde – a new instrument for atmospheric aerosol
 research, *Appl. Optics*, 30, 1552–1561, 1991.

Sadourny, R. and Laval, K.: January and July performance of the LMD general circulation
 model, in: *New perspectives in Climate Modeling*, edited by: Berger, A. and Nicolis, C., Else-
 vier, Amsterdam, 173–194, 1984.

Salawitch, R., Canty, T., Stimpfle, R., Wilmouth, D., Anderson, J. G., von Hobe, M., Stroh, F.,
 Rex, M., Schofield, R., Santee, M., Kinnison, D. E., Kurylo, M. J., and Sinnhuber, B. M.:
 Impact of recent laboratory measurements of the ClOOCl cross section on our understanding
 of polar ozone chemistry, *Abstr. Pap. Am. Chem. S.*, 237, 363–363, 2009.

Sander, S. P., Abbatt, J., Barker, J. R., Burkholder, J. B., Friedl, R. R., Golden, D. M., Huie, R. E.,
 Kolb, C. E., Kurylo, M. J., Moortgat, G. K., Orkin, V. L., and Wine, P. H.: Chemical kinetics and
 photochemical data for use in atmospheric studies, *Jet Propulsion Laboratory, Pasadena*,
 10–6, 2011.

Santee, M. L., Lambert, A., Read, W. G., Livesey, N. J., Cofield, R. E., Cuddy, D. T., Daffer, W. H.,
 Drouin, B. J., Froidevaux, L., Fuller, R. A., Jarnot, R. F., Knosp, B. W., Manney, G. L., Pe-
 run, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Waters, J. W., Mus-
 cari, G., de Zafra, R. L., Dibb, J. E., Fahey, D. W., Popp, P. J., Marcy, T. P., Jucks, K. W.,
 Toon, G. C., Stachnik, R. A., Bernath, P. F., Boone, C. D., Walker, K. A., Urban, J., and
 Murtagh, D.: Validation of the Aura Microwave Limb Sounder HNO₃ measurements, *J. Geo-
 phys. Res.*, 112, D24S40, doi:10.1029/2007JD008721, 2007a.

Santee, M. L., Lambert, A., Read, W. G., Livesey, N. J., Manney, G. L., Cofield, R. R.,
 Cuddy, D. T., Daffer, W. H., Froidevaux, L., Fuller, R. A., Jarnot, R. F., Knosp, B. W., Pe-
 run, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Waters, J. W., Con-
 nor, B., Urban, J., Murtagh, D., Ricaud, P., Barret, B., Kleinböhl, A. J. K., Kullmann, H., von

- Hobe, M., Toon, G. C., and Stachnik, R. A.: Validation of the Aura Microwave Limb Sounder CIO measurements, *J. Geophys. Res.*, 113, D15S22, doi:10.1029/2007JD008762, 2007b.
- Schiermeier, Q.: Chemists poke holes in ozone theory, *Nature*, 449, 382–383, 2007.
- Schofield, R., Frieler, K., Wohltmann, I., Rex, M., von Hobe, M., Stroh, F., Koch, G., Peter, T., Canty, T., Salawitch, R., and Volk, C. M.: Polar stratospheric chlorine kinetics from a self-match flight during SOLVE-II/EUPLEX, *Geophys. Res. Lett.*, 35, L01807, doi:10.1029/2007GL031740, 2008.
- Shi, Q., Jayne, J. T., Kolb, C. E., Worsnop, D. R., and Davidovits, P.: Kinetic model for reaction of ClONO₂ with H₂O and HCl and HOCl with HCl in sulfuric acid solutions, *J. Geophys. Res.*, 106, 24259–24274, 2001.
- Shindell, D. T., Miller, R. L., Schmidt, G. A., and Pandolfo, L.: Simulation of recent northern winter climate trends by greenhouse-gas forcing, *Nature*, 399, 452–455, 1999.
- Sigmond, M., Scinocca, J. F., and Kushner, P. J.: Impact of the stratosphere on tropospheric climate change, *Geophys. Res. Lett.*, 35, L12706, doi:10.1029/2008GL033573, 2008.
- Sitnikov, N. M., Yushkov, V. A., Afchine, A. A., Korshunov, L. I., Astakhov, V. I., Ulanovskii, A. E., Kraemer, M., Mangold, A., Schiller, C., and Ravegnani, F.: The FLASH instrument for water vapor measurements on board the high-altitude airplane, *Instrum. Exp. Tech.*, 50, 113–121, 2007.
- Solomon, S.: Stratospheric ozone depletion: a review of concepts and history, *Rev. Geophys.*, 37, 275–316, 1999.
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, *Nature*, 321, 755–758, 1986.
- Solomon, S., Portmann, R. W., Sasaki, T., Hofmann, D. J., and Thompson, D. W. J.: Four decades of ozonesonde measurements over Antarctica, *J. Geophys. Res.*, 110, D21311, doi:10.1029/2005JD005917, 2005.
- Sovde, O. A., Orsolini, Y. J., Jackson, D. R., Stordal, F., Isaksen, S. A., and Rognerud, B.: Estimation of Arctic ozone loss in winter 2006/2007 using a chemical transport model and data assimilation., *Q. J. Roy. Meteor. Soc.*, 137, 118–128, 2011.
- Spang, R., Grooß, J. U., Günther, G., Von Hobe, M., Kalicinsky, C., Riese, M., Stroh, F., Ungermann, J., Gerber, D., Moyna, B. P., Oldfield, M. L., Reburn, W. J., Siddans, R., Kerridge, B. J., Oelhaf, H., and Woiwode, W.: Final Report of the PremierEX Study, ESA, ESTEC, Netherlands, 2011.
- SPARC: SPARC Report No. 5, Chemistry-Climate Model Validation, SPARC, 2010.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Stolarski, R. S., Krueger, A. J., Schoeberl, M. R., McPeters, R. D., Newman, P. A., and Alpert, J. C.: Nimbus-7 satellite measurements of the springtime Antarctic ozone decrease, *Nature*, 322, 808–811, 1986.
- 5 Stroh, F., Grooß, J. U., Heinecke, F., von Hobe, M., Suminska-Ebersoldt, O., Volk, C. M., and Wegner, T.: Polar vortex ClO_x chemistry from RECONCILE: ClO and ClO dimer measurements, Quadrennial Ozone Symposium, Toronto, Canada, L08502, doi:10.1029/2009GL037524, 2012.
- 10 Struthers, H., Bodeker, G. E., Austin, J., Bekki, S., Cionni, I., Dameris, M., Giorgetta, M. A., Grewe, V., Lefèvre, F., Lott, F., Manzini, E., Peter, T., Rozanov, E., and Schraner, M.: The simulation of the Antarctic ozone hole by chemistry-climate models, *Atmos. Chem. Phys.*, 9, 6363–6376, doi:10.5194/acp-9-6363-2009, 2009.
- 15 Sumińska-Ebersoldt, O., Lehmann, R., Wegner, T., Grooß, J.-U., Hösen, E., Weigel, R., Frey, W., Griessbach, S., Mitev, V., Emde, C., Volk, C. M., Borrmann, S., Rex, M., Stroh, F., and von Hobe, M.: ClOOCl photolysis at high solar zenith angles: analysis of the RECONCILE self-match flight, *Atmos. Chem. Phys.*, 12, 1353–1365, doi:10.5194/acp-12-1353-2012, 2012.
- Tilmes, S., Müller, R., and Salawitch, R. J.: The sensitivity of polar ozone depletion to proposed geoeengineering schemes, *Science*, 320, 1201–1204, 2008.
- 20 Tsias, A., Wirth, M., Carslaw, K. S., Biele, J., Mehrtens, H., Reichardt, J., Wedekind, C., Weiss, V., Renger, W., Neuber, R., von Zahn, U., Stein, B., Santacesaria, V., Stefanutti, L., Fierli, F., Bacmeister, J., and Peter, T.: Aircraft lidar observations of an enhanced type Ia polar stratospheric clouds during APE-POLECAT, *J. Geophys. Res.*, 104, 23961–23969, 1999.
- Tuck, A. F., Watson, R. T., Condon, E. P., Margitan, J. J., and Toon, O. B.: The planning and execution of ER-2 and DC-8 aircraft flights over Antarctica, August and September 1987, *J. Geophys. Res.*, 94, 11181–11222, 1989.
- 25 Turco, R., Plumb, A., and Condon, E.: The Airborne Arctic Stratospheric Expedition – prologue, *Geophys. Res. Lett.*, 17, 313–316, 1990.
- Turner, J., Comiso, J. C., Marshall, G. J., Lachlan-Cope, T. A., Bracegirdle, T., Maksym, T., Meredith, M. P., Wang, Z., and Orr, A.: Non-annular atmospheric circulation change induced by stratospheric ozone depletion and its role in the recent increase of Antarctic sea ice extent, *Geophys. Res. Lett.*, 36, L08502, doi:10.1029/2009GL037524, 2009.
- 30

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ulanovsky, A. E., Yushkov, V. A., Sitnikov, N. M., and Ravegnani, F.: The FOZAN-II fast-response chemiluminescent airborne ozone analyzer, *Instrum. Exp. Tech.*, 44, 249–256, 2001.

Ungermann, J., Kalicinsky, C., Olschewski, F., Knieling, P., Hoffmann, L., Blank, J., Woiwode, W., Oelhaf, H., Hösen, E., Volk, C. M., Ulanovsky, A., Ravegnani, F., Weigel, K., Stroh, F., and Riese, M.: CRISTA-NF measurements with unprecedented vertical resolution during the RECONCILE aircraft campaign, *Atmos. Meas. Tech.*, 5, 1173–1191, doi:10.5194/amt-5-1173-2012, 2012.

Urban, J., Pommier, M., Murtagh, D. P., Santee, M. L., and Orsolini, Y. J.: Nitric acid in the stratosphere based on Odin observations from 2001 to 2009 – Part 1: A global climatology, *Atmos. Chem. Phys.*, 9, 7031–7044, doi:10.5194/acp-9-7031-2009, 2009.

van der A, R. J., Allaart, M. A. F., and Eskes, H. J.: Multi sensor reanalysis of total ozone, *Atmos. Chem. Phys.*, 10, 11277–11294, doi:10.5194/acp-10-11277-2010, 2010.

Vaughan, G., Schiller, C., MacKenzie, A. R., Bower, K., Peter, T., Schlager, H., Harris, N. R. P., and May, P. T.: SCOUT-03/ACTIVE – high-altitude aircraft measurements around deep tropical convection, *B. Am. Meteorol. Soc.*, 89, 647–662, 2008.

Viciani, S., D’Amato, F., Mazzinghi, P., Castagnoli, F., Toci, G., and Werle, P.: A cryogenically operated laser diode spectrometer for airborne measurement of stratospheric trace gases, *Appl. Phys. B*, 90, 581–592, 2008.

Vogel, B., Feng, W., Streibel, M., and Müller, R.: The potential impact of ClO_x radical complexes on polar stratospheric ozone loss processes, *Atmos. Chem. Phys.*, 6, 3099–3114, doi:10.5194/acp-6-3099-2006, 2006.

Voigt, C., Schlager, H., Luo, B. P., Dörnbrack, A., Roiger, A., Stock, P., Curtius, J., Vössing, H., Borrmann, S., Davies, S., Konopka, P., Schiller, C., Shur, G., and Peter, T.: Nitric Acid Trihydrate (NAT) formation at low NAT supersaturation in Polar Stratospheric Clouds (PSCs), *Atmos. Chem. Phys.*, 5, 1371–1380, doi:10.5194/acp-5-1371-2005, 2005.

Vömel, H., David, D. E., and Smith, K.: Accuracy of tropospheric and stratospheric water vapor measurements by the cryogenic frost point hygrometer: instrumental details and observations, *J. Geophys. Res.*, 112, D08305, doi:10.1029/2006JD007224, 2007.

von Hobe, M.: Atmospheric science – revisiting ozone depletion, *Science*, 318, 1878–1879, 2007.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- von Hobe, M., Grooß, J.-U., Müller, R., Hrechanyy, S., Winkler, U., and Stroh, F.: A re-evaluation of the ClO/Cl₂O₂ equilibrium constant based on stratospheric in-situ observations, *Atmos. Chem. Phys.*, 5, 693–702, doi:10.5194/acp-5-693-2005, 2005.
- von Hobe, M., Ulanovsky, A., Volk, C. M., Grooß, J. U., Tilmes, S., Konopka, P., Günther, G., Werner, A., Spelten, N., Shur, G., Yushkov, V., Ravegnani, F., Schiller, C., Müller, R., and Stroh, F.: Severe ozone depletion in the cold Arctic winter 2004–05, *Geophys. Res. Lett.*, 33, L17815, doi:10.1029/2006GL026945, 2006.
- von Hobe, M., Salawitch, R. J., Canty, T., Keller-Rudek, H., Moortgat, G. K., Grooß, J.-U., Müller, R., and Stroh, F.: Understanding the kinetics of the ClO dimer cycle, *Atmos. Chem. Phys.*, 7, 3055–3069, doi:10.5194/acp-7-3055-2007, 2007.
- von Hobe, M., Stroh, F., Beckers, H., Benter, T., and Willner, H.: The UV/Vis absorption spectrum of matrix-isolated dichlorine peroxide, ClOOCl, *Phys. Chem. Chem. Phys.*, 11, 1571–1580, 2009.
- von Hobe, M., Griessbach, S., and Wegner, T.: Impact of Arctic and Antarctic ozone depletion on UV radiation, *Quadrennial Ozone Symposium, Toronto, Canada, 2012a*.
- von Hobe, M., Suminska-Ebersoldt, O., Emde, C., and Stroh, F.: How well can we quantify the actinic flux driving catalytic ozone chemistry at high solar zenith angles?, in: *Geophys. Res. Abst., EGU General Assembly, Vienna, 8030, 22–27 April 2012, 2012b*.
- Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read, W. G., Siegel, P. H., Cofield, R. R., Filipak, M. J., Flower, D. A., Holden, J. R., Lau, G. K., Livesey, N. J., Manney, G. L., Pumphrey, H. C., Santee, M. L., Wu, D. L., Cuddy, D. T., Lay, R. R., Loo, M. S., Perun, C. S., Schwartz, M. J., Stek, P. C., Thurstans, R. P., Boyles, M. A., Chandra, K. M., Chavez, M. C., Chen, G. S., Chudasama, B. V., Dodge, R., Fuller, R. A., Girard, M. A., Jiang, J. H., Jiang, Y., Knosp, B. W., LaBelle, R. C., Lam, J. C., Lee, K. A., Miller, D., Oswald, J. E., Patel, N. C., Pukala, D. M., Quintero, O., Scaff, D. M., Snyder, W. V., Tope, M. C., Wagner, P. A., and Walch, M. J.: The Earth Observing System Microwave Limb Sounder (EOS MLS) on the Aura satellite, *IEEE T. Geosci. Remote*, 44, 1075–1092, 2006.
- Waugh, D. W. and Polvani, L. M.: Stratospheric polar vortices in the stratosphere: dynamics, chemistry, and transport, *AGU, Geophys. Monogr. Ser.*, 190, 43–57, 2010.
- Waugh, D. W., Oman, L., Kawa, S. R., Stolarski, R. S., Pawson, S., Douglass, A. R., Newman, P. A., and Nielsen, J. E.: Impacts of climate change on stratospheric ozone recovery, *Geophys. Res. Lett.*, 36, L03805, doi:10.1029/2008GL036223, 2009.

RECONCILE

M. von Hobe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Wegner, T., Grooß, J.-U., Müller, R., Stroh, F., Lehmann, R., Volk, C. M., Hösen, E., Vom Scheidt, M., Wintel, J., Riediger, O., Schlager, H., Scheibe, M., Stock, P., Ravegnani, F., Ulanovsky, A., Yushkov, V. A., and von Hobe, M.: Chasing air masses in the Arctic vortex: an evaluation of trajectory calculations using an active match, AGU, Fall Meeting 2010, San Francisco, USA, 2010.

Wegner, T., Grooß, J.-U., von Hobe, M., Stroh, F., Sumińska-Ebersoldt, O., Volk, C. M., Hösen, E., Mitev, V., Shur, G., and Müller, R.: Heterogeneous chlorine activation on stratospheric aerosols and clouds in the Arctic polar vortex, *Atmos. Chem. Phys.*, 12, 11095–11106, doi:10.5194/acp-12-11095-2012, 2012.

Wegner, T., Kinnison, D. E., Garcia, R. R., Madronich, S., and Solomon, S.: Polar stratospheric clouds in SD-WACCM~4, *J. Geophys. Res.*, submitted, 2012b.

Wegner, T., Kinnison, D. E., Garcia, R. R., Madronich, S., Solomon, S., and von Hobe, M.: On the depletion of HCl in the Antarctic polar vortex, *J. Geophys. Res.*, submitted, 2012c.

Weigel, R., Hermann, M., Curtius, J., Voigt, C., Walter, S., Böttger, T., Lepukhov, B., Belyaev, G., and Borrmann, S.: Experimental characterization of the COndensation PArticle counting System for high altitude aircraft-borne application, *Atmos. Meas. Tech.*, 2, 243–258, doi:10.5194/amt-2-243-2009, 2009.

Weigel, K., Riese, M., Hoffmann, L., Hofer, S., Kalicinsky, C., Knieling, P., Olschewski, F., Preusse, P., Spang, R., Stroh, F., and Volk, C. M.: CRISTA-NF measurements during the AMMA-SCOUT-O3 aircraft campaign, *Atmos. Meas. Tech.*, 3, 1437–1455, doi:10.5194/amt-3-1437-2010, 2010.

Weigel, R., Borrmann, S., Ebert, M., Kandler, K., Frey, W., Molleker, S., Volk, C. M., Günther, G., Schlager, H., Cairo, F., Di Donfrancesco, G., and Khaykin, S.: Non-volatile aerosol in the Arctic winter stratosphere and its role for PSC formation, European Aerosol Conference, Granada, Spain, 2012.

Werner, A., Volk, C. M., Ivanova, E. V., Wetter, T., Schiller, C., Schlager, H., and Konopka, P.: Quantifying transport into the Arctic lowermost stratosphere, *Atmos. Chem. Phys.*, 10, 11623–11639, doi:10.5194/acp-10-11623-2010, 2010.

Wetzel, G., Oelhaf, H., Kirner, O., Friedl-Vallon, F., Ruhnke, R., Ebersoldt, A., Kleinert, A., Maucher, G., Nordmeyer, H., and Orphal, J.: Diurnal variations of reactive chlorine and nitrogen oxides observed by MIPAS-B inside the January 2010 Arctic vortex, *Atmos. Chem. Phys.*, 12, 6581–6592, doi:10.5194/acp-12-6581-2012, 2012.

COBALD (Compact Optical Backscatter Aerosol Detector) Data Sheet: available at: http://www.iac.ethz.ch/groups/peter/research/Balloon_soundings/COBALD_data_sheet, last access: 8 October, 2012.

Wilmouth, D. M., Hanisco, T. F., Stimpfle, R. M., and Anderson, J. G.: Chlorine-catalyzed ozone destruction: Cl atom production from ClOOCl photolysis, *J. Phys. Chem.*, 113, 14099–14108, doi:10.1021/jp9053204, 2009.

WMO: Scientific assessment of ozone depletion 2010, World Meteorological Organization, Global ozone research and monitoring project, Geneva, Switzerland, 52, 2010.

Wohltmann, I. and Rex, M.: The Lagrangian chemistry and transport model ATLAS: validation of advective transport and mixing, *Geosci. Model Dev.*, 2, 153–173, doi:10.5194/gmd-2-153-2009, 2009.

Wohltmann, I., Lehmann, R., and Rex, M.: The Lagrangian chemistry and transport model ATLAS: simulation and validation of stratospheric chemistry and ozone loss in the winter 1999/2000, *Geosci. Model Dev.*, 3, 585–601, doi:10.5194/gmd-3-585-2010, 2010.

Wohltmann, I., Wegner, T., Müller, R., Lehmann, R., Rex, M., Manney, G. L., Santee, M. L., Bernath, P., Sumińska-Ebersoldt, O., Stroh, F., von Hobe, M., Volk, C. M., Hösen, E., Ravegnani, F., Ulanovsky, A., and Yushkov, V.: Uncertainties in modeling heterogeneous chemistry and Arctic ozone depletion in the winter 2009/2010, *Atmos. Chem. Phys. Discuss.*, 12, 26245–26295, doi:10.5194/acpd-12-26245-2012, 2012.

Woiwode, W., Oelhaf, H., Gulde, T., Piesch, C., Ebersoldt, A., and Orphal, J.: FTIR Limb-Sounding in the Arctic UTLS under cloudy conditions: the MIPAS-STR PSC flights during RECONCILE and ESSenCe, in: *Geophys. Res. Abst.*, EGU General Assembly, Vienna, Austria, EGU2012-11490, 2012a.

Woiwode, W., Oelhaf, H., Gulde, T., Piesch, C., Maucher, G., Ebersoldt, A., Keim, C., Höpfner, M., Khaykin, S., Ravegnani, F., Ulanovsky, A. E., Volk, C. M., Hösen, E., Dörnbrack, A., Ungermann, J., Kalicinsky, C., and Orphal, J.: MIPAS-STR measurements in the Arctic UTLS in winter/spring 2010: instrument characterization, retrieval and validation, *Atmos. Meas. Tech.*, 5, 1205–1228, doi:10.5194/amt-5-1205-2012, 2012b.

Wolff, M. A., Kerzenmacher, T., Strong, K., Walker, K. A., Toohey, M., Dupuy, E., Bernath, P. F., Boone, C. D., Brohede, S., Catoire, V., von Clarmann, T., Coffey, M., Daffer, W. H., De Mazière, M., Duchatelet, P., Glatthor, N., Griffith, D. W. T., Hannigan, J., Hase, F., Höpfner, M., Huret, N., Jones, N., Jucks, K., Kagawa, A., Kasai, Y., Kramer, I., Küllmann, H., Kuttippurath, J., Mahieu, E., Manney, G., McElroy, C. T., McLinden, C., Mébarki, Y.,

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Mikuteit, S., Murtagh, D., Piccolo, C., Raspollini, P., Ridolfi, M., Ruhnke, R., Santee, M., Senten, C., Smale, D., Tétard, C., Urban, J., and Wood, S.: Validation of HNO_3 , ClONO_2 , and N_2O_5 from the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS), *Atmos. Chem. Phys.*, 8, 3529–3562, doi:10.5194/acp-8-3529-2008, 2008.
- 5 Young, I. A. K., Murray, C., Blaum, C. M., Cox, R. A., Jones, R. L., and Pope, F. D.: Temperature dependent structured absorption spectra of molecular chlorine, *Phys. Chem. Chem. Phys.*, 13, 15318–15325, 2011.
- Young, I. A. K., Jones, R. L., and Pope, F. D.: The UV and visible spectrum of ClOOCl , in preparation, 2012.
- 10 Yushkov, V., Merkulov, S., and Astakhov, V.: Optical balloon hygrometer for upper stratosphere and stratosphere water vapour measurements, in: *Proc. SPIE, Optical Remote Sensing of the Atmosphere and Clouds*, 439–445, 1998.
- Yushkov, V., Oulanovsky, A., Lechenuk, N., Roudakov, I., Arshinov, K., Tikhonov, F., Stefanutti, L., Ravegnani, F., Bonafe, U., and Georgiadis, T.: A chemiluminescent analyzer for stratospheric measurements of the ozone concentration (FOZAN), *J. Atmos. Ocean. Tech.*, 15, 1345–1350, 1999.
- Zhou, T., Geller, M. A., and Lin, W.: An observational study on the latitudes where wave forcing drives Brewer–Dobson upwelling, *J. Atmos. Sci.*, 69, 1916–1935, 2012.
- Zobrist, B., Marcolli, C., Pedernera, D. A., and Koop, T.: Do atmospheric aerosols form glasses?, *Atmos. Chem. Phys.*, 8, 5221–5244, 2008, <http://www.atmos-chem-phys.net/8/5221/2008/>.
- 20 Zöger, M., Afchine, A., Eicke, N., Gerhards, M. T., Klein, E., McKenna, D. S., Morschel, U., Schmidt, U., Tan, V., Tuitjer, F., Woyke, T., and Schiller, C.: Fast in situ stratospheric hygrometers: a new family of balloon-borne and airborne Lyman alpha photofragment fluorescence hygrometers, *J. Geophys. Res.*, 104, 1807–1816, 1999.
- 25

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Overview of RECONCILE flights. All flights were carried out in 2010, take off and landing times are in UTC. The corresponding flight tracks are shown in Fig. 4.

No	date	take off	landing	Scientific targets
F1	17 Jan	11:20	15:33	PSC characterization
F2	20 Jan	08:13	11:55	PSC characterization and CALIPSO match
F3	22 Jan	10:01	13:35	PSC characterization
F4	24 Jan	13:30	16:56	PSCs and renitrification
F5	25 Jan	05:50	09:19	PSCs, de/renitrification, Cl active., self-match experiment ¹
F6	28 Jan	09:00	12:56	vortex survey (dynamics and mixing)
F7	30 Jan	06:36	10:15	self-match experiment ¹ , Cl activation, de/renitrification
F8	2 Feb	10:01	13:31	match experiment ² , Cl activation, de/renitrification, O ₃ loss
F9	27 Feb	12:06	15:41	vortex filament
F10 ³	2 Mar	02:51	06:31	vortex survey, Cl deactivation, de/renitrification, O ₃ loss
F11 ³	2 Mar	09:35	13:35	vortex survey, Cl deactivation, de/renitrification, O ₃ loss
F12	5 Mar	15:05	18:36	vortex filament
F13 ⁴	10 Mar	07:14	10:51	test of remote sensing instrumentation, O ₃ loss

¹ The self match flight is a particularly useful flight pattern to constrain kinetic parameters. Air masses are sampled twice during the same flight (Schofield et al., 2008). The two RECONCILE self match flights were carried out over sunrise to constrain the ClOOCl photolysis rate J and the ClO/ClOOCl thermal equilibrium constant K_{EQ} (Suminska-Ebersoldt et al., 2012).

² Three days after the second self-match flight, the air masses that had been sampled came within reach of the Geophysica again, and a match flight was carried out to test the accuracy of trajectory calculations and to check the consistency of observed and simulated chlorine activation and ozone loss (Wegner et al., 2012).

³ Flights F10 and 11 were carried out as relay flights with an intermediate stop in Spitsbergen.

⁴ Flight 13 was dedicated to test remote sensing instrumentation under the PremierEx project funded by ESA.

Table 2. Instrumental payload of M55 Geophysica during the RECONCILE field campaign.

instrument	parameter	time res.	accuracy	precision	technique	references
in situ						
FOZAN	O ₃	1 s	0.01 ppm	8 %	Dye chemiluminescence+ECC	(Ulanovsky et al., 2001; Yushkov et al., 1999)
FISH	H ₂ O (total)	1 s	0.2 ppm	4 %	Lyman- α	(Zöger et al., 1999)
FLASH	H ₂ O (gas phase)	8 s	0.2 ppm	6 %	Lyman- α	(Sitnikov et al., 2007)
SIOUX	NO	1 s	10 %	3 %	Chemiluminescence,	(Voigt et al., 2005)
	NO _y	1 s	15 %	5 %	+ Au-converter + Subisokinetic inlet	
HALOX	ClO	10 s	20 %	5 %	CCRF	(von Hobe et al., 2005)
	ClOOCl	30 s	30 %	15 %	+ thermal dissociation	
HAGAR	N ₂ O, CFC12	90 s	1.3 %	0.3 %	Gas Chromatography (GC)	(Homan et al., 2010;
	CFC 11	90 s	1.6 %	0.6 %	with electron capture detector (ECD)	Werner et al., 2010)
	CH ₄ , H ₂	90 s	1.8 %	0.8 %	IR absorption	
	SF ₆	90 s	2.3 %	1.3 %		
	Halon 1211	90 s	3.0 %	2.0 %		
COLD	CO ₂	5 s	0.1 %	0.05 %		
	CO	5 s	9 %	1 %	TDL	(Viciani et al., 2008)
COPAS	Condensation nuclei (CN-total, CN-non-volatile)	1 s	10 %	5 %	2-channel CN counter, one inlet heated	(Weigel et al., 2009)
	FSSP	Cloud particle size distrib. (0.4–47 μ m)	1 s	20 %	10 %	Laser-particle spectrometer
CCP	Cloud particle size distrib. (3–47 μ m)	1 s	20 %	10 %	Laser-particle spectrometer	
CIP	Cloud particle size distrib. (25–1600 μ m)	1 s	20 %	10 %	Laser-particle spectrometer	(Baumgardner et al., 2001)
Particle Images						
MAS	Aerosol optical properties	10 s	5 %	5 %	Multi-wavelength Scattering	(Buontempo et al., 2006)
WAS	Long lived trace gases and isotopo-logues	minutes	5 %	< 2 %	Whole air sampling with lab GC and MS analysis	(Kaiser et al., 2006; Laube et al., 2010a)
	Rosemount probe (TDC)	<i>T</i> , <i>P</i> horizontal wind	0.1 s 0.1 s	0.5 K 1 ms ⁻¹	0.1 K 0.1 ms ⁻¹	PT100, 5-hole probe
remote sensing						
MAL 1 & 2	Remote Aerosol Profile (2 km from aircraft altitude)	30–120 s	10 %	10 %	Microjoule-lidar	(Matthey et al., 2003)
miniDOAS	BrO, OClO				DOAS	
MIPAS-STR	Cloud Index, <i>T</i> , HNO ₃ , O ₃ , ClONO ₂ , CFCs, H ₂ O and minor species	typ. 2.4/3.8 min for 1 profile (depends on sampling programme)	<i>T</i> < 1 K/ vmr typ. 10–15 %	<i>T</i> < 1 K/ vmr typ. 4–12 %	FTIR limb sounder	(Woiwode et al., 2012b)
CRISTA-NF	H ₂ O, HNO ₃ , PAN, ClONO ₂ , CFCs, O ₃ , CCl ₄				MIR-Emission in limb viewing geometry	(Ungermann et al., 2012; Weigel et al., 2010)
MARSCHALS	O ₃ , H ₂ O, CO, HNO ₃ , N ₂ O				Millimetre Wave spectrometer in limb viewing geometry	(Moyna et al., 2006)

RECONCILE

M. von Hobe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Kinetic and photochemical parameters for the ClO-dimer cycle that are most consistent with the experiments and observations made within RECONCILE. Clearly, it does not represent a comprehensive evaluation of kinetic parameters determined in laboratory experiments such as given by Sander et al. (2011).

ClOOCl production rate constant k_f^*	(Nickolaisen et al., 1994) $2.1 \times 10^{-32} \cdot (T/300)^{-3.01}$	This is somewhat lower than the current JPL recommendation (Sander et al., 2011), but gives the best fit to field measurements of ClO made during the RECONCILE field campaign in 2010 (Suminska-Ebersoldt et al., 2012) and can be rationalized based on theoretical chemistry calculations (von Hobe et al., 2007)
ClO/ClOOCl equilibrium constant K_{EQ}	(Plenge et al., 2005) $1.92 \times 10^{-27} \cdot e^{8430/T}$	Field measurements of ClO before sunrise in air masses that were shown to be in thermal equilibrium rule out parameterizations that give higher values for K_{EQ} (Suminska-Ebersoldt et al., 2012). Again, this is in good agreement with theoretical considerations by von Hobe et al. (2007)
ClOOCl absorption cross sections σ_{ClOOCl}	(Papanastasiou et al., 2009)	These published cross sections are consistent with the ones measured in the laboratory as part of RECONCILE and also with ClO observations made during the RECONCILE aircraft campaign

* We only consider the low pressure limit for k_f , which is much more relevant for stratospheric conditions than the high pressure limit.

RECONCILE

M. von Hobe et al.




Fig. 1. Illustrative timeline of ozone research. More detailed discussions on the history of the discovery of the anthropogenic influence on stratospheric ozone are given by Solomon (1999), Brasseur (2008) and Müller (2009). Parts of this figure are from WMO (2010) and <http://www.nobelprize.org>, the Concorde photo is courtesy of K.-H. Nogai.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Conditions favouring the formation of an ozone hole in polar regions in spring. **(a)** The isolation of the air inside the polar vortex inhibits dilution of active chlorine and replenishment of ozone (McIntyre, 1989; Proffitt et al., 1989). **(b)** Fast heterogeneous reactions on particle surfaces activate chlorine from reservoir gases (Solomon et al., 1986). **(c)** Ozone is destroyed by catalytic cycles that are particularly efficient under cold temperature/high solar zenith angle conditions (McElroy et al., 1986; Molina and Molina, 1987).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. In RECONCILE, a fully integrated approach is adopted that joins detailed process studies including laboratory and field experiments with global chemistry-climate model (CCM) studies.

RECONCILE

M. von Hobe et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



RECONCILE

M. von Hobe et al.




Fig. 4. Geophysica flight paths during RECONCILE. In the left panel, all flights are marked by different colours (referring to flight date, yymmdd). In the right panel, flight sections in vortex air are marked red, flight sections outside the polar vortex are marked in blue. Two self-match flights were carried out on 25 and 30 January, a match flight on 2 February (matching the flight on 30 January).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

RECONCILE

M. von Hobe et al.




Fig. 5. Flight tracks of the second self match flight on 30 January and the match flight on 2 February with HAGAR N₂O measurements along the flight tracks shown by the coloured squares. The trajectories connecting the flights are colour coded according to the N₂O (in ppb) observed during the first flight. The grey contours in the background show PV from ERA-interim at the 425 K level on 30 January, 12:00 UTC, with white representing high PV and darker grey representing areas of low PV pointing to a significant contribution of non-vortex-air.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 6. Network of ozone sounding stations that participated in the RECONCILE match campaign (red). Gray dots indicate stations that were active in match campaigns in the past but which went out of operation during recent years due to lack of support.

RECONCILE

M. von Hobe et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



RECONCILE

M. von Hobe et al.




Fig. 7. Minimum temperatures T_{MIN} (K) between 65° N to 90° N on the 50 hPa (left panel) and 30 hPa (right panel) pressure surfaces. Black line: mean value from 1989–2011; red line the T_{MIN} evolution from July 2009 through June 2010, blue line from July 2010 through June 2011; shaded area encompasses the minimum/maximum T_{MIN} between 1 January 1989 and 30 June 2012. Source: ECMWF reanalyses interim (ERA Interim) data provided at 6 hourly temporal resolutions, see: <http://www.ecmwf.int>.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 8. Modified potential vorticity (PV, given in $10^{-6} \text{ K m}^2 \text{ kg}^{-1} \text{ s}^{-1}$, upper panel) and inert tracer P4 (indicating the relative contribution of air masses that were inside the vortex core at the time of initialization, lower panel) on 450 K isentropic level during the first vortex split, after reorganization of the vortex, during the second vortex split and at the end of the winter (order from left to right).




Fig. 9. Examples of CALIPSO PSC composition observations during the 2009–2010 Arctic winter. The panels on the left show CALIPSO PSC composition images for representative orbits from four periods during the winter: **(a)** 15–30 December, **(b)** 31 December–14 January, **(c)** 15–21 January, and **(d)** 22–28 January. GEOS5 temperature and geopotential height fields at 30 hPa for 12:00 UTC are shown at the right of each panel with the location of the corresponding CALIPSO orbit track indicated by the white line (red dot marks orbit track start).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 10. CALIPSO–COBALD comparison on the 17 January 2010. Upper panel: CALIPSO PSC composition observations. The black line denotes the location of the simultaneously performed COBALD sounding. Lower left panel: GEOS5 temperature and geopotential height fields at 30 hPa for 12:00 UTC with the location of the corresponding CALIPSO orbit track indicated by the white line and the position of the COBALD sonde by the yellow dot. Lower right panel: profiles of backscatter measurements from COBALD (red) and CALIPSO (black). According to the composition classification by Pitts et al. (2011) and the definition of ice, backscatter ratios larger than 5 are highlighted in blue. Grey profiles show maximum and minimum backscatter values from CALIPSO within a range of ± 25 km around the closest profile indicating the uncertainty between both measurements at 21 km altitude.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 11. Number size distribution obtained from FSSP-100 measurements during RECONCILE flight on 25 January 2010, averaged over 22 min (from 7:58 to 8:20 UTC). For comparison, the size distribution retrieved from NO_y observations in the Arctic winter 2000 by Fahey et al. (2001) is shown in red.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 12. PSC formation pathways as implemented into ZOMM with SBS = Supercooled Binary Solution ($\text{H}_2\text{SO}_4/\text{H}_2\text{O}$), STS = Supercooled Ternary Solution ($\text{HNO}_3/\text{H}_2\text{SO}_4/\text{H}_2\text{O}$), NAT = Nitric Acid Trihydrate ($\text{HNO}_3 \cdot 3\text{H}_2\text{O}$), \blacktriangle = solid inclusion, e.g. meteoritic dust. Grey arrows denote pathways included in previous schemes describing PSC formation pathways (going back to Peter, 1997). Black arrows show the heterogeneous nucleation pathways of NAT and ice on pre-existing solid particles supported by the new laboratory and field observations.




Fig. 13. MIPAS-STR standard retrieval results showing vertical redistribution of HNO₃ attributed to de/renitrification in Arctic polar winter/spring 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 14. Comparison of CIO simulated by CLaMS and observed by HALOX. Adapted from Wegner et al. (2012a).

RECONCILE

M. von Hobe et al.




Fig. 15. Results of ATLAS model runs, showing the reference (blue) and sensitivity (grey) runs for vortex-averaged ClO_x (left) and HCl (middle) for 18 January 2010, as well as ozone loss for 30 March 2010 as a function of potential temperature. For comparison, HCl observed by MLS and ozone loss inferred from MLS measurements (subtracting observed ozone measurements from the passive ozone tracer) are shown as black dashed lines in the respective panels. See Wohltmann et al. (2012) for details.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 16. Rates for the reaction $\text{Cl}^- + \text{ClONO}_2 \rightarrow \text{Cl}_2 + \text{NO}_3^-$ in/on the condensed phase (liquid or solid) calculated for pressure level 50 hPa and mixing ratio 1 ppb HCl, 10 ppb HNO_3 and 5 ppm H_2O . Rates for aerosols, ice clouds and “maximum NAT PSC” refer to NAT clouds characteristic for the Arctic, where the low NAT particle number densities with 2 orders of magnitude lower surface area densities hinder the gas phase to equilibrate. Transition from binary $\text{H}_2\text{SO}_4/\text{H}_2\text{O}$ aerosols to ternary $\text{HNO}_3/\text{H}_2\text{SO}_4/\text{H}_2\text{O}$ aerosols is indicated by the smooth transition from light green to black. Short time scales on the order of 1 day for chlorine activation appear to be possible by processing on binary aerosols, not requiring the formation of any PSCs (based on Biele et al., 2001; Carslaw et al., 1997; Dameris et al., 2007; Tsias et al., 1999).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 17. CIOOCI photolysis rate, J_{CIOOCI} , at different solar zenith angles for two different CIOOCI cross sections: Papa: determined by Papanastasiou et al. (2009) and currently recommended by JPL; vH/Lien: spectrum measured by von Hobe et al. (2009) scaled to the absolute measurement by Lien et al. (2009) at 248 nm wavelength. The grey area represents uncertainty due to variability in the actinic flux, with the black lines denoting a baseline scenario and the coloured error bars showing the sensitivities towards individual parameters affecting the actinic flux. This plot summarizes von Hobe et al. (2012b). Actinic fluxes were calculated using the radiative transfer model MYSTIC (Monte Carlo code for the physically correct Tracing of photons in Cloudy atmospheres, Mayer, 2009) in fully spherical geometry (Emde and Mayer, 2007) with the ALIS (Absorption Lines Importance Sampling) method to perform the high spectral resolution calculations efficiently (Emde et al., 2011). MYSTIC is part of the libRadtran radiative transfer package (Mayer and Kylling, 2005; see also <http://www.libradtran.org>).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.




Fig. 18. Difference in zonal mean annual mean ozone from a simulation of the SLIMCAT 3-D CTM with scaled J_{BrONO_2} (times 1.27) and $k_{\text{BrO}+\text{NO}_2}$ (times 0.75) compared to a run with standard JPL kinetics (from Kreycy et al., 2012).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 19. (a) Global geographic distribution of the DFA2 correlation exponent determined for the total column data base produced by Bodecker Scientific, and (b) for the LMDZrepro simulations. The reproduction of empirical correlation properties is very accurate in the overlapping region, spectroscopic measurements are not available over the poles. (c) Geographic distribution of the spectral weight of semiannual oscillations for the total column data base produced by Bodecker Scientific, and (d) for the LMDZrepro simulations. The agreement is satisfactory, however the simulations exhibit anomalously strong oscillations over the Indian ocean and the Tibetan plateau. Semiannual oscillations over the Antarctic is a well known feature in many atmospheric variables.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 20. Vertical profile of chemical ozone loss for 2011 and previous Arctic winters derived from match ozone sonde measurements.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RECONCILE

M. von Hobe et al.




Fig. 21. Modelled average column ozone north of 60° N for January, February and March 2010 (upper panel, dotted line) and 2011 (upper panel, full line) in simulations with no ozone-depleting chemistry north of 60° N. Relative difference in transport in 2010 compared to 2011 (lower panel, dotted line), and absolute ozone loss from heterogeneous chemistry in 2011 (lower panel, full line). Figure adapted from Isaksen et al. (2012).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)