

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges¹, D. E. Oram^{1,2}, J. C. Laube¹, C. E. Reeves¹, M. J. Newland¹,
C. Hogan¹, P. Martinerie³, E. Witrant⁴, C. A. M. Brenninkmeijer⁵, T. J. Schuck⁵,
and P. J. Fraser⁶

¹The School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK

²National Centre for Atmospheric Science, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK

³UJF – Grenoble 1/CNRS, Laboratoire de Glaciologie et Géophysique de l'Environnement (LGGE) UMR 5183, Grenoble, 38041, France

⁴Grenoble Image Parole Signal Automatique (GIPSA-lab), Université Joseph Fourier/CNRS, BP 46, 38 402 Saint Martin d'Hères, France

⁵Max Planck Institute for Chemistry, Air Chemistry Division, Mainz, Germany

⁶Centre for Australian Weather and Climate Research, Commonwealth Scientific and Industrial Research Organisation, Aspendale, Victoria 3195, Australia

Received: 22 December 2011 – Accepted: 3 January 2012 – Published: 10 January 2012

Correspondence to: W. T. Sturges (w.sturges@uea.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Long term measurements in background air (Cape Grim, Tasmania) and firn air (NEEM, Greenland) of the “super” greenhouse gas SF₅CF₃ show that emissions declined after the late 1990s, having grown since the 1950s, and became indistinguishable from zero after 2003. It is concluded that this is due to the cessation of global production of perfluorooctyl sulfonate-based fluorochemicals. An earlier observation of closely correlated atmospheric abundances of SF₅CF₃ and SF₆ are shown here to have likely been purely coincidental, as their respective trends diverged after 2002. Due to its long lifetime (ca. 900 yr), atmospheric concentrations of SF₅CF₃ have not declined, and it is now well mixed between hemispheres, as is also shown here from interhemispheric aircraft measurements. Total cumulative emissions of SF₅CF₃ amount to around 5 kT.

1 Introduction

In recent years the atmospheric growth of a number of fully-fluorinated and very long-lived greenhouse gases (“super” greenhouse gases), with lifetimes on the order of a thousand years or more, has been observed; e.g. CF₄ and C₂F₆ (Worton et al., 2007), C₃F₈ (Mühle et al., 2010), c-C₄F₈ (Oram et al., 2011), NF₃ (Weiss et al., 2008) and SF₆ (Levine et al., 2010). These gases have exceptionally high Global Warming Potentials (GWPs), in some cases on the order of several thousand times that of CO₂, and thus their unconstrained growth would be of considerable concern for the radiative forcing of climate.

In 2000 we reported our discovery of another, entirely unexpected, perfluorocarbon, namely SF₅CF₃ (Sturges et al., 2000). This was initially noticed as unidentified mass chromatographic peaks with *m/z* ratios representative of CF₃⁺ and SF₅⁺ ions during measurements of chlorofluorocarbons and SF₆ in air samples from the middle stratosphere. The appearance of this substance in air samples from above 32 km altitude

ACPD

12, 869–885, 2012

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



immediately implied that the “mystery” compound had to be long-lived. Two features marked this compound out as exceptional. Firstly, we determined that as a molecule it had the highest radiative forcing efficiency of any gas yet measured in the atmosphere ($0.57 \text{ W m}^{-2} \text{ ppb}^{-1}$). Secondly, it appeared to have grown in the atmosphere, between the mid-20th century and 1999, at almost exactly the same rate as SF_6 , to which it is closely chemically related. From this circumstantial evidence we concluded that its origins might be related to those of SF_6 , and we conjectured that it might be formed from high energy arc discharge in electrical equipment containing SF_6 as a high voltage dielectric. Indeed there has since been some laboratory evidence for this (Huang et al., 2005; Solovev et al., 2007). Immediately following publication of our article, however, an open letter to the publishing journal from the company 3M stated that “one source of this compound is as a by-product of the manufacture of certain 3M fluorochemicals” (Santoro et al., 2000). It transpired that the relevant process was electrochemical fluorination for the production of perfluorooctanyl sulphonate, (PFOS) and other fluorosurfactants, used in the manufacture of foams and stain-resist coatings. They went on to note that these production operations were to be imminently curtailed in the USA. In a subsequent personal communication, their bottom-up emission estimates were evidently close to the global emission rate that we had deduced from observations.

Our updated atmospheric measurements, presented here, clearly demonstrate that our original supposition that SF_5CF_3 originates from degradation of SF_6 was incorrect, and that all evidence points to 3M being correct in their assertion and that, as they had predicted, emissions of this potent greenhouse gas have subsequently reduced to the point where they are not distinguishable by observation from zero.

**Emissions halted of
the potent
greenhouse gas
 SF_5CF_3**

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Methodology

2.1 Sampling and analysis

The majority of the samples collected at Cape Grim, Tasmania (40.4° S, 144.4° E) since 1994 were pumped directly into 3-dm³ electrochemically passivated (Rasmussen) or Silcosteel-treated (Restek Corp.) stainless steel canisters at a pressure of ca. 2 atmospheres using a metal bellows air pump. Prior to 1994, the samples analysed were sub sampled from the parent Cape Grim air archive contained at high pressure (ca. 30 atmospheres) in large (35 dm³) stainless steel vessels. In all cases the air samples were collected under baseline (clean marine air) conditions. Details of the Cape Grim air archive and sampling techniques have been reported elsewhere (Oram et al., 1995; Fraser et al., 1999). Analyses were carried out using gas chromatography separation followed by detection with an EBE configuration triple-sector mass spectrometer (Waters/Micromass “Autospec”). Two slightly different techniques were employed using different capillary columns in the case of the Cape Grim air samples. In the first instance samples were preconcentrated on a glass bead trap cooled to around -185°C and separated on a Chrompack KCl-passivated alumina PLOT column. In the second case the samples were preconcentrated on a Hayesep D-packed trap cooled to -78°C. The two techniques are described in detail in Oram et al. (2011). Samples from NEEM and CARIBIC (see below) were exclusively analysed using the GasPro column. Analytical uncertainties were assessed from the reproducibility of duplicate samples and frequent calibration standard measurements.

To determine a corresponding time trend of SF₅CF₃ in the Northern Hemisphere we also made measurements on air pumped from deep snow (firn) in Northern Greenland (the North Greenland Eemian Ice Drilling project, NEEM; <http://www.neem.nbi.ku.dk>), which contains air dating from the early 20th century (Buizert et al., 2011). The sampling for trace gases at NEEM is described in Buizert et al. (2011) and Laube et al. (2010).

ACPD

12, 869–885, 2012

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Finally, measurements were also made on 49 air samples collected from a Lufthansa aircraft during four flights at cruising altitudes of 8.6 to 12.2 km between Frankfurt, Germany and Cape Town/Johannesburg, South Africa in the period October 2009 to March 2011 in the framework of the CARIBIC project (Brenninkmeijer et al., 2007; <http://www.caribic-atmospheric.com>).

2.2 2-D atmospheric modelling

A 2-D global atmospheric chemistry-transport model was used to derive “top-down” emission estimates of SF₅CF₃ from the measurements made at Cape Grim. The model grid is divided into 24 equal area, longitudinally-averaged bands and has 12 vertical layers each of 2 km depth. The model has been described elsewhere (Hough 1989, 1991) and has been used to study the temporal behaviour and global distribution of other halocarbons (Fraser et al., 1999; Reeves et al., 2005; Laube et al., 2010; Oram et al., 2011). The latitudinal distribution of emissions is based on an even split between the two main production sites of PFOS by 3M, Decatur, Alabama (34° N, 87° W) and Antwerp, Belgium (51° N, 4° E) (Paul et al., 2009). An atmospheric lifetime of 890 years was assumed for SF₅CF₃ (Montzka et al., 2010; Miller et al., 2002), and the annual emissions in the model were adjusted until the predicted concentrations matched the Cape Grim observations.

2.3 Modelling trace gas transport in firn

The relationship between past atmospheric time trends and concentration profiles in firn can be established using models of trace gas transport in firn, which is dominated by diffusion but is also subject to gravitational fractionation, firn sinking and gas trapping effects. Here we use a direct model of transport in firn as described by Witrant et al. (2011), which includes the main modelling principles proposed by Rommelaere et al. (1997), Fabre et al. (2000) and Martinerie et al. (2009). We use a firn diffusivity for the conditions at NEEM that has been evaluated using nine reference gases: CO₂,

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CH₄, SF₆, HFC-134a, CH₃CCl₃, ¹⁴CO₂, CFC-11, CFC-12 and CFC-113 as described in greater detail in Buizert et al. (2011). The diffusion coefficient for SF₅CF₃ was taken to be the same as that used in our earlier study (i.e. 0.0562 cm² s⁻¹, or 0.467 relative to that of CO₂; Sturges et al., 2000).

3 Results and discussion

The Cape Grim measurements are compiled in Fig. 1 as red diamonds for the measurements made with the GasPro capillary column, and green dashes for those made with the AI-PLOT capillary column. There is excellent agreement between the two sets of analyses for the overlapping time periods. The AI-PLOT measurements, however, had somewhat lower precision and were more scattered. The reason for this is not entirely clear (it might relate to slightly higher column bleed) as other gases showed generally excellent precision by the same method (e.g., Worton et al., 2007). Also shown, as blue circles, are the measurements from the original SF₅CF₃ paper. These were all from measurements in Antarctic firn air (Dome Concordia) and the dates have been estimated from co-measurements of SF₆ as described in Sturges et al. (2000). The tabulated dates so determined can be found at the Carbon Dioxide Information Analysis Center (<http://cdiac.ornl.gov/trends/otheratg/sturges/sturges.html>). The Dome C measurements usefully extend the southern hemispheric trend to dates before that of first Cape Grim measurements. The final set of measurements shown in Fig. 1 is the means of the measurements from the flights of the CARIBIC aircraft in 2009, 2010 and 2011 in the upper troposphere and lower stratosphere. These agree very well with the last few years of SF₅CF₅ measurements at Cape Grim, which is discussed further below.

Due to the greater scatter of the Cape Grim AI-PLOT measurements, we have based our subsequent model and curve fitting on the Cape Grim GasPro measurements, but we have also included the Dome C measurements prior to 1978 to define the earlier part of the trend. A five-parameter sigmoid expression (SPSS, now Systat Software, SigmaPlot Version 7) has been used to fit a smoothed line to the points (dashed blue-

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



green line), as this allows an envelope of uncertainty to be plotted based on the ± 1 sigma deviation of the fit (dashed black lines). The 2-D atmospheric model output for the surface grid box for the latitude of Cape Grim is shown as a solid red line. This is barely distinguishable from the sigmoidal line fit, and falls entirely within the deviation of the latter, as can be better seen in the expanded panel (inset). We therefore judge that the model fit represents well the observed trend.

The final curve plotted in Fig. 1, shown as a blue broken line, is a calculation of the SF₅CF₃ abundance expected had it exhibited the same relative molar abundance to SF₆ as was assumed in the modelling study from our original paper, i.e. a ratio to SF₆ of 3.05×10^{-2} . After 2002 it is clear that the apparent relationship between SF₅CF₃ and SF₆ began to diverge, with the latter continuing to grow whilst the SF₅CF₃ abundance began to stabilise. It can, therefore, be concluded that the emission of SF₅CF₃ is not linked to that of SF₆, contrary to our original hypothesis.

The measurements in Fig. 1 demonstrate the absence of significant emissions of SF₅CF₃ prior to the late 1960s, which is consistent with large-scale manufacture of PFOS having begun after 3M acquiring the patent for electrochemical fluorination in the 1950s. The first evidence for an unambiguous decrease in the atmospheric growth rate appears to have been around 2000, since which time the growth rate has declined rapidly. For the model to agree with the sigmoidal line fit, it was necessary to continue a small emission to the atmosphere after 2003 (see below). However, a model run (not shown) in which emissions were allowed to continue until 2003, and were then reduced to zero thereafter, resulted in a small decline in SF₅CF₃ (less than 1 % or 0.001 ppt) between 2003 and 2011, but which still fell within the uncertainty of the measurements. Equally, a longer atmospheric lifetime would imply emissions closer to zero, as noted in a sensitivity test with a lifetime of 3000 yr (not shown).

The observed trend is broadly consistent with Rosiek et al. (2007) who reported a continuing upward trend between 2001 and 2003 in Krakow, Poland, and an apparent subsequent slow-down in the growth rate of SF₅CF₃ reported by Busenberg and Plummer (2008) from various sites in North America.

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The annual emissions used in the atmospheric model to reproduce the observed atmospheric abundance of SF₅CF₃ are shown in Fig. 2. Emissions rose year-on-year from the 1960s to the 1990s, peaking around 1996 at around 300 t before progressively declining. The decline was very rapid from about 2000 to 2003. As discussed above, we cannot confidently distinguish the emissions shown after 2003 (around 10 t yr⁻¹) from zero.

The same model run can also be used to produce a temporal trend for the latitude and elevation relevant to the Greenland firn air sampling site. This trend can then be used as input to the firn diffusion model to derive an expected depth profile for SF₅CF₃. This is shown in Fig. 3 (line) along with the actual measurements made in the firn air (symbols). There is very good agreement suggesting excellent compatibility between the two sets of measurements and further validation of the modelled atmospheric distribution of SF₅CF₃.

The model computed no significant latitudinal gradient, and also no altitudinal gradient up to the tropopause, for the latter years (2005 onwards). This is confirmed by the agreement between the CARIBIC measurements in 2009, 2010 and 2011 in the upper troposphere/lower stratosphere, and the ground level measurements from NEEM (albeit made at about 3 km) and from Cape Grim (Fig. 1). Figure 4 further illustrates that, by the date of the aircraft flights, there was no evidence of a latitudinal gradient in SF₅CF₃, which also confirms the absence of further significant emissions to the atmosphere, such that the mixing ratio of this gas had equilibrated throughout the global atmosphere.

3M stated that a phase-out of PFOS manufacture was decided upon in 2000 (statement on “3M’s Phase Out and New Technologies” at http://solutions.3m.com/wps/portal/3M/en_US/PFOS/PFOA/Information/phase-out-technologies/), but also that overall greenhouse gas emissions (not just SF₅CF₃) had been reduced 40% since 1995 (Santoro, 2000). The PFOS phase-out was supposedly completed in 2002 (see “statement” above on 3M’s web site). Our results confirm that there was indeed a rapid decline in emissions between 2000 and 2003, although the uncertainties in

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

our observations preclude us from determining whether emissions actually ceased in 2003. Nevertheless, these findings strongly suggest that SF₅CF₃ emissions are indeed uniquely associated with such fluorochemical production. 3M state that associated fluorochemical production continued via a subsidiary (Dyneon), albeit using perfluorooctanoic acid (PFOA) not PFOS, but this too ceased in 2008 (see 3M “statement”). A number of companies worldwide continue to manufacture related perfluorocarboxylates (Rennner, 2006) but, it appears from our observations, without incurring substantial emissions of SF₅CF₃. 3M themselves have subsequently switched to processes that produce perfluorobutane sulfonate (PFBS) instead of the more toxic PFOS (3M “statement” and Renner, 2006).

4 Conclusions

The combined dataset from the Cape Grim atmospheric air archive, northern and southern hemispheric firn air, and interhemispheric aircraft flights results in a consistent atmospheric trend for SF₅CF₃ from the mid-20th century to 2011. Although the maximum atmospheric abundance of this molecule has never exceeded more than a small amount (0.15 ppt), and thus currently contributes only minutely to climate forcing (Sturges et al., 2000), it (along with the other fully-fluorinated species referred to in the Introduction) represents an essentially irreversible change to the composition of the Earth’s atmosphere and to the radiative forcing of the climate. The present cessation (or near-cessation), and presumed future avoidance of further emissions of this gas, is therefore to be welcomed.

Acknowledgement. WTS, DEO and JCL acknowledge the UK Natural Environment Research Council (NERC) for awards NE/F021194/1 (NEEM), R8/H12/38/012 (NCAS Composition) and NE/F015585/1 (NERC Fellowship), respectively. CH and MJN were both supported by NERC Doctoral Studentships. We thank Michael Leist and the staff of the Cape Grim Baseline Station and the CSIRO GASLAB for collection and maintenance of the Cape Grim air archive. The Cape Grim programme has previously received funding from the former UK Government

Department of Environment (GA/4/2/57). The CARIBIC programme has previously received funding from UK NERC (NER/T/S/2000/01040 and GST/02/2218) and the European Union (EVK2-GT2001-00101). We are grateful to the entire CARIBIC team and all its partners. We are indebted to Dr Jakob Schwander of the Physics Institute at the University of Berne, Switzerland for collecting the NEEM firn air samples. NEEM is directed and organized by the Center of Ice and Climate at the Niels Bohr Institute and US NSF, Office of Polar Programs. It is supported by funding agencies and institutions in Belgium (FNRS-CFB and FWO), Canada (GSC), China (CAS), Denmark (FIST), France (IPEV, CNRS/INSU, CEA and ANR), Germany (AWI), Iceland (RannIs), Japan (NIPR), Korea (KOPRI), The Netherlands (NWO/ALW), Sweden (VR), Switzerland (SNF), UK (NERC) and the USA (US NSF, Office of Polar Programs).

References

- Brenninkmeijer, C. A. M., Crutzen, P., Boumard, F., Dauer, T., Dix, B., Ebinghaus, R., Filippi, D., Fischer, H., Franke, H., Frieß, U., Heintzenberg, J., Helleis, F., Hermann, M., Kock, H. H., Koepfel, C., Lelieveld, J., Leuenberger, M., Martinsson, B. G., Miemczyk, S., Moret, H. P., Nguyen, H. N., Nyfeler, P., Oram, D., O'Sullivan, D., Penkett, S., Platt, U., Pucek, M., Ramonet, M., Randa, B., Reichelt, M., Rhee, T. S., Rohwer, J., Rosenfeld, K., Scharffe, D., Schlager, H., Schumann, U., Slemr, F., Sprung, D., Stock, P., Thaler, R., Valentino, F., van Velthoven, P., Waibel, A., Wandel, A., Waschitschek, K., Wiedensohler, A., Xueref-Remy, I., Zahn, A., Zech, U., and Ziereis, H.: Civil Aircraft for the regular investigation of the atmosphere based on an instrumented container: The new CARIBIC system, *Atmos. Chem. Phys.*, 7, 4953–4976, doi:10.5194/acp-7-4953-2007, 2007.
- Buizert, C., Martinerie, P., Petrenko, V. V., Severinghaus, J. P., Trudinger, C. M., Witrant, E., Rosen, J. L., Orsi, A. J., Rubino, M., Etheridge, D. M., Steele, L. P., Hogan, C., Laube, J. C., Sturges, W. T., Levchenko, V. A., Smith, A. M., Levin, I., Conway, T. J., Dlugokencky, E. J., Lang, P. M., Kawamura, K., Jenk, T. M., White, J. W. C., Sowers, T., Schwander, J., and Blunier, T.: Gas transport in firn: multiple-tracer characterisation and model intercomparison for NEEM, Northern Greenland, *Atmos. Chem. Phys. Discuss.*, 11, 15975–16021, doi:10.5194/acpd-11-15975-2011, 2011.
- Busenberg, E. and Plummer, L. N.: Dating groundwater with trifluoromethyl sulfurpentafluoride

ACPD

12, 869–885, 2012

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(SF₅CF₃), sulfur hexafluoride (SF₆), CF₃Cl (CFC-13), and CF₂Cl₂ (CFC-12), *Water Resour. Res.*, 44, W02431, doi:10.1029/2007WR006150, 2008.

Fabre, A., Barnola, J.-M., Arnaud, L., and Chappellaz, J.: Determination of gas diffusivity in polar firn: Comparison between experimental measurements and inverse modelling, *Geophys. Res. Lett.*, 27, 557–560, 2000.

Hough, A. M.: The development of a two-dimensional global tropospheric model, 1, the model transport, *Atmos. Environ.*, 23, 1235–1261, 1989.

Hough, A. M.: Development of a two-dimensional global tropospheric model: model chemistry, *J. Geophys. Res.*, 96, 7325–7362, 1991.

Huang, L., Zhu, L., Pan, X., Zhang, J., Ouyang, B., Hou, H.: One potential source of the potent greenhouse gas SF₅CF₃: The reaction of SF₆ with fluorocarbon under discharge, *Atmos. Environ.*, 39, 9, 1641–1653, 2005.

Laube, J. C., Martinerie, P., Witrant, E., Blunier, T., Schwander, J., Brenninkmeijer, C. A. M., Schuck, T. J., Bolder, M., Röckmann, T., van der Veen, C., Bönisch, H., Engel, A., Mills, G. P., Newland, M. J., Oram, D. E., Reeves, C. E., and Sturges, W. T.: Accelerating growth of HFC-227ea (1,1,1,2,3,3,3-heptafluoropropane) in the atmosphere, *Atmos. Chem. Phys.*, 10, 5903–5910, doi:10.5194/acp-10-5903-2010, 2010.

Levin, I., Naegler, T., Heinz, R., Osusko, D., Cuevas, E., Engel, A., Ilmberger, J., Langenfelds, R. L., Neining, B., Rohden, C. v., Steele, L. P., Weller, R., Worthy, D. E., and Zimov, S. A.: The global SF₆ source inferred from long-term high precision atmospheric measurements and its comparison with emission inventories, *Atmos. Chem. Phys.*, 10, 2655–2662, doi:10.5194/acp-10-2655-2010, 2010.

Martinerie, P., Nourtier-Mazauric, E., Barnola, J.-M., Sturges, W. T., Worton, D. R., Atlas, E., Gohar, L. K., Shine, K. P., and Brasseur, G. P.: Long-lived halocarbon trends and budgets from atmospheric chemistry modelling constrained with measurements in polar firn, *Atmos. Chem. Phys.*, 9, 3911–3934, doi:10.5194/acp-9-3911-2009, 2009.

Miller, T. M., Arnold, S. T., Viggiano, A. A., Knighton, W. B.: Electron attachment of SF₅CF₃ (296–563 K) and calculations of the neutral and anion thermochemistry, *J. Chem. Phys.*, 116, 6021–6027, 2002.

Montzka, S. A., Reimann, S. (Coordinating Lead Authors), Engel, A., Krüger, K., O'Doherty, S., Sturges, W., Blake, D., Dorf, M., Fraser, P., Froidevaux, L., Jucks, K., Kreher, K., Kurylo, M., Mellouki, A., Miller, J., Nielsen, O.-J., Orkin, V., Prinn, R., Rhew, R., Santee, M., Stohl, A., and Verdonik, D.: Scientific Assessment of Ozone Depletion: 2010, *Global Ozone Research*

and Monitoring Project, Report No. 52, 516 pp., Chapter 1, Ozone-Depleting Substances (ODSs) and Related Chemicals, World Meteorological Organization, Geneva, Switzerland, 2011.

5 Mühle, J., Ganesan, A. L., Miller, B. R., Salameh, P. K., Harth, C. M., Grealley, B. R., Rigby, M., Porter, L. W., Steele, L. P., Trudinger, C. M., Krummel, P. B., O'Doherty, S., Fraser, P. J., Simmonds, P. G., Prinn, R. G., and Weiss, R. F.: Perfluorocarbons in the global atmosphere: tetrafluoromethane, hexafluoroethane and octafluoropropane, *Atmos. Chem. Phys.*, 10, 5145–5164, doi:10.5194/acp-10-5145-2010, 2010.

10 Oram, D. E., Reeves, C. E., Fraser, P. J., and Penkett, S. A.: Measurements of HCFC-142b and HCFC-141b in the Cape Grim air archive: 1978–1993, *Geophys. Res. Lett.*, 22, 2741–2744, 1995.

15 Oram, D. E., Mani, F. S., Laube, J. C., Newland, M. J., Reeves, C. E., Sturges, W. T., Penkett, S. A., Brenninkmeijer, C. A. M., Röckmann, T., and Fraser, P. J.: Long-term tropospheric trend of octafluorocyclobutane (c-C4F8 or PFC-318), *Atmos. Chem. Phys.*, 12, 261–269, doi:10.5194/acp-12-261-2012, 2012.

Paul, A. G., Jones, K. C., and Sweetman, A. J.: A first global production, emission and environmental inventory for perfluorooctane sulfonate, *Environ. Sci. Technol.*, 43, 386–392, 2009.

20 Reeves, C. E., Sturges, W. T., Sturrock, G. A., Preston, K., Oram, D. E., Schwander, J., Mulvaney, R., Barnola, J.-M., and Chappellaz, J.: Trends of halon gases in polar firn air: implications for their emission distributions, *Atmos. Chem. Phys.*, 5, 2055–2064, doi:10.5194/acp-5-2055-2005, 2005.

Renner, R.: The long and the short of perfluorinated replacements, *Environ. Sci. Technol.*, 40, 12–13, doi:10.1021/es062612a, 2006.

25 Rommelaere, V., Arnaud, L., and Barnola, J.: Reconstructing recent atmospheric trace gas concentrations from polar firn and bubbly ice data by inverse methods, *J. Geophys. Res.*, 102, 30069–30083, 1997.

Rosiek, J., Lasa, J., Sliwka, I., and Róžański, K.: Application of GC and Modulated ECD for the Determination of SF₅CF₃ Mixing Ratios in the Atmosphere, *Chem. Anal. (Warsaw)*, 52, 235, 2007.

30 Santoro, M. A.: Clarifying the SF₅CF₃ record, *Science*, 290, 935, 2000.

Solovev, S., Palmentieri, A., Potekhina, N. D., and Madey, T. E.: Mechanism for electron-induced SF₅CF₃ formation in condensed molecular films, *J. Phys. Chem. C*, 111, 18271–18278, 2007.

ACPD

12, 869–885, 2012

Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sturges, W. T., Wallington, T. J., Hurley, M. D., Shine, K. P., Sihra, K., Engel, A., Oram, D. E., Penkett, S. A., Mulvaney, R., Brenninkmeijer, C. A. M.: A potent greenhouse gas identified in the atmosphere: SF₅CF₃, *Science*, 289, 611–613, 2000.

Weiss, R. F., Mühle, J., Salameh, P. K., and Harth, C. M.: Nitrogen trifluoride in the global atmosphere, *Geophys. Res. Lett.*, 35, L20821, doi:10.1029/2008GL035913, 2008.

Witrand, E., Martinerie, P., Hogan, C., Laube, J. C., Kawamura, K., Capron, E., Montzka, S. A., Dlugokencky, E. J., Etheridge, D., Blunier, T., and Sturges, W. T.: A new multi-gas constrained model of trace gas non-homogeneous transport in firn: evaluation and behavior at eleven polar sites, *Atmos. Chem. Phys. Discuss.*, 11, 23029–23080, doi:10.5194/acpd-11-23029-2011, 2011.

Worton, D. R., Sturges, W. T., Gohar, L. K., Shine, K. P., Martinerie, P., Oram, D. E., Humphrey, S. P., Begley, P., Gunn, L., Barnola, J.-M., Schwander, J., and Mulvaney, R.: Atmospheric trends and radiative forcings of CF₄ and C₂F₆ inferred from firn air, *Environ. Sci. Technol.*, 41, 2184–2189, doi:10.1021/es061710t, 2007.

ACPD

12, 869–885, 2012

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Emissions halted of the potent greenhouse gas SF₅CF₃

W. T. Sturges et al.




Fig. 1. The atmospheric record of SF₅CF₃ from measurements made in air collected at Cape Grim, Tasmania, in firn air collected at Dome Concordia, Antarctica, and in air collected between 8.6 and 12.2 km and 47° N–28° S from the CARIBIC aircraft. The dashed blue-green line is a least mean squares optimisation fit to a subset of the Cape Grim (see text) and Dome Concordia measurements (pre-1978) with the uncertainty of the fit shown as black dashed lines. The red line (almost indistinguishable from the blue-green line) is the output mixing ratio from the atmospheric model run for the location of Cape Grim using the emissions shown in Fig. 2. The blue broken line shows the “predicted” SF₅CF₃ assuming a constant ratio to SF₆ (see text). The inset figure is an expansion of the main graph between the dates 2000–2011.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.




Fig. 2. The model-derived annual and cumulative emissions of SF₅CF₃ used to produce the atmospheric model fit scenario presented in Fig. 1 (the red line in Fig. 1).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.




Fig. 3. The observed amounts (symbols) of SF₅CF₃ in samples collected at different depths of firn at the NEEM drilling site, Greenland. The purple line is the result of taking the long-term trend of the gas in the atmospheric model domain box appropriate for NEEM, based on the best fit to Cape Grim (red line in Fig. 1), and used as input to the firn diffusion model.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Emissions halted of
the potent
greenhouse gas
SF₅CF₃**

W. T. Sturges et al.




Fig. 4. Measurements of SF₅CF₃ from whole air samples collected from four interhemispheric CARIBIC flights flying at 8.6–12.2 km, as a function of latitude.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)