

Interactive comment on “Impact of the marine atmospheric boundary layer on VSL abundances in the eastern tropical and subtropical North Atlantic Ocean” by S. Fuhlbrügge et al.

R. Hossaini

chm3rh@leeds.ac.uk

Received and published: 30 January 2013

In the introduction the authors note that “oceanic emissions of VSLs might change in the future and in connection with an altering efficiency of the atmospheric upward transport might lead to significant future changes of the halogen budget of the troposphere/lower stratosphere”

I would like to draw the authors attention to the most recent study on future stratospheric VSLs loading and to note, along with potential transport changes, changes to the tropospheric oxidizing capacity are likely to be important also.

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



Hossaini, R., et al. (2012), Modelling future changes to the stratospheric source gas injection of biogenic bromocarbons, *Geophys. Res. Lett.*, 39, L20813, doi:10.1029/2012GL053401.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 12, 31205, 2012.

ACPD

12, C12158–C12159,
2013

Interactive
Comment

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)

