
Interactive
Comment

Interactive comment on “Volcanic SO₂ fluxes derived from satellite data: a survey using OMI, GOME-2, IASI and MODIS” by N. Theys et al.

Anonymous Referee #2

Received and published: 6 February 2013

General comments:

This paper presented a survey of four techniques for deriving volcanic SO₂ flux from satellite measurement of dispersed volcanic plumes, and provided an informative summary on the advantages and weaknesses of each technique. The authors analyzed the SO₂ column density maps of three selected (to represent different types) volcanic eruptions from multiple satellite sensors by applying some or all these loading-to-flux conversion techniques. They provided clear and easy-to-understand comparisons among the many combinations of volcanic events, satellite sensors, and conversion techniques, to illustrate main findings of this paper: SO₂ emission flux can be derived with < 50% error using just once a day full coverage of a volcanic plume, and the measured mass and the derived flux from the four satellite sensors (OMI, GOME2, IASI,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



and MODIS) are consistent even with the differences in measurement sensitivity, pixel sizes (spatial resolutions), and time of observations. These comparisons also highlighted the importance of volcanic plume height information in improving the accuracy of satellite SO₂ data. This is a well-written paper for publication in ACP.

Since this paper emphasized that these techniques mainly worked for 'dispersed and large-scale' plumes, the authors probably need to include the area sizes (in square kilometer) for the sample cases shown in this paper, and discuss how they are selected and if automatic processing is possible.

Also this paper needs to include the description of the SO₂ mass (or the total burden) calculation for a plume, whether a threshold SO₂ value is used to select the pixels or all the pixels inside a bounding box were included. How to treat the background SO₂ pixels may have a significant impact on the total burden, hence the derived flux, especially when a plume is well dispersed.

Specific comments:

1. Figures 5, 10, and 12 are too small see without enlargement.
2. Rightmost panel of figure 12 seems to be mislabeled.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 12, 31349, 2012.

ACPD

12, C12337–C12338,
2013

Interactive
Comment

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)

