

Interactive comment on “Tropospheric bromine chemistry: implications for present and pre-industrial ozone and mercury” by J. P. Parrella et al.

J. P. Parrella et al.

parrella@fas.harvard.edu

Received and published: 18 April 2012

The authors would like to thank P. Pongprueksa for bringing the updated rate constants to our attention. Here we will report the impact of the new Goodsite et al. (2012) rates on our results. The final version of our paper will include these updated results.

1. The present-day Hg(0) lifetime = 0.58 years
2. The pre-industrial Hg(0) lifetime = 0.35 years

Updating the kinetics does not affect our conclusions. Our estimated increase in the Hg(0) lifetime from pre-industrial to present is now 66%. And as a minor point, changes in the global Hg(0) lifetime with respect to bromine are now linear (instead of slightly

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



supralinear as we reported before the update), and this is due to the slower rate of HgBr thermal dissociation when using the Goodsite et al. (2012) kinetics.

Reference:

Goodsite M. E., Plane J. M. C., and Skov H., 2012, Correction to A Theoretical Study of the Oxidation of Hg⁰ to HgBr₂ in the Troposphere, Environ. Sci. Technol., DOI:10.1021/es301201c.

[Interactive comment on Atmos. Chem. Phys. Discuss., 12, 9665, 2012.](#)

ACPD

12, C1652–C1653, 2012

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C1653

