Atmos. Chem. Phys. Discuss., 12, C3046–C3047, 2012 www.atmos-chem-phys-discuss.net/12/C3046/2012/ © Author(s) 2012. This work is distributed under the Creative Commons Attribute 3.0 License.



ACPD

12, C3046-C3047, 2012

Interactive Comment

Interactive comment on " α -pinene photooxidation under controlled chemical conditions – Part 1: Gas-phase composition in low- and high-NO $_x$ environments" by N. C. Eddingsaas et al.

Anonymous Referee #2

Received and published: 29 May 2012

The manuscript reports high quality chamber experiments on photooxidation of α -pinene with various NOx concentrations. The gas phase composition was monitored by a CIMS system. Under low-NOx conditions α -pinene hydroxy hydroperoxides and pinonaldehyde are reported as major products and their formation pathways are discussed with focus on the RO2+HO2 reactions. A kinetic model was developed to calculate OH concentrations and branching ratios. Under high-NOx conditions pinonaldehyde , organonitrates and the isomerization products of alkoxy radicals are observed as the major products with implications on the understanding of OH radical regeneration. It was also found that especially the concentrations of PAN compounds are depending on the actual NO2 level. Consequently, the paper fits to the scope of ACP and it will

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



contribute to understand the gas phase chemistry of biogenic VOCs. The paper is well written and conclusions made are sound. We recommend publishing the manuscript in ACP, however, have some minor comments on the manuscript.

Minor revisions: Line 307: From a physico-chemical point of view, pinonaldehyde is not really nonpolar, it just contains no acidic hydrogen, therefore the positive mode needs to be used.

In Equations (4) + (5) O2 is missing on the right side of the equations

Line 371: α -pinene oxide has a relatively low polarity but it is again not nonpolar (see above). Arguing about this kind of ion formation should discuss gas phase acidities or gas phase basicities (or proton affinities).

Line 396f: "... forming an acyl peroxy radical" after O2 addition.....?

Line 405 and 422: Did the authors measure synthesized reference peroxyacids to be absolutely sure about these products or were they "just" tentatively identified based on the interpretation of the mass spectra?

Figures 1+2: To avoid confusion of the reader between molecular ions of complexes the authors should consider to mentioned the ion compositions (complexes of CF3 or HF) in the captions again.

Figure 6 and related text: Acronyms for the compounds would facilitate the reading.

Interactive comment on Atmos. Chem. Phys. Discuss., 12, 6447, 2012.

ACPD

12, C3046-C3047, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

