

Interactive comment on “Implementation of dust emission and chemistry into the Community Multiscale Air Quality modeling system and initial application to an Asian dust storm episode” by K. Wang et al.

Anonymous Referee #2

Received and published: 23 July 2012

General comments

This paper presents the implementation of two online dust emission parameterization schemes, the replacement of the thermodynamic module ISORROPIA with the newer version ISORROPIA II and also the inclusion of heterogeneous reactions of gaseous species at the surface of dust particles in the CMAQ model (US Environmental Protection Agency Community Multiscale Air Quality). The new system has been applied to simulate a dust storm event in Asia on April 2001, performing 9 simulations with different configurations, in order to reveal the impact of the new development to several

C4945

atmospheric chemical species and the improvement succeeded by this development. A small section is also devoted to the influence of Asian pollution on the air quality in the US. The title of the manuscript reflects the contents of the paper and is consider sufficient.

In terms of scientific quality and significance, the purpose of this work is in line with the current state-of-the-art in the field of atmospheric chemistry modelling research. The inclusion of the complex interactions between gases and aerosols into a regional atmospheric chemistry model is a significant effort towards understanding the uncertainties and the behaviour of aerosols in the atmospheric system. I am in favour of such research efforts and my comments on the manuscript are written with this in mind.

The authors have made a large effort to analyze and discuss a lot of aspects of the new development in the CMAQ model, from meteorology to chemical reactions. This has resulted in a text rather difficult to read with several inconsistencies that I will describe in detail in the specific comments section. Several errors have been made in the description of the dust emission schemes, and the authors must carefully correct them as this part of the manuscript is the core part of the new development. The effects of the new processes in the CMAQ model are multidimensional and the message from this work must be clear, stressing out the solid conclusions and the remaining uncertainties. The specific comments will help to clarify some issues that have arisen during the review process and strengthen the quality of the paper through recommendations on the presentation and the discussion of the results.

Specific comments

Abstract

Page 13459, lines 14-15: The total dust emission amount indicated in the abstract comes from the Zender or the Westphal scheme? Please include this information here.

Page 13459, lines 25-26: The increase of gases and aerosols due to Asian pollution

C4946

mainly affects the Western US as indicated in Section 5.4. Please include this information in the abstract.

Introduction

Page 13461, lines 3-5: I suggest you include the recent publication of Crowley et al. (2010) (see the full reference in the end of this document) where a full assessment of the heterogeneous processes on surfaces of solid particles is presented.

Page 13461, lines 7, 8: The modelling work of Cheng et al. (2008) and Huneus et al. (2011) did not include any chemistry on dust particles. You should not include these publications in the same list denoting the studies of heterogeneous chemistry on dust particles (line 9). The references on Tang et al. (2004), Fairlie et al. 2010 (already in your ref list), Astitha et al. (2010) could be part of this list.

Section 2.1

Page 13464, lines 8-17: I would suggest that the authors do not refer to the second emission scheme as the “Zender scheme” when discussing the physical concept and equations. Zender et al. (2003) clearly state that “We generally follow the microphysical and micrometeorological approach to dust mobilization developed by Marticorena and Bergametti [1995]”. The original work on the horizontal dust flux is done by White (1979) and also Kawamura (1964) and it should be mentioned here.

Page 13464, line 26: Please briefly explain how the Heaviside function H is used in the context of the dust emission flux (the physical meaning).

Page 13465, line 2: “ $RF=1.0$ ” will result in zero dust emission flux (Equation 1) for barren land? Please correct the values of RF used in Equation 1.

Page 13465, lines 2-9: The parameter EF , which is assumed to characterize the fraction of erodible land, is constant everywhere in the domain? Does it change depending on the soil properties in each arid area (roughness elements etc)? If not, then this parameter is mainly a tuning factor rather than a measure of the erodibility of the land.

C4947

Page 13465, line 10: I strongly disagree that a constant threshold velocity can be valid for any study on dust emissions, either regional or global. The dust emission process occurs on very small spatial scales and is directly related to the soil properties (mineralogy, roughness elements, smooth surface) and particle sizes (diameter of the particles capable to initiate saltation).

Page 13465, Equation 2: The value 121 should be 1.21.

Page 13466, Equation 7: This equation must be wrong. The last terms “ $H(u^*-u^*)$ ” are not included in the dust flux formulation described in Marticorena and Bergametti (1995), Zender et al. (2003) and elsewhere. Furthermore, why did you use the parameter EF in this equation of the horizontal flux? In the calculation of the vertical flux (Equation 11) you already include the “source erodibility factor S ”. I don’t see any physical meaning in adding one more constant factor in Eq.7 rather than a possible double-counting of the same concept.

Page 13466, line 21: There is no u^*l in the previous equation of the dust horizontal flux. The authors use this parameter also in Eqs. 8 and 9, which should be replaced by u^*t .

Page 13467, Eqs 8 and 9: In Iversen and White (1982) some of the values used in the empirical relationship are different from the ones shown here (instead of 0.1666681 the value is 0.129 etc). Please explain why you changed these values or refer to the appropriate publication. In addition, there is an error in Equation 9, at the last term of the right side ($gD2.5$ belongs to the denominator). Please correct it accordingly.

Page 13467, Eq 10: The Reynolds number calculated in equation 10 is a constant value since the diameter of the particle is fixed to $75\mu m$. This means that the threshold friction velocity is also a fixed number and there is no need for both Eqs 8 and 9. The use of Eq.10 could be employed only at the 1st time-step when there is no initial value for u^*t , and then $Re^*t = u^*t D/\nu$ providing a small change in the calculation of the threshold friction velocity.

C4948

Page 13467, line 12: Maybe you mean that u^*t is calculated from Eq2 with u^*t_l from Eqs 8 or 9 and then used in Eq.7? Of course if the previous comment is true, there is no need for both equations 8 and 9. If my comment is precise then the dust flux calculated by Zender's approach is subject to correction for soil moisture as done for the Westphal. If so, please indicate that explicitly in the text.

Page 13467, line 17: How do you calculate the "source erodibility factor" S (from an input table, a database, online calculation)? Add an explanation in the text.

Page 13467, Eq 12: This equation must be wrong. It is the same as used in Zender et al. (2003) but a careful comparison with the original equation in Marticorena and Bergametti (1995) will reveal the differences: The original equation is $\log(F/G)=0.134(\% \text{clay})-6$ coming from linear fitting of data in Marticorena and Bergametti (1995). This means that $\alpha=F/G=10^{(0.134(\% \text{clay})-6)}$ and for 10% of clay the sandblasting efficiency will be $\sim 2.18 \times 10^{-5} \text{ cm}^{-1}$ (shown in Fig.4 of Marticorena and Bergametti (1995)). For $M_{\text{clay}}=0.1$, equation 12 gives $\alpha=2.18 \times 10^{-3} \text{ cm}^{-1}$, two orders of magnitude different. If the authors have used this equation in their calculations they should carefully check the calculated sandblasting coefficient α with published values and the original work.

Page 13468, line 10: There is no Eq.14 in the text.

Section 2.2 (Heterogeneous chemistry)

A recent publication by Crowley et al. (2010) is dedicated to the heterogeneous processes on surfaces of solid particles present in the atmosphere, for which uptake coefficients and adsorption parameters have been presented on the IUPAC (website in 2009). In this publication, data of uptake coefficients is evaluated and a recommendation is made for each reaction, based on several arguments presented in the paper. A reference to this work must be included in the text, as it is a recent study based on experimental data and it is closely related to this publication. Some of the uptake coefficients suggested by Crowley et al. (2010) for the uptake on mineral dust particles are quite different from the values used herein. A brief discussion on the consequences

C4949

from using lower or higher values for the gamma coefficients compared to the ones in Table 2 would be very informative in the text.

Section 2.3 (Crustal species-ISORROPIA II)

The mineralogy of the soil that prescribes the chemical composition of atmospheric dust particles is very important and still not available for all the arid areas of the world. The inclusion of ISORROPIA II is a very good step towards improving the thermodynamics of the aerosols in the model.

1. How do you treat the products from the heterogeneous reactions in the model, i.e. the sulfates produced on dust are added to the sulfates produced from gas to particle conversion or they are kept separately in the thermodynamics and the other aerosol processes in the model?

2. The ratios between crustal species and dust (page 13470, line 17) is assigned to the dust emitted flux or the concentration?

Section 3 (Model configuration)

Page 13471, line 24: Please be more specific on the meteorological setup chosen for the 9 simulations described in Table 3. Is it WRF with FNL and nudging?

Page 13471, line 25: The WRF and the CMAQ models use the same geographical projection or a necessary recalculation is performed?

Page 13472, line 11: Which dust emission scheme did the authors use for the simulations, Zender or Westphal? This is not stated anywhere in the next sections and the description of both schemes in Section 1 is not justified by the performed simulations or a discussion in the text.

Section 4 (Model evaluation)

Page 13476: The evaluation of the dust emission scheme used for the 9 simulations is missing from this section. Figure 2 with the monthly AOD plots is not adequate to form

C4950

a solid evaluation of the dust scheme. This is very crucial as the main developments of this study rely on the dust emission online scheme embedded in the model. I would strongly suggest two ways the authors can deal with that: 1) use the data of dust concentration measurements in Asian sites published in Cheng et al. (2008). Data for April 2001 is included in that publication (10 stations out of the 16 give values for April 2001). 2) Use the daily AOD values from the AERONET Level 2 products (~13 AERONET sites in Asia have AOD values for April 2001). Selecting stations close to the desert areas will exclude the influence from anthropogenic sources and give a clear picture of the model performance. The AOD statistics presented in Table 7, (a spatial and temporal average on the domain), include all species with the new development and cannot account as an evaluation of the dust scheme.

Section 5.2

Figure 6: The spatial differences between DUST and CRUST_ONLY show increase of H₂O₂ but the differences between DUST and DUST_HIGH_UPTAKE show decrease in H₂O₂. The same occurs for NO_x and the opposite for NO₃. This means that changing the uptake coefficients γ into higher values can have the opposite effect in some species, indicating the high uncertainty of the γ values. The authors should mention this in the text and try to elaborate on the reasons for this behaviour.

Conclusions

It is not clear if the 9 simulations were performed with the Zender or the Westphal scheme. Which of the two schemes performed better in terms of efficiently producing the dust distribution in the selected domain and which scheme would the authors recommend to the modellers using CMAQ/Dust in the future?

Technical corrections Page 13459, line 18: “aerosol optical depth than the default CMAQ...”

Page 13460, line 22: “treatments for dust blowing up or uplifting processes...” should

C4951

be rephrased to “parameterizations of dust emission processes...”.

Page 13461, line 11: “For example,...”.

Page 13465, line 22: “In the current version of the emission scheme, only the ...”

Page 13466, line 1: There is a missing prime symbol in w (threshold gravimetric soil moisture).

Page 13468, line 1: “the mass faction” should be “mass fraction”.

Page 13483, line 28: “...altitudes is much larger...”

Tables 4-7: The values “Data Number” and “Mean Observed” must be clearly stated and independent of the model configuration in the first column. In addition, in tables 4-6 the different data networks must be clearly separated. These actions will facilitate the proper reading of the tables.

Figure 2: Indicate the satellite that gave the monthly AOD plot (first row). Also the title of each plot does not correspond to the respective simulation (1st plot: should be MODIS AOD, if MODIS data is used etc).

Figures 3-9: It would benefit the reader if a brief sentence on the scope of each figure was included in the legend. For example: In Figure 6 the sentence “Effects of heterogeneous chemistry” could be added.

Figures in the supplement: The figures in the supplement need refinement as some titles are not shown and the borders of the plots are not uniform.

References

1. Astitha, M., and CoAuthors, 2010: Chemically aged and mixed aerosols over the Central Atlantic Ocean - potential impacts, Atmos. Chem. Phys., 10, 5797-5822, doi:10.5194/acp-10-5797-2010.
2. Crowley, J. N., M. Ammann, R. A. Cox, R. G. Hynes, M. E. Jenkin, A. Mellouki,

C4952

M. J. Rossi, J. Troe and T. J. Wallington: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V - heterogeneous reactions on solid substrates. *Atmospheric Chemistry and Physics* 10, 18, 9059-9223 (2010).

3. White, B. R.: Soil transport by winds on Mars, *J. Geophys. Res.*, 84, 4643–4651, 1979.

4. Kawamura, R. (1964) Study of sand movement by wind, *Hydraul. Eng. Lab. Tech. Rep.*, HEL-2-8, pp. 99–108, Univ. of Calif., Berkeley.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 12, 13457, 2012.