

***Interactive comment on “Evaluation of  
HO<sub>x</sub> sources and cycling using measurement –  
constrained model calculations in a 2 – methyl – 3 –  
butene – 2 –  
ol (MBO) and monoterpene (MT) dominated ecosystem” by S. Kim et al.***

**Anonymous Referee #2**

Received and published: 13 August 2012

Last week I came across the following new article which describes a new interference on OH radical measurements made using the CIMS technique. The interference is reported to cause a positive artifact in "terpene +O<sub>3</sub>" rich environments. In the light of those interesting results, it would be nice if the authors include a discussion of how strongly the OH concentrations used in the present work would be affected by the newly reported interference and what bearing it would have on the conclusions of the current study.

R. L. Mauldin III, T. Berndt, M. Sipilä, P. Paasonen, T. Petäjä, S. Kim, T. Kurtén, F.

C5654

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Stratmann, V.M. Kerminen & M. Kulmala , A new atmospherically relevant oxidant of sulphur dioxide, Nature, 488,193–196, doi:10.1038/nature11278, 2012.

---

Interactive comment on Atmos. Chem. Phys. Discuss., 12, 15945, 2012.

ACPD

12, C5654–C5655, 2012

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C5655

