

## ***Interactive comment on “Cyclobutyl methyl ketone as a model compound for pinonic acid to elucidate oxidation mechanisms” by A. P. Praplan et al.***

**A. P. Praplan et al.**

arnaud.praplan@psi.ch

Received and published: 10 September 2012

We thank U. Neuenschwander for taking the time to read and comment on this manuscript and for his helpful and constructive comments. Comments are portrayed in regular font style, reply in italic.

### **1 Reply to U. Neuenschwander**

In terms of presented mechanisms (i.e. elementary reaction steps), I feel that an important reaction could be added to Figs. 5 and 10 in the manuscript: Whereas the authors correctly ignore the direct reaction between ozone and peroxy radicals under

C6753

the current experimental conditions, and invoke radical cross-reactions being the origin of alkoxy radicals, the intermediate radical prior to "alkoxy B", being a 1,5-dicarbonyl 4-peroxy species, can alternatively undergo intramolecular oxidation of the aldehyde moiety. Similar 1,6-H shifts have been published elsewhere (see e.g. Neuenschwander and Hermans (2010); Zhu et al. (2007)). The estimated rate constant for that 1,6-H shift at 298 K is about  $100\text{ s}^{-1}$ , using a TST approximation based on a literature prefactor ( $A = 10^{10}\text{ s}^{-1}$ ; Zhu et al. (2007)) and an ab initio activation energy, as specifically calculated for this reaction, on an extensive gaussian basis set within the unrestricted B3LYP level of theory (45 kJ/mol; Neuenschwander (2012)). Note that this unimolecular rearrangement is favored both entropically (viz. molecular preorganization) and enthalpically (viz. oxidation of weak aldehyde C-H bond). Thus, I think that it should be added to the presented framework of elementary reaction steps.

*The authors are very grateful for this comment and calculations supporting the discussion about "non-traditional" reactivity. This 1,6-H shift for the 1,5-dicarbonyl 4-peroxy marked in orange in Fig. 5 is now included in the section 3.5 as an example supporting our assumption that this compound may play a key role due to possible intramolecular reactions.*

### **References**

- Neuenschwander, Ulrich, and Hermans, Ive: Autoxidation of  $\alpha$ -pinene at high oxygen pressure, Phys. Chem. Chem. Phys., 12, 10542–10549, doi:10.1039/c0cp00010h, 2010.  
Neuenschwander, Ulrich: DFT calculation on UB3LYP/6-311++G(df,pd)//UB3LYP/6-31G(d,p) including ZPE, unpublished results, 2012.  
Zhu, Li and Bozzelli, Joseph W. and Kardos, Lisa M.: Thermochemical properties,  $\Delta_f H^\circ(298)$ ,  $S^\circ(298)$ , and  $C_p^\circ(T)$ , for n-butyl and n-pentyl hydroperoxides and the alkyl and peroxy radicals, transition states, and kinetics for intramolecular hydrogen shift reactions of the peroxy radicals, J. Phys. Chem. A, 111, 6361–6377, doi:10.1021/jp070342s, 2007.

C6755