

Interactive comment on “How relevant is the deposition of mercury onto snowpacks? – Part 2: A modeling study” by D. Durnford et al.

D. Durnford et al.

dorothy.durnford@ec.gc.ca

Received and published: 28 September 2012

Re: acp-2011-996

How relevant is the deposition of mercury onto snowpacks? – Part 2: A modeling study by D. Durnford et al.

Response to Reviewer 1

Specific comments 1. Page 2665, line 12: “Oceanic emissions were increased over Hudson Bay and polewards of 66.5 N ... in order to reproduce summertime atmospheric GEM concentrations ...” - There is no description of how the emissions from the ocean were modified (e.g. fit monthly mean summertime observations at some sites or increased by the magnitude of Hg in meltwater of appropriate catchment area).

C7565

It is principal as it affects the later conclusion on the importance of the ocean emission.

The ocean emissions of SORun were calculated as a remainder. Their calculation was guided by observations of atmospheric mercury. Text discussing the emissions has been added to the first paragraph of Sect. 2.3 from line 509-521.

2. Page 2667, Figs. 2a, 2b: As it follows from the figure there are snowpacks (seasonal - ?) in quite low-latitude regions (e.g. southern Europe – Italy, Spain; southern states of the US), where snow (if any) can hardly lie on the ground longer than over few days. So mercury in such snow rather originates from wet scavenging than from the air-snow exchange. Some discussion of this aspect could improve understanding of the results.

A paragraph discussing this issue, beginning “An interesting feature” has been added in Sect 3.1 at line 617.

3. Page 2669, Fig. 2c: Spatial distribution of the meltwater runoff looks very sporadic, whereas one can expect availability of meltwater during springtime wherever seasonal snowpack takes place. Some clarification of this is needed in the text.

The paragraph beginning “Figure 2c presents” in Sect. 3.1 has been expanded.

4. Page 2672, line 16: “... strongly suggests that the observed summertime maximum is caused by mercury emitted from the Arctic Ocean itself ...” It is interesting to note that the observed summertime maximum of GEM concentration is less pronounced at another high latitude site – Ny Ålesund – located at Spitsbergen (see, for example, Steffen et al., 2008 or measurement data at ebas.nilu.no). Both sites are located at the Arctic Ocean coast and should present similar behavior of GEM seasonality if we expect uniform increase of Hg emissions over the Arctic in summer. Probably the reason is location of Alert close to the Canadian Archipelago with the long coastline, which is more strongly affected by the meltwater runoff.

Text has been added in Sect. 3.2, lines 742-745, in the paragraph starting “From mid-June”.

C7566

5. Fig. 5: Units of the presented volatilization flux are not evident (ng m^{-2}). Probably, they should be $\text{ng m}^{-2} \text{h}^{-1}$.

The units have been corrected.

6. Page 2681, lines 3-11, Figs. 7b, 7d: The same for units of the net deposition flux and net accumulation (ug m^{-2}). Are they $\text{ug m}^{-2} \text{y}^{-1}$?

The units of Fig. 7c, 7d have been corrected.

7. Page 2681, line 26: "... Simulated concentrations of mercury in snowpacks and runoff agree well with observations ..." Really, one can hardly talk about a full-scale validation of the snowpack/meltwater model taking into account very limited amount of observations available for comparison. Besides, all of them relate to different time periods and represent very local conditions. So it should be mentioned that additional evaluation of the model is required in detailed case studies and under conditions of particular field measurements.

Text has been added in Sect. 4, lines 1021-1024 at the end of the paragraph beginning "The snowpack/meltwater model is seen".

Interactive comment on Atmos. Chem. Phys. Discuss., 12, 2647, 2012.