

Interactive  
Comment

## ***Interactive comment on “Modeling chemistry in and above snow at Summit, Greenland – Part 2: Impact of snowpack chemistry on the oxidation capacity of the boundary layer” by J. L. Thomas et al.***

**Anonymous Referee #2**

Received and published: 20 March 2012

### GENERAL COMMENTS

The coupled 1-D snow chemistry and atmospheric boundary layer model MISTRA-SNOW is used to study the impact of snow chemistry on the oxidising capacity of the boundary layer at Summit/Greenland. Model output is compared with observations in ambient and partially also in firn air over a three day period in summer 2008 (GSHOX campaign). Within the measurement uncertainties the model agrees well with observed oxidant levels, i.e. OH and the sum of HO<sub>2</sub>+RO<sub>2</sub> radicals and most of the time with O<sub>3</sub>. In order to investigate impact of snow chemistry on oxidant concentrations,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive  
Comment

runs without snow bromide and without snow nitrate are compared to the base case. The main result is that OH levels can only be matched by including snow emissions of NO<sub>x</sub> and halogens. The model indicates that NO<sub>x</sub> emissions contribute to ozone production, which is reduced but not offset by catalytic destruction via snow sourced bromine chemistry. The mismatch with HONO is interpreted as a potential missing HONO source.

This is a nice model study investigating the coupled snow-atmosphere system in order to understand better the chemical composition of air above snow. The finding that snow photochemistry influences the HO<sub>x</sub> radical budget at Summit is not a novelty as such, as this has been shown previously by Chen et al. (2007). However, difficulties in that study remained in reproducing radical levels. The present work is clearly a step forward in that it employs a truly coupled approach and matches observations of hydroxyl radicals with only temperature as a constraint. In addition, links and feedbacks between NO<sub>x</sub> and halogen chemistry and how they relate to the oxidant budget above snow are illustrated. The MISTRA-SNOW appears to be a valuable tool for the interpretation of atmospheric chemistry studies above snow.

The main concern, which needs to be addressed/discussed further before going ahead in ACP, is that the model probably underestimates the impact of snow emissions of HO<sub>x</sub> precursors on the respective atmospheric budget and on total oxidation capacity.

## SPECIFIC COMMENTS

The authors state that model predicted HO<sub>x</sub> precursors H<sub>2</sub>O<sub>2</sub> and CH<sub>2</sub>O compare well with observations in summer 2000. They find that H<sub>2</sub>O<sub>2</sub> snow emissions do not contribute much to a) the atmospheric budget of H<sub>2</sub>O<sub>2</sub> and b) to primary production of OH radicals. Both statements are quite different to what has been found previously at the same site or other polar sites. For example, regarding a) Chen et al. (2007) showed that in summer 2003 the net snow contribution of the gross chemical produc-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

tion of  $\text{H}_2\text{O}_2$  is about 25%. Re:b) the same study showed that primary  $\text{HO}_x$  sources were  $\text{O}(^1\text{D})+\text{H}_2\text{O}$  (41%) and the photolysis of snow emitted precursors  $\text{H}_2\text{O}_2$  and  $\text{CH}_2\text{O}$  (40%); for the latter  $\text{H}_2\text{O}_2$  dominated with 37% (3% from  $\text{CH}_2\text{O}$ ).

The conclusions regarding model performance & oxidant budget might be different had the authors not chosen to use  $\text{H}_2\text{O}_2$  and  $\text{CH}_2\text{O}$  from summer 2000 (due to lack of measurement during GSHOX). This was a summer with atypically low concentrations of both species at Summit, e.g. mean  $\text{H}_2\text{O}_2$  in the 16-Jun to 7-July 2000 period was 0.65 ppbv (Jacobi 2002) whereas average levels measured in 4-20 June 1996 (Hutterli, 2001) and 27-June to 12-July 2003 (Frey, 2009) were with 1.4 ppbv about twice as high. The same applies to  $\text{CH}_2\text{O}$  with 0.12 ppbv in summer 2000 (Jacobi et al., 2002) compared to 0.23 ppbv in June 1996 (Hutterli et al., 1999) or  $>0.15$  ppbv in July 2003 (Chen et al., 2007).

Thus, uncertainties in modelled oxidant levels & production/loss pathways from less accurate model description of  $\text{H}_2\text{O}_2/\text{CH}_2\text{O}$  snow-air exchange need further discussion & revision of corresponding statements.

As acknowledged by the authors, the model has some shortcomings such as lack of organic snow chemistry or still rather simplistic snow physics, e.g. the representation of the liquid-like layer (LLL). The main tuning parameter is the bulk-LLL ratio of nitrate and bromide and is adjusted to match NO and BrO in ambient air. How sensitive are modelled  $\text{NO}_x$  and BrO mixing ratios to this parameter?

And finally, a more detailed comparison between modelled and observed firn air concentrations would be beneficial for model validation, but certainly another study in its own right. However, discussing some more of the published firn air data at Summit would yield insight into where the model needs development, e.g. firn air concentrations of ozone (Peterson et al., 2001), OH radicals (Beyersdorf et al., 2007) or possibly organic peroxides (Frey et al., 2009).

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Further comments:

Abstract: I16-19 The impact of BrO is a lot smaller than that of NO<sub>x</sub>. It would be instructive to include the fractional contributions of NO<sub>x</sub> and BrO to the OH increase at noon. Credit should be given (here or in the summary of previous model work) that Chen et al. (2007) demonstrated previously the impact of snow chemistry on the HO<sub>x</sub> budget above snow at Summit, although not with a coupled snow-air model and with different conclusions (see above).

I24-25: but not with typical summer levels of H<sub>2</sub>O<sub>2</sub> & CH<sub>2</sub>O measured in other campaigns (e.g. 1996 and 2003, see above)

p5560, I16-23: A brief note would be helpful regarding H<sub>2</sub>O<sub>2</sub> & CH<sub>2</sub>O snow grain distribution and initial snow profiles.

p5562 I.12-15: BL physics seemingly well modelled as suggested by the match of observed vs. modelled T-profiles. So does this imply that wind-shear driven mixing plays a lesser role at Summit (during the model period)? How do wind speeds compare to observations (important for snowpack ventilation e.g. via wind pumping)?

p5564-5566 (section 3): A summary table would be helpful, e.g. parameter (mean, range), uncertainty & reference and also including a quantitative measure of model performance such as the rms error for each parameter & run (base, noNO, noBr).

p5570, I14-21 & Fig.6: The differences in O<sub>3</sub> mixing ratios between base case and noBro are surprisingly small and would not explain a difference of >15 ppbv between firn and ambient air previously observed by Peterson et al., (2001). Is this due to model inconsistencies or are perhaps other halogens at play?

p5571 I15-16: See above. In addition, the model underestimate of CH<sub>2</sub>O in firn air might be even more significant due to a low bias in measured CH<sub>2</sub>O from ventilation effects during sampling.

p5572,I8: see above, choice of H<sub>2</sub>O<sub>2</sub> data set for comparison

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

p5572,115: not Jacobi 2002,2004, the correct citation would be Hutterli et al., 2001, where June 1996 H<sub>2</sub>O<sub>2</sub> data were first described and interpreted

p5572, 118-19: here most of ambient H<sub>2</sub>O<sub>2</sub> production is from HO<sub>2</sub> self reaction; however, the model seems to underestimate the H<sub>2</sub>O<sub>2</sub> snow source if typical Summit values are taken into account and the budget would change accordingly.

p5572, 120-22: needs revision with above comments

p5572, 127: to avoid confusion define oxidation capacity, e.g. as the sum of HO<sub>x</sub>, H<sub>2</sub>O<sub>2</sub> and O<sub>3</sub> or similar.

p5574, 121-24: in contrast to the SUM03 model study (Chen et al., 2007), see previous comments.

p5575,124-27: An important terminal loss of HO<sub>2</sub> radicals identified by the model is the formation of organic peroxides (see also Fig.13). This could be further supported/evaluated by comparison with existing observations of ROOH at Summit (Frey et al., 2009).

p5579,111-22: revision according to comments above.

## TECHNICAL CORRECTIONS

Figure 12d. title should be 'HO<sub>2</sub> destruction'

## REFERENCES

Beyersdorf, A. J., Blake, N. J., Swanson, A. L., Meinardi, S., Dibb, J. E., Sjostedt, S., Huey, G., Lefer, B., Sherwood Rowland, F., and Blake, D. R.: Hydroxyl concentration estimates in the sunlit snowpack at Summit, Greenland, *Atmos. Environ.*, 41, 5101–5109, 2007.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Chen, G., Huey, L. G., Crawford, J. H., Olson, J. R., Hutterli, M. A., Sjostedt, S., Tanner, D., Dibb, J., Lefer, B., Blake, N., Davis, D., and Stohl, A.: An assessment of the polar HO<sub>x</sub> photochemical budget based on 2003 Summit Greenland field observations, *Atmos. Environ.*, 41, 7806–7820, 2007.

Frey, M. M., Hutterli, M. A., Chen, G., Sjostedt, S. J., Burkhart, J. F., Friel, D. K., and Bales, R. C.: Contrasting atmospheric boundary layer chemistry of methylhydroperoxide (CH<sub>3</sub>OOH) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) above polar snow, *Atmos. Chem. Phys.*, 9, 3261–3276, 2009.

Hutterli, Roethlisberger, and Bales] Hutterli, M. A., Roethlisberger, R., and Bales, R. C.: Atmosphere-to-snow-to-firn transfer studies of HCHO at Summit, Greenland, *Geophys. Res. Lett.*, 26, 1691–1694, 1999.

Hutterli, M. A., McConnell, J. R., Stewart, R. W., Jacobi, H. W., and Bales, R. C.: Impact of temperature-driven cycling of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) between air and snow on the planetary boundary layer, *J. Geophys. Res.*, 106, 15395–15 404, 2001.

Jacobi, H.-W., Frey, M. M., Hutterli, M. A., Bales, R. C., Schrems, O., Cullen, N. J., Steffen, K., and Koehler, C.: Measurements of hydrogen peroxide and formaldehyde exchange between the atmosphere and surface snow at Summit, Greenland, *Atmos. Environ.*, 36, 2619–2628, 2002.

Peterson, M. C. and Honrath, R. E.: Observations of rapid photochemical destruction of ozone in snowpack interstitial air, *Geophys. Res. Lett.*, 28, 511, 2001.

---

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 12, 5551, 2012.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)