

***Interactive comment on “Future methane, hydroxyl, and their uncertainties: key climate and emission parameters for future predictions” by C. D. Holmes et al.***

**M. Krol**

m.c.krol@uu.nl

Received and published: 13 October 2012

This comments is written also on behalf of S.A. Montzka and J. Lelieveld

This paper presents an analysis of methane lifetime variations calculated by three models. Overall, calculated variations in methane lifetime are rather small (1%, page 20944, line 13) and modeled variations can be reasonably well represented by a parametric function with only five parameters, (i) temperature (ii) water vapor (iii) column ozone (iv) biomass burning emissions, and (v) lightning  $\text{NO}_x$  emissions.

The paper is very well written, and provides an interesting synthesis of the main factors that are thought to influence the modeled methane lifetime.

C8141

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



## 1 Major comments

We have an objection, however, relating to a statement made in the paper that we feel is inaccurate. In section 3.3, modeled variations in the methane lifetime are compared to observed variations in the methyl chloroform (MCF) decay rate. Since 1998, MCF emissions have been small compared to the atmospheric burden, and variations in observed MCF decay rate may be interpreted as variations in the tropospheric oxidizing capacity, and hence in the methane lifetime. However, in Fig. 4 the authors show that the detailed interannual variations in the MCF decay rate depend on the observational network.

They conclude: “Given that differences in observed MCF decay rates between the two networks are as large as their difference from CTM  $\tau_{\text{CH}_4 \times \text{OH}}$  anomalies, we conclude that better understanding of the systematic differences between the observation networks is required before using them as a constraint on  $\tau_{\text{CH}_4 \times \text{OH}}$  and OH interannual variability.”

This statement suggests that an analysis of MCF data has no informative value with respect to mean OH concentrations, i.e. the lifetime of methane, and the interannual variability of OH. We think this sentence inaccurately reflects the value of MCF measurements for deriving information about OH. We agree that a determination of the phase of interannual OH variations from MCF data would be robust especially if the timing of those changes were consistently derived from both measurement networks, but poorly correlated variations derived from the two networks do not negate the fact that results from both networks provide a consistent estimate of mean annual OH (and, therefore,  $\tau_{\text{CH}_4 \times \text{OH}}$ ) and of the magnitude of interannual changes in OH since 1998. We also feel that this result does not negate the main message of Montzka et al. (2011), which is that the OH-variability derived from MCF since 1998 was remarkably small compared to that inferred in earlier years and that the tropospheric oxidizing capacity appears well buffered against perturbations.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Another point we would like to make is that modeled interannual OH-variability is likely an underestimate (1%), even when the main drivers of OH variability are accounted for. Possible reasons include unaccounted variability in VOC emissions, limited spatial resolution, and simplified chemistry schemes used in the analysis. In contrast, interannual OH-variability derived from MCF measurements (2.3%) is likely overestimated (as explicitly discussed in Montzka et al., 2011), due to potential calibration and measurement consistency issues and errors in translating measurements at a few stations to a global MCF decay rate, in addition to the potential errors associated with unknown emission variations.

We agree with the authors' statement that the differences between the two MCF measurement networks should be resolved, though this is a problem that has received attention in AGAGE and NOAA for many years. In the end, the problem may be unresolvable, since the discrepancies the authors discuss arise from differences in measured mixing ratios that are often only small fractions of a part per trillion. Even so, results of both networks are not uncorrelated as both show similar variability and a clear anomaly around 2003. Likewise, the different models agree approximately, though not perfectly. Ultimately, our goal should be to reduce errors in both models and measurements, while measurements remain imperative to evaluate models.

## 2 Minor Comments

Page 20936, line 13: "...which increases OH and biases  $\tau_{\text{CH}_4 \times \text{OH}}$  high by about 10%...". if OH increases one would expect the methane lifetime to be biased low.

Page 20944, line 29: "Fig. 2" should be "Fig. 4".

In the Supplement: **WLEF** tower is in Wisconsin.

NOAA data after Dec 2007 have been appropriately omitted from the analysis, but why

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

are NOAA data and not AGAGE data shown past that date in Figure S2?

---

Interactive comment on Atmos. Chem. Phys. Discuss., 12, 20931, 2012.

ACPD

12, C8141–C8144, 2012

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C8144

