

Interactive
Comment

Interactive comment on “Peroxy radical chemistry and OH radical production during the NO₃-initiated oxidation of isoprene” by A. J. Kwan et al.

Anonymous Referee #2

Received and published: 6 February 2012

This paper presents a detailed analysis of a chamber experiment on NO₃+isoprene, conducted with excess isoprene to force HO₂+RO₂ and RO₂+RO₂ chemistry to dominate over NO₃+RO₂. This enables the authors to study in greater detail the product channels of HO₂+RO₂ and RO₂+RO₂, which were hypothesized in a companion paper (Ng et al., 2008) to be responsible for higher SOA yields. This work addresses important scientific questions in the production of atmospheric nitrates and SOA, within the scope of ACP. The use of CIMS data to constrain RO₂/HO₂/NO₃ reaction productions adds a novel contribution to the literature on NO₃ + isoprene. The manuscript is well-written. The product branching ratios and modeling are subject to large uncertainties that must be clearly stated, but the publication of this qualitative picture is likely to be of great value to the research community in identifying important chemistry for follow-up study. I recommend publication after the authors consider the comments & questions

C83

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



below:

Specific comments: 1) The production of OH seems to be central to discussion of results and implications in this manuscript but is not highlighted in the title or introduction. I suggest adding background on this question of additional oxidant sources upfront.

2) (related to point 1) In the discussion of Figure 4 and in the figure itself: clearly you were tweaking the model to get more OH production. What determined the degree of your tweaks? Can you put any, even approximate, constraint on the likely amount of OH produced based on observed OH-isoprene products? If so it would be great to include the apparent branching ratio of OH production in the manuscript.

3) Table 1 and last paragraph of p. 2266: I'd like to be able to tell if this slow equilibration with surfaces in tubing /instrumentation is a big effect or not – is the peak mixing ratio likely a big underestimate of actual product yield or not? Can you provide any kind of uncertainty estimate due to this effect, e.g. based on other experiments with mass balance? In the heading in Table 1, I suggest labeling as “maximum/peak concentration” to remind/highlight this.

4) Please indicate if the yields are determined on a molar or mass basis.

5) Last paragraph on p. 2268: why would the possibility of radical interconversion make the distribution of isomers more sensitive to initial conditions?

6) Last paragraph of section 3.1, p. 2269: here you say nitrate yields should be considered upper limits, while in the abstract you list yield as “>~80%”, should this be “<~80%”?

7) Section 3.2: box model is introduced early – do you really need it to demonstrate than HO₂+NO cannot possibly be important? Seems you can say this without the model, but it really is needed for the later RO₂+HO₂ branching – I would move its discussion down there.

8) last paragraph on p. 2272: RO₂ /HO₂ reactions from isomerized nitrooxy RO₂ are

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



ignored, but earlier you pointed out that this isomerization was facile – how to reconcile this?

9) Last sentence above section 3.2.1 p. 2273: Elaborate please – why do you expect resonance stabilization of the radical reactant to preference the radical propagation product channel?

10) last paragraph, p. 2275: “There remain many uncertainties regarding the mechanism of $\text{RO}_2 + \text{RO}_2$ reactions (Dibble, 2008), so it is difficult to assess whether reported ROOR formation (or lack thereof) is a result of the particular radicals studied or the analytical techniques employed to study their reaction.” Sounds like you’re pointing out discrepancies in previous measurements of RO_2/RO_2 to yours for isoprene? I don’t see a huge discrepancy as you’ve set it up – perhaps you can start this paragraph in a way that highlights the difference you’re trying to point out? Is it the difference between 38% and 50%?

11) Top of 2277: Why would O_2 abstraction be faster for these radicals? Is there a structural explanation?

12) At end of section 3.5: add a wrap-up statement to make clear what you’re saying: you’re proposing that $\text{RO}_2 + \text{isoprene}$ might be an important reaction pathway here

13) Top paragraph on p. 2280: $\text{RO}_2\text{-RO}_2$ reactions are sped up by NO_3 electron-withdrawing group: what about your resonance-stabilized radical

14) Since you emphasize a few times the sensitivity to initial conditions, maybe a comment is warranted on how much greater than ambient conditions this experiment is. Do you expect the sensitivity to be primarily in the RATIO of NO_3 to isoprene?

Technical corrections: a) Caption to Fig. 1 : suggest “are detected by the CIMS instrument as $\text{CF}_3\text{O-}$ adducts at the indicated. . .” b) Caption to Table 2: suggest “for box model assessment”

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

