

Interactive comment on “Nested-grid simulation of mercury over North America” by Y. Zhang et al.

Anonymous Referee #2

Received and published: 27 March 2012

This study describes the development of a higher resolution nested version of the well known GEOS-Chem model, and describes the results obtained simulating the wet deposition of Hg over continental North America, and compares these results with the extensive observations made in the Mercury Deposition Network. The article also discusses the speciation of emissions from major anthropogenic sources, and assesses the quality of the simulation results, representing in plume reduction by using different speciation profiles for certain types of anthropogenic emission source. Overall the paper is clearly written and well presented. The subject is relevant, in 2013 UNEP hopes to complete negotiations for a legally binding instrument on mercury before the twenty-seventh regular session of the Governing Council / Global Ministerial Environment Forum in 2013, see, <http://www.unep.org/hazardoussubstances/Mercury/Negotiations/tabid/3320/Default.aspx>, which is something the authors may wish to add to their Introduction.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



The approach used by the authors is sound; using the same model (physics parametrizations, and chemistry scheme) to perform both the global and regional simulations makes more sense than taking boundary and initial conditions from a global model, as input for a regional model which calculates atmospheric chemistry and physics differently. The results certainly seem to give a greater insight into the processes influencing the spatial and temporal distribution of the wet deposition of H_g over North America. There is no doubt that the article deserves publication, although there are a few general and a couple of specific points which the authors should address by the authors before final publication in ACP.

Generally speaking throughout the article there seems to me to be a tendency to overestimate the amount of confidence that can be placed in the results obtained from model simulations, and to present model results as if they were fact, some of the instances where a caveat might be useful are noted in the following comments.

In the abstract the authors use the term high-resolution. In terms of regional modelling, would a half by two-thirds degree resolution be classified as as high, although certainly it is higher than the global model obviously. Higher resolution, maybe? Also in the abstract the last three sentences read as facts rather than findings from model simulations. The last sentence regarding dry deposition should certainly be qualified as there are very few dry deposition measurements with which to compare the simulation results, and the methods used for the measurements are still under development.

In the introduction the authors could add the following references: regarding emissions; one or both of AMAP/UNEP (2008); Pirrone et al. (2009), and concerning the influence of boundary conditions Pongprueksa et al. (2008), which showed very clearly the correlation between boundary conditions and deposition fluxes in regional models.

In the description of the model (p2607 onwards) the authors give an annual total of
C992

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

220 Mg yr⁻¹ for the contribution of Hg emissions from biomass burning, this is noticeably less than the 300 Mg yr⁻¹ in Holmes et al. (2010), which the authors cite, and significantly less than the 675 ± 240 Mg yr⁻¹ in Friedli et al. (2009) which is not cited. Is this because the authors prefer to use an enrichment factor $\Delta[\text{Hg}]/\Delta[\text{CO}]$ at least in part derived from measurements over North America?

On page 2609, the parametrisation process to describe polar AMDEs is described, however there are no references, and the figure of 5 ppt is given as the concentration of BrO in the boundary layer when a series of conditions is met. What period do the authors consider as springtime, how much sea-ice per model cell is necessary, is there a cut off value for the incoming solar radiation at which there is no BrO, how are stable conditions defined, and why below 268 K? The reader is left with rather a lot of questions and the authors should expand a little, or provide a reference for the source of the parametrisation.

In the description of the IPR simulation, p2611, the authors describe the studies that show that the oxidised Hg content of coal fired power plant plumes is significantly less downwind of the plant than it is when measured in close proximity to the stack, which suggests that there is an in plume reduction of Hg^{II}. If I recall correctly it was proposed that SO₂ is involved in the reduction mechanism. The authors adjust the proportions of Hg species in the emission inventory for CFPPs. The authors then apply similar changes to the emissions from incinerators (is there evidence of high concentrations of SO₂ from incinerator stacks?), and use 96% Hg(0), 0% Hg^{II}, and 4% HgP; this speciation profile is cited as from Streets et al. (2009). Streets et al. (2009) actually cites Streets et al. (2005), which for Hg speciation from waste and residue burning in the table 7 of their article cites three articles by Friedli et al. namely Friedli et al. (2001, 2003a,b), which refer to biomass burning and agricultural waste burning. Therefore I am not convinced that the application of the in plume reduction parametrisation used for CFPPs is entirely valid. Given the underestimation of Hg wet deposition in Florida at certain times of the year, it may well be a good idea to rerun the model removing this

assumption.

The total emissions from natural and anthropogenic sources are described on page 2612 and illustrated in Figure 1. I am surprised that there are no (or extremely low) emissions from the sea all along the West Coast of the United States, and also that the emissions from parts of the North Atlantic appear to be as high or higher than they are from the most part of the continental US (middle row, left, figure 1). It also seems from the two figures in the middle row that the area of intense emissions in the vicinity of the Great Lakes in the global model is not present in the regional model. Could the authors comment on this please? Also on page 2612, the sensitivity study with primary anthropogenic emissions turned off, this refers only to Hg emissions I assume?

On page 2616, last line. Is 'very well' perhaps a slight overstatement?

On page 2617, the last sentences of section 3.2. I wonder if these underestimations would improve if the authors remove their assumption about the speciation profile from waste incineration and rerun the model. Or perhaps they tried and it led to poor results in other areas. Would it be possible to include a little more discussion here, perhaps mentioning what happens in the waste incineration in-plume reduction assumption is removed?

Page 2717, Section 3.3. In the discussion of the simulated mean surface concentrations of atmospheric Hg there are a number of comments in the text which seem unusual. Temme et al. (2003) and Lindberg et al. (2007) both give the hemispheric background of Hg⁰ as 1.5 - 1.7 ng m⁻³, (northern), and 1.1 - 1.3 ng m⁻³ (southern). Therefore values of 1.7 - 1.8 ng m⁻³ for TGM in the Ohio River Valley cannot be considered 'high', nor can 1.5 - 1.6 ng m⁻³ in Nevada and Utah be considered 'elevated' and 1.3 - 1.5 ng m⁻³ over the Great Plains, southern Canada and northern Mexico is not 'typical' background. The concentrations in the MBL seem anomalous as well, mean values of the measurements in the Atlantic (northern hemisphere) are higher

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



than $1.2 - 1.4 \text{ ng m}^{-3}$ (see Sprovieri et al. (2009)), and the authors ascribe the low MBL values to low emissions (fig 1, middle panels) which as mentioned before hardly seems to be the case looking closely at the figure. These concentrations seem low and require explanation.

On page 2618, the authors state that at higher altitudes, Hg^{II} concentrations are high due to faster oxidation and slower removal processes. This is another instance where modelling studies suggest that this is indeed the case, however there are not enough experimental results of certain enough quality to state this definitively. The authors should qualify this statement.

On pages 2619-20 the authors discuss the seasonal cycle of TGM and refer to the summertime increase in oxidation and subsequent deposition, citing a number of previous studies. However the studies by Bergan and Rodhe (2001) and by Selin et al. (2007) used a different chemical mechanism based on oxidation by OH (and O_3 in the case of Bergan and Rodhe (2001)), therefore the authors should explain why they think this comparison is valid, or point out the fact that whatever the atmospheric oxidation pathway for Hg^0 is, observations indicate that it is more rapid in summer and any attempt at modelling the atmospheric Hg cycle needs to be able to reproduce this phenomena.

p2622 section 3.6. Do the authors mean all anthropogenic Hg emissions?

On p2624 the authors mention that the model resolution cannot capture the local scale enhancements in Hg deposition which occur close to point sources. Is it possible that the IPR simulations therefore underestimate the contribution of north American sources to north American deposition? And therefore that the 12% estimated in the last line of the Conclusions may not be as robust as it seems?

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

References

- AMAP/UNEP: Technical Background Report to the Global Atmospheric Mercury Assessment, Tech. rep., Arctic Monitoring and Assessment Programme / UNEP Chemicals, 2008.
- Bergan, T. and Rodhe, H.: Oxidation of Elemental Mercury in the Atmosphere; Constraints Imposed by Global Scale Modelling, *Journal of Atmospheric Chemistry*, 40, 191–212, 2001.
- Friedli, H., Radke, L., and Lu, J.: Mercury in smoke from biomass fires, *Geophysical Research Letters*, 28, 3223–3226, 2001.
- Friedli, H., Radke, L., Lu, J., Banic, C.M., Leitch, W., and MacPherson, J.: Mercury emissions from burning of biomass from temperate North American forests: Laboratory and airborne measurements, *Atmospheric Environment*, 37, 253–267, cited By (since 1996) 82, 2003a.
- Friedli, H., Arellano, A., Cinnirella, S., and Pirrone, N.: Initial Estimates of Mercury Emissions to the Atmosphere from Global Biomass Burning, *Environ. Sci. Technol.*, 43, 3507–3513, 2009.
- Friedli, H. R., Radke, L. F., Prescott, R., Hobbs, P. V., and Sinha, P.: Mercury emissions from the August 2001 wildfires in Washington State and an agricultural waste fire in Oregon and atmospheric mercury budget estimates, *Global Biogeochem. Cycles*, 17, 1039–, 2003b.
- Holmes, C. D., Jacob, D. J., Corbitt, E. S., Mao, J., Yang, X., Talbot, R., and Slemr, F.: Global atmospheric model for mercury including oxidation by bromine atoms, *Atmospheric Chemistry and Physics*, 10, 12037–12057, 2010.
- Lindberg, S., Bullock, R., Ebinghaus, R., Engstrom, D., Feng, X., Fitzgerald, W., Pirrone, N., Prestbo, E., and Seigneur, C.: A Synthesis of Progress and Uncertainties in Attributing the Sources of Mercury in Deposition, *AMBIO: A Journal of the Human Environment*, 36, 19–33, 2007.
- Pirrone, N., Cinnirella, S., Feng, X., Finkelman, R., Friedli, H., Leaner, J., Mason, R., Mukherjee, A., Stracher, G., Streets, D., and Telmer, K.: Global Mercury Emissions to the Atmosphere from Natural and Anthropogenic Sources, in: *Mercury Fate and Transport in the Global Atmosphere*, edited by Mason, R. and Pirrone, N., pp. 1–47, Springer US, 2009.
- Pongprueksa, P., Lin, C.-J., Lindberg, S. E., Jang, C., Braverman, T., Jr., O. R. B., Ho, T. C., and Chu, H.-W.: Scientific uncertainties in atmospheric mercury models III: Boundary and initial conditions, model grid resolution, and Hg(II) reduction mechanism, *Atmospheric Environment*, 42, 1828 – 1845, 2008.
- Selin, N. E., Jacob, D. J., Park, R. J., Yantosca, R. M., Strode, S., Jaeglé, L., and Jaffe, D.: Chemical cycling and deposition of atmospheric mercury: Global constraints from observa-

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



- tions, *J. Geophys. Res.*, 112, D02 308–, 2007.
- Sprovieri, F., Pirrone, N., Mason, R. P., and Andersson, M.: Spatial coverage and temporal trends of over-water, air-surface exchange, surface and deep sea water mercury measurements, in: *Mercury Fate and Transport in the Global Atmosphere*, edited by Mason, R. and Pirrone, N., pp. 323–380, Springer US, 2009.
- Streets, D. G., Hao, J., Wu, Y., Jiang, J., Chan, M., Tian, H., and Feng, X.: Anthropogenic mercury emissions in China, *Atmospheric Environment*, 39, 7789 – 7806, 2005.
- Streets, D. G., Zhang, Q., and Wu, Y.: Projections of Global Mercury Emissions in 2050, *Environmental Science & Technology*, 43, 2983–2988, 2009.
- Temme, C., Slemr, F., Ebinghaus, R., and Einax, J.: Distribution of mercury over the Atlantic Ocean in 1996 and 1999–2001, *Atmospheric Environment*, 37, 1889 – 1897, 2003.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)