



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak<sup>1,\*</sup> and P. J. Adams<sup>1,2</sup>

<sup>1</sup>Department of Civil and Environmental Engineering, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

<sup>2</sup>Department of Engineering and Public Policy, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

\* now at: The Department of Highways, Ministry of Transport, Bangkok, Thailand

Received: 27 February 2013 – Accepted: 14 March 2013 – Published: 22 April 2013

Correspondence to: P. J. Adams (petera@andrew.cmu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

A model of carbonaceous aerosols has been implemented into the TwO-Moment Aerosol Sectional (TOMAS) microphysics module in the GEOS-Chem CTM, a model driven by assimilated meteorology. Inclusion of carbonaceous emissions alongside pre-existing treatments of sulfate and sea-salt aerosols increases the number of emitted primary aerosol particles by a factor of 2.5 and raises annual-average global CCN(0.2 %) concentrations by a factor of two. Compared to the prior model without carbonaceous aerosols, this development improves the model prediction of CN10 number concentrations significantly from  $-45$  to  $-7$  % bias when compared to long-term observations. However, similar to other OC/EC models, the model underpredicts OC and EC mass concentrations by a factor of 2–5 when compared to EMEP observations. Because primary OA and secondary OA affect aerosol number size distributions differently, we assess the sensitivity of CCN production, for a fixed source of OA mass, to the assumed POA-SOA split in the model. For a fixed OA budget, we found that CCN(0.2 %) decreases nearly everywhere as the model changes from a world dominated by POA emissions to one dominated by SOA condensation. POA is about twice as effective per unit mass at CCN production compared to SOA. Changing from a 100 % POA scenario to a 100 % SOA scenario, CCN(0.2 %) concentrations in the lowest model layer decrease by about 20 %. In any scenario, carbonaceous aerosols contribute significantly to global CCN. The SOA-POA split has a significant effect on global CCN and the microphysical implications of POA emissions versus SOA condensation appear to be at least as important as differences in chemical composition as expressed by the hygroscopicity of OA. These findings stress the need to better understand carbonaceous aerosols loadings, the global SOA budget, microphysical pathways of OA formation (emissions versus condensation) as well as chemical composition to improve climate modeling.

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

Cloud condensation nuclei (CCN) are the fraction of aerosol particles that activate to become cloud droplets. CCN concentrations and cloud processes play a critical role in determining the aerosol indirect radiative forcings, including increased cloud reflectivity and changes in cloud cover (Albrecht, 1989; Twomey, 1974), which are caused by anthropogenic increases in CCN levels. The Intergovernmental Panel on Climate Change (IPCC) has estimated the globally and annually averaged indirect aerosol radiative forcing (only the cloud brightness effect) to be between  $-1.8$  to  $-0.3 \text{ W m}^{-2}$ , compared to  $+2.6 \text{ W m}^{-2}$  by changes in long-lived greenhouse gases and pointed out that the aerosol indirect forcing remains the dominant uncertainty in the net anthropogenic forcing estimate (IPCC, 2007). Improving the representation of CCN in global models, including mechanisms leading to their formation, enhances our understanding of CCN with the long-term goal of reducing uncertainty in aerosol indirect radiative forcing estimates.

Organic aerosol (OA) is found to be a significant fraction of aerosol mass in the accumulation mode that accounts for most CCN (Hitzenberger et al., 1999). Many organics in aerosols have been shown to be CCN-active as OA contains a significant fraction of water-soluble organic carbon (WSOC) (Raymond and Pandis, 2002, 2003; Kerminen, 1997). Global modeling has also demonstrated the important role that OA plays in the global CCN budget (Pierce et al., 2007; Merikanto et al., 2009).

Organic aerosol or, synonymously, organic matter (OM) is composed of many compounds and has traditionally been categorized by the way it enters the atmosphere: primary emission or secondary production. OA that is emitted directly from fossil fuel and biomass combustion is termed “primary organic aerosol” (POA). In contrast, secondary organic aerosol (SOA) is formed in the atmosphere by condensation of low volatility products from oxidation of volatile organic compounds. The relative contributions of POA and SOA to the total OA remains an open issue as different studies report wide ranges of estimates (Kanakidou et al., 2005; Hallquist et al., 2009). Traditionally,

ACPD

13, 10561–10601, 2013

### Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

models have tended to predict a predominance of POA over SOA (Chung and Seinfeld, 2002; Kanakidou et al., 2005; Pun et al., 2003; Vutukuru et al., 2006), but measurement studies show striking evidence of SOA dominance observed at various locations, even in heavily urbanized locations (Zhang et al., 2007; de Gouw et al., 2005; Volkamer et al., 2006; Zhang et al., 2005), and recent modeling efforts have addressed this discrepancy (Jathar et al., 2011; Pye and Seinfeld, 2010). To complicate the matter even further, different measurement techniques result in different SOA/OA fractions, as was the case for the Pittsburgh Air Quality Study, for which SOA/OA values were estimated to be 35–73 % (Subramanian et al., 2007; Cabada et al., 2004; Shrivastava et al., 2007; Zhang et al., 2005). Therefore, there exists large uncertainty regarding the POA-SOA split.

Apart from obvious differences in the chemical composition of POA and SOA, they form via different microphysical pathways. POA emissions add to both aerosol mass and number concentrations, in both the ultrafine and CCN size ranges. Primary ultrafine particles compete for condensable gas, if available, and often grow to CCN (Pierce and Adams, 2007; Pierce et al., 2007). Alternatively, these ultrafines undergo high coagulation scavenging loss, resulting in short atmospheric lifetimes. In contrast, condensation of SOA adds mass to existing particles without changing aerosol number concentrations. Condensation of SOA grows particles and can lead to CCN formation, but this depends on what size particle the SOA condenses onto (and, therefore, the pre-existing aerosol size distribution) and whether the additional SOA results in a particle activating that would not otherwise activate. Previous work has demonstrated that uncertainty in the total SOA source has a significant impact on global CCN concentrations (Pierce and Adams, 2009).

Despite large uncertainty in the POA-SOA split, total organic aerosol sources, on the other hand, may be better quantified than those of POA or SOA individually. This leads to the question of how much, for a fixed total OA source, this POA-SOA split uncertainty translates into uncertainty in CCN predictions. This work will explore the sensitivity of

CCN concentrations to different POA-SOA split assumptions and ask whether POA or SOA, per unit mass, is better at forming CCN.

Additionally, this paper documents the implementation of carbonaceous aerosols in the highly size-resolved Two-Moment Aerosol Sectional (TOMAS) microphysics model (Adams and Seinfeld, 2002), which was integrated in the GEOS-Chem CTM (Bey et al., 2001) by Trivitayanurak et al. (2008). Previous work had developed a treatment of carbonaceous aerosols in TOMAS hosted by the GISS GCM II-prime (Pierce et al., 2007). Using TOMAS hosted by GEOS-Chem, we present CCN predictions with and without carbonaceous aerosols to demonstrate the contribution of carbonaceous aerosols to CCN at the global scale. Comparison of model results to aerosol observations is performed to evaluate the model performance. Finally, we perform sensitivity simulations to test different POA-SOA split assumptions. Results from these simulations are used to answer the question whether POA or SOA is more effective at CCN formation.

Section 2 provides model overview and describes development of the carbonaceous aerosol in GEOS-CHEM with TOMAS. Section 3 contains model results, discusses contribution of carbonaceous aerosols and presents comparison to observations. Section 4 discusses the effects of POA-SOA split on the CCN results. Finally, Sect. 5 concludes this study.

## 2 Model description

### 2.1 Overview

We represent carbonaceous aerosols in the GEOS-Chem global CTM with the Two-Moment Aerosol Sectional (TOMAS) microphysics model (Adams and Seinfeld, 2002). TOMAS tracks two independent moments, number and mass, of the aerosol size distribution for each size bin. Previous work implemented TOMAS in the host model GEOS-Chem as described in Trivitayanurak et al. (2008) and simulated sulfate and sea-salt aerosols.

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The GEOS-Chem model is a 3-D tropospheric chemistry model driven by assimilated meteorological observations from the Goddard Earth Observing System (GEOS) of the NASA Global Modeling and Assimilation Office (GMAO) (Bey et al., 2001). We implemented our aerosol capabilities in the GEOS-Chem version 7.04.11 ([http://acmg.seas.harvard.edu/geos/geos\\_versions.html](http://acmg.seas.harvard.edu/geos/geos_versions.html)). The GEOS-3 GMAO meteorological fields are used to run the model with horizontal resolution of 4 degrees latitude by 5 degrees longitude and a 30-level sigma-coordinate vertical grid between the surface and 0.01 hPa.

TOMAS is configured to track 30 size bins defined by dry particle mass, which span the size range of roughly 10 nm to 10  $\mu$ m. In each bin, TOMAS tracks 8 quantities: sulfate mass, sea salt mass, mass of externally mixed EC, mass of internally mixed EC, hydrophobic OM mass, hydrophilic OM mass, mass of water and the number of aerosol particles in the bin. Besides the aerosol microphysics, GEOS-Chem simulates photochemistry with online oxidant, i.e. OH, H<sub>2</sub>O<sub>2</sub>, O<sub>3</sub>, concentrations and also calculates sulfur chemistry, including DMS, MSA, SO<sub>2</sub>, and H<sub>2</sub>SO<sub>4</sub>. We use the binary nucleation scheme described in Adams and Seinfeld (2002), which calculates new particle formation based on a critical threshold value for rapid nucleation and available sulfuric acid concentrations (Wexler et al., 1994).

Advection is calculated every 30 min using the TPCORE algorithm (Lin and Rood, 1996). Size-dependent dry deposition of aerosols is modeled using the scheme of Zhang et al. (2001). Wet deposition includes in-cloud scavenging, below-cloud scavenging, and scavenging in convective updrafts. The large-scale clouds and convective clouds are assumed to experience maximum supersaturations of 0.19% and 0.75%, respectively, for purposes of nucleation scavenging. The supersaturation at which particles in each size bin activate is determined from modified Köhler theory based on the current composition of that size bin and model grid cell (Seinfeld and Pandis, 1998; Hanel, 1976; Laaksonen et al., 1998; Raymond and Pandis, 2003). We neglect interstitial scavenging in clouds. First-order removal by below-cloud scavenging is calculated using size-dependent washout rate constants as in Adams and Seinfeld (2002).

Convective updrafts remove aerosol by a first-order loss that affects only activated particles. A more comprehensive description of TOMAS implementation in GEOS-Chem model can be found in Trivitayanurak et al. (2008).

## 2.2 Improvements in particulate emissions

5 Primary particulate emissions contribute significantly to aerosol number. “Primary” sulfate emissions, representing plume processing of power plant emissions, consist of an assumed fraction of total sulfur emission with a bimodal lognormal size distribution with number geometric mean diameters of 10 and 70 nm and standard deviations of 1.6 and 2.0, respectively (Adams and Seinfeld, 2002). In Trivitayanurak et al. (2008), particulate  
10 sulfate was prescribed to be 5 % by mass of sulfur emission in Europe and 3 % elsewhere. In this work, we reduce the fraction of “primary” particulate sulfate to be 1 % everywhere following the development of Pierce and Adams (2009), who found that 3 % of total sulfur consistently overpredicted the measured aerosol number in polluted regions even when carbonaceous aerosols were omitted.

15 Another improvement related to particulate emission is the development of a sub-grid coagulation parameterization based on Pierce et al. (2009). In this parameterization, primary particles undergo coagulation with the pre-existing aerosol size distribution in the source grid cell for 10 h to determine their “effective” regional emission and number size distributions; this helps the model to simulate appropriate regional background  
20 aerosol number concentrations.

## 2.3 Carbonaceous aerosols implementation

Size-resolved carbonaceous aerosols, namely organic matter (OM) and elemental carbon (EC), are the new aerosol species implemented in this work. This implementation for TOMAS in the GEOS-Chem CTM largely follows the development of carbonaceous  
25 aerosols for TOMAS in GISS GCM II-Prime by Pierce et al. (2007).

### Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 2.3.1 Emissions

Combustion sources of carbonaceous aerosols are categorized into fossil fuel, biofuel, and biomass burning. Note that oceanic emissions of organic aerosol are not considered here. Fossil fuel and biofuel emissions are taken from the Bond et al. (2004) global inventory. Seasonality of these emissions over North America follows Cooke et al. (1999) and Park et al. (2003) for fossil fuel and biofuel, respectively. Biomass burning emission is from the Global Fire Emissions Database version 2 (GFED2: on the web at <http://ess1.ess.uci.edu/~jranders/data/GFED2/>), which provides monthly variability as well as interannual variability from 1997 to 2004 (Giglio et al., 2006; van der Werf et al., 2006). Annual emission rates of OM and EC are summarized in Table 1. To convert from OC mass normally reported in emission inventories to OM mass presented in our model, we assume a globally uniform and constant OM : OC ratio of 1.8 (El-Zanan et al., 2005; Yu et al., 2005; Zhang et al., 2005). For both OM and EC, fossil fuel combustion is assumed to emit particles with a lognormal size distribution with number median diameter of 30 nm with a geometric standard deviation of 2; biofuel and biomass burning emissions are assumed to follow a lognormal distribution with a number median diameter of 100 nm and a geometric standard deviation of 2.

### 2.3.2 Secondary organic aerosol

This work incorporates a simple treatment of SOA for several reasons: (1) to observe the global aerosol prediction with complete organic aerosol sources, (2) to demonstrate effects of varying SOA contributions on CCN production, and (3) given rapidly changing knowledge of SOA and TOMAS's focus on aerosol microphysics, a simple treatment is more flexible and allows for easier sensitivity studies. In GEOS-Chem, SOA is produced by condensation of low-volatility products of the photooxidation of biogenic VOCs; traditionally, global models have predicted that SOA produced from anthropogenic organic compounds makes a small contribution (about 10%) to global OA sources (Tsigaridis and Kanakidou, 2003) and is not considered here. Following the

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



assumption used by Park et al. (2003), we assume a 10% yield from monoterpenes emissions to produce SOA in our model BASE case. In the model, SOA products are assumed to be produced immediately after emission of monoterpenes and are thereafter treated as non-volatile. Biogenic emissions in GEOS-Chem are calculated from MEGAN model (Guenther et al., 2006). Annual-average SOA production from this calculation is  $18.4 \text{ Tgyr}^{-1}$ , in agreement with previous estimates (Griffin et al., 1999).

Previous SOA global modeling studies have adopted different choices regarding what constitutes the absorbing media for SOA partitioning into the aerosol phase. These choices include SOA itself (Kanakidou et al., 2000), entire condensed-phase organic (POA+SOA) (Pun et al., 2003; Chung and Seinfeld, 2002; Tsigaridis and Kanakidou, 2003; Kanakidou et al., 2000; Johnson et al., 2006), organic + inorganic aerosols (Tsigaridis and Kanakidou, 2003), and aqueous portion of the aerosol for the case of water soluble organic compounds (Seinfeld et al., 2001). Absorbing media still remains an uncertain factor. Regardless of what media constitutes the absorbing phase, which governs the thermodynamic equilibrium of SOA, there is still a question of kinetics and mass transfer that governs the size distribution of the condensed SOA. For simplicity here, we assume that the produced SOA irreversibly condenses onto the size distribution in proportion to the pre-existing aerosol surface area under ambient (hydrated) conditions, an assumption that has produced results in reasonable agreement with observations (Riipinen et al., 2011). Use of total aerosol surface area from all species is consistent with smog chamber studies in which SOA is observed to condense onto inorganic seed particles. This assumption also guarantees that SOA will reside primarily in the accumulation mode that contributes most aerosol surface area, consistent with ambient observations.

### 2.3.3 Hygroscopicity, mixing state, CCN activation, and aging

Carbonaceous aerosols are presented in the model as four categories: externally mixed EC, internally mixed EC, hydrophobic OM, and hydrophilic OM. Aerosols are divided into two populations for the purpose of CCN activation calculations. The first

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

population is solely the externally mixed EC or “pure EC”, and the other population is an internal mixture of all other aerosols, including mixed EC, hydrophilic OM, hydrophobic OM, sulfate and sea salt. The externally mixed EC population does not activate and, therefore, is not subject to nucleation scavenging. CCN activation for the mixed aerosol population is calculated using modified Köhler theory (Pierce et al., 2007), which allows for calculation of the activation diameter of particles containing various soluble and insoluble species. We ignore changes in surface tension by surfactant effects of the organic aerosols. EC is insoluble in both populations, but the mixed EC may occur in activated particles and, therefore, be subject to nucleation scavenging. Hydrophobic and hydrophilic OM are assumed to be insoluble and completely soluble, respectively. Hydrophilic OM is assumed to have a critical dry diameter for activation of 140 nm at 0.2% supersaturation similar to pinonic acid and adipic acid (Raymond and Pandis, 2002). In terms of the  $\kappa$  parameter (Petters and Kreidenweis, 2007), hydrophilic OM has  $\kappa = 0.18$  and hydrophobic OM has  $\kappa = 0$  at 273 K. The assumed density of hydrophilic OM is  $1.4 \text{ g cm}^{-3}$  and hydrophobic OM is  $1.8 \text{ g cm}^{-3}$ .

Our emissions of OM and EC use hygroscopicity and mixing state assumptions following Cooke et al. (1999). Fifty percent of OM emissions are assumed to be hydrophobic and the other 50% are hydrophilic. As for EC, 80% is assumed to be emitted as pure EC and 20% as mixed EC. Secondary organic aerosol formed in the model is assumed to be completely hydrophilic. SOA water solubility and its relation to WSOC has been discussed in a number of studies (Saxena and Hildemann, 1996; Kumagai et al., 2009; Kondo et al., 2007).

Aging of carbonaceous particles includes the conversion of hydrophobic OM to hydrophilic OM and the conversion of pure EC to mixed EC via mixing processes. Both processes are modeled in our work using an e-folding time of 1.15 days. This conversion timescale affects the wet deposition lifetime of carbonaceous aerosol but remains uncertain (Cooke et al., 1999; Koch et al., 1999; Park et al., 2005). The crudeness of this parameter may be a source of uncertainty but since this timescale is shorter than the mean lifetime of particles in the atmosphere, uncertainties in the aging timescale

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



should not significantly affect the carbonaceous aerosol burden. Because the aging timescale is relatively short, hydrophilic OM tends to dominate the organic aerosol composition. Although the treatment of aging is crude, this result may be seen as broadly consistent with ambient data that show a predominance of oxygenated organic aerosol (OOA) over hydrophobic organic aerosol (HOA) (Zhang et al., 2007) as well as data that show that typically 60 % of OA is water soluble (Kerminen, 1997).

### 2.4 Overview of simulations

Table 2 summarizes the base case and different sensitivity simulations used in this work. All simulations are carried out for 14 months starting from 1 November 2000 with the first two months discarded from the analysis to allow for model initialization. Each simulation differs in the aerosol species included or the treatment of organic aerosol sources.

The NOCARB simulation has only sulfate and sea salt aerosols. It should be noted that the NOCARB simulation is different from the results presented in Trivitayanurak et al. (2008) due to the recent developments discussed in Sect. 2.3.1 that lower the predicted number concentrations. The BASE simulation includes sulfate, sea-salt, EC and OM with SOA production calculated as 10 % of monoterpene emissions.

In the POA-SOA split experiments, which include the 0SOA, 10SOA, 50SOA, 90SOA, and 100SOA simulations, the only change from the BASE simulation is that we omit the SOA production from monoterpenes. Instead, we take the current POA emission rates to be the total OA source and artificially vary the contribution of SOA production to the total OA by deducting  $f$  percent of mass from the POA source and condensing it as SOA. The 5 experimental simulations have SOA/OA source ratios of 0 %, 10 %, 50 %, 90 %, and 100 %.

As we increase the SOA contribution in these experiments, the hydrophobic/hydrophilic split is shifted since SOA is assumed to form as soluble species unlike POA emission that also has the insoluble fraction. However, due to relatively fast aging assumed in the model, hydrophilic OM dominates relatively quickly and thus the

composition of the 0SOA and 100SOA cases are not dramatically different. Therefore, the major differences between the simulations result from different aerosol size distributions stemming from the different microphysical pathways that form POA and SOA. Influence of composition to global CCN will be discussed again in Sect. 4.

## 3 Results

### 3.1 Global aerosol distributions

Figure 1a and b show the annual-average predicted aerosol number (CN10) and CCN(0.2%) concentrations for the model surface layer from the BASE simulation. CN10 is defined as condensation nuclei with dry diameter larger than 10 nm and CCN(0.2%) is cloud condensation nuclei at 0.2% supersaturation; all concentrations are  $\text{cm}^{-3}$  at 298 K and 1 atm. The predictions follow expected features with higher number concentrations over land than ocean. Predicted CN10 concentrations reach  $10\,000\text{ cm}^{-3}$  in the most polluted areas. Continental CN10 concentrations outside the most polluted regions range from 300 to  $3000\text{ cm}^{-3}$ . Marine boundary layer CN10 concentrations are roughly  $60\text{--}300\text{ cm}^{-3}$ , comparable with observed values (Andreae et al., 1995; Clarke et al., 1987; Covert et al., 1996; Fitzgerald, 1991; Pandis et al., 1995; Raes et al., 2000).

CCN(0.2%) predictions at the lowest model layer follow similar trends as CN10 predictions. CCN(0.2%) concentrations over the most polluted regions exceed  $1000\text{ cm}^{-3}$ . CCN(0.2%) concentrations typically range from 100 to  $1000\text{ cm}^{-3}$  over the continents, while they range from 25 to  $100\text{ cm}^{-3}$  over oceans in agreement with observations (Andreae et al., 1995).

### 3.2 Contribution of carbonaceous aerosols to CN and CCN

Figure 1c and d show the annual-average model surface layer CCN(0.2%) concentrations of the NOCARB simulation and the CCN(0.2%) enhancement ratios due to

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the additional carbonaceous aerosols in the BASE simulation. CCN(0.2%) concentrations increase over all continental and near-continental regions by  $\sim 20\%$  over polluted marine areas up to a tenfold increase in the biomass burning source regions. The increased CCN(0.2%) concentrations are found throughout the troposphere but are strongest within the Northern hemisphere where the majority of combustion sources are located. At the 300 hPa level, the CCN(0.2%) increase is spatially homogenous within hemispheres and around 20% over the S.H. and a  $\sim 60\%$  increase over the N.H. (not shown).

Table 3 presents globally and annually averaged aerosol number budget calculations for the NOCARB and BASE simulations. For these budget calculations, the burden includes only aerosols below the annual mean tropopause level. For ease of interpretation, source, growth, and sink terms are normalized by a fixed tropospheric volume assuming a 12 km tropopause. The budgets are categorized by size modes as ultrafine (UF), defined as particles with dry diameter below 80 nm, and CCN mode, defined as those with dry diameter above 80 nm. Note that we categorize budget components as source, microphysical growth, and sink. Coagulation can be either a sink for small particles or a microphysical growth term for larger particles. Carbonaceous aerosol increases annual-average primary emissions from 14 to  $35 \text{ cm}^{-3} \text{ day}^{-1}$ , almost all in the ultrafine mode. This increase in emissions raises the contribution of primary emissions to UF sources from 14 to 26%, the rest being nucleation. Among processes responsible for CCN production, increased emissions in the CCN-mode enlarges the contribution of primary emission relative to microphysical growth from 16 to 31%. SOA condensation accounts for only 5% of CCN sources in the BASE case, but it should be remembered that SOA in the BASE case is only  $18 \text{ Tg yr}^{-1}$ . The global UF number burden increases by 15% while CCN numbers double. Primary carbonaceous aerosol plays important roles in global CCN increase, both by direct emission at CCN sizes as well as by UF emissions that subsequently grow to CCN sizes by condensation. These latter processes can be seen in Table 3 in which the BASE simulation has higher CCN formation rates from  $\text{SO}_4$  condensation and aqueous oxidation of sulfate despite the

fact that sulfur emissions are the same in both simulations; emissions of UF carbonaceous aerosols increase the amount of sulfate condensing onto smaller particles and the number of UF particles that grow to CCN sizes. Lifetimes of UF- and CCN-mode aerosols change slightly from NOCARB to BASE simulations with similar contributions of each sink, e.g. coagulation being the largest sink for UF-mode particles, wet deposition as the major sink of CCN-mode.

### 3.3 Carbonaceous model evaluation

#### 3.3.1 Aerosol number

We compare the annual-average predicted CN10 concentrations in the surface layer with long-term CN observations shown in Table 4. The data is limited to sites outside of urban areas with a minimum sample of about one year. The sites included are part of a European network of sites presented in Van Dingenen et al. (2004), the Global Monitoring Division (GMD) of the Earth Systems Research Laboratory (Schnell, 2003) (<http://www.esrl.noaa.gov/gmd/>) and the Thompson Farm site of AIRMAP (<http://airmap.unh.edu/>). Locations of the CN observation sites are shown in Fig. 2.

Figure 3 presents the comparison of simulated CN10 against the observed CN10 at various sites for the NOCARB and BASE simulations. Types of observational sites are color-coded in the data points as blue for European sites, red for North American sites, green for remote sites, and cyan for free tropospheric sites. Before the implementation of carbonaceous aerosols, the model underpredicted aerosol number concentrations at most locations in the NOCARB simulation and most especially in more polluted locations. Without primary emissions of carbonaceous aerosols, model predictions had a log mean normalized bias (LMNB) of  $-0.26$ , meaning that it underpredicted on average by a factor of  $10^{-0.26} = 0.55$  ( $-45\%$  bias). The log mean normalized error (LMNE) was  $0.37$ , translated to an overall absolute error of a factor of  $10^{0.37} = 2.34$ . With carbonaceous aerosols, model predictions compare better with observations, resulting in LMNB of  $-0.032$  ( $-7\%$  bias). The LMNE is calculated to be  $0.19$ , which equals an

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



overall absolute error of a factor of 1.55. Introduction of carbonaceous aerosol markedly improved model predictions for European and North American sites while having minor impacts on the remote and free troposphere locations.

### 3.3.2 Carbonaceous aerosol mass

5 We compare the predicted organic carbon and elemental carbon with the measurements from the European Monitoring and Evaluation Program (EMEP) (<http://www.emep.int/index.html>) shown in Table 5. All measurements were made as part of the OC/EC campaign during year 2002 and 2003. Locations of monitoring sites are shown in Fig. 4.

10 Figure 5 summarizes the annual-average comparison of OC and EC concentrations predicted at the surface level in the BASE simulation comparing with the measurements. The model consistently underpredicts OC and EC mass with LMNB values of  $-0.67$  ( $-79\%$  bias) and  $-0.41$  ( $-61\%$  bias) for OC and EC, respectively. This degree of underprediction is not surprising for a model with a “traditional” (e.g. SOA from biogenic precursors only) treatment of organic aerosol. For example, similar global models have exhibited underpredictions of a factor of 2 to 10 (Chung and Seinfeld, 2002; Park et al., 2003; Liousse et al., 1996; Tsigaridis and Kanakidou, 2003; de Gouw et al., 2005; Volkamer et al., 2006; Heald et al., 2005, 2006) whereas some models have had only minor biases of  $\sim 20\%$  (Park et al., 2006). Overall, these comparisons suggest that  
20 our model has approximately the correct number of primary carbonaceous particles but insufficient carbonaceous mass on average. Because the ultimate contribution of carbonaceous aerosol to CCN depends on both number and (soluble) mass, we expect that our model underpredicts the CCN contribution somewhat.

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## 4 Effects of POA-SOA split on CCN(0.2%)

To explore the effect of varying the POA-SOA split on global CCN(0.2%) predictions, we conducted 5 POA-SOA split sensitivity runs as described in Sect. 2.5 ranging from treating OA as pure POA (0SOA simulation) to all OA as SOA (100SOA simulation).

5 These runs predict what the tropospheric aerosol would look like in a POA-dominant world versus a SOA-dominant world. As the model has a relatively fast aging (hydrophobic – hydrophilic conversion), the composition of the OA in all scenarios is similar and mostly dominated by hydrophilic OA. Therefore, these sensitivity results highlight the microphysical effects of POA versus SOA on CCN formation as reflected in the  
10 aerosol number distributions. CCN(0.2%) concentrations in the model surface layer are the focus here because boundary layer CCN are thought to be most important to altering stratiform clouds, which contribute most to global albedo.

### 4.1 Effect on global CCN distribution

Figure 6 presents annual-average CCN(0.2%) concentrations in the surface layer predicted from the 10SOA simulation and their changes in the 90SOA simulation, which are presented as ratios of CCN(0.2%) concentrations in the 90SOA to 10SOA simulations. In general, the result of increasing SOA in the POA-SOA split is reduction of  
15 CCN(0.2%) by 10–30% away from biomass burning source regions and up to 80% decreases over the biomass burning source regions. Over the oceans, where carbonaceous aerosols make a small contribution as shown in Fig. 1d, the changes in  
20 CCN(0.2%) are less than 10%, mostly decreases, due to differences in background aerosol concentrations in the two simulations.

There are three effects occurring to the CCN(0.2%) population when we increase SOA in the split: (1) direct decrease of CCN(0.2%) by reducing CCN-mode primary  
25 emissions and (2) reduction of UF-mode primary emission that might grow to CCN, and (3) additional condensation of SOA that may or may not contribute to CCN production, depending on what size of pre-existing particles that SOA condenses onto.

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To contribute to CCN(0.2%) formation, SOA must condense onto UF-mode particles. The overall tendency to decrease CCN when shifting POA to SOA indicates that the first two effects combined are stronger than the third and highlights the importance of primary particles to the global CCN budget (Adams and Seinfeld, 2003).

Comparing the 90SOA to 10SOA simulations, major CCN(0.2%) reductions are localized in biomass burning source regions, and more modest decreases occur in the fossil fuel combustion source regions such as North America or East Asia. This is largely because there are available ultrafine sulfate aerosols (resulting either from assumed in-plume nucleation events or regional-scale nucleation events) in polluted source regions to compensate for the reduced UF-mode POA emission whereas primary carbonaceous particles are the dominant source of aerosol number in biomass burning regions. This will be discussed more in the next section.

## 4.2 Effect on size distributions

Figure 7a and b show number size distributions from the 10SOA and 90SOA simulations for Congo (12.5–32.5° E and 0 to 8° S) biomass burning region and a North American (107.5–112.5° W and 48–52° N) location away from urban areas; these two locations are displayed on the map in Fig. 6b as insets number 1 and 2, respectively. We show aerosol surface area distributions for these two locations in Fig. 7c and d to demonstrate the distribution of SOA condensation. Generally, as the fraction of SOA increases, the number of particles decreases and the size distribution shifts to larger sizes. This is to be expected since the number of particles decreases while the mass remains relatively constant. In the Congo region, as OM dominates aerosol mass in the submicron mode (83%) and ultrafine mode (62%) in the 10SOA case, reducing primary emission of OM results in decreased surface area below ~500 nm diameter. We define the ultrafine surface fraction (USF) as the fraction of aerosol surface area in the ultrafine mode ( $D_p < 0.1 \mu\text{m}$ ) with respect to the total surface; this represents the potential of SOA condensation to grow UF particles to CCN sizes. Figure 7c for the Congo shows a reduction of USF from 11% in 10SOA case to 6% in 90SOA case.

### Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





more than chemical composition. Doing a similar calculation on the global tropospheric CCN(0.2%) burden shows that POA is 2.5 times as effective at producing CCN(0.2%) per unit mass relative to SOA.

In terms of absolute concentrations, the reduction of surface CCN(0.2%) is from 160 to 125 cm<sup>-3</sup> for 0SOA and 100SOA case, respectively, which is about a 20% reduction. The reduction of the global tropospheric CCN(0.2%) burden is also approximately 20% going from 0 to 100% SOA/OA. The 20% change of global CCN(0.2%) concentration due to the change in microphysical path of OA formation is comparable to the influence of chemical composition/hygroscopicity. Pierce et al. (2007) found less than a 20% reduction of global CCN(0.2%) when changing carbonaceous aerosols from highly soluble scenario to a completely insoluble scenario.

Thus far, our discussion has focused on CCN at 0.2% supersaturation. At lower in-cloud supersaturations, CCN is limited to larger particles. In Fig. 7, we see that higher SOA fractions shift the size distribution to larger sizes, increasing the number of CCN active at lower supersaturations. These potentially “giant” cloud condensation nuclei (GCCN) may be important for cloud evolution and precipitation formation. We looked at the change in predicted CCN at different supersaturations down to 0.01% for different POA-SOA splits. We found that POA enhanced CCN concentrations at all supersaturations greater than 0.05%, below which SOA is more effective at producing such “giant” CCN. However, even at very high SOA fractions (90SOA case), the enhancement in CCN(0.01%) is still modest (within 4%) compared to the 10SOA case, so we conclude that there is no strong impact of SOA on the formation of giant CCN.

## 5 Conclusions

Carbonaceous aerosols have been implemented in the TOMAS microphysics module inside the global chemical transport model, GEOS-Chem. This version of TOMAS tracks aerosol, both number and mass, in 30 size sections covering the sizes from 10 nm to 10 μm. Hydrophobic OM, hydrophilic OM, pure EC and mixed EC are tracked

### Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in the model. At this point, aerosol mass species in the model include sulfate, sea salt, EC and OM.

Contributions of carbonaceous aerosol to the CN10 and CCN(0.2%) predictions have been examined in comparison to a simulation with only sulfate and sea salt aerosols. In the model surface layer, CN10 increases localized around the carbonaceous source regions. With the introduction of carbonaceous aerosols, primary number emissions increase by a factor of 2.5, and annual-average global CCN(0.2%) burden increases by a factor of 2. Inclusion of carbonaceous aerosols improves comparison of CN10 predictions to a set of long-term observations at various locations with a change from 45% underprediction on average to 7% underprediction on average. The total source of OA in this model, 55 to 73 Tg yr<sup>-1</sup> depending on the simulation, is low compared to more recent observationally driven estimates (Spracklen et al., 2011; Heald et al., 2010). As a result, comparison of predicted EC and OC to measurements shows that the model underpredicts carbonaceous mass concentrations in the European boundary layer by a factor of 2–4, similar to several other “traditional” global models.

We performed simulations to assess the sensitivity of CCN production to the POA-SOA split by artificially shifting OA sources from primary POA emissions to SOA condensation while holding the total OA source constant. SOA and POA contribute to CCN via different microphysical pathways. Because the model assumes that rapid (1.15 days) aging of OM from hydrophobic to hydrophilic occurs, the overall CCN activity of OM in these simulations are quite similar, and differences primarily reflect the different microphysical pathways. The main finding is that CCN(0.2%) decreases nearly everywhere as the model changes from a POA-dominated world to an SOA-dominated world. This is because SOA condenses mostly onto the accumulation mode, thus not promoting any CCN(0.2%) growth. In contrast, POA emissions add significant numbers of particles, either directly to the CCN mode or to the ultrafine mode, a fraction of which grow to become CCN. Not surprisingly, the reductions in CCN(0.2%) when shifting from POA to SOA are strongest in biomass burning regions, which are dominated

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by carbonaceous aerosol. In contrast, CCN(0.2%) reductions are less pronounced in locations with abundant inorganic ultrafine emissions because these provide condensational sink for SOA leading to CCN growth, which compensates for the missing POA.

We find that, compared to SOA, POA is about twice as effective per unit mass at CCN production at the model surface. By changing from completely POA to completely SOA, the CCN(0.2%) in the lowest model layer differ by about 20% compared to the values in the completely POA case. In contrast, Pierce et al. (2007) found that global CCN(0.2%) changed by less than 20% on average for a drastic change in assumed OM solubility (a change from highly CCN active to completely insoluble OM). We conclude that the SOA-POA split has a significant effect on global CCN, and that the microphysical implications of POA versus SOA appear to be at least as important as differences in chemical composition. We conclude that carbonaceous aerosol overall makes a significant contribution to global CCN; therefore, a better understanding of its total source will aid climate change simulations.

## References

- Adams, P. J. and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, *J. Geophys. Res.-Atmos.*, 107, 4370–4392, doi:10.1029/2001JD001010, 2002.
- Adams, P. J. and Seinfeld, J. H.: Disproportionate impact of particulate emissions on global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 30, 1239–1242, doi:10.1029/2002GL016303, 2003.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, 1989.
- Andreae, M. O., Elbert, W., and Demora, S. J.: Biogenic Sulfur Emissions and Aerosols over the Tropical South-Atlantic. 3. Atmospheric Dimethylsulfide, Aerosols and Cloud Condensation Nuclei, *J. Geophys. Res.-Atmos.*, 100, 11335–11356, 1995.
- Bey, I., Jacob, D. J., Logan, J. A., and Yantosca, R. M.: Asian chemical outflow to the Pacific in spring: Origins, pathways, and budgets, *J. Geophys. Res.-Atmos.*, 106, 23097–23113, 2001.

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.-Atmos.*, 109, D14203, doi:10.1029/2003JD003697, 2004.
- Cabada, J. C., Pandis, S. N., Subramanian, R., Robinson, A. L., Polidori, A., and Turpin, B.: Estimating the secondary organic aerosol contribution to PM<sub>2.5</sub> using the EC tracer method, *Aerosol Sci. Technol.*, 38, 140–155, 2004.
- Chung, S. H. and Seinfeld, J. H.: Global distribution and climate forcing of carbonaceous aerosols, *J. Geophys. Res.-Atmos.*, 107, 4407, doi:10.1029/2001JD001397, 2002.
- Clarke, A. D., Ahlquist, N. C., and Covert, D. S.: The Pacific Marine Aerosol-Evidence for Natural Acid Sulfates, *J. Geophys. Res.-Atmos.*, 92, 4179–4190, 1987.
- Cooke, W. F., Liousse, C., Cachier, H., and Feichter, J.: Construction of a 1 degrees × 1 degrees fossil fuel emission data set for carbonaceous aerosol and implementation and radiative impact in the ECHAM4 model, *J. Geophys. Res.-Atmos.*, 104, 22137–22162, 1999.
- Covert, D. S., Kapustin, V. N., Bates, T. S., and Quinn, P. K.: Physical properties of marine boundary layer aerosol particles of the mid-Pacific in relation to sources and meteorological transport, *J. Geophys. Res.-Atmos.*, 101, 6919–6930, 1996.
- de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M., Fehsenfeld, F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C., Marchewka, M., Bertman, S. B., and Bates, T. S.: Budget of organic carbon in a polluted atmosphere: Results from the New England Air Quality Study in 2002, *J. Geophys. Res.-Atmos.*, 110, D16305, doi:10.1029/2004JD005623, 2005.
- El-Zanan, H. S., Lowenthal, D. H., Zielinska, B., Chow, J. C., and Kumar, N.: Determination of the organic aerosol mass to organic carbon ratio in IMPROVE samples, *Chemosphere*, 60, 485–496, 2005.
- Fitzgerald, J. W.: Marine Aerosols – a Review, *Atmos. Environ. A-Gen.*, 25, 533–545, 1991.
- Giglio, L., van der Werf, G. R., Randerson, J. T., Collatz, G. J., and Kasibhatla, P.: Global estimation of burned area using MODIS active fire observations, *Atmos. Chem. Phys.*, 6, 957–974, doi:10.5194/acp-6-957-2006, 2006.
- Griffin, R. J., Cocker, D. R., Seinfeld, J. H., and Dabdub, D.: Estimate of global atmospheric organic aerosol from oxidation of biogenic hydrocarbons, *Geophys. Res. Lett.*, 26, 2721–2724, 1999.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and



## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.

Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.

Hanel, G.: Single-Scattering Albedo of Atmospheric Aerosol-Particles as a Function of Relative Humidity, *J. Atmos. Sci.*, 33, 1120–1124, 1976.

Heald, C. L., Jacob, D. J., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H., and Weber, R. J.: A large organic aerosol source in the free troposphere missing from current models, *Geophys. Res. Lett.*, 32, L18809, doi:10.1029/2005GL023831, 2005.

Heald, C. L., Jacob, D. J., Turquety, S., Hudman, R. C., Weber, R. J., Sullivan, A. P., Peltier, R. E., Atlas, E. L., de Gouw, J. A., Warneke, C., Holloway, J. S., Neuman, J. A., Flocke, F. M., and Seinfeld, J. H.: Concentrations and sources of organic carbon aerosols in the free troposphere over North America, *J. Geophys. Res.-Atmos.*, 111, D23S47, doi:10.1029/2006JD007705, 2006.

Heald, C. L., Ridley, D. A., Kreidenweis, S. M., and Drury, E. E.: Satellite observations cap the atmospheric organic aerosol budget, *Geophys. Res. Lett.*, 37, L24808, doi:10.1029/2010GL045095, 2010.

Hitzenberger, R., Berner, A., Giebl, H., Kromp, R., Larson, S. M., Rouc, A., Koch, A., Marischka, S., and Puxbaum, H.: Contribution of carbonaceous material to cloud condensation nuclei concentrations in European background (Mt. Sonnblick) and urban (Vienna) aerosols, *Atmos. Environ.*, 33, 2647–2659, 1999.

IPCC: Intergovernmental Panel on Climate Change, *Climate Change 2007: The Physical Science Basis. Summary for Policymakers*, Cambridge Univ. Press, New York, 2007.

Jathar, S. H., Farina, S. C., Robinson, A. L., and Adams, P. J.: The influence of semi-volatile and reactive primary emissions on the abundance and properties of global organic aerosol, *Atmos. Chem. Phys.*, 11, 7727–7746, doi:10.5194/acp-11-7727-2011, 2011.

Johnson, D., Utembe, S. R., Jenkin, M. E., Derwent, R. G., Hayman, G. D., Alfarra, M. R., Coe, H., and McFiggans, G.: Simulating regional scale secondary organic aerosol formation

---

**Does the POA-SOA split matter for global CCN formation?**

---

W. Trivitayanurak and  
P. J. Adams

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

during the TORCH 2003 campaign in the southern UK, *Atmos. Chem. Phys.*, 6, 403–418, doi:10.5194/acp-6-403-2006, 2006.

Kanakidou, M., Tsigaridis, K., Dentener, F. J., and Crutzen, P. J.: Human-activity-enhanced formation of organic aerosols by biogenic hydrocarbon oxidation, *J. Geophys. Res.-Atmos.*, 105, 9243–9254, 2000.

Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.

Kerminen, V. M.: The effects of particle chemical character and atmospheric processes on particle hygroscopic properties, *J. Aerosol Sci.*, 28, 121–132, 1997.

Koch, D., Jacob, D., Tegen, I., Rind, D., and Chin, M.: Tropospheric sulfur simulation and sulfate direct radiative forcing in the Goddard Institute for Space Studies general circulation model, *J. Geophys. Res.-Atmos.*, 104, 23799–23822, 1999.

Kondo, Y., Miyazaki, Y., Takegawa, N., Miyakawa, T., Weber, R. J., Jimenez, J. L., Zhang, Q., and Worsnop, D. R.: Oxygenated and water-soluble organic aerosols in Tokyo, *J. Geophys. Res.-Atmos.*, 112, D01203, doi:10.1029/2006JD007056, 2007.

Kumagai, K., Iijima, A., Tago, H., Tomioka, A., Kozawa, K., and Sakamoto, K.: Seasonal characteristics of water-soluble organic carbon in atmospheric particles in the inland Kanto plain, Japan, *Atmos. Environ.*, 43, 3345–3351, 2009.

Laaksonen, A., Korhonen, P., Kulmala, M., and Charlson, R. J.: Modification of the Kuehler equation to include soluble trace gases and slightly soluble substances, *J. Atmos. Sci.*, 55, 853–862, 1998.

Lin, S. J. and Rood, R. B.: Multidimensional flux-form semi-Lagrangian transport schemes, *Mon. Weather Rev.*, 124, 2046–2070, 1996.

Liousse, C., Penner, J. E., Chuang, C., Walton, J. J., Eddleman, H., and Cachier, H.: A global three-dimensional model study of carbonaceous aerosols, *J. Geophys. Res.-Atmos.*, 101, 19411–19432, 1996.

Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, *Atmos. Chem. Phys.*, 9, 8601–8616, doi:10.5194/acp-9-8601-2009, 2009.

**Does the POA-SOA split matter for global CCN formation?**W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pandis, S. N., Wexler, A. S., and Seinfeld, J. H.: Dynamics of Tropospheric Aerosols, *J. Phys. Chem.*, 99, 9646–9659, 1995.
- Park, R. J., Jacob, D. J., Chin, M., and Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, *J. Geophys. Res.-Atmos.*, 108, 4355, doi:10.1029/2002JD003190, 2003.
- Park, R. J., Jacob, D. J., Palmer, P. I., Clarke, A. D., Weber, R. J., Zondlo, M. A., Eisele, F. L., Bandy, A. R., Thornton, D. C., Sachse, G. W., and Bond, T. C.: Export efficiency of black carbon aerosol in continental outflow: Global implications, *J. Geophys. Res.-Atmos.*, 110, D11205, doi:10.1029/2004JD005432, 2005.
- Park, R. J., Jacob, D. J., Kumar, N., and Yantosca, R. M.: Regional visibility statistics in the United States: Natural and transboundary pollution influences, and implications for the Regional Haze Rule, *Atmos. Environ.*, 40, 5405–5423, 2006.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.
- Pierce, J. R. and Adams, P. J.: Efficiency of cloud condensation nuclei formation from ultrafine particles, *Atmos. Chem. Phys.*, 7, 1367–1379, doi:10.5194/acp-7-1367-2007, 2007.
- Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation and primary emission rates, *Atmos. Chem. Phys.*, 9, 1339–1356, doi:10.5194/acp-9-1339-2009, 2009.
- Pierce, J. R., Chen, K., and Adams, P. J.: Contribution of primary carbonaceous aerosol to cloud condensation nuclei: processes and uncertainties evaluated with a global aerosol microphysics model, *Atmos. Chem. Phys.*, 7, 5447–5466, doi:10.5194/acp-7-5447-2007, 2007.
- Pierce, J. R., Theodoritsi, G., Adams, P. J., and Pandis, S. N.: Parameterization of the effect of sub-grid scale aerosol dynamics on aerosol number emission rates, *J. Aerosol Sci.*, 40, 385–393, 2009.
- Pun, B. K., Wu, S. Y., Seigneur, C., Seinfeld, J. H., Griffin, R. J., and Pandis, S. N.: Uncertainties in modeling secondary organic aerosols: Three-dimensional modeling studies in Nashville/Western Tennessee, *Environ. Sci. Technol.*, 37, 3647–3661, 2003.
- Pye, H. O. T. and Seinfeld, J. H.: A global perspective on aerosol from low-volatility organic compounds, *Atmos. Chem. Phys.*, 10, 4377–4401, doi:10.5194/acp-10-4377-2010, 2010.

**Does the POA-SOA split matter for global CCN formation?**W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Raes, F., Van Dingenen, R., Vignati, E., Wilson, J., Putaud, J. P., Seinfeld, J. H., and Adams, P.: Formation and cycling of aerosols in the global troposphere, *Atmos. Environ.*, 34, 4215–4240, 2000.

Raymond, T. M. and Pandis, S. N.: Cloud activation of single-component organic aerosol particles, *J. Geophys. Res.-Atmos.*, 107, 4787, doi:10.1029/2002JD002159, 2002.

Raymond, T. M. and Pandis, S. N.: Formation of cloud droplets by multicomponent organic particles, *J. Geophys. Res.-Atmos.*, 108, 4469, doi:10.1029/2003JD003503, 2003.

Riipinen, I., Pierce, J. R., Yli-Juuti, T., Nieminen, T., Häkkinen, S., Ehn, M., Junninen, H., Lehtipalo, K., Petäjä, T., Slowik, J., Chang, R., Shantz, N. C., Abbatt, J., Leaitch, W. R., Kerminen, V.-M., Worsnop, D. R., Pandis, S. N., Donahue, N. M., and Kulmala, M.: Organic condensation: a vital link connecting aerosol formation to cloud condensation nuclei (CCN) concentrations, *Atmos. Chem. Phys.*, 11, 3865–3878, doi:10.5194/acp-11-3865-2011, 2011.

Saxena, P. and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical review of the literature and application of thermodynamics to identify candidate compounds, *J. Atmos. Chem.*, 24, 57–109, 1996.

Schnell, R. C.: Chapter 3: Aerosols and Radiation, in: *Climate Monitoring and Diagnostics Laboratory Summary Report*, edited by: McComisky, A., 2002–2003, 2003.

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons, New York, 1998.

Seinfeld, J. H., Erdakos, G. B., Asher, W. E., and Pankow, J. F.: Modeling the formation of secondary organic aerosol (SOA). 2. The predicted effects of relative humidity on aerosol formation in the  $\alpha$ -pinene-,  $\beta$ -pinene-, sabinene-,  $\Delta^3$ -carene-, and cyclohexene-ozone systems *Environ. Sci. Technol.*, 35, 3272–3272, 2001.

Shrivastava, M. K., Subramanian, R., Rogge, W. F., and Robinson, A. L.: Sources of organic aerosol: Positive matrix factorization of molecular marker data and comparison of results from different source apportionment models, *Atmos. Environ.*, 41, 9353–9369, 2007.

Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, A., and Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos. Chem. Phys.*, 11, 12109–12136, doi:10.5194/acp-11-12109-2011, 2011.

Subramanian, R., Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F., and Robinson, A. L.: Insights into the primary-secondary and regional-local contributions to organic aerosol and PM<sub>2.5</sub> mass in Pittsburgh, Pennsylvania, *Atmos. Environ.*, 41, 7414–7433, 2007.

**Does the POA-SOA split matter for global CCN formation?**W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Trivitayanurak, W., Adams, P. J., Spracklen, D. V., and Carslaw, K. S.: Tropospheric aerosol microphysics simulation with assimilated meteorology: model description and intermodel comparison, *Atmos. Chem. Phys.*, 8, 3149–3168, doi:10.5194/acp-8-3149-2008, 2008.
- Tsigaridis, K. and Kanakidou, M.: Global modelling of secondary organic aerosol in the troposphere: a sensitivity analysis, *Atmos. Chem. Phys.*, 3, 1849–1869, doi:10.5194/acp-3-1849-2003, 2003.
- Twomey, S.: Pollution and Planetary Albedo, *Atmos. Environ.*, 8, 1251–1256, 1974.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.
- Van Dingenen, R., Raes, F., Putaud, J. P., Baltensperger, U., Charron, A., Facchini, M. C., Decesari, S., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Hüglin, C., Jones, A. M., Laj, P., Lorbeer, G., Maenhaut, W., Palmgren, F., Querol, X., Rodriguez, S., Schneider, J., ten Brink, H., Tunved, P., Torseth, K., Wehner, B., Weingartner, E., Wiedensohler, A., and Wahlin, P.: A European aerosol phenomenology-1: physical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2561–2577, 2004.
- Volkamer, R., Jimenez, J. L., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L. T., Worsnop, D. R., and Molina, M. J.: Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophys. Res. Lett.*, 33, L17811, doi:10.1029/2006GL026899, 2006.
- Vutukururu, S., Griffin, R. J., and Dabdub, D.: Simulation and analysis of secondary organic aerosol dynamics in the South Coast Air Basin of California, *J. Geophys. Res.-Atmos.*, 111, D10S12, doi:10.1029/2005JD006139, 2006.
- Wexler, A. S., Lurmann, F. W., and Seinfeld, J. H.: Modelling urban and regional aerosols – I. Modeling development, *Atmos. Environ.*, 28, 531–546, 1994.
- Yu, L. E., Shulman, M. L., Kopperud, R., and Hildemann, L. M.: Fine organic aerosols collected in a humid, rural location (Great Smoky Mountains, Tennessee, USA): Chemical and temporal characteristics, *Atmos. Environ.*, 39, 6037–6050, 2005.
- Zhang, L. M., Gong, S. L., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environ.*, 35, 549–560, 2001.

Zhang, Q., Worsnop, D. R., Canagaratna, M. R., and Jimenez, J. L.: Hydrocarbon-like and oxygenated organic aerosols in Pittsburgh: insights into sources and processes of organic aerosols, *Atmos. Chem. Phys.*, 5, 3289–3311, doi:10.5194/acp-5-3289-2005, 2005.

5 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M.,  
10 and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007GL029979, 2007.

**Does the POA-SOA split matter for global CCN formation?**

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Emission and production rates of OM and EC in BASE scenario.

Sources	OM sources (Tgyr <sup>-1</sup> )	EC sources (Tgyr <sup>-1</sup> )	Reference
Primary Emissions			
Fossil fuel	5	3.5	Bond et al. (2004)
Biofuel	13	1.6	Bond et al. (2004)
Biomass burning	37	2.6	GFED2, Giglio et al. (2006)
Secondary production			
Biogenic SOA	18	–	10% of monoterpenes from MEGAN, Guenther et al. (2006)



## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Overview of simulations.

Name	EC Emission Rate (Tgyr <sup>-1</sup> )	POA Emission Rate (Tgyr <sup>-1</sup> )	SOA Production Rate (Tgyr <sup>-1</sup> )
NOCARB	None	None	None
BASE	8	55	18
0SOA	8	55	0
10SOA	8	50	6
50SOA	8	28	28
90SOA	8	6	50
100SOA	8	0	55

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 3.** Global annual-average aerosol number budgets comparing NOCARB and BASE simulations. Size mode for ultrafine (UF) covers 0.01–0.08  $\mu\text{m}$  and CCN-mode covers 0.08–10  $\mu\text{m}$ .

Size mode	NOCARB		BASE	
	UF	CCN	UF	CCN
Sources ( $\text{cm}^{-3} \text{day}^{-1}$ )				
Primary emissions	13	1.2	31	4
Nucleation ( $J_{10}$ )	78	0	89	0
Total	91	1.2	120	4
Microphysical growth ( $\text{cm}^{-3} \text{day}^{-1}$ )				
SO <sub>4</sub> Condensation	−2.1	2.1	−2.5	2.5
Aqueous oxidation	−4.2	4.2	−5.8	5.8
SOA condensation	−	−	−0.6	0.6
Total	−6.3	6.3	−8.9	8.9
Sinks ( $\text{cm}^{-3} \text{day}^{-1}$ )				
Dry deposition	8.8	1.2	15	2.6
Wet deposition	6.7	6.2	8.5	10
Coagulation	69	0	87	0.6
Total	85	7.4	111	13
Burden ( $\text{cm}^{-3}$ )	750	29	871	61
Lifetime (days)	9	4	8	5

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams

**Table 4.** Locations of number concentration measurements used for comparison.


Location	Region	Reference	Time	Long.	Lat.	Elevation (m)	CN10 (cm <sup>-3</sup> )
A Aspvereten, Sweden	Europe	Van Dingenen et al. (2004)	Jan–Dec 2001	17.4° E	58.8° N	20	2000
B Harwell, United Kingdom	Europe	Van Dingenen et al. (2004)	May 1998–Nov 2000	1.3° W	51.6° N	125	3000
C Hohenpeissenberg, Germany	Europe	Van Dingenen et al. (2004)	Apr 1998–Aug 2000	11° E	47.8° N	988	2500
D Melpitz, Germany	Europe	Van Dingenen et al. (2004)	Dec 1996–Nov 1997	12.9° E	51.5° N	86	5600
E Ispra, Italy	Europe	Van Dingenen et al. (2004)	Feb 2000–Dec 2000	8.6° E	45.8° N	209	9000
F Thompson Farm, New Hampshire, US	North America	http://airmap.unh.edu	2001–2005	71° W	43.1° N	75	7250
G Lamont, Oklahoma, US	North America	http://www.cmdl.noaa.gov/aero/data/	1996–2004	97.5° W	36.5° N	318	5200
H Bondville, Illinois, US	North America	http://www.cmdl.noaa.gov/aero/data/	1994–2005	88.3° W	40.1° N	230	3700
I Sable Island, Nova Scotia, Canada	North America	http://www.cmdl.noaa.gov/aero/data/	1992–1999	60° W	43.9° N	5	850
J Trinidad Head, California, US	North America	http://www.cmdl.noaa.gov/aero/data/	2002–2005	124.2° W	41.1° N	107	590
K American Samoa	Remote	http://www.cmdl.noaa.gov/aero/data/	1995–2005	170.5° W	14.2° S	42	220
L South Pole	Remote	http://www.cmdl.noaa.gov/aero/data/	1995–2005	102° E	90° S	2810	100
M Point Barrow, Alaska, US	Remote	http://www.cmdl.noaa.gov/aero/data/	1995–2005	156.6° W	71.3° N	11	110
N Mauna Loa, Hawaii, US	Free Troposphere	http://www.cmdl.noaa.gov/aero/data/	1995–2005	155.6° W	19.5° N	3397	330
O Jungfraujoch, Switzerland	Free Troposphere	Van Dingenen et al. (2004)	Jun 1997–May 1998	8° E	47.6° N	3580	525

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams




**Fig. 1.** Annual-average number concentrations in the surface layer ( $\text{cm}^{-3}$  at 298 K, 1 atm) for **(a)** CN10 from BASE simulation; **(b)** CCN(0.2%) from BASE simulation; **(c)** CCN(0.2%) from NOCARB simulation, and **(d)** ratios of predicted annual-average CCN(0.2%) concentrations in the surface layer as  $(\text{CCN BASE})/(\text{CCN NOCARB})$ .

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[⏴](#)
[⏵](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams




**Fig. 2.** Locations of aerosol number concentration (CN10) measurements used for model evaluation. Letters correspond to those listed in Table 4. Dark blue letters denote European boundary layer, red letters denote polluted North American boundary layer, green is for remote boundary layer, and cyan denote free troposphere.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams




**Fig. 3.** Comparison of simulated CN10 concentrations to observed CN10 ( $\text{cm}^{-3}$  at 298 K and 1 atm) for **(a)** NOCARB and **(b)** BASE simulations. Solid line shows a 1 : 1 ratio and dashed lines show ratios of 10 : 1 and 1 : 10. The letters refer to the locations presented in Table 4 and Fig. 2. Blue letters refer to European sites. Red letters refer to North Americans sites. Green letters refer to remote sites. Cyan letters refer to free tropospheric sites. Log-mean normalized bias (LMNB) and log-mean normalized error (LMNE) are listed in each panel.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[⏴](#)
[⏵](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams




**Fig. 4.** Locations of carbonaceous aerosol mass concentration measurements used in model evaluation. Letters correspond to the list in Table 5.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams



**Fig. 5.** Comparison of simulated concentrations to observed concentrations ( $\mu\text{gC m}^{-3}$  at 298 K and 1 atm) for **(a)** organic carbon and **(b)** elemental carbon. Solid line shows a 1 : 1 ratio and dashed lines show ratios of 1 : 10. The letters refer to the locations presented in Table 5 and Fig. 4. Log-mean normalized bias (LMNB) and log-mean normalized error (LMNE) are listed in each panel.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 6.** (a) Predicted annual-average CCN(0.2%) concentrations ( $\text{cm}^{-3}$  at 298 K and 1 atm) in the surface layer for the 10SOA simulation; (b) Ratios of predicted annual-average surface CCN(0.2%) concentrations:  $(\text{CCN } 90\text{SOA})/(\text{CCN } 10\text{SOA})$ .

## Does the POA-SOA split matter for global CCN formation?

W. Trivittayanurak and  
P. J. Adams




**Fig. 7.** Annual-average number size distributions for the surface layer of the **(a)** Congo (12.5–32.5° E and 0 to 8° S) and **(b)** North American (107.5–112.5° W and 48–52° N), regions, which are displayed as insets on Fig. 6 **(b)**; annual-average surface area distribution for **(c)** Congo and **(d)** North America, respectively. The distributions from the 10SOA simulations are shown as black lines and those from the 90SOA simulations as grey lines.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Does the POA-SOA split matter for global CCN formation?

W. Trivitayanurak and  
P. J. Adams



**Fig. 8.** Global-average CCN(0.2%) concentrations predicted in different POA-SOA sensitivity simulations for the model surface layer as well as the entire troposphere. The 10SOA run corresponds to the 10% SOA/OA point on the x-axis. Black circles and lines denote surface CCN(0.2%) concentrations with values displayed on the left vertical axis. Grey triangles and lines denote tropospheric average with values displayed on the right vertical axis. The corresponding CCN(0.2%) concentrations from the NOCARB simulation are also shown for reference as black and grey horizontal dashed lines.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)