

Gravity wave
influence on NLC at
ALOMAR

H. Wilms et al.

Gravity wave influence on NLC: experimental results from ALOMAR, 69° N

H. Wilms¹, M. Rapp^{1,2}, P. Hoffmann³, J. Fiedler³, and G. Baumgarten³

¹Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

²Meteorologisches Institut München, Ludwig-Maximilians-Universität München, Munich, Germany

³Leibniz Institute of Atmospheric Physics, Schlosstr. 6, 18225 Kühlungsborn, Germany

Received: 6 June 2013 – Accepted: 14 July 2013 – Published: 31 July 2013

Correspondence to: H. Wilms (henrike.wilms@dlr.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The influence of gravity waves on noctilucent clouds (NLC) at ALOMAR (69° N) is analyzed by relating gravity wave activity to NLC occurrence from common-volume measurements. Gravity wave kinetic energies are derived from MF-radar wind data and filtered into different period ranges by wavelet transformation. From the data set covering the years 1999–2011 a direct correlation between gravity wave kinetic energy and NLC occurrence is not found. I.e., NLC appear independently of the simultaneously measured gravity wave kinetic energy. In addition, gravity wave activity is divided into weak and strong activity as compared to a 13 yr mean. The NLC occurrence rates during strong and weak activity are calculated separately for a given wave period and compared to each other. Again, for the full data set no dependence of NLC occurrence on relative gravity wave activity is found. However, concentrating on 12 h of NLC detections during 2008, we do find an NLC-amplification with strong long-period gravity wave occurrence. Our analysis hence confirms previous findings that in general NLC at ALOMAR are not predominantly driven by gravity waves while exceptions to this rule are at least possible.

1 Introduction

Noctilucent clouds (NLC) are a remarkable feature of the polar summer mesopause. These clouds, first documented by Leslie (1885), Jesse (1885) and Backhouse (1885), are the direct indication of extremely low temperatures enabling ice particle growth to visible sizes. Owing to the high altitude of about 83 km NLC can be observed by naked eye when the sun is 6° to 15° below the horizon and illuminates the thin clouds from below (Gadsden and Schröder, 1989).

NLC are subject to a high degree of variability on scales ranging from minutes to decades. Long-term trends have been identified in the brightness of polar mesospheric clouds (DeLand et al., 2007) as well as their occurrence frequency (Shettle

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2009). According to model results these trends are mainly due to temperature changes of the mesopause, induced by the shrinking of the stratosphere, and solar cycle modulations of mesospheric water vapor content (Lübken et al., 2009; Lübken and Berger, 2011). On the shorter end of scales, namely minutes to hours, NLC are highly influenced by dynamical features of the mesopause. For example, diurnal and semi-diurnal variations of NLC occurrence frequency, brightness and central height due to tides have been detected in lidar measurements at several sites, including South Pole (Chu et al., 2001), Rothera, Antarctica (Chu et al., 2006) and ALOMAR, Norway (von Zahn et al., 1998; Fiedler et al., 2005, 2011). Gravity waves can induce temperature fluctuations as high as 10 K (e.g., Rapp et al., 2002), which influence in a complex interplay with vertical winds and sedimentation the growth of ice particles. The signature of wave-like structures in NLC have been used to determine gravity wave parameters from ground-based (e.g., Witt, 1962; Pautet et al., 2011) and satellite-based (e.g., Chandran et al., 2009) images and recently also from high resolution lidar data (Käfler et al., 2013). NLC therefore open a possibility to study small scale dynamics in the mesopause which are not accessible by other remote sensing methods. However, for the correct interpretation of NLC variability in terms of gravity wave activity it is of crucial importance to clearly understand all processes modifying NLC parameters.

Model simulations of NLC under the influence of gravity waves have shown that short period gravity waves tend to destroy NLC, whereas longer period gravity waves enhance NLC (Turco et al., 1982; Jensen and Thomas, 1994; Rapp et al., 2002; Chandran et al., 2012), with the transition period lying at 6.5 h (Rapp et al., 2002). This anticorrelation for short period gravity waves has been confirmed by lidar measurements at the South Pole (Chu et al., 2009) and in Sondrestrom, Greenland (Thayer et al., 2003; Gerrard et al., 2004). On the contrary, no such anticorrelation was found in Rothera, Antarctica (Chu et al., 2009), Davis, Antarctica (Innis et al., 2008) and ALOMAR, Norway (Schöch, 2007). The applied methodology was basically the same at all locations: Density fluctuations in the stratosphere serve as a measure for gravity wave activity, which is then correlated to mesospheric cloud parameters. The underlying assumption

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion thereby is that gravity waves in the stratosphere are closely related to those in the mesosphere. Consequently, modifications of the gravity wave spectrum due to filtering processes by the background wind (Eckermann, 1995) are not taken into account, as well as non-vertically propagating gravity waves (Preusse et al., 2009). Altogether it is not known, how and to what extend the gravity wave field changes during 40 km of upward propagation. So far, it has not been identified, whether the absence of an anticorrelation between gravity wave activity and NLC brightness at Rothera, Davis and ALOMAR can be attributed to probing different volumes for gravity waves and NLC, or whether gravity waves are indeed only a minor contributor to NLC brightness at those sites.

The aim of this work is to closer examine this question by determining gravity wave parameters in a common volume with the NLC observations. We combine lidar measurements from ALOMAR, the arctic atmospheric research site in Andenes, Norway, with mesospheric wind measurements by a closely located radar. The data used for the following analysis is briefly described in Sect. 2. Gravity wave activity in different period ranges is determined by filtering of the wind data by means of a wavelet transformation, which is explained in Sect. 3. The results of two different methods analysing the relation of NLC and gravity wave occurrence are presented in Sect. 4 and discussed in Sect. 5. The concluding Sect. 6 summarizes our results.

2 Data base

For the following analysis of the correlation between NLC occurrence rates and the presence of gravity waves, data sets obtained by lidar and radar in Andenes, Northern Norway (69.3° N, 16.0° E) are combined.

Wind data used for determining gravity wave activity was obtained with the Andenes MF radar which operates at 1.98 MHz and continuously provides wind measurements between 70 km and 94 km. A vertically directed wide beam antenna transmits electromagnetic pulses of 4 km length with a peak power of 50 kW. Three crossed dipoles

**Gravity wave
influence on NLC at
ALOMAR**

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



arranged in an equilateral triangle collect the atmospheric return signal (see Singer et al., 1997 for details). For determining the horizontal wind speed the spaced-antenna method (Briggs, 1984) is applied to the return signal and the height bins are oversampled to yield a height resolution of 2 km. The data is collected with a time resolution of 3 min and is then averaged over 1 h for the following analysis. The uncertainty of the wind velocities is 2 m s^{-1} (Gonzalez et al., 1994). Gravity wave activity in different period bands is derived by means of a wavelet transformation, as previously applied by Hoffmann et al. (2010).

The Rayleigh/Mie/Raman- (RMR-) lidar at the Arctic Lidar Observatory for Middle Atmosphere Research (ALOMAR) is optimized for the detection of NLC by using a spectral highly stable laser and strong spatial and spectral filtering in the detection unit (von Zahn et al., 2000). This allows lidar measurements throughout the Arctic summer at all local times. Lidar count rate profiles are converted to volume backscatter coefficients $\beta(z)$ depending on the altitude z and NLC are identified where $\beta(z)$ is greater than the measurement uncertainty $\Delta\beta(z)$. Details about the data processing are described by Fiedler et al. (2009). For our analysis the maximum backscatter coefficient β_{max} and the centroid altitude z_c were extracted from 14 min averaged profiles. In the year 2006 a major improvement of the detection limit was achieved which reduced the uncertainty to $\Delta\beta \approx 1 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ from previously $\Delta\beta \approx 4 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ (Fiedler et al., 2011), thus allowing the detection of weaker NLC. To ensure that the analysis is performed on the same brightness classes of NLC in every year and that the analysis is unaffected by the change of the detection threshold, we will consider the NLC data in the long-term detection limit (Fiedler et al., 2009) which only includes backscatter coefficients with $\beta_{\text{max}} > 4 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$. The use of the high-sensitivity data ($\beta_{\text{max}} > 1 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$) will be stated explicitly.

3 Data processing

3.1 Wavelet analysis

As model simulations predict that the impact of gravity waves on mesospheric ice layers depends on the waves' period (Turco et al., 1982; Jensen and Thomas, 1994; Rapp et al., 2002; Chandran et al., 2012), we performed our analysis period-dependently. A suitable tool to extract the time evolution of different period components of a signal is the wavelet transformation. Based on the description of Torrence and Compo (1998), the wavelet transformation $W(s, \tau)$ of the wind time series at a certain scale s and time step τ have been calculated. As mother wavelet functions the so-called Morlet wavelet, a cosine function with the frequency ω_0 as free parameter and damped with a Gaussian window (e.g., Torrence and Compo, 1998), has been employed as well as the Paul-Wavelet (Farge, 1992) of order 4. Following the advice of Liu et al. (2007), each wavelet coefficient is weighted by the square root of its corresponding scale $\tilde{f}(s, \tau) = \frac{W(s, \tau)}{\sqrt{s}}$. This allows the direct comparison of different peaks in the wavelet spectrum in the sense that equal amplitudes in a superposition of harmonic functions lead to equally strong peaks in the spectrum. The local wavelet spectrum or wavelet power is given by $|\tilde{f}(s, \tau)|^2$. The sum of the wavelet power over a range of scales, henceforth referred to as a band, results in a time series of the variance in this band (see Eq. 24 of Torrence and Compo, 1998). In this manner we calculate the wind variance in a certain band centered around the mean scale S by using the following relation

$$\chi'^2(S, \tau) = \frac{\delta j \delta t}{C_\delta} \sum_{j=j_1}^{j_2} |\tilde{f}(s_j, \tau)|^2 \quad (1)$$

where s_{j_1} and s_{j_2} denote the scales equivalent to the lower and upper period of the considered band. The time and scale resolution is given by δt and δj , whereas C_δ is the reconstruction factor depending on the wavelet function. From the variances of the u and v component of the wind the kinetic energy density in the chosen band is

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calculated as $E_{\text{kin}}(S, \tau) = \frac{1}{2}(u'^2(S, \tau) + v'^2(S, \tau))$ and averaged over the altitude range of 80 km to 88 km. The band width was chosen to be 2 h, which is a compromise between performing the analysis for different periods and taking the width of a peak produced by a single frequency signal into account.

5 3.2 Normalization of gravity wave kinetic energy

In the second part of our analysis we will divide the gravity wave activity into strong and weak events. The threshold dividing the two domains is derived for each band and altitude bin from the global wavelet spectrum (the time averaged wavelet power). By analogy with the summation of different scales to calculate the variance in a band (Eq. 1), the same scales of the global wavelet spectrum are summed to calculate the reference variance

$$x'^2_{\text{ref}}(S) = \frac{\delta j \delta t}{C_\delta} \sum_{j=j_1}^{j_2} |\tilde{f}_{\text{gl}}(s_j)|^2 \quad (2)$$

with the global wavelet spectrum $\tilde{f}_{\text{gl}}(s) = \frac{1}{N} \sum_{\tau=1}^N \tilde{f}(s, \tau)$. The global wavelet spectrum is based on the time series from 1 June to 15 August, which covers most of the NLC season at ALOMAR. To ensure comparability between different years, the averaged global wavelet spectrum from the years 1999 to 2011 is used as reference. Normalized wind variances are then obtained by dividing the wind variances by their corresponding reference variance $x'^2_{\text{norm}}(S, \tau) = x'^2(S, \tau) / x'^2_{\text{ref}}(S)$ and accordingly, the normalized kinetic energy is calculated as $E_{\text{kin, norm}}(S, \tau) = \frac{1}{2}(u'^2_{\text{norm}}(S, \tau) + v'^2_{\text{norm}}(S, \tau))$. This procedure is repeated for all height bins and the normalized energies are averaged over the altitude range of 80 km to 88 km. Enhanced gravity wave activity will be indicated by $E_{\text{kin, norm}} > 1$, whereas $E_{\text{kin, norm}} < 1$ will denote reduced gravity wave activity. Figure 1 shows in the top panel the global wavelet spectrum used to calculate the reference variance displayed in the bottom panel for the u wind component. The global wavelet

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spectrum exhibits a $f^{-5/3}$ frequency dependency (not shown) and the dominant peaks belong to the diurnal and semidiurnal tide at 24 h and 12 h respectively. The power spectral density (PSD) of the global wavelet spectrum and also the reference variances increases with increasing altitude in the gravity wave range as expected from energy conservation. By using the reference variance as normalization factor this amplitude growth is eliminated and the different altitude bins are comparable. The mean value of the different altitude bins will therefore be equally weighted and not biased by the larger variances at higher altitudes.

4 Results

4.1 Relation of NLC and gravity wave strength

A direct comparison of NLC backscatter coefficients with the simultaneously occurring gravity wave kinetic energy is shown in Fig. 2a for the complete data set of 1999–2011 and the 3 h wave (meaning the kinetic energy in the 2 h to 4 h band). Most NLC and the strongest NLC occurred during times of lower kinetic energy, whereas less and mostly weak NLC occurred during times of high gravity wave kinetic energy. This distribution is more clearly seen in panel b, which shows the histogram of NLC detections in different energy ranges. Figure 2c shows the histogram of gravity wave energies independent of NLC detection during the complete time of lidar measurements. While this figure includes all data points from simultaneous lidar and gravity wave measurements, the data of panel b is limited to those data points of NLC detection and is consequently a subset of the complete data set of panel c. From the two histograms in panel b and c the NLC occurrence rate in different energy ranges is calculated and displayed in panel d. The occurrence rate is defined as the ratio of NLC detection time (panel b) to the total measurement time (panel c), in this specific case limited to certain energy bins of the 3 h wave. In the higher energy range the NLC occurrence rates are based on data sets containing a factor of 100 less data points than in the maximum, so that single data

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



points have a much greater influence. Occurrence rates based on comparatively small data sets of less than 5 h (20 data points) of NLC detection are indicated by empty symbols in Fig. 2d. This indicates their high unreliability compared to the occurrence rates at the maximum and these data points will not be considered further.

In this manner the data sets of waves with larger periods (4 h to 12 h) were analysed. As the results strongly resemble each other, the NLC occurrence rates of the 5 h, 7 h, 9 h and 11 h wave are presented exemplarily in Fig. 3.

The NLC occurrence rates are uniformly distributed over the different energy bins and show no preference to either higher or lower gravity wave activity. As a consequence we conclude that there is no explicit dependence of NLC occurrence on gravity wave kinetic energy.

4.2 NLC occurrence rates during strong and weak gravity wave activity

We further developed a second method to analyse the relation of NLC occurrence and gravity wave activity, where we do not focus on the absolute gravity wave energy during NLC events as in the previous method, but classify the gravity wave activity in weak and strong events, relative to a 13 yr summer season mean condition. By this classification, the data set can be divided into two subsets, one containing all times with enhanced gravity wave strength (indicated by $E_{kin,norm} > 1$) and the other containing all times of attenuated gravity wave strength ($E_{kin,norm} < 1$). For each subset, the NLC occurrence rate (OR) is calculated and compared with the other. A simple way to compare the two occurrence rates is to consider the ratio

$$\eta = \frac{OR_{NLC}(E_{kin,norm} > 1)}{OR_{NLC}(E_{kin,norm} < 1)}. \quad (3)$$

Figure 4 schematically visualizes the calculation of η . The rectangle depicts all points in time where wind measurements are available, respectively the oval all points in time where lidar measurements were performed. In our analysis we include all points in the overlapping area (not to size). The data points are then further sorted according to

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



also the NLC occurrence rates. The finest scale resolution of our three types is obtained by the Morlet wavelet with parameter $\omega_0 = 10$, the finest time resolution by the Paul wavelet of order 4. An additional uncertainty of η is constituted by the uncertainty of the raw wind data of about 2 m s^{-1} . To estimate the sensitivity of η to errors of the wind data, we added random fluctuations following a Gaussian distribution with standard deviation $\sigma = 2 \text{ m s}^{-1}$ to the raw data and then recalculated η . This procedure was repeated 100 times and the minimum and maximum values for η are marked by the errorbars. In this sense the errorbars span the range of values possibly obtained by the analysis when varying the raw data within its uncertainty.

Only in the year 2008 η features a clear increasing trend for all three mother wavelet functions in the period range from 6 h to 12 h and values distinctly different from $\eta = 1$ at periods larger than 9 h. This trend is consistent with model prediction of NLC enhancement due to long period gravity waves and is supported by the scatter plots for the year 2008 in Fig. 6: The brightest NLC occur during comparatively high gravity wave activity in the long period range (8 h and 11 h), whereas in the short period range (3 h) the brightest NLC occur during weak gravity wave activity. It is important to note, that the NLC distribution does not simply follow the gravity wave distribution as in Figs. 2 and 3 and that the accumulation of NLC events at the kinetic energy density of $50 \text{ m}^2 \text{ s}^{-2}$ in the right panel cannot be explained by an increased gravity wave activity in this energy range.

However, further analysis of the data of 2008 reveals that the period dependency of η significantly depends on only two NLC events. One of these events occurred on 5 August 2008 and is shown in greater detail in Fig. 7. It gives an overview over NLC height and brightness and the normalized kinetic energy in the different period ranges. The NLC event marked by the box is accompanied by enhanced activity of gravity waves with periods longer than 8 h and greatly reduced activity of the 3 h and 4 h wave.

If the two events are removed from the data set, i.e. only 12 h out of 378 h of lidar measurements in 2008, η still features an increase with increasing period, but less substantial and only over a reduced period range as shown in Fig. 8 (compare to upper

right panel in Fig. 5). Considering the small amount of data points removed from the data set, this change is remarkable and indicates that the analysis is highly influenced by the sampling of the data. Drawing general conclusions about a robust correlation between enhanced long period gravity waves and NLC occurrence based on single year data is therefore not possible.

In order to increase the statistical robustness we extended the analysis to the time period from 1999 to 2011 and recalculated η for the combined data set. The result is represented in Fig. 9a. So far, only NLC brighter than the long term detection limit of $\beta_{\max} = 4 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ were taken into account, but starting from 2006 high sensitivity data with $\beta_{\max} > 1 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ is available. These weak NLC are included in the second analysis presented in Fig. 9b where η is calculated on the basis of the 2006–2011 high sensitivity data set. Both data sets, the long term and the high sensitivity data, exhibit values of η close to 1 with deviations of smaller magnitude than in the single year data (compare to Fig. 5). The long term data set in panel a features a small decrease in η for periods between 6 h and 9 h. This decrease is least prominent in the Paul wavelet data, as the Paul wavelet has the coarsest period resolution and therefore variations of η are smoothed out over a larger period range. A comparable decrease occurs in the 2006–2011 data in the long term limit (not shown), but it almost completely vanishes in the high sensitivity data (panel b), where $\eta = 1$ lies within the range of possible values of η for all periods except the 8 h wave. The mean value of η determined by the three wavelet types at the 8 h wave is $\eta = 0.87$, which means that the NLC occurrence rate during reduced gravity wave activity was roughly 15 % higher than during enhanced gravity wave activity. However values of η as high as 0.94 lie within the possible range and indicate, that if the 8 h wave has a special influence at all, it is only of minor importance as η does not considerably differ from 1. The overall conclusion of the analysis of η is that η does not vary significantly over the period range from 3 h to 12 h and within the uncertainty the NLC occurrence rates during weak and strong gravity wave activity are equal.

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Discussion

So far, all previous studies on the correlation of NLC and gravity wave activity basically employed the same method by taking stratospheric density perturbations as a proxy for mesospheric gravity wave strength. Our analysis fundamentally differs in three points.

5 Firstly, we derive gravity wave activity from wind instead from density perturbations and secondly the gravity wave activity is determined in the same altitude range as NLC occur. Density perturbations lead to the potential energy of a gravity wave, whereas wind perturbations result in the kinetic energy. According to Geller and Gong (2010) the kinetic energy is proportional to the potential energy, which has recently been confirmed
10 for the mesopause by Placke et al. (2013). Thus, for a given period it does not make a difference whether the potential or kinetic energy is used to determine the gravity wave strength and our energy scale is proportional to those used in previous studies. Determining gravity wave activity in the mesopause instead of the stratosphere yields the advantage that we probe the same waves that influence the NLC formation and no variation of the gravity wave field due to filtering processes has to be considered.

15 Another difference, attributed to the almost continuously available wind data, is that short period waves as well as long period gravity waves can be identified and for the first time the predicted enhancement of NLC due to long period gravity waves can be studied. Additionally, from the large data set of gravity wave energies we are able to perform a detailed analysis of their distribution. It is important to address the question
20 how this energy distribution is possibly distorted by reducing the data set to only those times when lidar measurements were performed. During the NLC season the lidar measurement time covers between 25 % and 7 % of the total time, varying from year to year (Fiedler et al., 2011). To verify that no artificial bias is introduced due to taking only
25 a small subset, the distribution of E_{kin} during the whole NLC season and during lidar measurements is examined. Figure 10 gives an exemplary overview over the kinetic energy distribution prior to normalization for the 3 h wave, 6 h and 9 h respectively. No fundamental differences can be found between the two data sets, which is supported by

a fairly constant ratio between them both. We therefore conclude that the lidar subset is large enough to adequately cover all energy ranges.

The distribution of NLC in Fig. 2a strongly resembles the schematic illustration of Gerrard et al. (2004, Fig. 5) with the pronounced triangular shape in the NLC strength/gravity wave strength plane. Their illustration displayed the theorized relationship between gravity wave activity and NLC brightness for short period gravity waves. At a first sight the resemblance of our data to the illustration could be interpreted as another experimental support of the destructing effect of short period gravity waves on NLC. However, as we have shown, NLC occurrence is independent of gravity wave kinetic energy as indicated by the constant NLC occurrence rates in different energy ranges and for different period bands. The absence of strong NLC during high gravity wave activity in Fig. 2a can simply be explained by the generally rare occurrence of strong NLC. For example, only 1 % of all NLC are brighter than $\beta_{\max} = 30 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ and given only 10 NLC detections at energy densities around $50 \text{ m}^2 \text{ s}^{-2}$ we would expect less than one to be as bright as $30 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ in this energy range. While these numbers hold for the 3 h wave, the arguments can be generalized for larger periods. Thus, concluding from this analysis no enhancement or attenuation of NLC by gravity waves is supported by our data.

The analysis of η , the ratio of NLC occurrence rates during high and low gravity wave activity, results in the same conclusion. While the data limited to single NLC seasons partly show large deviations from $\eta = 1$, the combined data set of the years 1999–2011 is not indicative of any destructing or enhancing characteristic of gravity waves. Nevertheless, the analysis of the single year data of 2008 revealed two interesting NLC events which were accompanied by enhanced long period gravity wave activity (Fig. 7 shows one of them). The transition period between reduced and enhanced gravity wave activity falls into the range of 6 h to 8 h. This is in good agreement to model studies by Rapp et al. (2002) who found NLC enhancement due to gravity waves with periods of 6.5 h and more. Furthermore it is worth remarking that this event is preceded by 36 h of enhanced activity of the 8 h wave. During the following NLC event one of the

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



highest backscatter coefficients ($\beta_{\max} = 57 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$) of the whole data set was measured. This exceptionally strong NLC might suggest that the conditions prior to NLC detection have highly favored ice particle growth. We can only speculate whether the gravity waves prior and during NLC detection were one of the factors enabling NLC of this brightness, but it seems possible that there is a physical connection. To entirely answer questions of this type, the history of air parcels with temperature and humidity in the mesopause region would be needed.

Not finding any correlation of gravity wave activity and NLC occurrence in Andenes agrees with the results of Schöch (2007). One of his explanations for the absence of an anticorrelation refers to a strong thermal tidal influence on NLC occurrence over ALOMAR which is reflected by the semi-diurnal variation of NLC height and brightness. This is confirmed by Fiedler et al. (2011) who found an almost perfect anticorrelation of the diurnal variation of NLC occurrence rates and model mesopause temperatures during times of lidar measurements. In the early morning hours, the cold phase of the wave, NLC occurrence rates as high as 40 % are reached, whereas in the afternoon and evening NLC occurrence rates are less than 20 %. Temperature variations, which are one of the main factors controlling NLC properties (e.g., Rapp and Thomas, 2006) can be induced by gravity waves as well as thermal tides, so that in general a superposition of both exist in the mesopause region. In the context of previous studies we conclude that at ALOMAR gravity waves play only a minor role in controlling NLC occurrence compared to the tidal influence. For this reason we do not find any relation of NLC occurrence to gravity wave kinetic energy. Note that we do not find a positive correlation with wave activity in the 11 h to 13 h period range because we relate NLC occurrence to the overall strength of a wave and not to a certain phase (like the cold phase) of a wave. Of course we need to stress that our results are only valid for the location of ALOMAR and cannot be generalized.

Gravity wave strength might vary considerably between different locations. Chandran et al. (2010) have derived gravity wave occurrence rates from polar mesospheric clouds which feature a strong longitudinal variability. ALOMAR is situated in one of the minima

of gravity wave occurrence rates whereas Rothera and Sondrestrom are located at a maximum. From this point of view the presence or absence of an anticorrelation of NLC brightness and gravity wave activity can likely be explained by the different levels of gravity wave activity.

6 Summary

We combined the 1999–2011 ALOMAR-RMR lidar data set with wind measurements by MF radar to study the relationship of NLC occurrence and gravity wave activity. Gravity wave kinetic energy in different period bands, derived from the horizontal wind data by means of a wavelet transformation, was obtained in the same volume as NLC were detected. The direct comparison of NLC brightness and gravity wave kinetic energy showed no correlation of these two parameters. After normalization to the occurrence of gravity waves themselves, we found that NLC occurred equally frequently in all energy ranges of all periods. This implies, that the occurrence of NLC is independent of the simultaneously determined gravity wave kinetic energy. In a second approach we divided the gravity wave energy data into two subsets, one of reduced and one of enhanced energy as compared to a 13 yr summer season mean condition. For each subset the NLC occurrence rate was determined. We analysed whether NLC occurred more frequently during enhanced gravity wave activity by looking at the ratio η of the NLC occurrence rate during strong and weak gravity wave activity. For the period range of 3 h to 12 h this ratio was calculated for the years 2007, 2008, 2009 and 2011 separately and for combined data sets of 1999–2011 and 2006–2011. For the single year data (except 2008) η exhibited values in the order of 1, which means that NLC occurred equally frequently during high and low gravity wave activity, but with quite large fluctuations from one period to the next as well as quite large differences between the three applied mother wavelet functions. The data of 2008 showed a remarkably clear increase of η with increasing period, which is the expected trend from microphysical model results (e.g., Rapp et al., 2002). However, it appeared that the increasing trend

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of η was mainly due to two NLC events of 12 h duration. One of the events was accompanied by enhanced gravity wave activity in periods larger than 8 h and preceded by 36 h of enhanced activity in the 7 h to 9 h period band. As this NLC was one of the brightest NLC of the whole data set, it seems possible that gravity waves have positively influenced the growth of ice particles during this specific NLC. For the combined data set of the years 1999–2011 η shows a slight decrease in the period range from 6 h to 9 h. This decrease however vanishes when using the high-sensitivity data set available since 2006 with a detection limit of $\beta_{\text{thres}} = 1 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$ and η attains values close to 1 for all periods.

From both analysing methods we come to the conclusion that gravity waves above ALOMAR neither negatively nor positively influence NLC occurrence rates. Our findings are in accordance to earlier studies by Schöch (2007). They also agree with the interpretation of the diurnal variation of NLC occurrence rates by Fiedler et al. (2011) who conclude that tidal temperature variations are the primary driver for diurnal NLC occurrence variability. In future, the relative importance of gravity waves and tides influencing NLC parameters should be analyzed in dependence of their geographic position.

The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

References

- Backhouse, T. W.: The luminous cirrus cloud of June and July, *Meteorol. Mag.*, 20, 133, 133–133, 1885. 20050
- Briggs, B. H.: The analysis of spaced sensor records by correlation techniques, in: *Middle Atmosphere Program Handbook*, edited by: Vincent, R., vol. 13, SCOSTEP, 168–188, 1984. 20053

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chandran, A., Rusch, D. W., Palo, S. E., Thomas, G. E., and Taylor, M. J.: Gravity wave observations in the summertime polar mesosphere from Cloud Imaging and Particle Size (CIPS) experiment on the AIM spacecraft, *J. Atmos. Sol.-Terr. Phys.*, 71, 392–400, doi:10.1016/j.jastp.2008.09.041, 2009. 20051

5 Chandran, A., Rusch, D. W., Merkel, A. W., Palo, S. E., Thomas, G. E., Taylor, M. J., Bailey, S. M., and Russell III, J. M.: Polar mesospheric cloud structures observed from the cloud imaging and particle size experiment on the Aeronomy of Ice in the Mesosphere spacecraft: Atmospheric gravity waves as drivers for longitudinal variability in polar mesospheric cloud occurrence, *J. Geophys. Res.*, 115, D13102, doi:10.1029/2009JD013185, 2010. 20063

10 Chandran, A., Rusch, W., Thomas, G. E., Palo, S. E., Baumgarten, G., Jensen, E. J., and Merkel, A. W.: Atmospheric gravity wave effects on polar mesospheric clouds: a comparison of numerical simulations from CARMA 2D with AIM observations, *J. Geophys. Res.*, 117, D20104, doi:10.1029/2012JD017794, 2012. 20051, 20054

15 Chu, X., Gardener, C. S., and Papen, G.: Lidar observations of polar mesospheric clouds at South Pole: diurnal variations, *Geophys. Res. Lett.*, 28, 1937–1940, doi:10.1029/2000GL012524, 2001. 20051

20 Chu, X., Espy, P., Nott, G. J., Dietrich, J. C., and Gardner, C. S.: Polar mesospheric clouds observed by an iron Boltzmann lidar at Rothera (67.5° S, 68.0° W), Antarctica from 2002 to 2005: properties and implications, *J. Geophys. Res.*, 111, D20213, doi:10.1029/2006JD007086, 2006. 20051

Chu, X., Yamashita, C., Espy, P. J., Nott, G. J., Jensen, E. J., Liu, H.-L., Huang, W., and Thayer, J. P.: Responses of polar mesospheric cloud brightness to stratospheric gravity waves at the South Pole and Rothera, Antarctica, *J. Atmos. Sol.-Terr. Phys.*, 71, 434–445, doi:10.1016/j.jastp.2008.10.002, 2009. 20051

25 DeLand, M. T., Shettle, E. P., Thomas, G. E., and Olivero, J. J.: Latitude-dependent long-term variations in polar mesospheric clouds from SBUV version 3 PMC data, *J. Geophys. Res.*, 112, D10315, doi:10.1029/2006JD007857, 2007. 20050

Eckermann, S. D.: Effect of background winds on vertical wavenumber spectra of atmospheric gravity waves, *J. Geophys. Res.*, 100, 14097–14112, doi:10.1029/95JD00987, 1995. 20052

30 Farge, M.: Wavelet transforms and their applications to turbulence, *Annu. Rev. Fluid Mech.*, 24, 395–457, doi:10.1146/annurev.fluid.24.1.395, 1992. 20054

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fiedler, J., Baumgarten, G., and von Cossart, G.: Mean diurnal variations of noctilucent clouds during 7 years of lidar observations at ALOMAR, *Ann. Geophys.*, 23, 1175–1181, doi:10.5194/angeo-23-1175-2005, 2005. 20051

Fiedler, J., Baumgarten, G., and Lübken, F.-J.: NLC observations during one solar cycle above ALOMAR, *J. Atmos. Sol.-Terr. Phys.*, 71, 424–433, doi:10.1016/j.jastp.2008.11.010, 2009. 20053

Fiedler, J., Baumgarten, G., Berger, U., Hoffmann, P., Kaifler, N., and Lübken, F.-J.: NLC and the background atmosphere above ALOMAR, *Atmos. Chem. Phys.*, 11, 5701–5717, doi:10.5194/acp-11-5701-2011, 2011. 20051, 20053, 20061, 20063, 20065

Gadsden, M. and Schröder, W.: *Noctilucent Clouds*, Springer-Verlag, 1989. 20050

Geller, M. A. and Gong, J.: Gravity wave kinetic, potential, and vertical fluctuation energies as indicators of different frequency gravity waves, *J. Geophys. Res.*, 115, D11111, doi:10.1029/2009JD012266, 2010. 20061

Gerrard, A. J., Kane, T. J., Thayer, J. P., and Eckermann, S. D.: Concerning the upper stratospheric gravity wave and mesospheric cloud relationship over Sondrestrom, Greenland, *J. Atmos. Sol.-Terr. Phys.*, 66, 229–240, doi:10.1016/j.jastp.2003.12.005, 2004. 20051, 20062

Gonzalez, N., Hauchecorne, A., Kirkwood, S., Lübken, F.-J., Manson, A. H., Mourier, A., Schmidlin, F. J., Schminder, R., Kürschner, D., Singer, W., and Widdel, H.-U.: Intercomparisons of simultaneous remote and in situ wind measurements, *J. Atmos. Terr. Phys.*, 56, 1985–2001, doi:10.1016/0021-9169(94)90024-8, 1994. 20053

Hoffmann, P., Becker, E., Singer, W., and Placke, M.: Seasonal variation of mesospheric waves at northern middle and high latitudes, *J. Atmos. Sol.-Terr. Phys.*, 72, 1068–1079, doi:10.1016/j.jastp.2010.07.002, 2010. 20053

Innis, J. L., Klekociuk, A. R., Morris, R. J., Cunningham, A. P., Graham, A. D., and Murphy, D. J.: A study of the relationship between stratospheric gravity waves and polar mesospheric clouds at Davis Antarctica, *J. Geophys. Res.*, 113, D14102, doi:10.1029/2007JD009031, 2008. 20051

Jensen, E. J. and Thomas, G. E.: Numerical simulations of the effects of gravity waves on noctilucent clouds, *J. Geophys. Res.*, 99, 3421–3430, doi:10.1029/93JD01736, 1994. 20051, 20054

Jesse, O.: Auffallende Abenderscheinung am Himmel, *Meteorol. Z.*, 2, 311–312, 1885. 20050

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kaifler, N., Baumgarten, G., Fiedler, J., and Lübken, F.-J.: Quantification of waves in lidar observations of noctilucent clouds at scales from seconds to minutes, *Atmos. Chem. Phys. Discuss.*, 13, 7397–7429, doi:10.5194/acpd-13-7397-2013, 2013. 20051
- Leslie, R. C.: Sky glows, *Nature*, 245, 16, 245–245, 1885. 20050
- 5 Liu, Y., Liang, X. S., and Weisberg, R. H.: Rectification of the bias in the wavelet power spectrum, *J. Atmos. Ocean. Tech.*, 24, 2093–2102, doi:10.1175/2007JTECHO511.1, 2007. 20054
- Lübken, F.-J. and Berger, U.: Latitudinal and interhemispheric variation of stratospheric effects on mesospheric ice layer trends, *J. Geophys. Res.*, 116, D00P03, doi:10.1029/2010JD015258, 2011. 20051
- 10 Lübken, F.-J., Berger, U., and Baumgarten, G.: Stratospheric and solar cycle effects on long-term variability of mesospheric ice clouds, *J. Geophys. Res.*, 114, D00I06, doi:10.1029/2009JD012377, 2009. 20051
- Pautet, P.-D., Stegman, J., Wrasse, C. M., Nielsen, K., Takahashi, H., Taylor, M. J., Hopfel, K. W., and Eckermann, S. D.: Analysis of gravity waves structures visible in noctilucent cloud images, *J. Atmos. Sol.-Terr. Phys.*, 73, 2082–2090, doi:10.1016/j.jastp.2010.06.001, 2011. 20051
- 15 Placke, M., Hoffmann, P., Gerding, M., Becker, E., and Rapp, M.: Testing linear gravity wave theory with simultaneous wind and temperature data from the mesosphere, *J. Atmos. Sol.-Terr. Phys.*, 93, 57–69, doi:10.1016/j.jastp.2012.11.012, 2013. 20061
- 20 Preusse, P., Eckermann, S. D., Ern, M., Oberheide, J., Picard, R. H., Roble, R. G., Riese, M., Russell III, J. M., and Mlynczak, M. G.: Global ray tracing simulations of SABER gravity wave climatology, *J. Geophys. Res.*, 114, D08126, doi:10.1029/2008JD011214, 2009. 20052
- Rapp, M. and Thomas, G. E.: Modeling the microphysics of mesospheric ice particles: assessment of current capabilities and basic sensitivities, *J. Atmos. Sol.-Terr. Phys.*, 68, 715–744, doi:10.1016/j.jastp.2005.10.015, 2006. 20063
- 25 Rapp, M., Lübken, F.-J., Müllemann, A., Thomas, G. E., and Jensen, E. J.: Small-scale temperature variations in the vicinity of NLC: experimental and model results, *J. Geophys. Res.*, 107, 4392, doi:10.1029/2001JD001241, 2002. 20051, 20054, 20062, 20064
- 30 Schöch, A.: Thermal structure and gravity waves in the Arctic middle atmosphere above ALOMAR (69.3° N, 16.0° E), Ph. D. thesis, Leibniz-Institut für Atmosphärenphysik, Universität Rostock, Kühlungsborn, Germany, 151 pp., available at: http://www.iap-kborn.de/fileadmin/user_upload/MAIN-abteilung/optik/Forschung/Doktorarbeiten/schoech_diss.pdf, 2007. 20051, 20063, 20065

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Shettle, E. P., DeLand, M. T., Thomas, G. E., and Olivero, J. J.: Long term variations in the frequency of polar mesospheric clouds in the Northern Hemisphere from SBUV, *Geophys. Res. Lett.*, 36, L02803, doi:10.1029/2008GL036048, 2009. 20050

5 Singer, W., Keuer, D., and Eriksen, W.: The Alomar MF radar: technical design and first results, in: *Proceedings of the 13th ESA Symposium on European Rocket and Balloon Programmes and Related Research*, Oeland, Sweden (ESA SP-397), edited by: Kaldeich-Schürmann, B., 101–103, 1997. 20053

10 Thayer, J. P., Rapp, M., Gerrard, A. J., Gudmundsson, E., and Kane, T. J.: Gravity-wave influences on Arctic mesospheric clouds as determined by a Rayleigh lidar at Sondrestrom, Greenland, *J. Geophys. Res.*, 108, 8449, doi:10.1029/2002JD002363, 2003. 20051

Torrence, C. and Compo, G. P.: A Practical Guide to Wavelet Analysis, *B. Am. Meteorol. Soc.*, 79, 61–78, doi:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2, 1998. 20054

15 Turco, R. P., Toon, O. B., Whitten, R. C., Keesee, R. G., and Hollenbach, D.: Noctilucent clouds: Simulations studies of their genesis, properties and global influences, *Planet. Space Sci.*, 30, 1147–1181, doi:10.1016/0032-0633(82)90126-X, 1982. 20051, 20054

von Zahn, U., von Cossart, G., and Fiedler, J.: Tidal variations of noctilucent clouds measured at 69° N latitude by groundbased lidar, *Geophys. Res. Lett.*, 25, 1289–1292, doi:10.1029/98GL00546, 1998. 20051

20 von Zahn, U., von Cossart, G., Fiedler, J., Fricke, K. H., Nelke, G., Baumgarten, G., Rees, D., Hauchecorne, A., and Adolfsen, K.: The ALOMAR Rayleigh/Mie/Raman lidar: objectives, configuration, and performance, *Ann. Geophys.*, 18, 815–833, doi:10.1007/s00585-000-0815-2, 2000. 20053

Witt, G.: Height, structure and displacements of noctilucent clouds, *Tellus A*, 14, 1–18, doi:10.1111/j.2153-3490.1962.tb00115.x, 1962. 20051

Gravity wave
influence on NLC at
ALOMAR

H. Wilms et al.




Fig. 1. (a) Global wavelet spectrum based on wind u data from 1 June to 15 August of the years 1999 to 2011. (b) Reference variances obtained from the global wavelet spectrum by summation of overlapping period bands of 2 h width (see brackets below upper panel).

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. (a) NLC backscatter coefficient β_{\max} versus gravity wave kinetic energy density in the 2 h to 4 h period band. (b) Total time of NLC detection during simultaneously identified gravity wave activity of the 3 h wave in different energy bins (histogram of panel a). (c) Total time of gravity wave occurrence of the 3 h wave in different energy bins during the complete lidar measurement time independent of NLC detection. (d) NLC occurrence rate (ratio of NLC detection time to total measurement time) in different energy bins. Open symbols indicate occurrence rates based on less than 5 h of NLC detection.

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. The same as Fig. 2d, but for the 5 h, 7 h, 9 h and 11 h wave.




Fig. 4. Schematic visualization of the available lidar and radar data. From the radar data times of high and low gravity wave activity in a certain period range are identified ($E_{kin,norm} > 1$ and $E_{kin,norm} < 1$ respectively) and from the lidar data times of NLC detection. The NLC occurrence rate during high gravity wave activity is the fraction of the shaded area with respect to the framed area.

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gravity wave influence on NLC at ALOMAR

H. Wilms et al.




Fig. 5. Ratio η (see Eq. 3) for the years 2007, 2008, 2009 and 2011 and three different mother wavelet functions (Morlet with parameter $\omega_0 = 6$ and $\omega_0 = 10$ and Paul of order $m = 4$). The errorbars result from variation of the raw data within its uncertainty.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.




Fig. 6. NLC brightness versus kinetic energy density for data of 2008 and three different periods (3 h, 8 h, 11 h). The step plot shows the histogram of gravity wave occurrence as in Fig. 2c.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. NLC and gravity wave data for an exceptionally bright NLC on 5 August 2008. The top panel shows the NLC height and color-coded the maximum backscatter coefficient β_{\max} . Times of lidar measurement are indicated by the black line above the NLC data. The panels below show the height-time cross-section of normalized kinetic energy for periods between 3 h and 12 h.

**Gravity wave
influence on NLC at
ALOMAR**

H. Wilms et al.




Fig. 8. Same as Fig. 5 for 2008, but with 12 h of lidar data removed from the data set.

Gravity wave influence on NLC at ALOMAR

H. Wilms et al.




Fig. 9. **(a)** Ratio η as in Fig. 5 for the combined data set of 1999–2011 with long term detection limit of $\beta_{\max} = 4 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$. **(b)** Ratio η for the combined data set of 2006–2011 with NLC detection limit of $\beta_{\max} = 1 \times 10^{-10} \text{ m}^{-1} \text{ sr}^{-1}$.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Gravity wave influence on NLC at ALOMAR

H. Wilms et al.




Fig. 10. Gravity wave occurrence of the 3 h, 6 h and 9 h wave in different energy bins. Red: complete summer season data set of the years 1999–2011. Blue: data set limited to times of lidar measurements.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)