



Technical Note:
**Temporal change in
averaging kernels**

J. Yoon et al.

Technical Note: Temporal change in averaging kernels as a source of uncertainty in trend estimates of carbon monoxide retrieved from MOPITT

J. Yoon¹, A. Pozzer¹, P. Hoor², D. Y. Chang¹, S. Beirle¹, T. Wagner¹, S. Schloegl², J. Lelieveld^{1,3}, and H. M. Worden⁴

¹Atmospheric Chemistry Department, Max-Planck Institute of Chemistry, P.O. Box 3060, 55020 Mainz, Germany

²Institute of Atmospheric Physics, Johannes Gutenberg University, 55099 Mainz, Germany

³The Cyprus Institute, Energy, Environment and Water Research Center, P.O. Box 27456, 1645 Nicosia, Cyprus

⁴National Center for Atmospheric Research (NCAR), Boulder, CO, USA

Received: 18 June 2013 – Accepted: 26 July 2013 – Published: 5 August 2013

Correspondence to: J. Yoon (jongmin.yoon@mpic.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

It is now possible to monitor the global and long-term trends of trace gases that are important to atmospheric chemistry, climate, and air quality with satellite data records that span more than a decade. However, many of the remote sensing techniques used by satellite instruments produce measurements that have variable sensitivity to the vertical profiles of atmospheric gases. In the case of constrained retrievals like optimal estimation, this leads to a varying amount of a priori information in the retrieval and is represented by an averaging kernel. In this study, we investigate to what extent such trends can be biased by temporal changes of averaging kernels used in the retrieval algorithm. In particular, the surface carbon monoxide data retrieved from the Measurements Of Pollution In The Troposphere (MOPITT) instrument from 2001 to 2010 were analysed. As a practical example based on the MOPITT data, we show that if the true atmospheric mixing ratio is continuously 50 % higher or lower than the a priori state, the temporal change of the averaging kernel at the surface level gives rise to an artificial trend in retrieved surface carbon monoxide, ranging from -10.71 to $+13.21$ ppbv yr⁻¹ (-5.68 to $+8.84$ % yr⁻¹) depending on location. Therefore, in the case of surface (or near-surface level) CO derived from MOPITT, the AKs trends multiplied by the difference between true and a priori states must be quantified in order to estimate trend biases.

1 Introduction

Since the Industrial Revolution human activity has brought about a significant change in atmospheric trace gas concentrations (Hansen et al., 2007), which has important consequences for global air quality and the Earth's atmosphere radiative budget, thus causing climate change (Forster et al., 2007 (IPCC AR4)). One of the most important trace gases is carbon monoxide (CO). It is produced by the oxidation of methane (CH₄) and non-methane hydrocarbons (NMHC), and directly emitted by biomass burning and

ACPD

13, 20319–20340, 2013

Technical Note:
**Temporal change in
averaging kernels**

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



combustion of fossil fuels (Wallace and Hobbs, 2006). It acts as a major sink for tropospheric hydroxyl radical (OH) (Crutzen, 1974). OH, in turn, regulates the lifetimes of various tropospheric trace gases (Khalil and Rasmussen, 1990; Wallace and Hobbs, 2006). The oxidation of CO also leads to the formation of carbon dioxide (CO₂) and tropospheric ozone (O₃) (Crutzen and Gidel, 1983; Fishman and Crutzen, 1978; Burrows et al., 1995). Because of its moderately long lifetime (weeks to months) and inhomogeneous distribution in the troposphere, CO is an ideal tracer for monitoring the air pollution sources and transports (Cicerone, 1988; Logan et al., 1981; Shindell et al., 2006; Lelieveld et al., 2001; Hoor et al., 2005; Edwards et al., 2004, 2006; Pan et al., 1995; Worden et al., 2013a; Liu et al., 2011, 2013).

Remote sensing from satellites provides the unique opportunity to study regional and global changes in atmospheric composition (Burrows et al., 2011). Since its launch on board the EOS-Terra satellite in 1999, MOPITT has provided global and long-term CO products, which may be used to estimate trends of tropospheric CO and the influence of anthropogenic emissions on atmospheric chemistry (Worden et al., 2013a). To retrieve CO vertical profiles using MOPITT observations, the theory developed by Rodgers (1976) and Rodgers et al. (2000) is employed in the retrieval algorithm (Pan et al., 1998; Deeter et al., 2003). The algorithm is designed to use a priori profiles derived from a climatology based on simulations by the Model for Ozone and Related chemical Tracers-4 (MOZART-4) model (Pan et al., 1998; Deeter et al., 2003; Deeter, 2009; Worden et al., 2013a). The retrieval sensitivity is generally evaluated through averaging kernels (AKs) (Pan et al., 1998; Rodgers, 2000; Deeter et al., 2003; Eskes and Boersma, 2003). The AKs are essential for meaningful comparisons of satellite products against in situ data and model results (Rodgers, 2000; Rodger and Connor, 2003; Eskes and Boersma, 2003). However, we will show that their temporal change can lead to a significant uncertainty in the trend estimation of retrieved CO using AKs.

The goal of this paper is to investigate the uncertainty caused by temporal AKs change in the trend estimation of retrieved state. It is shown that without adequate knowledge of the differences between the a priori and true profiles of the observed trace

gases, AKs uncertainty cannot be removed in the trend estimation of retrieved state. Although MOPITT monthly surface CO data were used in this study, we emphasize that the uncertainties in trend estimations apply to any remote sensing observations based on similar AK concepts.

In Sect. 2, the MOPITT CO data and the standard method used for trend estimation are described. Section 3 presents the analysis of uncertainties and a practical example based on MOPITT CO surface data. Section 4 analyses possible issues associated with statistically merging Level 3 MOPITT data, and Sect. 5 summarizes this work and presents conclusions.

2 MOPITT surface CO and linear trend model

The MOPITT instrument on the EOS-Terra spacecraft provides tropospheric CO and CH₄ distributions on a global scale (Deeter et al., 2003). The temperature constraints of the MOPITT radiometric calibration (University of Toronto and NCAR MOPITT Team, 1995 (MOPITT ATBD)) are ± 0.5 K for the CO profile channels and ± 1 K for the CO and CH₄ column channels for the expected precisions, being 10 % for CO and 1 % for CH₄ products (Pan et al., 1995, 1998). The global MOPITT retrieved CO data have been used for estimating CO distributions, atmospheric transport, sources, and sinks (e.g., Drori et al., 2012; Fortems-Cheiney et al., 2011; Hooghiemstra et al., 2011; Kanakidou et al., 2011; Arellano and Hess, 2010; Choi and Chang, 2010; Stroppiana et al., 2010). Recently the MOPITT Version 5 (V5) Level 3 (L3) thermal infrared (TIR) retrievals (in daytime) lead to an improved sensitivity for the lower tropospheric CO retrieval, a reduced retrieval bias drift associated with long-term instrumental degradation, and a more exact representation of the effects of random errors in the radiances (Clerbaux et al., 2009; Worden et al., 2010, 2013a; Deeter et al., 2007a, 2011, 2012, 2013), and are used in this study. Figure 1 shows the total mean distribution of global surface CO from the products (spatial resolution: $1^\circ \times 1^\circ$ and temporal resolution: 1 month) from 2001 to 2010.

The linear trends (ω) in unit of ppbv yr⁻¹ are analysed using a linear least squares fit based on monthly data as follows:

$$y_t = \omega x_t + \tilde{y}^m + n_t + \mu \quad (1)$$

where x_t , μ , n_t , and \tilde{y}^m denote the time index term ($t/12$), the constant term, the monthly noise for the analysed periods ($t = 1 \dots T$), and the seasonal mean of y_t for each month ($m = 1 \dots 12$), respectively. This linear trend model is applied to the data time series for each point that represents complete yearly datasets defined as encompassing at least 9 months of data per year. Otherwise, the location is discarded to avoid statistical inconsistencies. In general, the noise term can include random and systematic measurement errors and autocorrelation due to atmospheric variability (Weatherhead et al., 1998, 2002). In this study, we focus on the contribution from changes in the averaging kernel to this noise term.

Figure 2 presents the linear trends of surface CO from MOPITT V5 TIR L3 products during daytime from 2001 to 2010. Here we have included all the MOPITT V5 data, including scenes that used water vapour climatology when the NCEP (National Centers for Environmental Prediction) Global Data Assimilation System (GDAS) product was not available. The use of water vapour climatology can produce anomalous results in the CO retrievals, as found by the MOPITT team, in particular, off the coast of California. Therefore, we do not have confidence in the higher CO values or trends that can be seen in this region in Figs. 1 and 2.

An overall decrease of the surface CO from biomass burning over South America/Africa (Wallace and Hobbs, 2006; Liu et al., 2011) and fossil-fuel combustion over East USA, West Europe and South/East Asia (Wallace and Hobbs, 2006; Liu et al., 2011; Buchwitz et al., 2007) is indicated. This tendency is consistent with the decreasing trends in total CO columns retrieved from MOPITT, the Atmospheric InfraRed Sounder (AIRS), the Tropospheric Emission Spectrometer (TES), and the Infrared Atmospheric Sounding Interferometer (IASI), which are reported in Worden et al. (2013a).

Technical Note:
Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, the trends in Fig. 2 are affected by intrinsic uncertainty due to temporal changes of averaging kernels (AKs).

3 Uncertainty from temporal change of averaging kernels

The V5 MOPITT CO retrieval algorithm is based on the maximum a posteriori (MAP) that is an iterative optimal estimation technique (Pan et al., 1998; Rodgers, 2000; Deeter et al., 2003, 2010). The V5 algorithm employs the log(VMR: Volume Mixing Ratio) retrievals (Deeter et al., 2013) that are the basis of this study. The statistical technique to solve the inversion problem is employed to find the optimal CO retrieval using the MOPITT measured radiances, employing a priori defined CO vertical profiles. If the forward model error is neglected, the relation equation between retrieved and true states can be expressed as follows (Rodgers, 1990; Pan et al., 1998; Deeter et al., 2003, 2010):

$$\hat{\mathbf{x}} \cong \mathbf{x}_0 + \mathbf{A}(\mathbf{x} - \mathbf{x}_0) = \mathbf{A}\mathbf{x} + (\mathbf{I} - \mathbf{A})\mathbf{x}_0 \quad (2)$$

where $\hat{\mathbf{x}}$, \mathbf{x}_0 , and \mathbf{x} represent the retrieved, the a priori, and the true states of the atmosphere, respectively. \mathbf{A} and \mathbf{I} denote the AKs and identity matrices, respectively. The AKs matrix, which reflects the sensitivity of the retrieved state to the true state, is the most important diagnostics of the retrieval (Pan et al., 1998; Rodgers, 2000; Deeter et al., 2003; Eskes and Boersma, 2003). When the AKs matrix is exactly equal to the identity matrix, the retrieved state is identical to the representation of the true atmospheric state on the altitude grid of the retrieval. However, in practice, the AKs matrix does not match up with \mathbf{I} . For example, Fig. 3 shows the mean AK for surface CO, a priori, and retrieved CO profiles over the grid cell defined by 30° E, 45° E, 110° N, and 120° N in East China, where large quantities of CO are emitted into the atmosphere from the combustion of fossil fuels (see Fig. 1) (Liu et al., 2011; Buchwitz et al., 2007). The AK theoretically represents how much the true and a priori profiles contribute to

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the retrieved CO at the surface as shown in Eq. (2). The surface retrieved CO is therefore the weighted mean contribution of the true and a priori profiles in multiple layers. However, there is a clear difference between the a priori and retrieved CO profiles as shown in Fig. 3. For example, the retrieved CO at the surface is higher by about 52 % compared to the a priori CO.

Most importantly, we are interested in the temporal change of the retrieved state, and its possible connection to the true trend. The retrieved trend can be expressed by the partial differential of Eq. (2) as follows:

$$\frac{\partial \hat{x}}{\partial t} \cong \frac{\partial x_0}{\partial t} + \frac{\partial \mathbf{A}}{\partial t} (\mathbf{x} - \mathbf{x}_0) + \mathbf{A} \frac{\partial}{\partial t} (\mathbf{x} - \mathbf{x}_0) = \frac{\partial x_0}{\partial t} (\mathbf{I} - \mathbf{A}) + \frac{\partial \mathbf{A}}{\partial t} (\mathbf{x} - \mathbf{x}_0) + \mathbf{A} \frac{\partial \mathbf{x}}{\partial t} \quad (3)$$

No temporal change is present in the a priori state ($\frac{\partial x_0}{\partial t} = 0$), as it is obtained from a monthly climatology of the Model for Ozone and Related chemical Tracers-4 (MOZART-4), and does not represent interannual variability (Pan et al., 1998; Deeter et al., 2003; Deeter, 2009; Worden et al., 2013a). Therefore, Eq. (3) can be rewritten as follows:

$$\frac{\partial \hat{x}}{\partial t} \cong \frac{\partial \mathbf{A}}{\partial t} (\mathbf{x} - \mathbf{x}_0) + \mathbf{A} \frac{\partial \mathbf{x}}{\partial t}. \quad (4)$$

Again, if the AKs matrix is the identity matrix ($\mathbf{A} = \mathbf{I}$), the temporal change of the retrieved state is the same as the change of the true state ($\frac{\partial \hat{x}}{\partial t} = \frac{\partial \mathbf{x}}{\partial t}$). Even if the AKs matrix is not the identity matrix, but the AKs have no temporal change ($\mathbf{A} \neq \mathbf{I}$, but $\frac{\partial \mathbf{A}}{\partial t} = 0$), then the change of the retrieved state is an averaged contribution of the true changes in multiple layers ($\frac{\partial \hat{x}}{\partial t} = \mathbf{A} \frac{\partial \mathbf{x}}{\partial t}$). In other words, no temporal change of AKs ($\frac{\partial \mathbf{A}}{\partial t} = 0$) is an ideal condition for the trend estimation. However, in fact the AKs do have a significant variability because they are influenced by all variables as inputs to the forward radiative transfer model (e.g., atmospheric temperature profile, surface pressure, and surface temperature) (Deeter et al., 2003). In addition, the changes in AK values over time are likely due to the changes in CO itself since retrieval sensitivity depends on CO

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



amount as shown in Deeter et al. (2007b) and Worden et al. (2013b). Therefore, the trend of the retrieved state ($\frac{\partial \hat{x}}{\partial t}$) can be biased by the term ($\frac{\partial A}{\partial t} (x - x_0)$), whose value (and even sign) is unknown, as the true state (and hence its difference with the a priori state) is unknown.

To estimate the general uncertainties caused by temporal varying AKs, we make the assumption that the true state is 50 % more (less) than the a priori state (i.e. $x = x_0 \pm 50\% \times x_0$). This 50 % difference could be estimated from the difference between retrieved and a priori states at the surface as shown in Fig. 3. In addition, Fig. 4 shows the global distributions of maximum and minimum differences between them, and demonstrates that the assumption is reasonable globally. Eventually, the AKs uncertainty in the trend analysis of MOPITT CO at the surface is balancing between these extremes.

Following this assumption, the trend of the constructed “true state” is the same as the trend of the a priori state, which theoretically equals to zero (i.e. $\frac{\partial x}{\partial t} = \frac{\partial x_0}{\partial t} = 0$) (Pan et al., 1998; Deeter et al., 2003; Deeter, 2009; Worden et al., 2013a). Nevertheless some significant trends of the retrieved CO at the surface are present, ranging from -10.71 to $+13.21$ ppbvyr $^{-1}$ (-5.68 to $+8.84\%$ yr $^{-1}$) in Fig. 5. The largest deviation from a null trend is found over China where large surface CO emissions are reported (Liu et al., 2011; Buchwitz et al., 2007). Figure 5a contrasts to Fig. 5b, as the difference between true and a priori state (i.e. $(x - x_0)$) is constant in both figures, but with different sign. In general, the magnitude of the uncertainty depends on the temporal AKs change ($\frac{\partial A}{\partial t}$). For instance, in Fig. 5a (5b) when the difference is $-50\% \times x_0$ ($+50\% \times x_0$), the temporal increase of the AK for surface CO state force an under- (over-) estimation of the trends of retrieved state. Therefore, for both assumed conditions of true state (i.e., $x = x_0 \pm 50\% \times x_0$), the trend of the retrieved data is determined by the AKs trends multiplied by the difference between true and a priori states ($\frac{\partial A}{\partial t} (x - x_0)$).

Since the true state (and hence its difference with the a priori state) is unknown, it is impossible to estimate the actual AKs uncertainty. Furthermore, this issue is relevant for all instrumentation and algorithms that rely on temporally varying AKs. Therefore,

future trend studies using similar atmospheric sounding data like MOPITT retrievals, need to first check the temporal changes of AKs. Furthermore, because the difference between true and a priori states could be insignificant at higher level as shown in Fig. 3, the AKs uncertainty could be less than the expectation.

4 Temporal variation of level 3 a priori

In Eq. (4), we assumed that no trends are present in the a priori profiles (i.e. $\frac{\partial x_0}{\partial t} = 0$). In fact, the a priori profiles are obtained from the MOZART-4 model, which does not present a inter-annual variability (Pan et al., 1998; Deeter et al., 2003; Deeter, 2009; Worden et al., 2013a). Nevertheless, this is strictly valid only for Level 2 (daily) products, where the model climatology is spatially and temporally interpolated to the exact location and day of the observation for each retrieval (Deeter et al., 2010). In the Level 3 (monthly) data, instead, the a priori profiles can have small temporal variation globally, as shown in Fig. 6a, even though it is negligible compared to the trend of retrieved surface CO (see Fig. 2). This small trend can be attributed to the mathematical/statistical aggregation (e.g. cloud filtering-out process) to monthly (Level 3) $1^\circ \times 1^\circ$ data. Figure 6b shows the MOPITT a priori CO at the surface between 2001 and 2010 for the location where the highest trend of the a priori CO is present (i.e. $+4.08 \text{ ppbvyr}^{-1}$ at 89.5° E and 43.5° N , red star symbol in Fig. 6a). The anomaly of surface a priori CO includes an inter-annual variability and a net distinction in 2005. Hence not only does the temporal variation of Level 3 a priori data make it more difficult to predict the difference between true and a priori states (i.e. $\mathbf{x} - \mathbf{x}_0$), it also creates the additional uncertainty ($\frac{\partial \mathbf{x}_0}{\partial t} (\mathbf{I} - \mathbf{A})$) in the trend estimation as shown in Eq. (3).

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Summary and conclusions

Time varying AKs have been shown to be a significant source of uncertainties/errors in the estimation of temporal variation in retrieved data. The uncertainty test applied in this study, based on the MOPITT V5 TIR Level 3 CO products at the surface, demonstrates that the temporal changes of AKs and the differences between true and a priori states can cause significant uncertainties in the trend estimation. Unfortunately, it is not possible to eliminate the AKs uncertainty solely based on satellite observations, as the true state is unknown. Retrieved total column CO product might also be affected, since it is the vertical integral of the retrieved CO profiles (Worden et al., 2013a). However, since at higher altitudes (a) the averaging kernels are higher, and (b) the uncertainties are lower, the related uncertainties are expected to be smaller than for the near-surface layers. Future studies should investigate this aspect. Furthermore, trend studies based on satellite- and ground-based data products, which have been derived using similar atmospheric sounding methods like the MOPITT retrieval, need to carefully estimate uncertainties and errors that may be caused by time varying AKs.

Acknowledgements. The authors would like to thank the NASA Langley Research Center (Atmospheric Science Data Center, https://eosweb.larc.nasa.gov/project/mopitt/mopitt_table) for providing the MOPITT Version 5 TIR Level 3 products used in the study.

The service charges for this open access publication have been covered by the Max Planck Society.

References

- Arellano, A. F. and Hess, P. G.: Sensitivity of top-down estimates of CO sources to GCTM transport, *Geophys. Res. Lett.*, 33, L21807, doi:10.1029/2006GL027371, 2006.
- Buchwitz, M., Khlystova, I., Bovensmann, H., and Burrows, J. P.: Three years of global carbon monoxide from SCIAMACHY: comparison with MOPITT and first results related to the

ACPD

13, 20319–20340, 2013

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

detection of enhanced CO over cities, *Atmos. Chem. Phys.*, 7, 2399–2411, doi:10.5194/acp-7-2399-2007, 2007.

Burrows, J. P., Hölzle, E., Goede, A. P. H., Visser, H., and Fricke, W.: SCIAMACHY – Scanning Imaging Absorption Spectrometer for Atmospheric Cartography, *Acta Astronaut.*, 35, 445–451, 1995.

Burrows, J. P., Goede, A. P. H., Muller, C., and Bovensmann, H.: SCIAMACHY – The need for atmospheric research from space, in: *SCIAMACHY – Exploring the Changing Earth's Atmosphere*, edited by: Gottwald, M. and Bovensmann, H., Springer Science + Business Media B. V., Dordrecht, the Netherlands, 1–17, doi:10.1007/978-90-481-9896-2_1, 2011.

Choi, S. D. and Chang, Y. S.: Carbon monoxide monitoring in Northeast Asia using MOPITT: effects of biomass burning and regional pollution in April 2000, *Atmos. Environ.*, 40, 686–697, doi:10.1016/j.atmosenv.2005.09.081, 2006.

Cicerone, R. J.: How has the atmospheric concentration of CO changed, in: *The Changing Atmosphere*, edited by: Rowland, F. S. and Isaksen, I. S. A., Wiley, Chichester, UK, 49–61, 1988.

Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, 9, 6041–6054, doi:10.5194/acp-9-6041-2009, 2009.

Crutzen, P. J.: Photochemical reactions initiated by and influencing ozone in unpolluted tropospheric air, *Tellus*, 26, 47–57, 1974.

Crutzen, P. J. and Gidel, L. T.: A two-dimensional photochemical model of the atmosphere, 2: The tropospheric budgets of anthropogenic chlorocarbons CO, CH₄, CH₃Cl and the effect of various, NO_x sources on tropospheric ozone, *J. Geophys. Res.*, 88, 6641–6661, 1983.

Deeter, M. N.: MOPITT (Measurements of Pollution in the Troposphere) Validated Version 4 Product User's Guide, available at: http://www.acd.ucar.edu/mopitt/v4_users_guide_val.pdf, (last access: May 2013), 1–21, 2009.

Deeter, M. N., Emmons, L. K., Francis, G. L., Edwards, D. P., Gille, J. C., Warner, J. X., Khattatov, B., Ziskin, D., Lamarque, J.-F., Ho, S.-P., Yudin, V., Attié, J.-L., Packman, D., Chen, J., Mao, D., and Drummond, J. R.: Operational carbon monoxide retrieval algorithm and selected results for the MOPITT instrument, *J. Geophys. Res.*, 108, 4399, doi:10.1029/2002JD003186, 2003.

**Technical Note:
Temporal change in
averaging kernels**

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Deeter, M. N., Edwards, D. P., Gille, J. C., and Drummond, J. R.: Sensitivity of MOPITT observations to carbon monoxide in the lower troposphere, *J. Geophys. Res.*, 112, 24306, doi:10.1029/2007JD008929, 2007a.

Deeter, M. N., Edwards, D. P., and Gille, J. C.: Retrievals of carbon monoxide profiles from MOPITT observations using lognormal a priori statistics, *J. Geophys. Res.*, 112, D11311, doi:10.1029/2006JD007999, 2007b.

Deeter, M. N., Edwards, D. P., Gille, J. C., Emmons, L. K., Francis, G., Ho, S.-P., Mao, D., Masters, D., Worden, H., Drummond, J. R., and Novelli, P. C.: The MOPITT version 4 CO product: algorithm enhancements, validation, and long-term stability, *J. Geophys. Res.*, 115, D07306, doi:10.1029/2009JD013005, 2010.

Deeter, M. N., Worden, H. M., Gille, J. C., Edwards, D. P., Mao, D., and Drummond, J. R.: MOPITT multispectral CO retrievals: origins and effects of geophysical radiance errors, *J. Geophys. Res.*, 116, D15303, doi:10.1029/2011JD015703, 2011.

Deeter, M. N., Worden, H. M., Edwards, D. P., Gille, J. C., and Andrews, A. E.: Evaluation of MOPITT retrievals of lower tropospheric carbon monoxide over the United States, *J. Geophys. Res.*, 117, D13306, doi:10.1029/2012JD017553, 2012.

Deeter, M. N., Martínez-Alonso, S., Edwards, D. P., Emmons, L. K., Gille, J. C., Worden, H. M., Pittman, J. V., Daube, B. C., and Wofsy, S. C.: Validation of MOPITT Version 5 thermal-infrared, near-infrared, and multispectral carbon monoxide profile retrievals for 2000–2011, *J. Geophys. Res.*, 118, 1–16, doi:10.1002/jgrd.50272, 2013.

Drori, R., Dayan, U., Edwards, D. P., Emmons, L. K., and Erlick, C.: Attributing and quantifying carbon monoxide sources affecting the Eastern Mediterranean: a combined satellite, modelling, and synoptic analysis study, *Atmos. Chem. Phys.*, 12, 1067–1082, doi:10.5194/acp-12-1067-2012, 2012.

Edwards, D. P., Emmons, L. K., Hauglustaine, D. A., Chu, A., Gille, J. C., Kaufman, Y. J., Pétron, G., Yurganov, L. N., Giglio, L., Deeter, M. N., Yudin, V., Ziskin, D. C., Warner, J., Lamarque, J.-F., Francis, G. L., Ho, S. P., Mao, D., Chan, J., and Drummond, J. R.: Observations of carbon monoxide and aerosol from the terra satellite: Northern Hemisphere variability, *J. Geophys. Res.*, 109, D24202, doi:10.1029/2004JD004727, 2004.

Edwards, D. P., Emmons, L. K., Gille, J. C., Chu, A., Attié, J.-L., Giglio, L., Wood, S. W., Haywood, J., Deeter, M. N., Massie, S. T., Ziskin, D. C., and Drummond, J. R.: Satellite-observed pollution from Southern Hemisphere biomass burning, *J. Geophys. Res.*, 111, 14312, doi:10.1029/2005JD006655, 2006.

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Eskes, H. J. and Boersma, K. F.: Averaging kernels for DOAS total-column satellite retrievals, *Atmos. Chem. Phys.*, 3, 1285–1291, doi:10.5194/acp-3-1285-2003, 2003.
- Fishman, J. and Crutzen, P. J.: The origin of ozone in the troposphere, *Nature*, 274, 855–858, 1978.
- 5 Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in atmospheric constituents and in radiative forcing, in: *Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D.,
- 10 Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK, New York, NY, USA, 129–234, 2007.
- Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Szopa, S., Deeter, M. N., and Clerbaux, C.: Ten years of CO emissions as seen from Measurements of Pollution in the Troposphere (MOPITT), *J. Geophys. Res.*, 116, D05304, doi:10.1029/2010JD014416, 2011.
- 15 Hansen, J., Sato, M., Kharecha, P., Russell, G., Lea, D. W., and Siddall, M.: Climate change and trace gases, *Philos. T. Roy. Soc. A*, 365, 1925–1954, doi:10.1098/rsta.2007.2052, 2007.
- Hooghiemstra, P. B., Krol, M. C., Meirink, J. F., Bergamaschi, P., van der Werf, G. R., Novelli, P. C., Aben, I., and Röckmann, T.: Optimizing global CO emission estimates using a four-dimensional variational data assimilation system and surface network observations, *Atmos. Chem. Phys.*, 11, 4705–4723, doi:10.5194/acp-11-4705-2011, 2011.
- 20 Hoor, P., Fischer, H., and Lelieveld, J.: Tropical and extratropical tropospheric air in the lowermost stratosphere over Europe: a CO based budget, *Geophys. Res. Lett.*, 32, L07802, doi:10.1029/2004GL022018, 2005.
- Kanakidou, M., Mihalopoulos, N., Kindap, T., Im, U., Vrekoussis, M., Gerasopoulos, E., Dermizaki, E., Unal, A., Koçak, M., Markakis, K., Melas, D., Kouvarakis, G., Youssef, A. F., Richter, A., Hatzianastassiou, N., Hilboll, A., Ebojje, F., Wittrock, F., von Savigny, C., Burrows, J. P., Ladstaetter-Weissenmayer, A., and Moubasher, H.: Megacities as hot spots of air pollution in the East Mediterranean, *Atmos. Environ.*, 45, 1223–1235, doi:10.1016/j.atmosenv.2010.11.048, 2011.
- 25 Khalil, M. A. K. and Rasmussen, R. A.: The global cycle of carbon monoxide: trends and mass balance, *Chemosphere*, 20, 227–242, 1990.
- Lelieveld, J., Crutzen, P. J., Andreae, M. O., Brenninkmeijer, C. A. M., Campos, T., Cass, G. R., Dickerson, G. R., Fischer, H., de Gouw, J. A., Hansel, A., Jefferson, A., Kley, D., de

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Laat, A. T. J., Lal, S., Lawrence, M. G., Lobert, J. M., Mayol-Bracero, O., Mitra, A. P., Novakov, T., Oltmans, S. J., Prather, K. A., Ramanathan, V., Reiner, T., Rodhe, H., Scheeren, H. A., Sikka, D., and Williams, J.: The Indian Ocean Experiment: widespread air pollution from South and South-East Asia, *Science*, 291, 1031–1036, 2001.

5 Liu, C., Beirle, S., Butler, T., Liu, J., Hoor, P., Jöckel, P., Penning de Vries, M., Pozzer, A., Frankenberg, C., Lawrence, M. G., Lelieveld, J., Platt, U., and Wagner, T.: Application of SCIAMACHY and MOPITT CO total column measurements to evaluate model results over biomass burning regions and Eastern China, *Atmos. Chem. Phys.*, 11, 6083–6114, doi:10.5194/acp-11-6083-2011, 2011.

10 Liu, C., Beirle, S., Butler, T., Hoor, P., Frankenberg, C., Jöckel, P., Penning de Vries, M., Platt, U., Pozzer, A., Lawrence, M. G., Lelieveld, J., Tost, H., and Wagner, T.: CO profiles from SCIAMACHY observations using cloud slicing and comparison with model simulations, *Atmos. Chem. Phys. Discuss.*, 13, 11659–11688, doi:10.5194/acpd-13-11659-2013, 2013.

Logan, J. A., Prather, M. J., Wofsy, S. C., and Mcelroy, M. B.: Tropospheric chemistry – a global perspective, *J. Geophys. Res.*, 86, 7210–7254, 1981.

15 Pan, L., Edwards, D. P., Gille, J. C., Smith, M. W., and Drummond, J. R.: Satellite remote sensing of tropospheric CO and CH₄: forward model studies of the MOPITT instrument, *Appl. Optics*, 34, 6976–6988, doi:10.1364/AO.34.006976, 1995.

20 Pan, L., Gille, J. C., Edwards, D. P., Bailey, P. L., and Rodgers, C. D.: Retrieval of tropospheric carbon monoxide for the MOPITT experiment, *J. Geophys. Res.*, 103, D24, 32277–32290, doi:10.1029/98JD01828, 1998.

Rodgers, C. D.: Retrieval of atmospheric temperature and composition from remote measurements of thermal radiation, *Rev. Geophys.*, 14, 609–624, doi:10.1029/RG014i004p00609, 1976.

25 Rodgers, C. D.: Characterization and error analysis of profiles retrieved from remote sounding measurements, *J. Geophys. Res.*, 95, D5, 5587–5595, doi:10.1029/JD095iD05p05587, 1990.

Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *J. Geophys. Res.*, 108, 4116, doi:10.1029/2002JD002299, 2003.

30 Shindell, D. T., Faluvegi, G., Stevenson, D. S., Krol, M. C., Emmons, L. K., Lamarque, J. F., Petron, G., Dentener, F. J., Ellingsen, K., Schultz, M. G., Wild, O., Amann, M., Atherton, C. S., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isak-

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

sen, I. S. A., Lawrence, M. G., Montanaro, V., Muller, J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Strahan, S. E., Sudo, K., Szopa, S., Unger, N., van Noije, T. P. C., and Zeng, G.: Multimodel simulations of carbon monoxide: comparison with observations and projected near-future changes, *J. Geophys. Res.*, 111, D19306, doi:10.1029/2006jd007100, 2006.

Stroppiana, D., Brivio, P. A., Grégoire, J.-M., Lioussse, C., Guillaume, B., Granier, C., Mieville, A., Chin, M., and Pétron, G.: Comparison of global inventories of CO emissions from biomass burning derived from remotely sensed data, *Atmos. Chem. Phys.*, 10, 12173–12189, doi:10.5194/acp-10-12173-2010, 2010.

University of Toronto and NCAR MOPITT Team: MOPITT (Measurement of Pollution in the Troposphere) Level 1 Algorithm Theoretical Basis Documents, ATBD-MOP-01, University of Toronto and NCAR MOPITT Team, <http://eosps0.gsfc.nasa.gov/sites/default/files/atbd/atbd-mop-01.pdf>, 1–45, August, 1996.

Wallace, J. M. and Hobbs, P. V.: Atmospheric chemistry, in: *Atmospheric Science: An Introductory Survey*, 2nd edn., edited by: Dmowska, R., Hartmann, D., and Rossby, H. T., Elsevier, MA, USA, California, USA, and London, UK, 153–207, 2006.

Weatherhead, E. C., Reinsel, G. C., Tiao, G. C., Meng, X.-L., Choi, D., Cheang, W.-K., Keller, T., DeLuisi, J., Wuebbles, D. J., Kerr, J. B., Miller, A. J., Oltmans, S. J., and Frederick, J. E.: Factors affecting the detection of trends: statistical considerations and applications to environmental data, *J. Geophys. Res.*, 103, 17149–17161, doi:10.1029/98JD00995, 1998.

Weatherhead, E. C., Stevermer, A. J., and Schwartz, B. E.: Detecting environmental changes and trends, *Phys. Chem. Earth*, 27, 399–403, doi:10.1016/S1474-7065(02)00019-0, 2002.

Worden, H. M., Deeter, M. N., Edwards, D. P., Gille, J. C., Drummond, J. R., and Nédélec, P. P.: Observations of near-surface carbon monoxide from space using MOPITT multispectral retrievals, *J. Geophys. Res.*, 115, D18314, doi:10.1029/2010JD014242, 2010.

Worden, H. M., Deeter, M. N., Frankenberg, C., George, M., Nichitiu, F., Worden, J., Aben, I., Bowman, K. W., Clerbaux, C., Coheur, P. F., de Laat, A. T. J., Detweiler, R., Drummond, J. R., Edwards, D. P., Gille, J. C., Hurtmans, D., Luo, M., Martínez-Alonso, S., Massie, S., Pfister, G., and Warner, J. X.: Decadal record of satellite carbon monoxide observations, *Atmos. Chem. Phys.*, 13, 837–850, doi:10.5194/acp-13-837-2013, 2013a.

Worden, H. M., Edwards, D. P., Deeter, M. N., Fu, D., Kulawik, S. S., Worden, J. R., and Arellano, A.: Averaging kernel prediction from atmospheric and surface state parameters based

on multiple regression for nadir-viewing satellite measurements of carbon monoxide and ozone, Atmos. Meas. Tech., 6, 1633–1646, doi:10.5194/amt-6-1633-2013, 2013b.

ACPD

13, 20319–20340, 2013

Technical Note: Temporal change in averaging kernels

J. Yoon et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Technical Note:
Temporal change in
averaging kernels**

J. Yoon et al.




Fig. 1. Total mean distribution of global surface CO from MOPITT Version 5 TIR Level 3 product, referring to daytime from 2001 to 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Technical Note:
Temporal change in
averaging kernels**

J. Yoon et al.




Fig. 2. The linear trends of surface CO from MOPITT (Version 5 TIR, Level 3, daytime) from 2001 to 2010. “Not Enough Data” represents the data series discarded to avoid statistical inconsistencies. Complete yearly datasets encompass at least 9 months of data over the year.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Mean MOPITT averaging kernel for surface CO, a priori, and retrieved CO profiles over the grid cell defined by 30° E, 45° E, 110° N, and 120° N over East China on October 2002. The relative difference between retrieved CO with a priori profiles is shown to estimate the difference of true with a priori states ($x - x_0$). The retrieved CO at the surface is about 52 % higher than the a priori CO mixing ratio.

**Technical Note:
Temporal change in
averaging kernels**

J. Yoon et al.




Fig. 4. Global distributions of **(a)** maximum and **(b)** minimum relative difference between retrieved and a priori CO at surface.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technical Note: Temporal change in averaging kernels

J. Yoon et al.




Fig. 5. The linear trends of retrieved states under the assumption that the true state is **(a)** -50% or **(b)** $+50\%$ higher than the a priori state (i.e. $x = x_0 \pm 50\% \times x_0$). The trends of retrieved states range from -10.71 to $+13.21$ ppbv yr⁻¹ (-5.68 to $+8.84\%$ yr⁻¹).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Technical Note:
Temporal change in
averaging kernels**

J. Yoon et al.




Fig. 6. (a) The linear trends of surface a priori CO from MOPITT (V5 TIR Level 3) (daytime) from 2001 to 2010; the coordinates 89.5° E and 43.5° N are marked by a red star symbol; (b) MOPITT surface a priori and the seasonal CO are depicted. The anomaly of surface a priori CO shows that it varies considerably with time.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion