



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Observation and modelling of HO<sub>x</sub> radicals in a boreal forest

K. Hens<sup>1</sup>, A. Novelli<sup>1</sup>, M. Martinez<sup>1</sup>, J. Auld<sup>1,\*\*</sup>, R. Axinte<sup>1</sup>, B. Bohn<sup>4</sup>, H. Fischer<sup>1</sup>, P. Keronen<sup>3</sup>, D. Kubistin<sup>1,\*</sup>, A. C. Nölscher<sup>1</sup>, R. Oswald<sup>2</sup>, P. Paasonen<sup>3</sup>, T. Petäjä<sup>3</sup>, E. Regelin<sup>1</sup>, R. Sander<sup>1</sup>, V. Sinha<sup>5</sup>, M. Sipilä<sup>3</sup>, D. Taraborrelli<sup>1</sup>, C. Tatum Ernest<sup>1</sup>, J. Williams<sup>1</sup>, J. Lelieveld<sup>1</sup>, and H. Harder<sup>1</sup>

<sup>1</sup>Dept. Airchemistry, Max Planck Institute for Chemistry, 55128 Mainz, Germany

<sup>2</sup>Dept. Biogeochemistry, Max Planck Institute for Chemistry, 55128 Mainz, Germany

<sup>3</sup>Dept. Phys., P.O. Box 64. 00014 University of Helsinki, Finland

<sup>4</sup>Institut für Energie- und Klimaforschung IEK-8: Troposphäre Forschungszentrum Jülich GmbH, 52428 Jülich, Germany

<sup>5</sup>Department of Earth and Environmental Sciences, Indian Institute of Science Education and Research Mohali, Sector 81, S.A.S. Nagar, Manauli PO, Mohali 140 306, Punjab, India

\*now at: University of Wollongong, School of Chemistry, Wollongong, NSW, Australia

\*\*now at: Department of Chemistry and Biochemistry, University of Windsor, Windsor, Ontario N9B3P4, Canada

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 9 October 2013 – Accepted: 15 October 2013 – Published: 1 November 2013

Correspondence to: H. Harder (hartwig.harder@mpic.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

13, 28561–28629, 2013

## HO<sub>x</sub> photochemistry in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Measurements of OH and HO<sub>2</sub> radicals were conducted in a pine dominated forest in Southern Finland during the HUMPPA-COPEC-2010 (Hyytiälä United Measurements of Photochemistry and Particles in Air – Comprehensive Organic Precursor Emission and Concentration study) field campaign in summer 2010. Simultaneous side-by-side measurements of hydroxyl radicals were conducted with two instruments using chemical ionization mass spectrometry (CIMS) and laser-induced fluorescence (LIF), indicating good agreement. Subsequently, the LIF instrument was moved to the top of a 20 m tower, just above the canopy, to investigate the radical chemistry at the ecosystem–atmosphere interface. Comprehensive measurements including observations of many VOCs and the total OH reactivity were conducted and analysed using steady-state calculations as well as an observationally constrained box model.

Production rates of OH calculated from measured OH precursors are consistent with those derived from the steady state assumption and measured total OH loss under conditions of moderate OH reactivity. The primary photolytic sources of OH contribute up to one third to the total OH production. OH recycling, which occurs mainly by HO<sub>2</sub> reacting with NO and O<sub>3</sub>, dominates the total hydroxyl radical production in this boreal forest. Box model simulations agree with measurements for hydroxyl radicals ( $\text{OH}_{\text{mod.}}/\text{OH}_{\text{obs.}} = 1.04 \pm 0.16$ ), while HO<sub>2</sub> mixing ratios are significantly underpredicted ( $\text{HO}_2^{\text{mod.}}/\text{HO}_2^{\text{obs.}} = 0.3 \pm 0.2$ ) and simulated OH reactivity does not match the observed OH reactivity. The simultaneous underprediction of HO<sub>2</sub> and OH reactivity in periods in which OH concentrations were simulated well, suggests that the missing OH reactivity is an unaccounted source of HO<sub>2</sub>.

Detailed analysis of the HO<sub>x</sub> production, loss, and recycling pathways suggests that in periods of high total OH reactivity there are additional recycling processes forming OH directly, not via reaction of HO<sub>2</sub> with NO or O<sub>3</sub>. Nevertheless, a major fraction of the OH recycling occurs via the reaction of HO<sub>2</sub> with NO and O<sub>3</sub> in this terpene dominated environment.

## HO<sub>x</sub> photochemistry in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



## 1 Introduction

Removal of pollutants and trace gases from the atmosphere is mostly initiated by oxidation processes. The most important oxidant during daytime under most atmospheric conditions is the hydroxyl radical (OH), which dominates the cleansing capacity of the atmosphere.

In the lower troposphere, the main primary source of OH on a global scale is the photolysis of ozone (O<sub>3</sub>) and the subsequent reaction of the excited oxygen atom with water vapour (Levy II, 1971):



Other sources of OH, e.g. photolysis of peroxides or ozonolysis of alkenes, are known as well (Jaegle et al., 2000; Ren et al., 2008, and references therein). Once formed, OH reacts rapidly – with a lifetime of typically less than one second. Many atmospheric compounds (e.g. CO or O<sub>3</sub>) subsequently produce hydroperoxyl radicals (HO<sub>2</sub>). The oxidation of hydrocarbons by OH leads to the formation of peroxy radicals (RO<sub>x</sub> = RO<sub>2</sub> + HO<sub>2</sub>). In the presence of nitric oxide (NO) RO<sub>2</sub> is converted to HO<sub>2</sub>, which reacts with O<sub>3</sub> or NO forming OH (e.g. Martinez et al., 2003). The main HO<sub>x</sub> (= OH + HO<sub>2</sub>) sinks are self-reactions with radicals and the formation of acids and peroxides.

Forests cover about one third of the earth's total land surface (FAO, 2010). They are known to be an important global source of biogenic volatile organic compounds (BVOCs) that are very reactive towards OH (Williams, 2004). A major contribution to BVOCs, such as isoprene, monoterpenes (e.g.  $\alpha$ -pinene), and sesquiterpenes (e.g. farnesene), is emission by vegetation (Fehsenfeld et al., 1992; Guenther et al., 1995; Kesselmeier and Staudt, 1999). Therefore, forest emissions are expected to significantly influence the oxidation capacity of the atmosphere. Measurement of OH and HO<sub>2</sub> radicals in forest environments, however, unveils serious deficits in our understanding of the underlying processes. Different types of forests are known to emit various characteristic BVOCs influencing the oxidation chemistry. Several studies in

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and above isoprene-rich forest environments have been conducted in the past. Aircraft measurements of atmospheric trace gases performed over the pristine Amazon rainforest during the GABRIEL campaign (Lelieveld et al., 2008; Kubistin et al., 2010) unexpectedly showed higher OH concentrations than predicted from box and global models. Furthermore, Kubistin et al. (2010) reported that isoprene acts as a buffer for the hydroxyl radicals, which is related to an increased HO<sub>x</sub> recycling in the oxidation mechanism of BVOCs at low NO<sub>x</sub> conditions, when the OH recycling reaction of HO<sub>2</sub> with NO is inhibited. Similarly, other studies from tropical forest regions (Stone et al., 2011; Whalley et al., 2011) reported that increased recycling would be necessary to explain the measured HO<sub>x</sub> values. An increase of the deposition rates for OH reactants, like methyl vinyl ketone (MVK) and methacrolein (MACR), can help explain the measurement results for ground-based observations during the OP3 campaign in Borneo (Pugh et al., 2010). It is unclear if measurements of the total OH reactivity are available as the observations are not compared with observations in the latter study. The authors conclude that the inadequacies apparent in the model are related to the representation of detailed physical and micrometeorological effects rather than errors in the chemical scheme. Measurements conducted in a deciduous forest (Tan et al., 2001) also showed an underestimation of observed OH concentrations from box model simulations. Although two follow up studies at the same site showed model predicted OH generally being in reasonable agreement with the measured OH, the model did underestimate the OH concentrations observed by a factor of about three in the afternoon during warmer periods (Griffith et al., 2013). It remains unclear if higher biogenic VOC emissions within those periods caused an instrumental interference as recently suggested by Mao et al. (2012).

Further, Kim et al. (2013) reported reasonable agreement between observed OH and that from steady-state calculations using observations, particularly measurements of HO<sub>2</sub>, in a monoterpene dominated forest environment. Constrained box model calculations underpredicted the observed HO<sub>2</sub> by as much as a factor of eight and underestimated the observed OH by a factor of four. The authors conclude that the OH

recycling occurs mainly via the reaction of HO<sub>2</sub> with NO in this forest, which is characterized by high 2-methyl-3-buten-2-ol (MBO) and monoterpene emissions (Kim et al., 2013).

Studies on oxidation processes in monoterpene dominated environments are rare. Direct OH reactivity measurements in a boreal forest, conducted by Sinha et al. (2008) and a box model study investigating the OH reactivity budget (Mogensen et al., 2011), revealed a significant fraction of “unknown OH reactivity”. Comprehensive measurements in the same boreal forest were conducted during the HUMPPA-COPEC-2010 campaign (**H**yytiälä **U**nited **M**easurements of **P**hotochemistry and **P**articles in **A**ir – **C**omprehensive **O**rganic **P**recursor **E**mission and **C**oncentration study) (Williams et al., 2011). HUMPPA-COPEC-2010 included observations of many VOCs, HO<sub>x</sub> and the total OH reactivity to increase our understanding of atmospheric oxidation processes based on detailed analysis of production and loss processes of the hydroxyl radical and exploring the HO<sub>x</sub> budget in a coniferous forest, using direct calculations from measured species as well as an observationally constrained chemical box model in steady state.

The “Mainz Isoprene Mechanisms,” a set of reduced chemical reaction schemes considering only isoprene as the predominant primary BVOC (Taraborrelli et al., 2009 based on Pöschl et al., 2000; Taraborrelli et al., 2012), is compared to a preliminary terpene mechanism, also taking the most abundant terpenes measured during HUMPPA-COPEC-2010 (e.g.  $\alpha$ -pinene,  $\beta$ -pinene,  $\beta$ -myrcene, farnesene, and  $\Delta^3$ -carene) and their oxidation products into account. The modification of the chemical mechanism is analysed and discussed.

## 2 Methodology

In order to measure OH and HO<sub>2</sub> radicals the HORUS instrument (**H**yd**O**xyl **R**adical **M**easurement **U**nit based on fluorescence **S**pectroscopy) was deployed during the intensive HUMPPA-COPEC field experiment in summer 2010. HORUS uses the laser-

### HO<sub>x</sub> photochemistry in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



induced fluorescence based on fluorescent assay by gas expansion (LIF-FAGE) technique to detect OH and HO<sub>2</sub>.

A comprehensive dataset including measurements of the main oxidants OH, O<sub>3</sub>, and NO<sub>3</sub>; important trace gases such as CO, NO<sub>x</sub>, H<sub>2</sub>O<sub>2</sub>, HCHO, HONO; anthropogenic and biogenic VOCs, and their oxidation products, inorganic chemical constituents, aerosol properties, aerosol size distributions, as well as photolysis frequencies and other meteorological parameters was collected at the field station SMEAR II (Station for Measuring Forest Ecosystem-Atmosphere Relations). An overview can be found in Williams et al. (2011).

## 2.1 Measurement site and instrumentation

The field site is located in a boreal forest in Hyytiälä, Southern Finland (61°51' N, 24°17' E, 181 m a.s.l.) (Vesala et al., 1998). The largest city near the station is Tampere (about 200 000 inhabitants), located about 60 km S–SW of the measurement site (Hari and Kulmala, 2005). The SMEAR II station is equipped with several masts and towers surrounded by a more than 40 yr-old pine dominated forest (*Pinus Sylvestris* L.). The canopy height during the measurement period was about 20 m (for more detailed information about the continuous measurements and the infrastructure see Vesala et al., 1998; Kulmala et al., 2001; Hari and Kulmala, 2005).

During the intensive measurement period of the HUMPPA-COPEC-2010 campaign an additional scaffolding tower was set up at the field site within a clearing (the HUMPPA tower, ~ 24 m high), reaching about 2–3 m above the canopy top. The sensors and inlet lines of instruments measuring reactive species like OH, HO<sub>2</sub>, O<sub>3</sub>, NO<sub>x</sub>, NO<sub>3</sub>, HONO, H<sub>2</sub>O<sub>2</sub>, organic peroxides, HCHO, monoterpenes, as well as the associated photolysis frequencies ( $J_{\text{NO}_2}$ ,  $J_{\text{O}(^1\text{D})}$ ), and the total OH reactivity were collocated on the top platform of the HUMPPA tower to investigate the photochemistry at the ecosystem–atmosphere interface.

Since the campaign instrumentation and meteorological conditions are described elsewhere (Williams et al., 2011), only a brief description of the instruments used in

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



this study is given here; time resolution and lower limits of detection are summarized in Table 1.

Ozone was measured using a UV instrument, sharing an inlet line with the chemiluminescence detection system (CLD) for measuring NO and NO<sub>2</sub>. NO<sub>2</sub> was measured indirectly by conversion to NO using a blue light converter. The instrument setup was similar to the well-established one described by Hosaynali Beygi et al. (2011) for a different field campaign. NO<sub>3</sub> and N<sub>2</sub>O<sub>5</sub> were measured by cavity-ring-down spectroscopy (Schuster et al., 2009; Crowley et al., 2010). Two long path absorption photometer systems (LOPAP) were set up within the forest and on the HUMPPA tower, measuring HONO (Kleffmann et al., 2002). Hydrogen peroxide and the sum of organic peroxides were observed with a wet chemical system based on derivatisation and fluorescence enzyme (DEF) described by Klippel et al. (2011). Measurements of carbon monoxide were conducted using a commercial vacuum-UV resonance fluorescence CO instrument (AeroLaser GmbH, Garmisch-Partenkirchen, Germany). Detection of HCHO was based on the Hantzsch reaction and subsequent quantification of the reaction product via fluorescence detection. BVOCs were measured by several mass spectrometers. Gas chromatograph mass spectrometry (GC-MS) was used to investigate alkanes, alkenes, and particularly isoprene and monoterpenes (Yassaa et al., 2012). Proton transfer reaction mass spectrometry (PTR-MS) was applied for the observation of methanol, acetone, acetonitrile, isoprene, total monoterpenes, benzene and toluene. In the case of isoprene measurements, GC-MS data is used for further analysis, due to higher uncertainty in PTR-MS measurements, probably caused by a cold-trap connected to the PTR-MS instrument. Furthermore, the PTR-MS isoprene data might be affected by an interference due to 2-methyl-3-buten-2-ol (MBO) which is detected at the same mass to charge ratio as isoprene using conventional H<sub>3</sub>O<sup>+</sup> ionization (Goldan et al., 1997; Williams et al., 2001; Karl et al., 2012).

Photolysis frequencies ( $J_{\text{NO}_2}$ ,  $J_{\text{O}(1D)}$ ) were measured within the clearing at the forest ground, as well as on top of the HUMPPA tower with a set of filter radiometers at each position (Junkermann et al., 1989). Water vapour, temperature, and pressure

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

were recorded at several different levels on a 75 m meteorological mast located about 100 m away. The total reactivity towards OH was determined by a comparative reactivity method (Sinha et al., 2008; Nölscher et al., 2012). This method is based on the competitive scavenging of OH by a reference gas (pyrrole) and atmospheric trace gases.

5 A chemical ionization mass spectrometry (CIMS) instrument was deployed to measure OH on ground (Petäjä et al., 2009).

## 2.2 OH and HO<sub>2</sub> measurements by LIF-FAGE

Observations of OH and HO<sub>2</sub> concentrations were conducted using the Max Planck Institute for Chemistry (Mainz, Germany) HO<sub>x</sub> instrument based on laser-induced fluorescence, HORUS. The HORUS instrument is originally based on the design of GTHOS (Ground-based Tropospheric Hydrogen Oxides Sensor), the Penn State HO<sub>x</sub> instrument described by Faloon et al. (2004). HORUS is described in detail by Martinez et al. (2010).

15 OH molecules are selectively excited at low pressure by pulsed UV light at around 308 nm on resonance with the Q<sub>1</sub>(2) transition line ( $A^2\Sigma^+ - X^2\Pi$ ,  $v' = 0 \leftarrow v'' = 0$ ), and fluorescence is detected using gated micro channel plate (MCP) detectors. The UV light for excitation of the hydroxyl radicals is provided by a Nd:YAG pumped, pulsed, tunable dye laser system (Martinez et al., 2010; Wennberg et al., 1994) operated at a pulse repetition frequency of 3 kHz. During the HUMPPA-COPEC-2010 campaign, 20 optical fibers about 4 m in length were used to transfer the 308 nm light to the detection system. The sample air is drawn through a critical orifice (pinhole size of about 1.2 mm) into the low pressure detection cell, achieving a constant volume flow of about 10 slm ( $p = 1013$  hPa,  $T = 273.15$  K). In order to maximize the fluorescence signal at a certain radical level (i.e. maximizing the instrument sensitivity) a White Cell setup 25 crossing the detection volume with 32 light paths is used (White, 1942). The detection of HO<sub>2</sub> is achieved via chemical conversion to OH by adding NO in excess to the total flow of sample air downstream of the OH detection. The sum of remaining ambient

OH plus OH originating from HO<sub>2</sub> conversion is measured in a second detection axis. Calibrations of the HORUS instrument were performed on a regular basis.

## 2.2.1 Calibration

For calibration of the HORUS instrument, OH and HO<sub>2</sub> radicals were produced by photolysis of water vapour in calibration air at constant levels of absolute humidity, as described by Martinez et al. (2010). Exact knowledge of the actinic flux density of the mercury lamp (Pen-ray line source, LOT-Oriel, Europe) used for the photolytic radical production is crucial for this method. The mercury lamp used for calibration of the HORUS instrument during HUMPPA-COPEC-2010 was calibrated using the actinometry method by N<sub>2</sub>O photolysis, as described by Martinez et al. (2010), immediately before and after the campaign. The resulting actinic flux density at different N<sub>2</sub>O mixing ratios is shown in Fig. 1. Both measurements of the actinic flux density,  $\Phi_0$ , before and after the field campaign, agree within uncertainties and indicate the long-term stability of the actinic flux density. From the overall fit we obtained  $\Phi_0 = (3.68 \pm 0.04) \times 10^{12} \text{ cm}^{-2} \text{ s}^{-1}$  with a reasonable quality of fit. The systematic error of the actinic flux density measurement is calculated to be about 17 %, similar to the value by Martinez et al. (2010).

Furthermore, the instrument sensitivity is limited by quenching of the fluorescence, which happens mainly due to water vapour. To minimize this effect, the HORUS system is operated at low pressure ( $\sim 3\text{--}5$  hPa), but still high enough to keep wall losses small and provide enough molecules for excitation. The quenching effect by H<sub>2</sub>O is considered in our calculations. An additional dependency on water vapour of about 12 % and 17 % decrease in sensitivity per 1 % increase in water vapour mixing ratio was observed for the first and second fluorescence cell, respectively. This additional water effect indicates further losses at higher water mixing ratios possibly due to formation of OH-water clusters during the cold-adiabatic expansion of the sample air while entering the low pressure detection system. Changing instrument sensitivity with respect to water vapour mixing ratio, which is not caused by quenching, was reported by others (Hofzumahaus et al., 1996; Holland et al., 1998) hypothesizing that condensation

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



processes during the cold-adiabatic expansion are causing this effect. The above mentioned quantification of the additional water dependency during instrument calibration was used to correct for this dependency.

Unfortunately, the MCP detectors changed in sensitivity during the HUMPPA-COPEC-2010 field campaign, decreasing over time. Calibrations of the instrument were conducted about every second day to keep track of changes in sensitivity. Within some limitations, the behaviour of the laser scattering inside the system is an indicator of the sensitivity changes in the system. For the laser scatter signal, factors such as laser power fluctuations and background reflections have to be taken into account. Differing amounts and composition of aerosols in the sample air therefore might cause variable scatter. However, comparison of the laser scatter signal in ambient air with the quasi-simultaneous measured sensitivity during calibration shows a linear dependency. This functional dependency was applied to correct for the changing sensitivity (see Fig. 2).

### 2.2.2 Interferences

Knowledge about possible interferences and avoiding those when indicated is required in order to measure OH and HO<sub>2</sub> reliably. Interferences can be caused by the instrument itself (e.g. laser generated OH) and/or atmospheric substances which fluoresce at wavelengths similar to the hydroxyl radical. Laboratory studies testing the effect of sulfur dioxide, formaldehyde, nitrous acid, nitric acid, acetone, hydrogen peroxide, and various hydrocarbons on the OH signal did not show any significant interference for measurements in the atmosphere (Kubistin, 2009; Faloon et al., 2000; Ren et al., 2004). A negative interference on the OH signal by naphthalene was observed in polluted urban environments and can be used for the specific measurement of this compound (Martinez et al., 2004). It was recently reported that LIF measurements can be affected from internally generated OH (Mao et al., 2012; Novelli et al., 2012). Therefore, a chemical modulation method to determine the background signal for the measurement of atmospheric OH (as proposed by Brune et al., 2010) was applied to the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HORUS-LIF for the first time during HUMPPA-COPEC-2010. A new injection unit IPI (Inlet PreInjector) was developed and optimized to scavenge more than 95 % of atmospheric OH by periodic injection of a chemical reactant in front of the standard inlet. In order to minimize wall losses, a bypass flow, large compared to the sample flow into the detection system, was established. Optimization of this injection system included tests using different reactants, e.g. propylene and hexafluoropropene, and varying injection and bypass flow conditions to determine the best parameters for continuous operation. A different publication on the characterisation of the OH scavenger injection device IPI is in preparation by Novelli et al. (2013).

Measurements of HO<sub>2</sub> concentrations are conducted using chemical conversion into OH via the reaction with NO and the subsequent detection of the hydroxyl radicals by the LIF technique (Reaction R3):



In the atmosphere, RO<sub>2</sub> is also converted into HO<sub>2</sub> in the presence of NO (Reactions R4 and R5). These reactions were expected to be negligible in low pressure FAGE detection systems due to the reduced oxygen concentration and the short reaction time between injection of NO and detection of OH within the system (Heard and Pilling, 2003).



We observed a small interference (< 10 %) originating from methyl peroxy radicals in the laboratory for the HORUS instrument, comparable to the results reported by Holland et al. (2003), who detected an interference smaller than 5 % for their LIF-FAGE instrument. Significant interferences in HO<sub>2</sub> measurements by the LIF-FAGE technique from RO<sub>2</sub> species resulting from the OH-initiated oxidation of alkenes and aromatics have been observed by Fuchs et al. (2011) and others (Dillon, 2011). In contrast to alkane-based peroxy radicals, which are formed in the reaction of VOC + OH via H-atom abstraction, alkene-based peroxy radicals (mainly produced via OH-addition on

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the C-C double bond) form specific RO by reaction with NO, that can decompose under the low pressure conditions forming HO<sub>2</sub> rapidly. The conversion efficiency is limited by Reaction R4. Thus, any potential interference from alkenes on HO<sub>2</sub> can be reduced by reducing reaction time and/or NO concentration. During HUMPPA-COPEC-2010 NO mixing ratios of about 400 ppmV occurred inside the detection system of HORUS by injection of pure NO into sample air, yielding a conversion efficiency from HO<sub>2</sub> to OH of more than 95 % at the operating conditions of the instrument. This could also have caused a high conversion efficiency for interfering RO<sub>2</sub> species as described above. Nevertheless, this interference depends strongly on the available VOCs for the production of RO<sub>2</sub> in the investigated environment as well as on the instrumental setup which defines the conversion efficiency of those peroxy radicals inside the instrument. The observed HO<sub>2</sub> can be interpreted as the sum of atmospheric HO<sub>2</sub> and a contribution from the effective interference by specific RO<sub>2</sub>:

$$[\text{HO}_2]_{\text{obs.}} = [\text{HO}_2]_{\text{atm.}} + \sum_i (\alpha_{\text{RO}_2}^i \times [\text{RO}_2]_i) \quad (1)$$

The relative detection sensitivities  $\alpha_{\text{RO}_2}^i$  for the specific RO<sub>2</sub> compared to HO<sub>2</sub> are strongly dependent on the instrumental setup and can reach values up to about 0.9 (Fuchs et al., 2011; Lu et al., 2012; Whalley et al., 2013). However, the speciation of RO<sub>2</sub> in the observed environment strongly influences the effective interference. Based on model simulations the magnitude of the RO<sub>2</sub> interference on HO<sub>2</sub> measurements by some LIF instruments in different environments was estimated between 10 % up to about 30 % (Lu et al., 2012; Fuchs et al., 2011; Mao et al., 2012; Whalley et al., 2013; Griffith et al., 2013).

NO titration experiments during calibration and ambient air measurements in two forest environments in Germany have been conducted after HUMPPA-COPEC-2010, quantifying the maximum interference by RO<sub>2</sub> in the HORUS instrument in these environments to be less than 20 % (Tatum Ernest et al., 2012).

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To further investigate the effect of RO<sub>2</sub> interference on the measurement of HO<sub>2</sub> concentrations during HUMPPA-COPEC-2010, we calculated the H<sub>2</sub>O<sub>2</sub> budget taking the production by the self-reaction of HO<sub>2</sub> and the loss via photolysis and deposition into account and compared the calculated hydrogen peroxide with the measurements (Fig. 3). The production of H<sub>2</sub>O<sub>2</sub> by HO<sub>2</sub> shows a quadratic dependency and is therefore highly sensitive to the HO<sub>2</sub> concentration. The decay of hydrogen peroxide in the afternoon is mainly determined by the deposition process. A reasonable deposition rate of  $4 \times 10^{-5} \text{ s}^{-1}$ , corresponding to a deposition velocity of  $4 \text{ cm s}^{-1}$  in a 1 km high boundary layer, was used for the calculation. This is comparable to values reported by Hall and Claiborn (1997) for a boreal forest, ranging from 1 to  $5 \text{ cm s}^{-1}$ . Removal of hydrogen peroxide by photolysis makes up to 10% of the total H<sub>2</sub>O<sub>2</sub> loss. Hydrogen peroxide concentrations derived using the measured HO<sub>2</sub> is in reasonable agreement with the measured H<sub>2</sub>O<sub>2</sub>. In some cases measured HO<sub>2</sub> is still not sufficient to explain the measured H<sub>2</sub>O<sub>2</sub> concentrations. Even though under some conditions hydrogen peroxide measurements might be affected by mixing with different air masses, e.g. from the residual layer, which have not been considered in our calculation, the comparison gives confidence that the measurements of HO<sub>2</sub> are not subject to a major interference. Although the magnitude of the RO<sub>2</sub> interference during HUMPPA-COPEC-2010 cannot be conclusively derived there is no evidence for an extraordinary large interference in the HORUS instrument compared to other LIF systems. Therefore, a contribution of 30% to the observed HO<sub>2</sub> signal (presently the maximum value observed in LIF instruments) is considered as an upper limit estimate of the RO<sub>2</sub>-interference for further analysis.

### 2.3 OH budget – calculated based on observations

Subsequent to the ground-level comparison with the OH-CIMS (1–8 August), the HORUS instrument was moved to the top of the HUMPPA tower to investigate the photochemistry at the ecosystem–atmosphere interface. Making use of the comprehensive measurements of atmospherically relevant species, described in Sect. 2.1, the known

daytime production rates of OH can be calculated. Besides the photolytic sources (Reactions R6 and R7), ozonolysis of different biogenic VOCs (Reaction R8) contributes to the primary production of OH. Secondary sources, e.g. the recycling of HO<sub>2</sub> by NO and O<sub>3</sub> (Reactions R11 and R12) and peroxide photolysis (Reactions R9 and R10), additionally play an important role in OH radical production. We define:

Primary production



Secondary production/Recycling



Under steady-state conditions, which can be assumed for short-lived compounds like OH, the sum of these production rates should equal the total loss of hydroxyl radicals, which can be derived from the product of the measured OH concentration and the total OH reactivity measurements ( $k'_{\text{OH}}$ ).

$$P_{\text{OH}}^{\text{total}} = \sum_i P_{\text{OH},i} \quad (2)$$

$$L_{\text{OH}}^{\text{total}} = \sum_i L_{\text{OH},i} = k'_{\text{OH}}[\text{OH}] = \tau^{-1}[\text{OH}] \quad (3)$$

$$P_{\text{OH}}^{\text{total}} \stackrel{!}{=} L_{\text{OH}}^{\text{total}} \quad (4)$$

Comparing the sum of production rates from known pathways with the measured total loss of hydroxyl radicals can be used as a tool to unveil gaps in our understanding of the underlying processes. Furthermore, the importance of the different contributions to the total production of OH can be determined. A more detailed analysis was also conducted taking into account unmeasured BVOCs and higher order oxidation products using a measurement constrained box model (see Sect. 2.4).

## 2.4 Box model

To investigate the influence of unmeasured intermediate reaction products, the observations were compared to simulations with version 3.0 of the chemical box model CAABA/MECCA (**C**hemistry **A**s **A** Boxmodel **A**pplication/**M**odule **E**fficiently **C**alculating the **C**hemistry of the **A**tmosphere) by Sander et al. (2011). We constrained the model with observed, complete 5 min datasets. Each simulation was terminated when both OH and HO<sub>2</sub> reached steady-state, typically after about 48 h.

MECCA contains a comprehensive atmospheric chemistry reaction scheme. However, since we focus on organics, we switched off halogen and sulfur chemistry, as well as heterogeneous and aqueous phase reactions. A list of the chemical reactions used in this study, including rate coefficients and references, is available in the Supplement. In the base configuration, version 2 of the isoprene chemistry from the “Mainz Isoprene Mechanism” (MIM2) was used, considering 68 species and 195 reactions (Taraborrelli et al., 2009 based on Pöschl et al., 2000). For sensitivity studies, the recently developed isoprene mechanism MIM3 (Taraborrelli et al., 2012), which includes additions to the isoprene chemistry; such as the photo-oxidation of unsaturated hydroperoxy-aldehydes; and a preliminary version of the monoterpene mechanism MTM (Taraborrelli et al., in preparation) were also used. The latter is based on a MIM2-like version of MIM3 (MIM3\*), i.e. hydroperoxy-aldehyde chemistry, H-shifts, and RO<sub>2</sub> + HO<sub>2</sub> reactions are considered the way of MIM2, while updated estimates of rate constants from MIM3 were retained. Furthermore, it has a representation for the oxidation of the major terpenes during HUMPPA-COPEC-2010 that being  $\alpha$ -pinene,  $\beta$ -pinene,  $\beta$ -

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



myrcene,  $\Delta^2$ -carene,  $\Delta^3$ -carene, and  $\alpha$ -farnesene chemistry. The oxidation of the first two monoterpenes is taken from the MCM<sup>1</sup>, with updates based on recent literature. Carene is assumed to yield the same products as  $\alpha$ -pinene. Finally the oxidation of  $\beta$ -myrcene and  $\alpha$ -farnesene is simplified and partially follows an isoprene-like oxidation.

5 Deposition was included for the species listed in Table 2, according to values given in literature or derived from measurements. For the numerical integration of the resulting set of differential ordinary equations (ODEs), the KPP software (Sandu and Sander, 2006) with a positive definite Rosenbrock solver and automatic time-step control was used. For the photolysis frequencies of NO<sub>2</sub> and O(<sup>1</sup>D), measured values (as described

10 above) were used. Photolysis frequencies for other observed photolabile species are calculated based on measured  $J_{O(^1D)}$  and  $J_{NO_2}$  using parameterisations from spectroradiometer measurements at other field sites (Bohn et al., 2008, B. Bohn, personal communication, 2012). Therefore effects of cloud coverage and aerosols are considered implicitly. In case of missing methane (CH<sub>4</sub>) data, box model simulations were

15 conducted using the median value of CH<sub>4</sub> observed, 1.79  $\mu\text{mol mol}^{-1}$ . The 1 $\sigma$  variability in CH<sub>4</sub> measurements during the entire campaign was 0.03  $\mu\text{mol mol}^{-1}$ . Measured photolysis frequencies below the lower limit of detection were set to zero, which affects the derived photolysis frequencies from other species accordingly. The uncertainty of simulated HO<sub>x</sub> caused by these assumptions is less than 10%. Datasets missing any

20 other of the species used as model input were omitted from this study.

### 2.4.1 Significance and sensitivity

The significance of the discrepancies between observed and simulated HO<sub>x</sub> concentrations depends on the uncertainty of the HO<sub>x</sub> measurements as well as on the uncertainty of the model simulation. The latter comprises the uncertainties of all measured

25 input variables such as trace gas concentrations and meteorological parameters as well as the uncertainties due to the chemical mechanism applied.

<sup>1</sup>Master Chemical Mechanism, MCM v3.2

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sensitivity analysis on measured species by varying constraint parameters within their uncertainty can be used to identify the impact on the model result. In the case of unmeasured species, e.g. oxidation products and intermediates, the uncertainties depend on the uncertainty of their production and destruction pathways as well as the uncertainty of the chemical mechanism. The uncertainties of the simulated OH and HO<sub>2</sub> concentrations related to the reaction rate coefficients were estimated by Monte-Carlo simulations as described by Sander et al. (2011). The result of one Monte-Carlo simulation in case of OH applying the isoprene mechanism (MIM2) is shown as an example in Fig. 4. The histogram shows a binned frequency distribution of the model OH resulting from the individual Monte-Carlo runs, simulated with diverse sets of rate coefficients. This yields a 1 $\sigma$ -uncertainty of the simulated OH resulting from the uncertainty of rate coefficients of 15%.

### 3 Results and discussion

#### 3.1 Observations

During the HUMPPA-COPEC-2010 field experiment gradients of the hydroxyl radical were measured using the CIMS instrument from University of Helsinki on ground (Petäjä et al., 2009) and the HORUS-LIF instrument above the canopy. To assure the comparability of both instruments and techniques they had to be compared side-by-side under ambient conditions. The HORUS-LIF instrument already participated in the international HO<sub>x</sub>Comp 2005 project, a ground-based intercomparison of six OH instruments (4 LIF, 1 CIMS, 1 DOAS<sup>2</sup>) performing measurements in the atmosphere simulation chamber SAPHIR as well as in ambient air. The HORUS-LIF showed good agreement with CIMS measurements ( $R^2 = 0.96$ ) during daytime. However, we were

<sup>2</sup>Differential Optical Absorption Spectroscopy

expecting from that study to see a nighttime signal in ambient air whereas the CIMS method usually does not detect nighttime OH (Schlosser et al., 2009).

During HUMPPA-COPEC-2010 the two OH instruments were intercompared at the beginning of the field experiment (27–30 July) before starting the gradient measurement of OH concentrations. The inlet system of the LIF was placed next to the CIMS inlet on ground. The meteorological conditions at the field site within the four days of instrument intercomparison were dominated by above average temperatures, exceeding 25 °C during noon, and mainly south-easterly winds, without rainy periods.

The result of these four days of instrument intercomparison is shown in Fig. 5. OH radical concentrations observed by both techniques are generally in agreement. The LIF data shows stronger fluctuations compared to the measurements by CIMS on the same averaging interval timescale of 30 min. Daytime maximum OH concentrations reached about  $(1.5\text{--}2) \times 10^6 \text{ molec cm}^{-3}$ . During nighttime, both instruments observed OH concentrations below  $5 \times 10^5 \text{ molec cm}^{-3}$ , but still well above the lower limit of detection of the CIMS, which was  $5 \times 10^4 \text{ molec cm}^{-3}$  at a time resolution of 30 s. For the LIF the detection limit was  $4.8 \times 10^5 \text{ molec cm}^{-3}$  at a time resolution of 60 min. The larger limit of detection as well as the increased fluctuations in the LIF observation compared to CIMS is mainly caused by application of the chemical modulation method. To determine atmospheric OH the interference signal is subtracted from the total observed signal. The detection limit is therefore not only given by instrument properties, like laser power, optical properties, and detector efficiency. It is also prone to the atmospheric variability both of OH and of the species causing the interference inside the detection unit. Nevertheless, the good agreement during the intercomparison (Fig. 6) provides confidence in the chemical modulation method and that the LIF measurements are not affected by additional unknown OH interferences resulting from the laser fluorescence technique (Novelli et al., 2012).

Finally, the LIF instrument was moved to the top of the HUMPPA tower to investigate the radical chemistry at the interface between atmosphere and ecosystem while

## HO<sub>x</sub> photochemistry in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the CIMS stayed on the ground. The resulting hydroxyl radical measurements are presented in Fig. 7.

Simultaneous OH measurements on ground and above canopy revealed a factor of 2–3 difference in observed concentration maxima, reaching values up to about  $3 \times 10^6$  molec $\text{cm}^{-3}$  on the tower. During nighttime, the hydroxyl radical observations above canopy and on ground showed similar values. Linear correlation of OH and photolysis frequency  $J_{\text{O}(^1\text{D})}$  was previously found during ground-based campaigns (Rohrer and Berresheim, 2006). However, the slope varies with location, depending on the abundance of VOCs and  $\text{NO}_x$ . The comparison of the correlation on ground and above canopy shows that higher OH values on tower are driven by higher radiation but does not suggest differences in the chemical regime (Fig. 8).

The nighttime OH, which was occasionally well above the detection limit of both instruments, can not be explained by the supporting observations, with known sources accounting for about 20 % of the total OH production necessary to explain the measured concentrations.  $\text{NO}_3$  was always below the lower limit of detection of about 1 pptV (at 5 min. time resolution) for the CRD instrument (Rinne et al., 2012). However, the production rate for  $\text{NO}_3$  from the reaction of  $\text{NO}_2$  and  $\text{O}_3$  is of the order of  $1 \times 10^6$  molec $\text{cm}^{-3} \text{s}^{-1}$ . Assuming this would directly cause an equally high OH production in the oxidation process of VOCs, which is an unlikely high upper estimate, this would still only explain an additional 10 to 15 % of the missing production during nighttime. The ozonolysis of unmeasured VOCs and their oxidation products, as well as enhanced  $\text{HO}_x$  recycling are potential candidates to explain the missing OH production. A more detailed analysis of the most relevant OH production terms is given in Sect. 3.1.1.

### 3.1.1 Radical budget based on observations

Hydroxyl radical production was calculated taking the primary production by  $\text{O}_3$  photolysis, as well as ozonolysis of observed hydrocarbons and recycling via  $\text{HO}_2$  by reaction with NO and  $\text{O}_3$  into account (see Sects. 2.3, R6–R12). It reaches a maximum of about

28580

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$1 \times 10^7$  molec $\text{cm}^{-3} \text{s}^{-1}$  on ground and  $1.4 \times 10^7$  molec $\text{cm}^{-3} \text{s}^{-1}$  above canopy around local solar noon as shown in Fig. 9. Since NO and NO<sub>2</sub> were not measured at ground level, they were derived from the above canopy measurements assuming constant NO<sub>x</sub> with height, using the filter radiometer measurements of the photolysis frequency  $J_{\text{NO}_2}$  on ground and above the canopy to calculate partitioning between NO<sub>2</sub> and NO. The enhanced radical production above canopy is caused by higher  $J_{\text{O}(^1\text{D})}$  observed on the tower and the enhanced recycling (Fig. 9). The dominant primary source of OH (21 %) is the reaction of O(<sup>1</sup>D) with water. HONO photolysis, with a contribution during noon time of about 7 %, is also significant. However, the conversion of HO<sub>2</sub> to OH via the reactions with NO and O<sub>3</sub> dominates the total production of hydroxyl radicals (60–80 %). The ozonolysis of measured biogenic VOCs plays a minor role as a source of OH during daytime but becomes more important during nighttime.

The budget of steady-state OH was calculated as described in Sect. 2.3, using the available measurements of the chemical species contributing to the production rates and the total OH reactivity observed above canopy as constraints. Mean OH reactivity measured during this period was  $11.5 \text{ s}^{-1}$  varying typically between the lower limit of detection ( $3\text{--}4 \text{ s}^{-1}$ ) and about  $30 \text{ s}^{-1}$ , with some peaks reaching above  $70 \text{ s}^{-1}$  (Nölscher et al., 2012).

The known OH sources are almost sufficient to close the budget above the canopy. Isoprene contributes less than 10 % at maximum to the total OH reactivity measured during HUMPPA-COPEC-2010 due to its low mixing ratio (typically below 200 pptV). Observed terpenes, being similar or slightly less reactive than isoprene, e.g.  $\alpha$ -pinene (up to 5 %),  $\beta$ -pinene (up to 4 %),  $\beta$ -myrcene (up to 6 %), and  $\Delta^3$ -carene (up to 9 %) were more abundant, thus providing in sum a higher reactivity towards OH.



Nevertheless, assuming steady-state conditions for HO<sub>2</sub> and taking the hydroperoxy radical measurements into account, the RO<sub>2</sub> concentration can be estimated.

$$\begin{aligned} \frac{d[\text{HO}_2]}{dt} &= 0 \\ &= P_{\text{HO}_2}^{\text{total}} - L_{\text{HO}_2}^{\text{total}} \\ &= P_{\text{HO}_2}^{\text{known}} + b k_{\text{R19}}[\text{NO}][\text{RO}_2] - L_{\text{HO}_2}^{\text{known}} - k_{\text{R24}}[\text{HO}_2][\text{RO}_2] \\ &= P_{\text{HO}_2}^{\text{known}} - L_{\text{HO}_2}^{\text{known}} + (b k_{\text{R19}}[\text{NO}] - k_{\text{R24}}[\text{HO}_2])[\text{RO}_2] \end{aligned} \quad (5)$$

where  $b$  is a branching ratio and  $k_i$  denotes the reaction rate constant associated with Reaction  $R_i$ . From this it follows

$$[\text{RO}_2] = \frac{L_{\text{HO}_2}^{\text{known}} - P_{\text{HO}_2}^{\text{known}}}{b k_{\text{R19}}[\text{NO}] - k_{\text{R24}}[\text{HO}_2]} \quad (6)$$

Typically there was about twice as much RO<sub>2</sub> as HO<sub>2</sub> (RO<sub>2</sub>/HO<sub>2</sub>: median 1.8; min. 0.5; max. 4.5). This result was confirmed by calculation of the modified Leighton ratio (RO<sub>2</sub>/HO<sub>2</sub>: median 1.8; min. 0.1; max. 6.3) (Leighton, 1961).

$$\Phi := \frac{J_{\text{NO}_2}[\text{NO}_2]}{[\text{NO}](k_{\text{NO}+\text{O}_3}[\text{O}_3] + k_{\text{R19}}([\text{RO}_2] + [\text{HO}_2]))} \quad (7)$$

$$[\text{RO}_2]_{\Phi=1} = \frac{J_{\text{NO}_2}[\text{NO}_2]}{k_{\text{R19}}[\text{NO}]} - \frac{k_{\text{NO}+\text{O}_3}[\text{O}_3] - [\text{HO}_2]}{k_{\text{R19}}} \quad (8)$$

Thus organic peroxy radicals seem to play a major role in the radical photochemistry in the observed boreal forest environment, providing a strong link between OH and HO<sub>2</sub>.

The importance of radical cycling via HO<sub>2</sub> shown in the budget demands careful examination of the reliability of HO<sub>2</sub> measurements. The measurement of HO<sub>2</sub> by conversion to OH using NO can be severely affected by RO<sub>2</sub> reacting with NO also producing hydroxyl radicals (Fuchs et al., 2011). However, this interference depends on

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the instrumental setup (Whalley et al., 2013). It is to be quantified specifically for each instrument and depends on the composition of RO<sub>2</sub> in ambient air. The HO<sub>2</sub> observed during HUMPPA-COPEC-2010 therefore gives an upper limit for the atmospheric HO<sub>2</sub> including an unknown contribution from RO<sub>2</sub>. To put limits on the atmospheric HO<sub>2</sub>, the budget of H<sub>2</sub>O<sub>2</sub> was analyzed. The H<sub>2</sub>O<sub>2</sub> intercomparison (Fig. 3) does not suggest that the HO<sub>2</sub> measurements by HORUS during HUMPPA-COPEC-2010 were affected by a major RO<sub>2</sub> interference (see Sect. 2.2.2). Nevertheless, assuming an interference by RO<sub>2</sub> species on the HO<sub>2</sub> measurements during HUMPPA-COPEC-2010 of 30% would increase the gap between total OH production and total loss of OH. Recycling via HO<sub>2</sub> would still be the predominant source of OH.

### 3.1.2 HO<sub>x</sub> recycling pathways

Production, loss, and recycling pathways of HO<sub>x</sub> above the canopy under various conditions are shown in Table 3. Average rates calculated from observations highlight once more the importance of OH recycling via HO<sub>2</sub>. Ozonolysis of BVOCs is an important primary radical source at low radiation levels. Photolytic OH production from O<sub>3</sub> as well as HONO photolysis are more relevant primary radical sources during daytime. The contribution of acetone photolysis to HO<sub>2</sub> formation is small at the surface, typically less than 1% of the total direct radical sources. The main sink terms are radical self-reactions, particularly HO<sub>2</sub> + RO<sub>2</sub>. Since RO<sub>2</sub> was not measured during HUMPPA-COPEC-2010, it is estimated from HO<sub>2</sub> steady-state (see Sect. 3.1.1). The total OH production,  $P_{\text{OH}}^{\text{total}}$ , is calculated from total OH reactivity data and the hydroxyl radical measurements assuming steady-state (Eqs. 3 and 4). The loss rate of OH via reaction with VOCs and oxygenated VOCs (OVOCs) is derived by subtracting the known OH loss rates caused by CO, O<sub>3</sub>, HCHO, H<sub>2</sub>O<sub>2</sub>, NO<sub>2</sub>, NO, SO<sub>2</sub>, and HO<sub>2</sub> from the total loss observed ( $k'_{\text{OH}}[\text{OH}]$ ).

The dataset can be grouped by conditions of low and high radiation, and moderate and high observed total OH reactivity. The resulting four cases (see Table 3) are described in more detail in the following paragraphs.



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Assuming steady-state, the total OH production is equal to the product of the measured total OH reactivity and the OH concentration (Eqs. 3 and 4). Under *daylight* conditions ( $J_{O(^1D)} > 3 \times 10^{-6} \text{ s}^{-1}$ ), at *moderate* observed total OH reactivity ( $k'_{OH} \leq 15 \text{ s}^{-1}$ ) (Table 3, upper left)  $L_{OH}^{\text{total}} \approx 3.3 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  is calculated from observations.

Primary production of OH radicals is on average  $1.2 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ . The reactions of HO<sub>2</sub> with NO ( $\chi(\text{NO}) \approx 46 \text{ pptV}$ ) and O<sub>3</sub> produce about  $3.3 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  OH, which is 73 % of the known summed OH production. The known OH sources are more than sufficient to close the OH budget. An OH loss rate of about  $1.2 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  is missing under these conditions, which is close to the combined uncertainty of 40 %. OH is directly lost at a rate of  $0.1 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  by radical termination reactions with NO<sub>2</sub>, NO, and HO<sub>2</sub>. OH reacts with CO, O<sub>3</sub>, HCHO, SO<sub>2</sub>, and H<sub>2</sub>O<sub>2</sub>, yielding HO<sub>2</sub> radicals at a rate of  $0.8 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ . The remaining observed total OH reactivity (equivalent to an average OH loss rate of  $2.4 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ ) constrains the maximum production of RO<sub>2</sub> from the reaction of OH with VOCs. However, this is only 75 % of the RO<sub>2</sub> loss rate via reaction with NO, which is the dominant source of HO<sub>2</sub>. This indicates a missing RO<sub>2</sub> source. Besides the ozonolysis of unmeasured BVOCs, reservoir species such as PAN, which thermally decompose (showing a strong temperature dependency), could contribute to the missing RO<sub>2</sub> production rate of the order of  $1.2 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ . The photolytic HO<sub>2</sub> production by HCHO +  $h\nu$  contributes with a rate of  $0.2 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ . Radical-radical termination reactions govern the HO<sub>2</sub> loss (on average about  $0.9 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ ).

Under conditions of *low radiation* ( $J_{O(^1D)} \leq 3 \times 10^{-6} \text{ s}^{-1}$ ) and *moderate* observed total OH reactivity ( $k'_{OH} \leq 15 \text{ s}^{-1}$ ) (Table 3, upper right) the known primary production of OH radicals is only  $0.3 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ . Also, the classical recycling of HO<sub>2</sub> to OH by reaction with NO ( $\chi(\text{NO}) \approx 3 \text{ pptV}$ ) and O<sub>3</sub> is inhibited ( $0.9 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ ), mainly due to the low NO concentration. Comparison with the derived total production rate (Eq. 4) reveals a missing fraction of 40 % ( $P_{OH}^{\text{missing}} \approx 1 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ ).

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This is in the same order of magnitude of the observed NO<sub>3</sub> production rate of  $1 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup>. Since the NO<sub>3</sub> concentration always remained below the lower limit of detection its reactivity had to be fast, potentially to some extent producing OH. Also ozonolysis of unmeasured BVOCs could directly produce OH and possibly close the OH budget.

Under *daylight* conditions ( $J_{O(1D)} > 3 \times 10^{-6}$  s<sup>-1</sup>), with *high* observed total OH reactivity ( $k'_{OH} > 15$  s<sup>-1</sup>) (Table 3, lower left) the primary production of OH radicals is on average  $1.5 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup>. Less than 10% of these primary produced OH radicals are directly lost by radical termination reactions with NO<sub>2</sub>, NO, or HO<sub>2</sub>, forming acids and H<sub>2</sub>O. The largest known source of OH is again HO<sub>2</sub> recycling by reaction with NO ( $\chi(\text{NO}) \approx 28$  pptV) and O<sub>3</sub> with an average OH production rate of  $6.3 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup>. However, only about 30% of the total OH production is known. The excess OH reactivity (equivalent to an average OH loss rate of  $27 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup>) is available for production of RO<sub>2</sub> radicals. RO<sub>2</sub> radicals react with NO forming HO<sub>2</sub> with a yield of about 90%, representing the main source of HO<sub>2</sub>. In comparison to this, the direct production of HO<sub>2</sub> radicals from HCHO photolysis is negligible ( $0.2 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup>). Through the relationship between the RO<sub>2</sub> loss (forming HO<sub>2</sub>) and the potential production rate of RO<sub>2</sub> by unaccounted observed total OH reactivity it can be estimated that about 50% of the remaining OH reactivity, equivalent to a rate of  $14 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup>, is from other processes, e.g. direct OH recycling, not via the reaction of HO<sub>2</sub> + NO/O<sub>3</sub>. Furthermore, an OH production rate of the order of  $20 \times 10^6$  molec cm<sup>-3</sup> s<sup>-1</sup> is missing, to close the OH budget under these conditions. The reaction of specific alkylperoxy radicals with HO<sub>2</sub> could represent an additional RO<sub>2</sub> sink and OH source. Laboratory studies investigating the OH production from the reactions of ethyl peroxy, acetyl peroxy, and acetonyl peroxy radicals with HO<sub>2</sub> revealed OH yields up to 70% (Thornton et al., 2002; Hasson et al., 2004; Jenkin et al., 2007; Dillon and Crowley, 2008). Assuming that all RO<sub>2</sub> species, as an upper limit, would react with HO<sub>2</sub> forming OH, this would contribute an addi-

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tional  $10 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  to the OH production. However, this would still not be sufficient to close the OH budget under these conditions of high radiation and high OH reactivity. A direct recycling mechanism, returning OH from RO<sub>2</sub> in the absence of NO, by a 1,5-H-shift, was proposed by Peeters et al. (2009) for the OH-initiated degradation of isoprene. A recent chamber study suggests a reduced reaction rate constant by about a factor of two with respect to the value given by Peeters et al. (2009) for this 1,5-H-shift (Fuchs et al., 2013). Nevertheless, at low NO levels ( $\chi(\text{NO}) \approx 100 \text{ pptV}$ ) these isomerization reactions for isoprene-related RO<sub>2</sub> radicals forming OH directly become competitive with the traditional reactions of RO<sub>2</sub> with HO<sub>2</sub>, RO<sub>2</sub>, and NO. Similarly, other RO<sub>2</sub> radicals, specifically all conjugated alkadienes, e.g. myrcene, many mono-cyclic monoterpenes, and also sesquiterpenes, can undergo a fast H-shift isomerization (Peeters et al., 2001; Vereecken et al., 2007, 2012). Monoterpenes and sesquiterpenes have many more pathways to oxidise than isoprene, and the channels that allow for H-migration are therefore a smaller percentage than in isoprene. Still, an OH production rate of the order of  $6 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  is missing. A direct OH source or additional RO<sub>2</sub> production, balanced by an equivalent loss of RO<sub>2</sub> forming OH via mechanisms like the ones mentioned above could close the OH budget.

Similarly, at *low radiation* ( $J_{\text{O}(^1\text{D})} \leq 3 \times 10^{-6} \text{ s}^{-1}$ ) and *high observed total OH reactivity* ( $k'_{\text{OH}} > 15 \text{ s}^{-1}$ ) (Table 3, lower right) the known production of OH radicals is on average only about 20 % of the total OH production. HO<sub>2</sub> reacts with NO ( $\chi(\text{NO}) \approx 17 \text{ pptV}$ ) and O<sub>3</sub> producing about  $3.9 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  OH, the largest known contribution to  $P_{\text{OH}}^{\text{total}}$ . For the production of RO<sub>2</sub> radicals the remaining observed total OH reactivity suggests a rate of  $20.6 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ . About  $7 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  of the RO<sub>2</sub> radicals react with NO forming HO<sub>2</sub>. An RO<sub>2</sub> loss rate of about  $12 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$  remains, e.g. for direct recycling mechanisms, recycling OH as proposed above, i.e. alkylperoxy radicals reacting with HO<sub>2</sub> and/or H-shift isomerization or similar mechanisms are likely, potentially providing a large fraction of the missing OH production rate of  $17 \times 10^6 \text{ molec cm}^{-3} \text{ s}^{-1}$ .

## 3.2 Box Model simulations

The previous discussion indicates, that under conditions of low reactivity and high radiation, recycling of OH in this environment occurs mainly by HO<sub>2</sub>. In order to identify the influence of unmeasured oxidation products on the HO<sub>x</sub> budget and to examine the current understanding of the underlying processes box model simulations have been conducted.

The CAABA/MECCA box model was applied in steady-state mode with concentrations of NO, CO, O<sub>3</sub>, H<sub>2</sub>O<sub>2</sub>, organic peroxides, isoprene (C<sub>5</sub>H<sub>8</sub>), terpenes, HONO, HCHO, and H<sub>2</sub>O as well as photolysis frequencies constrained to measured values. Simulations were only done when data of all key constraints were available (see Sect. 2.4).

Applying the chemistry scheme from the “Mainz Isoprene Mechanism” (MIM2), OH concentrations are overestimated by the model by about 40 % on average (Fig. 10). HO<sub>2</sub> is underestimated in the simulation. The model-measurement discrepancy can be divided into two groups: The ratio of simulated to observed HO<sub>2</sub> concentrations is about 0.7 for a minor part of the dataset, while the simulation significantly underestimates the observed total OH reactivity for the rest of the dataset. The ratio between simulated and observed HO<sub>2</sub> concentrations in this case is only about 0.3. A recently published chemical reaction scheme (MIM3) including new additions to the isoprene chemistry, such as the photo-oxidation of unsaturated hydroperoxy-aldehydes (Taraborrelli et al., 2012), produces even more hydroxyl radicals leading to an overestimation by a factor of up to three in the simulation as compared to the observations. The ratio between simulated and observed hydroperoxyl radicals increases to  $\text{HO}_2^{\text{mod.}}/\text{HO}_2^{\text{obs.}} = 0.6$  on average, still separating into two regimes with a minor part of the dataset showing a ratio of  $\text{HO}_2^{\text{mod.}}/\text{HO}_2^{\text{obs.}} = 1.5$ , while most of the data shows a ratio of  $\text{HO}_2^{\text{mod.}}/\text{HO}_2^{\text{obs.}} = 0.6$ .

Since isoprene was not the predominant biogenic VOC during HUMPPA-COPEC-2010, contributing less than 10% to the total OH reactivity measured, the terpene mechanism MTM was added to the chemistry mechanism. Inclusion of terpene chem-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



istry helped to reproduce the OH reasonably in the simulation (Fig. 11). The ratio between simulated and observed HO<sub>2</sub> slightly increased compared to the isoprene chemistry reference run, but still separates into two regimes. At lower observed HO<sub>2</sub> levels, a ratio of up to HO<sub>2</sub><sup>mod.</sup>/HO<sub>2</sub><sup>obs.</sup> = 0.8 is reached for a minor fraction of the dataset.

5 Only about 30 % of the observed HO<sub>2</sub> concentration can be reproduced by the model on average, when the simulated total OH reactivity does not match the observed reactivity.

The underprediction of HO<sub>2</sub>, when OH reactivity is missing in the model while OH is reproduced accurately, indicates that the missing reactivity is an unaccounted source of HO<sub>2</sub>. Furthermore, the recycling reaction of NO and O<sub>3</sub> with the missing HO<sub>2</sub> has the potential under most conditions to compensate for the additional OH loss, which has not been considered yet. This preserves the good agreement of simulated and observed OH even when the model accounts for all the observed total OH reactivity. However, when high total OH reactivities (> 20 s<sup>-1</sup>) were observed, the enhanced HO<sub>2</sub> recycling is not sufficient to reproduce the observed OH. Additional mechanisms, e.g. direct OH recycling as discussed for isoprene-dominated, low-NO<sub>x</sub> environments, or an additional primary source of OH is required to explain the observed hydroxyl radical concentrations under these conditions.

To investigate the hypothesis that the OH reactivity missing in the model using the terpene mechanism acts as a source of HO<sub>2</sub>, additional reactivity towards OH was introduced by adding the surrogate molecule X which behaves chemically identically to α-pinene and its concentration was iteratively tuned for each dataset in such a way that the total OH reactivity in the model agreed with the observed OH reactivity. To match the observed total OH reactivity, up to 3 ppbV of X (typically a factor of 5 to 10 times the measured α-pinene) had to be added. Tuning the simulation to the observed total OH reactivity improves the HO<sub>2</sub> agreement slightly (HO<sub>2</sub><sup>mod.</sup>/HO<sub>2</sub><sup>obs.</sup> = 0.4), as shown in Fig. 12. The production of OH from the additional HO<sub>2</sub>, however, is not sufficient to compensate for the enhanced reactivity in the model, leading to underprediction of hydroxyl radicals in the simulation. The unaccounted OH reactivity in the simulation

might still be linked to the missing source of HO<sub>2</sub>, nevertheless, this shows that  $\alpha$ -pinene-like chemistry for the compound X does not suffice to provide this connection.

### 3.2.1 Significance and uncertainties

To quantify the significance of model-measurement discrepancies the uncertainties in both, observations and simulations, have to be considered. The uncertainties of HO<sub>x</sub> measurements are listed in Table 1. The uncertainties of simulated HO<sub>x</sub> related to the reaction rate coefficients were determined via Monte-Carlo simulations as described in Sect. 2.4.1. Monte-Carlo-Analysis using the terpene mechanism yields uncertainties of 14 % and 11 % for the simulated OH and HO<sub>2</sub>, respectively. Measured and simulated HO<sub>x</sub> concentrations including their 1 $\sigma$ -uncertainties are compared in Fig. 13. Modelled and observed OH within their uncertainties partly agree within a factor of two. Significant discrepancies by more than a factor of two are present for a minor part of the dataset (22 %). For HO<sub>2</sub> the underestimation by more than a factor of two in the model is significant for a large part of the dataset (52 %).

The uncertainties in observed trace gas species and photolysis frequencies used to constrain the model can also have a direct impact on the uncertainty of simulated HO<sub>x</sub>. Even less obviously these parameters which are kept fixed at the observed values within the simulation, influence the abundance of unmeasured trace gases and intermediate products, therefore also showing an indirect effect on the resulting calculation of HO<sub>x</sub> concentrations. Sensitivity studies were conducted by varying the observed trace gas concentrations separately by a factor between 0.5–10 to investigate the overall influence on the simulated HO<sub>x</sub> (Table 4) with the terpene mechanism.

Substances relevant for primary production of HO<sub>x</sub> show the highest impact in this sensitivity study. Doubling the O<sub>3</sub> or water vapour concentration leads to about 25 % increase in simulated OH and about 15 % enhanced HO<sub>2</sub>. An even larger effect (OH: +37 %, HO<sub>2</sub>: +26 %) is found for photolysis elevated by a factor of two. Constraining the model with twice the amount of NO observed causes 24 % more hydroxyl radicals in the simulation. The simulated HO<sub>2</sub> decreases by 23 %. Reducing the CO concentration

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by a factor of two leads to a 8 % increase in simulated OH and a 8 % decrease in simulated HO<sub>2</sub> while doubling the observed CO yields a 13 % decrease in OH and a 13 % increase in HO<sub>2</sub>. Doubling formaldehyde or H<sub>2</sub>O<sub>2</sub> only has a minor influence on the simulated HO<sub>x</sub>. Changing the deposition rates by a factor of two causes at most a 12 % change in HO<sub>x</sub>. None of these input parameter variations yield a change in OH and/or HO<sub>2</sub> of a factor of two or more. All photolysis frequencies multiplied by a factor of ten could cause such changes in simulated HO<sub>x</sub>. Unless the uncertainties of the species used to constrain the simulation (Table 1) exceed the variation intervals, these uncertainties individually can neither explain the discrepancies between observation and modelled HO<sub>2</sub>, nor cause a significant disagreement for OH.

Free running HCHO, i.e. not constraining the simulation to the HCHO observations, leads to enhanced OH and HO<sub>2</sub> concentrations (up to 30 %), which is for a minor part of the dataset sufficient to produce good agreement between measured and simulated HO<sub>2</sub>, while OH is overestimated in these simulations. Formaldehyde concentrations in the free running model are up to four times higher than observed levels. This suggests a formaldehyde sink not represented in the box model, possibly due to transport processes or uptake by plants (Lowe and Schmidt, 1983; Benning and Wahner, 1998). However, it was shown that changes in the deposition velocity have only a small impact on the simulated HO<sub>x</sub>. The overprediction of formaldehyde in this case is more likely due to inadequate representation of the HCHO sources in the simulation. Another sensitivity simulation by also constraining the model to estimated 2-methyl-3-buten-2-ol (MBO) concentrations was conducted. Unfortunately, MBO, which is often referred to as “the isoprene of coniferous forests” was not directly measured during HUMPPA-COPEC-2010, though PTR-MS measurements of isoprene were affected by an MBO interference, as described in Sect. 2.1. Previous measurements in this forest showed that MBO accounts only for 1–3 % of the total monoterpene emission rate (Tarvainen et al., 2005). In American pine forests it can be more important. However, the PTR-MS signal was used as an upper limit estimate of MBO for this simulation to investigate the influence on the simulated HO<sub>x</sub>. This MBO causes, on average, about 0.5 s<sup>-1</sup> additional



5 reactivity towards OH, similar to the reactivity by methane. Only about 20 % of the observed HO<sub>2</sub> can be explained by the model and the observed OH is underpredicted by 40–60 % in this simulation. Considering the results from the above mentioned sensitivity tests and significance analysis, the discrepancy between measured and simulated HO<sub>2</sub> is significant, thus, the chemical mechanism applied in the simulations needs further improvement to reproduce the HO<sub>2</sub> for the observed boreal forest environment.

### 3.2.2 Sources and sinks of HO<sub>x</sub>

10 The production rates of the hydroxyl radical in the box model using the terpene mechanism are shown in Fig. 14, with respect to their contribution to the total OH production derived from measurements. The dataset is again grouped by conditions of different radiation and total OH reactivity (see Table 3). The total production rate of hydroxyl radicals in the simulation accounts only for about 10 % of the observed production when the total OH reactivity is high ( $k'_{\text{OH}} > 15 \text{ s}^{-1}$ ). This is mainly due to the strong underestimation of HO<sub>2</sub> by the model under these conditions (see Sects. 3.1.1 and 15 3.1.2). Photolytic production accounts for one third of the OH formation in the simulation. A similar contribution is found by HO<sub>2</sub> recycling via reaction with NO and O<sub>3</sub>. Ozonolysis of BVOCs and a minor contribution by other species produced in the model account for the remaining OH production. At lower radiation ( $J_{\text{O}(1\text{D})} \leq 3 \times 10^{-6} \text{ s}^{-1}$ ), the ozonolysis of BVOCs becomes relatively more important and the recycling reaction of 20 HO<sub>2</sub> with O<sub>3</sub> more relevant in comparison to the reaction with NO. Under conditions of moderate total OH reactivity ( $k'_{\text{OH}} \leq 15 \text{ s}^{-1}$ ) and low radiation ( $J_{\text{O}(1\text{D})} \leq 3 \times 10^{-6} \text{ s}^{-1}$ ) about 33 % of the observed total OH production is represented by the box model simulation. Ozonolysis of observed biogenic VOCs accounts for on average 15 % of the total production rate. Even though the hydroxyl radical production rates due to recycling of HO<sub>2</sub> via the reactions with NO and O<sub>3</sub> are based on the underestimated HO<sub>2</sub> by the box model, they are relevant source terms contributing 13 % to the total OH production. 25 Photolytic sources and production due to other species play a minor role. By assuming observed rather than model calculated HO<sub>2</sub>, this contribution would increase to 30 %.

28592

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Almost one quarter of the OH during periods characterized by high radiation values ( $J_{O(^1D)} > 3 \times 10^{-6} \text{ s}^{-1}$ ) and moderate total OH reactivity ( $k'_{OH} \leq 15 \text{ s}^{-1}$ ) is produced by photolytic sources. Ozonolysis of BVOCs and other sources in the simulation contribute 4 % each to the total observed OH production. HO<sub>2</sub> recycling via NO dominates the OH production (37 %). Nonetheless, about 30 % of the total OH observed production is not represented in the simulation, due to the underestimation of HO<sub>2</sub> and the associated recycling pathways.

OH reactivity contributions calculated from individually measured compounds compared to directly measured OH reactivity revealed 58 % missing OH reactivity under “normal” boreal conditions and up to about 90 % under “stressed” boreal conditions (i.e. prolonged high temperature) during HUMPPA-COPEC-2010 (Nölscher et al., 2012). About 50 % of missing OH reactivity was reported from the same site during summer in 2008 by Sinha et al. (2010) when more typical boreal conditions at lower temperatures than in 2010 prevailed. The loss of OH in terms of OH reactivity in the simulation is presented in Fig. 15. Similar to the findings of another modeling study based on different measurements conducted at the field station SMEAR II (Mogensen et al., 2011), the inorganic contribution (CO, O<sub>3</sub>, H<sub>2</sub>, H<sub>2</sub>O<sub>2</sub>, NO, NO<sub>2</sub>, HO<sub>2</sub>, and HONO) is significant (15 %). A large sink for OH is due to organic compounds, most importantly monoterpenes (5 %), methane (4 %), and isoprene (2 %). In contrast to the study by Mogensen et al. (2011), second and higher order organic reaction products by the model contribute an additional 42 % to the total OH reactivity. On average, almost one third of the observed total OH reactivity remains unexplained. The large contribution by second and higher order organic reaction products in the model is mainly caused by aldehydes coming from monoterpene oxidation (~ 30 %) and secondary products of isoprene oxidation (MVK + MACR ~ 4 %). Primary biogenic emissions, e.g. unmeasured monoterpenes and sesquiterpenes, photooxidation products, as well as anthropogenic pollutants occasionally transported to the measurement site are likely candidates causing the “missing” fraction of total OH reactivity.

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HO<sub>2</sub> radicals are lost by reaction with other radicals as well as recycled towards OH by reaction with NO and O<sub>3</sub> (Fig. 16). The biggest contribution to the loss of HO<sub>2</sub> under all conditions is the yielding peroxyntic acid (HNO<sub>4</sub>). However, under standard conditions, HNO<sub>4</sub> is rather unstable and decomposes typically within a few seconds. This creates an equilibrium between HNO<sub>4</sub> and HO<sub>2</sub> + NO<sub>2</sub> which is strongly temperature dependent (Dentener et al., 2002). In the simulation the production and decomposition of HNO<sub>4</sub> occur at the same rate, thus HNO<sub>4</sub> acts as a reservoir species for the hydroperoxyl radicals.

At conditions of low radiation ( $J_{O(1D)} \leq 3 \times 10^{-6} \text{ s}^{-1}$ ) hydroperoxyl radicals reacting with RO<sub>2</sub> and HO<sub>x</sub> contribute to the total loss of HO<sub>2</sub> in a similar amount as the recycling reactions with NO and O<sub>3</sub>. This is still the case at conditions of enhanced observed OH reactivity and low radiation. The higher absolute HO<sub>2</sub> loss is caused by enhanced recycling as well as enhanced loss by radical-radical reactions.

At high photolysis frequencies ( $J_{O(1D)} > 3 \times 10^{-6} \text{ s}^{-1}$ ) both the total HO<sub>2</sub> production and loss increase by about a factor of two compared to nighttime. The HO<sub>2</sub> loss contribution from the recycling reaction with NO is predominant (41 %) at a total OH reactivity  $k'_{OH} \leq 15 \text{ s}^{-1}$ . This is followed by the reaction with RO<sub>2</sub> contributing about 12 %. Other reactions, e.g. HO<sub>2</sub> + NO<sub>3</sub> or HO<sub>2</sub> + HO<sub>x</sub>, contribute only a few percent. When the total OH reactivity is high,  $k'_{OH} > 15 \text{ s}^{-1}$ , the OH recycling reaction of HO<sub>2</sub> with NO becomes less important (23 %) in the simulation, whereas the reaction with RO<sub>2</sub> contributes about 20 % to the total HO<sub>2</sub> loss. In addition to HNO<sub>4</sub> decomposition, the reaction of OH with carbon monoxide and the reaction of RO<sub>2</sub> with NO are the main sources of HO<sub>2</sub> in the model. The photolysis of formaldehyde contributes less than 1 % to the total HO<sub>2</sub> production under all conditions. Ozonolysis of monoterpenes yields up to 8 % when photolysis is low but is rather unimportant at higher  $J$  values. OH reactions with O<sub>3</sub>, HCHO, and H<sub>2</sub>O<sub>2</sub> are minor contributors (typically in total about 4 %) in HO<sub>2</sub> production. Differences between conditions of moderate and high total OH reactivity show up in the reactions of OH + CO and RO<sub>2</sub> + NO, forming HO<sub>2</sub>. The absolute HO<sub>2</sub> production under conditions of enhanced radiation and high observed total OH reactivity is

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



smaller compared to the case with enhanced radiation and moderate total OH reactivity. Under these conditions, the simulation underestimates the observed HO<sub>2</sub> most, indicating that important species and HO<sub>2</sub> production pathways are missing in the chemical mechanism, very likely including unmeasured BVOCs and their oxidation products. Unfortunately, there were no direct observations of RO<sub>2</sub> during HUMPPA-COPEC-2010. Thus, we are lacking the possibility to constrain the model by RO<sub>2</sub> observations, which would improve our understanding of the underlying processes.

#### 4 Summary and conclusions

Our understanding of the chemistry of hydroxyl and hydroperoxyl radicals in forest environments is limited, and although we are making progress, the present work could not fully elucidate the observations using state-of-the-art models. In contrast to well-known OH recycling pathways via HO<sub>2</sub> by reaction with NO and O<sub>3</sub>, various mechanisms recycling OH directly in the absence of NO have been proposed and tested in different studies to improve the understanding of HO<sub>x</sub>-chemistry influenced by forest emissions (Paulot et al., 2009; Peeters et al., 2009; Crouse et al., 2011). Mao et al. (2012), addressing the impact of measurement artefacts in LIF measurements, showed reasonable agreement between model predictions and observations using their new “chemical removal method” for detection of OH in a California forest during the BEARPEX09<sup>3</sup> campaign. Griffith et al. (2013) found good model-to-measurement agreement during the CABINEX campaign, when colder temperatures led to lower observed mixing ratios of isoprene, its oxidation products MVK and MACR, and formaldehyde. As discussed in the paper by Griffith et al. (2013) it is not clear whether the instrument was sensitive to a similar interference as shown by Mao et al. (2012), although measurements conducted later in the same environment using an external scrubbing technique did not reveal a significant interference.

<sup>3</sup>BEARPEX09 = Biosphere Effects on AeRosols and Photochemistry EXperiment

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The HORUS instrument was operated for the first time using the IPI-LIF-FAGE technique to measure OH during HUMPPA-COPEC-2010. To identify possible measurement interferences an instrument intercomparison between the CIMS and HORUS-LIF instruments was conducted at the beginning of the campaign on the ground. The hydroxyl radical measurements by the two independent techniques show generally good agreement. During daytime, OH concentrations up to  $3.5 \times 10^6$  molec $\text{cm}^{-3}$  were observed above canopy, a factor of two to three times higher than the measurements on ground. During nighttime these differences in OH vanished, but still during some nights both instruments, CIMS (on ground) and IPI-LIF-FAGE (above canopy), detected significant amounts of OH that cannot be fully explained by known production rates calculated from observations. The missing OH source during nighttime of the order of  $1 \times 10^6$  molec $\text{cm}^{-3} \text{s}^{-1}$  could partly be related to NO<sub>3</sub>. The NO<sub>3</sub> mixing ratio during HUMPPA-COPEC-2010 was always below the lower limit of detection of the CRD instrument (Rinne et al., 2012). However, it could nevertheless be produced at a rate of up to  $1 \times 10^6$  molec $\text{cm}^{-3} \text{s}^{-1}$  from the reaction of NO<sub>2</sub> with O<sub>3</sub>. RO<sub>2</sub> radicals from NO<sub>3</sub>-initiated VOC oxidation potentially undergo isomerization reactions or react with HO<sub>2</sub> as well, forming OH. Thus, NO<sub>3</sub> might, at least under some conditions, contribute to the missing OH source during nighttime. Ozonolysis of unmeasured VOCs is also likely to contribute.

Comparison of the calculated total OH production rate with the total loss rate calculated from OH concentrations and total OH reactivity measurements showed that the known OH sources are almost sufficient to close the budget above the canopy. Detailed analysis of the radical production, loss, and recycling pathways revealed that OH recycling in the observed boreal forest environment occurs mainly by recycling via HO<sub>2</sub> under conditions of high radiation and moderate observed total OH reactivity. In addition, recycling mechanisms of OH, not via reaction of HO<sub>2</sub> with NO/O<sub>3</sub>, are likely under conditions of enhanced total OH reactivity.

The role of potentially undetected VOCs and oxidation products in missing OH reactivity can not be understood from observations alone. Hence, box model simulations

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



have been considered for further investigation. The chemistry of the condensed isoprene mechanism (“Mainz Isoprene Mechanism”) is shown to be deficient for the observed monoterpene-dominated boreal forest environment. Isoprene levels of typically less than 200 pptV contributed at most 10 % to the total observed OH reactivity thus leading to an overprediction of OH due to the missing sinks by a factor of up to 3 and HO<sub>2</sub> being significantly underpredicted ( $\text{HO}_2^{\text{mod.}}/\text{HO}_2^{\text{obs.}} = 0.3$ ). Inclusion of the Mainz Terpene Mechanism (MTM) to account for the reactivity towards OH due to observed terpene species and their oxidation products leads to much better agreement between observed and simulated OH concentrations. However, this is due to two compensating effects. On average about one third of the observed total OH reactivity is not reproduced in the simulation, thus leading to underestimation of the total sink of OH. On the other hand, the production of OH is significantly underestimated due to the underpredicted HO<sub>2</sub> available for recycling by reaction with NO and O<sub>3</sub>, which was shown to be an important source of OH by direct calculation from observations. Tuning the simulation on the observed total OH reactivity by adding an unknown compound which behaves like  $\alpha$ -pinene produced a somewhat better  $\text{HO}_2^{\text{mod.}}/\text{HO}_2^{\text{obs.}}$  agreement. Nevertheless, the additional HO<sub>2</sub> being recycled towards OH was not sufficient to compensate for the resulting enhanced OH loss. The biggest model-observation discrepancies for HO<sub>2</sub> occurred when “missing” OH reactivity was highest. Therefore, this provides evidence that the “missing” OH reactivity in the simulation is a source of HO<sub>2</sub>. However, a single terpene following the chemical mechanism of  $\alpha$ -pinene as described in the preliminary terpene mechanism cannot account for this alone.

The OH production in this forest environment seems to be understood reasonably well under moderate total OH reactivity conditions (Table 3). The loss of OH cannot be fully explained, even through inclusion of higher order oxidation products by application of a box model. Similarly to the study of Kim et al. (2013), HO<sub>2</sub> is underestimated in simulations, indicating missing BVOCs, which could also account for the missing reactivity in the model. Additional recycling processes of OH, not via HO<sub>2</sub> + NO/O<sub>3</sub> are indicated under conditions of high observed total OH reactivity by detailed analysis of the HO<sub>x</sub>



## References

- Benning, L. and Wahner, A.: Measurements of atmospheric formaldehyde (HCHO) and acetaldehyde (CH<sub>3</sub>CHO) during POPCORN 1994 using 2.4-DNPH coated silica cartridges, *J. Atmos. Chem.*, **31**, 105–117, doi:10.1023/A:1005884116406, 1998. 28591
- 5 Bohn, B., Corlett, G. K., Gillmann, M., Sanghavi, S., Stange, G., Tensing, E., Vrekoussis, M., Bloss, W. J., Clapp, L. J., Kortner, M., Dorn, H.-P., Monks, P. S., Platt, U., Plass-Dülmer, C., Mihalopoulos, N., Heard, D. E., Clemmshaw, K. C., Meixner, F. X., Prevot, A. S. H., and Schmitt, R.: Photolysis frequency measurement techniques: results of a comparison within the ACCENT project, *Atmos. Chem. Phys.*, **8**, 5373–5391, doi:10.5194/acp-8-5373-2008, 2008. 28577
- 10 Brune, W. H., Mao, J., and Ren, X.: OH and HO<sub>2</sub> Measurements in Blodgett Forest, CA during BEARPEX 2009, abstract A51G-02 presented at 2010 Fall Meeting, AGU, San Francisco, Calif., 13–17 December, 2010. 28571
- Crouse, J. D., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: Peroxy radical isomerization in the oxidation of isoprene, *Phys. Chem. Chem. Phys.*, **13**, 13607–13613, doi:10.1039/C1CP21330J, 2011. 28595
- 15 Crowley, J. N., Ammann, M., Cox, R. A., Hynes, R. G., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V – heterogeneous reactions on solid substrates, *Atmos. Chem. Phys.*, **10**, 9059–9223, doi:10.5194/acp-10-9059-2010, 2010. 28568
- 20 Dentener, F., Williams, J., and Metzger, S.: Aqueous phase reaction of HNO<sub>4</sub>: the impact on tropospheric chemistry, *J. Atmos. Chem.*, **41**, 109–133, doi:10.1023/A:1014233910126, 2002. 28594
- Dillon, T.: Comment on “Detection of HO<sub>2</sub> by laser-induced fluorescence: calibration and interferences from RO<sub>2</sub> radicals” by: Fuchs, H. et al., *Atmos. Meas. Tech. Discuss.*, **4**, 1255, 2011. 28572
- 25 Dillon, T. J. and Crowley, J. N.: Direct detection of OH formation in the reactions of HO<sub>2</sub> with CH<sub>3</sub>C(O)O<sub>2</sub> and other substituted peroxy radicals, *Atmos. Chem. Phys.*, **8**, 4877–4889, doi:10.5194/acp-8-4877-2008, 2008. 28586
- 30 Evans, M., Shallcross, D., Law, K., Wild, J., Simmonds, P., Spain, T., Berrisford, P., Methven, J., Lewis, A., McQuaid, J., Pilling, M., Bandy, B., Penkett, S., and Pyle, J.: Evaluation of a Lagrangian box model using field measurements from EASE (Eastern Atlantic Summer Exper-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





iment) 1996, *Atmos. Environ.*, 34, 3843–3863, doi:10.1016/S1352-2310(00)00184-9, 2000. 28611

Faloona, I., Tan, D., Brune, W. H., Jaegle, L., Jacob, D. J., Kondo, Y., Koike, M., Chatfield, R., Poeschel, R., Ferry, G., Sachse, G., Vay, S., Anderson, B., Hannon, J., and Fuelberg, H.: Observations of HO<sub>x</sub> and its relationship with NO<sub>x</sub> in the upper troposphere during SONEX, *J. Geophys. Res.-Atmos.*, 105, 3771–3783, doi:10.1029/1999JD900914, 2000. 28571

Faloona, I., Tan, D., Leshner, R., Hazen, N., Frame, C., Simpas, J., Harder, H., Martinez, M., Di Carlo, P., Ren, X., and Brune, W.: A laser-induced fluorescence instrument for detecting tropospheric OH and HO<sub>2</sub>: characteristics and calibration, *J. Atmos. Chem.*, 47, 139–167, doi:10.1023/B:JOCH.0000021036.53185.0e, 2004. 28569

FAO: Global forest resources assessment 2010, FAO forestry paper, FAO, Rome, available at: <http://www.fao.org/docrep/013/i1757e/i1757e.pdf> (last access: 24 october 2013), 2010. 28564

Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A. B., Hewitt, C. N., Lamb, B., Liu, S., Trainer, M., Westberg, H., and Zimmerman, P.: Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry, *Global Biogeochem. Cy.*, 6, 389–430, doi:10.1029/92GB02125, 1992. 28564

Fuchs, H., Bohn, B., Hofzumahaus, A., Holland, F., Lu, K. D., Nehr, S., Rohrer, F., and Wahner, A.: Detection of HO<sub>2</sub> by laser-induced fluorescence: calibration and interferences from RO<sub>2</sub> radicals, *Atmos. Meas. Tech.*, 4, 1209–1225, doi:10.5194/amt-4-1209-2011, 2011. 28572, 28573, 28583

Fuchs, H., Hofzumahaus, A., Rohrer, F., Bohn, B., Brauers, T., Dorn, H.-P., Haseler, R., Holland, F., Kaminski, M., Li, X., Lu, K., Nehr, S., Tillmann, R., Wegener, R., and Wahner, A.: Experimental evidence for efficient hydroxyl radical regeneration in isoprene oxidation, *Nat. Geosci.*, doi:10.1038/ngeo1964, 2013. 28587

Goldan, P. D., Kuster, W. C., and Fehsenfeld, F. C.: Nonmethane hydrocarbon measurements during the Tropospheric OH Photochemistry Experiment, *J. Geophys. Res.-Atmos.*, 102, 6315–6324, doi:10.1029/96JD01868, 1997. 28568

Griffith, S. M., Hansen, R. F., Dusanter, S., Stevens, P. S., Alaghmand, M., Bertman, S. B., Carroll, M. A., Erickson, M., Galloway, M., Grossberg, N., Hottle, J., Hou, J., Jobson, B. T., Kammrath, A., Keutsch, F. N., Lefer, B. L., Mielke, L. H., O'Brien, A., Shepson, P. B., Thurlow, M., Wallace, W., Zhang, N., and Zhou, X. L.: OH and HO<sub>2</sub> radical chemistry during PROPHET 2008 and CABINEX 2009 – Part 1: Measurements and model comparison, *At-*

## HO<sub>x</sub> photochemistry in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mos. Chem. Phys., 13, 5403–5423, doi:10.5194/acp-13-5403-2013, 2013. 28565, 28573, 28595

Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdaun, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *J. Geophys. Res.-Atmos.*, 100, 8873–8892, doi:10.1029/94JD02950, 1995. 28564

Hall, B. D. and Claiborn, C. S.: Measurements of the dry deposition of peroxides to a Canadian boreal forest, *J. Geophys. Res.-Atmos.*, 102, 29343–29353, doi:10.1029/97JD01113, 1997. 28574

Hari, P. and Kulmala, M.: Station for Measuring Ecosystem–Atmosphere Relations (SMEAR II), *Boreal Environ. Res.*, 10, 315–322, 2005. 28567

Hasson, A. S., Tyndall, G. S., and Orlando, J. J.: A product yield study of the reaction of HO<sub>2</sub> radicals with ethyl peroxy (C<sub>2</sub>H<sub>5</sub>O<sub>2</sub>), acetyl peroxy (CH<sub>3</sub>C(O)O<sub>2</sub>), and acetonyl peroxy (CH<sub>3</sub>C(O)CH<sub>2</sub>O<sub>2</sub>) radicals, *J. Phys. Chem. A*, 108, 5979–5989, doi:10.1021/jp048873t, 2004. 28586

Heard, D. E. and Pilling, M. J.: Measurement of OH and HO<sub>2</sub> in the troposphere, *Chem. Rev.*, 103, 5163–5198, doi:10.1021/cr020522s, 2003. 28572

Hofzumahaus, A., Aschmutat, U., Hessling, M., Holland, F., and Ehhalt, D. H.: The measurement of tropospheric OH radicals by laser-induced fluorescence spectroscopy during the POPCORN Field Campaign, *Geophys. Res. Lett.*, 23, 2541–2544, doi:10.1029/96GL02205, 1996. 28570

Holland, F., Aschmutat, U., Hessling, M., Hofzumahaus, A., and Ehhalt, D.: Highly time resolved measurements of OH during POPCORN using laser-induced fluorescence spectroscopy, *J. Atmos. Chem.*, 31, 205–225, doi:10.1023/A:1005868520002, 1998. 28570

Holland, F., Hofzumahaus, A., Schäfer, J., Kraus, A., and Pätz, H.-W.: Measurements of OH and HO<sub>2</sub> radical concentrations and photolysis frequencies during BERLIOZ, *J. Geophys. Res.-Atmos.*, 108, 2–23, doi:10.1029/2001JD001393, 2003. 28572

Hosaynali Beygi, Z., Fischer, H., Harder, H. D., Martinez, M., Sander, R., Williams, J., Brookes, D. M., Monks, P. S., and Lelieveld, J.: Oxidation photochemistry in the Southern Atlantic boundary layer: unexpected deviations of photochemical steady state, *Atmos. Chem. Phys.*, 11, 8497–8513, doi:10.5194/acp-11-8497-2011, 2011. 28568

Jaegle, L., Jacob, D. J., Brune, W. H., Faloon, I., Tan, D., Heikes, B. G., Kondo, Y., Sachse, G. W., Anderson, B., Gregory, G. L., Singh, H. B., Poeschel, R., Ferry, G.,

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Blake, D. R., and Shetter, R. E.: Photochemistry of HO<sub>x</sub> in the upper troposphere at northern midlatitudes, *J. Geophys. Res.-Atmos.*, 105, 3877–3892, doi:10.1029/1999JD901016, 2000. 28564

Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel of the CH<sub>3</sub>C(O)O<sub>2</sub> + HO<sub>2</sub> reaction in the gas phase, *Phys. Chem. Chem. Phys.*, 9, 3149–3162, doi:10.1039/B702757E, 2007. 28586

Junkermann, W., Platt, U., and Volz-Thomas, A.: A photoelectric detector for the measurement of photolysis frequencies of ozone and other atmospheric molecules, *J. Atmos. Chem.*, Kluwer Academic Publishers, 8, 203–227, 1989. 28568

Karl, T., Hansel, A., Cappellin, L., Kaser, L., Herdinger-Blatt, I., and Jud, W.: Selective measurements of isoprene and 2-methyl-3-buten-2-ol based on NO<sup>+</sup> ionization mass spectrometry, *Atmos. Chem. Phys.*, 12, 11877–11884, doi:10.5194/acp-12-11877-2012, 2012. 28568

Kesselmeier, J. and Staudt, M.: Biogenic volatile organic compounds (VOC): an overview on emission, physiology and ecology, *J. Atmos. Chem.*, 33, 23–88, doi:10.1023/A:1006127516791, 1999. 28564

Kim, S., Wolfe, G. M., Mauldin, L., Cantrell, C., Guenther, A., Karl, T., Turnipseed, A., Greenberg, J., Hall, S. R., Ullmann, K., Apel, E., Hornbrook, R., Kajii, Y., Nakashima, Y., Keutsch, F. N., DiGangi, J. P., Henry, S. B., Kaser, L., Schnitzhofer, R., Graus, M., Hansel, A., Zheng, W., and Flocke, F. F.: Evaluation of HO<sub>x</sub> sources and cycling using measurement-constrained model calculations in a 2-methyl-3-butene-2-ol (MBO) and monoterpene (MT) dominated ecosystem, *Atmos. Chem. Phys.*, 13, 2031–2044, doi:10.5194/acp-13-2031-2013, 2013. 28565, 28566, 28597

Kleffmann, J., Heland, J., Kurtenbach, R., Lorzer, J., and Wiesen, P.: A new instrument (LOPAP) for the detection of nitrous acid (HONO), *Environ. Sci. Pollut. R.*, 9, 48–54, 2002. 28568

Klippel, T., Fischer, H., Bozem, H., Lawrence, M. G., Butler, T., Jöckel, P., Tost, H., Martinez, M., Harder, H., Regelin, E., Sander, R., Schiller, C. L., Stickler, A., and Lelieveld, J.: Distribution of hydrogen peroxide and formaldehyde over Central Europe during the HOOVER project, *Atmos. Chem. Phys.*, 11, 4391–4410, doi:10.5194/acp-11-4391-2011, 2011. 28568

Kubistin, D.: OH- und HO<sub>2</sub>-Radikale über dem tropischen Regenwald, Ph. D. thesis, Johannes Gutenberg-Universität, Mainz, Germany, 2009. 28571

Kubistin, D., Harder, H., Martinez, M., Rudolf, M., Sander, R., Bozem, H., Eerdekens, G., Fischer, H., Gurk, C., Klüpfel, T., Königstedt, R., Parchatka, U., Schiller, C. L., Stickler, A., Taraborrelli, D., Williams, J., and Lelieveld, J.: Hydroxyl radicals in the tropical troposphere

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



over the Suriname rainforest: comparison of measurements with the box model MECCA, Atmos. Chem. Phys., 10, 9705–9728, doi:10.5194/acp-10-9705-2010, 2010. 28565

Kulmala, M., Hämeri, K., Aalto, P. P., Mäkelä, J. M., Pirjola, L., Nilsson, E. D., Buzorius, G., Rannik, U., Maso, M. D., Seidl, W., Hoffman, T., Janson, R., Hansson, H.-C., Viisanen, Y., Laaksonen, A., and O'Dowd, C. D.: Overview of the international project on biogenic aerosol formation in the boreal forest (BIOFOR), Tellus B, 53, 324–343, doi:10.1034/j.1600-0889.2001.530402.x, 2001. 28567

Leighton, P.: Photochemistry of Air Pollution, Academic Press, New York and London, 1961. 28583

Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a tropical forest, Nature, 452, 737–740, 2008. 28565

Levy II, H.: Normal atmosphere: large radical and formaldehyde concentrations predicted, Science, 173, 141–143, 1971. 28564

Lowe, D. C. and Schmidt, U.: Formaldehyde (HCHO) measurements in the nonurban atmosphere, J. Geophys. Res.-Oceans, 88, 10844–10858, doi:10.1029/JC088iC15p10844, 1983. 28591

Lu, K. D., Rohrer, F., Holland, F., Fuchs, H., Bohn, B., Brauers, T., Chang, C. C., Häseler, R., Hu, M., Kita, K., Kondo, Y., Li, X., Lou, S. R., Nehr, S., Shao, M., Zeng, L. M., Wahner, A., Zhang, Y. H., and Hofzumahaus, A.: Observation and modelling of OH and HO<sub>2</sub> concentrations in the Pearl River Delta 2006: a missing OH source in a VOC rich atmosphere, Atmos. Chem. Phys., 12, 1541–1569, doi:10.5194/acp-12-1541-2012, 2012. 28573

Mao, J., Ren, X., Zhang, L., Van Duin, D. M., Cohen, R. C., Park, J.-H., Goldstein, A. H., Paulot, F., Beaver, M. R., Crouse, J. D., Wennberg, P. O., DiGangi, J. P., Henry, S. B., Keutsch, F. N., Park, C., Schade, G. W., Wolfe, G. M., Thornton, J. A., and Brune, W. H.: Insights into hydroxyl measurements and atmospheric oxidation in a California forest, Atmos. Chem. Phys., 12, 8009–8020, doi:10.5194/acp-12-8009-2012, 2012. 28565, 28571, 28573, 28595

Martinez, M., Harder, H., Kovacs, T. A., Simpas, J. B., Bassis, J., Leshner, R., Brune, W. H., Frost, G. J., Williams, E. J., Stroud, C. A., Jobson, B. T., Roberts, J. M., Hall, S. R., Shetter, R. E., Wert, B., Fried, A., Alicke, B., Stutz, J., Young, V. L., White, A. B., and Zamora, R. J.: OH and HO<sub>2</sub> concentrations, sources, and loss rates during the South-

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ern Oxidants Study in Nashville, Tennessee, summer 1999, *J. Geophys. Res.*, 108, 4617, doi:10.1029/2003JD003551, 2003. 28564

Martinez, M., Harder, H., Ren, X., Leshner, R. L., and Brune, W. H.: Measuring atmospheric naphthalene with laser-induced fluorescence, *Atmos. Chem. Phys.*, 4, 563–569, doi:10.5194/acp-4-563-2004, 2004. 28571

Martinez, M., Harder, H., Kubistin, D., Rudolf, M., Bozem, H., Eerdeken, G., Fischer, H., Klüpfel, T., Gurk, C., Königstedt, R., Parchatka, U., Schiller, C. L., Stickler, A., Williams, J., and Lelieveld, J.: Hydroxyl radicals in the tropical troposphere over the Suriname rainforest: airborne measurements, *Atmos. Chem. Phys.*, 10, 3759–3773, doi:10.5194/acp-10-3759-2010, 2010. 28569, 28570

Mogensen, D., Smolander, S., Sogachev, A., Zhou, L., Sinha, V., Guenther, A., Williams, J., Nieminen, T., Kajos, M. K., Rinne, J., Kulmala, M., and Boy, M.: Modelling atmospheric OH-reactivity in a boreal forest ecosystem, *Atmos. Chem. Phys.*, 11, 9709–9719, doi:10.5194/acp-11-9709-2011, 2011. 28566, 28593

Nölscher, A. C., Williams, J., Sinha, V., Custer, T., Song, W., Johnson, A. M., Axinte, R., Bozem, H., Fischer, H., Pouvesle, N., Phillips, G., Crowley, J. N., Rantala, P., Rinne, J., Kulmala, M., Gonzales, D., Valverde-Canossa, J., Vogel, A., Hoffmann, T., Ouwersloot, H. G., Vilà-Guerau de Arellano, J., and Lelieveld, J.: Summertime total OH reactivity measurements from boreal forest during HUMPPA-COPEC 2010, *Atmos. Chem. Phys.*, 12, 8257–8270, doi:10.5194/acp-12-8257-2012, 2012. 28569, 28581, 28593

Novelli, A., Hens, K., Tatum Ernest, C., Trawny, K., Rudolf, M., Kubistin, D., Hosaynali Beygi, Z., Fischer, H., Williams, J., and Paasonen, P., Sipilä, M., Keronen, P., Petäjä, T., Elste, T., Plass-Duelmer, C., Vereecken, L., Martinez, M., Lelieveld, J., and Harder, H.: The role of stabilised Criegee intermediate in gas phase H<sub>2</sub>SO<sub>4</sub> formation, abstract A24D-04 presented at 2012 Fall Meeting, AGU, San Francisco, Calif., 3–7 December, 2012. 28571, 28579

Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kürten, A., St. Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325, 730–733, doi:10.1126/science.1172910, 2009. 28595

Peeters, J., Vereecken, L., and Fantechi, G.: The detailed mechanism of the OH-initiated atmospheric oxidation of  $\alpha$ -pinene: a theoretical study, *Phys. Chem. Chem. Phys.*, 3, 5489–5504, doi:10.1039/B106555F, 2001. 28587

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Peeters, J., Nguyen, T. L., and Vereecken, L.: HO<sub>x</sub> radical regeneration in the oxidation of isoprene, *Phys. Chem. Chem. Phys.*, 11, 5935–5939, doi:10.1039/B908511D, 2009. 28587, 28595, 28612
- 5 Petäjä, T., Mauldin III, R. L., Petäjä, T., Mauldin, III, R. L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, *Atmos. Chem. Phys.*, 9, 7435–7448, doi:10.5194/acp-9-7435-2009, 2009. 28569, 28578, 28618, 28619
- 10 Pöschl, U., von Kuhlmann, R., Poisson, N., and Crutzen, P. J.: Development and intercomparison of condensed isoprene oxidation mechanisms for global atmospheric modeling, *J. Atmos. Chem.*, 37, 29–52, doi:10.1023/A:1006391009798, 2000. 28566, 28576
- 15 Pugh, T. A. M., MacKenzie, A. R., Hewitt, C. N., Langford, B., Edwards, P. M., Furneaux, K. L., Heard, D. E., Hopkins, J. R., Jones, C. E., Karunaharan, A., Lee, J., Mills, G., Misztal, P., Moller, S., Monks, P. S., and Whalley, L. K.: Simulating atmospheric composition over a South-East Asian tropical rainforest: performance of a chemistry box model, *Atmos. Chem. Phys.*, 10, 279–298, doi:10.5194/acp-10-279-2010, 2010. 28565
- 20 Ren, X., Harder, H., Martinez, M., Faloona, I., Tan, D., Leshner, R., Di Carlo, P., Simpas, J., and Brune, W.: Interference testing for atmospheric HO<sub>x</sub> measurements by laser-induced fluorescence, *J. Atmos. Chem.*, 47, 169–190, doi:10.1023/B:JOCH.0000021037.46866.81, 2004. 28571
- 25 Ren, X., Olson, J. R., Crawford, J. H., Brune, W. H., Mao, J., Long, R. B., Chen, Z., Chen, G., Avery, M. A., Sachse, G. W., Barrick, J. D., Diskin, G. S., Huey, L. G., Fried, A., Cohen, R. C., Heikes, B., Wennberg, P. O., Singh, H. B., Blake, D. R., and Shetter, R. E.: HO<sub>x</sub> chemistry during INTEX-A 2004: Observation, model calculation, and comparison with previous studies, *J. Geophys. Res.-Atmos.*, 113, 2156–2202, doi:10.1029/2007JD009166, 2008. 28564
- 30 Rinne, J., Markkanen, T., Ruuskanen, T. M., Petäjä, T., Keronen, P., Tang, M.J., Crowley, J. N., Rannik, Ü., and Vesala, T.: Effect of chemical degradation on fluxes of reactive compounds – a study with a stochastic Lagrangian transport model, *Atmos. Chem. Phys.*, 12, 4843–4854, doi:10.5194/acp-12-4843-2012, 2012. 28580, 28596
- Rohrer, F. and Berresheim, H.: Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet radiation, *Nature*, 442, 184–187, doi:10.1038/nature04924, 2006. 28580
- Sander, R., Baumgaertner, A., Gromov, S., Harder, H., Jöckel, P., Kerkweg, A., Kubistin, D., Regelin, E., Riede, H., Sandu, A., Taraborrelli, D., Tost, H., and Xie, Z.-Q.: The at-

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ospheric chemistry box model CAABA/MECCA-3.0, *Geosci. Model Dev.*, 4, 373–380, doi:10.5194/gmd-4-373-2011, 2011. 28576, 28578

Sandu, A. and Sander, R.: Technical note: Simulating chemical systems in Fortran90 and Matlab with the Kinetic PreProcessor KPP-2.1, *Atmos. Chem. Phys.*, 6, 187–195, doi:10.5194/acp-6-187-2006, 2006. 28577

Schlosser, E., Brauers, T., Dorn, H.-P., Fuchs, H., Häseler, R., Hofzumahaus, A., Holland, F., Wahner, A., Kanaya, Y., Kajii, Y., Miyamoto, K., Nishida, S., Watanabe, K., Yoshino, A., Kubistin, D., Martinez, M., Rudolf, M., Harder, H., Berresheim, H., Elste, T., Plass-Dülmer, C., Stange, G., and Schurath, U.: Technical Note: Formal blind intercomparison of OH measurements: results from the international campaign HO<sub>x</sub>Comp, *Atmos. Chem. Phys.*, 9, 7923–7948, doi:10.5194/acp-9-7923-2009, 2009. 28579

Schuster, G., Labazan, I., and Crowley, J. N.: A cavity ring down/cavity enhanced absorption device for measurement of ambient NO<sub>3</sub> and N<sub>2</sub>O<sub>5</sub>, *Atmos. Meas. Tech.*, 2, 1–13, doi:10.5194/amt-2-1-2009, 2009. 28568

Sinha, V., Williams, J., Crowley, J. N., and Lelieveld, J.: The Comparative Reactivity Method – a new tool to measure total OH Reactivity in ambient air, *Atmos. Chem. Phys.*, 8, 2213–2227, doi:10.5194/acp-8-2213-2008, 2008. 28566, 28569

Sinha, V., Williams, J., Lelieveld, J., Ruuskanen, T., Kajos, M., Patokoski, J., Hellen, H., Hakola, H., Mogensen, D., Boy, M., Rinne, J., and Kulmala, M.: OH reactivity measurements within a boreal forest: evidence for unknown reactive emissions, *Environ. Sci. Technol.*, 44, 6614–6620, doi:10.1021/es101780b, 2010. 28593

Stone, D., Evans, M. J., Edwards, P. M., Commane, R., Ingham, T., Rickard, A. R., Brookes, D. M., Hopkins, J., Leigh, R. J., Lewis, A. C., Monks, P. S., Oram, D., Reeves, C. E., Stewart, D., and Heard, D. E.: Isoprene oxidation mechanisms: measurements and modelling of OH and HO<sub>2</sub> over a South-East Asian tropical rainforest during the OP3 field campaign, *Atmos. Chem. Phys.*, 11, 6749–6771, doi:10.5194/acp-11-6749-2011, 2011. 28565

Tan, D., Faloon, I., Simpas, J., Brune, W., Shepson, P., Couch, T., Sumner, A., Carroll, M., Thornberry, T., Apel, E., Riemer, D., and Stockwell, W.: HO<sub>x</sub> budgets in a deciduous forest: results from the PROPHET summer 1998 campaign, *J. Geophys. Res.*, 106, 24407–24427, doi:10.1029/2001JD900016, 2001. 28565

Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric

modelling, *Atmos. Chem. Phys.*, 9, 2751–2777, doi:10.5194/acp-9-2751-2009, 2009. 28566, 28576

Taraborrelli, D., Lawrence, M. G., Crowley, J. N., Dillon, T. J., Gromov, S., Grosz, C. B. M., Vereecken, L., and Lelieveld, J.: Hydroxyl radical buffered by isoprene oxidation over tropical forests, *Nat. Geosci.*, 5, 190–193, doi:10.1038/ngeo1405, 2012. 28566, 28576, 28588

Tarvainen, V., Hakola, H., Hellén, H., Bäck, J., Hari, P., and Kulmala, M.: Temperature and light dependence of the VOC emissions of Scots pine, *Atmos. Chem. Phys.*, 5, 989–998, doi:10.5194/acp-5-989-2005, 2005. 28591

Tatum Ernest, C., Novelli, A., Trawny, K., Hens, K., Rudolf, M., Elste, T., Werner, A., Englert, J., Gilge, S., Plass-Dülmer, C., Lelieveld, J., Martinez, M., and Harder, H.: OH and HO<sub>2</sub> Measurements in a Rural Mid-Latitude Environment during HOPE2012, paper presented at the Atmospheric Chemical Mechanisms Conference, Davis, CA, 10–13 December, available at: <https://sites.google.com/site/atmoschemmech2012/> (last access: 24 October 2013), 2012. 28573

Thornton, J. A., Wooldridge, P. J., Cohen, R. C., Martinez, M., Harder, H., Brune, W. H., Williams, E. J., Roberts, J. M., Fehsenfeld, F. C., Hall, S. R., Shetter, R. E., Wert, B. P., and Fried, A.: Ozone production rates as a function of NO<sub>x</sub> abundances and HO<sub>x</sub> production rates in the Nashville urban plume, *J. Geophys. Res.-Atmos.*, 107, 7–17, doi:10.1029/2001JD000932, 2002. 28586

Vereecken, L., Muller, J.-F., and Peeters, J.: Low-volatility poly-oxygenates in the OH-initiated atmospheric oxidation of  $\alpha$ -pinene: impact of non-traditional peroxy radical chemistry, *Phys. Chem. Chem. Phys.*, 9, 5241–5248, doi:10.1039/B708023A, 2007. 28587

Vereecken, L., Harder, H., and Novelli, A.: The reaction of Criegee intermediates with NO, RO<sub>2</sub>, and SO<sub>2</sub>, and their fate in the atmosphere, *Phys. Chem. Chem. Phys.*, 14, 14682–14695, doi:10.1039/C2CP42300F, 2012. 28587

Vesala, T., Haataja, J., Aalto, P., Altimir, N., Buzorius, G., Garam, E., Hämeri, K., Ilvesniemi, H., Jokinen, V., Keronen, P., Lahti, T., Markkanen, T., Mäkelä, J. M., E., N., Palmroth, S., Palva, L., Pohja, T., Pumpanen, J., Rannik, U., Siivola, E., Ylitalo, H., Hari, P., and Kulmala, M.: Long-term field measurements of atmosphere-surface interaction in boreal forest combining forest ecology, micrometeorology, aerosol physics and atmospheric chemistry, *Trends in Heat, Mass Moment. Transf.*, 4, 17–35, 1998. 28567

Wennberg, P., Cohen, R., Hazen, N., Lapson, L., Allen, N., Hanisco, T., Oliver, J., Lanham, N., Demusz, J., and Anderson, J.: An aircraft-borne, laser-induced fluorescence instrument for

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the in situ detection of hydroxyl and hydroperoxyl radicals, Rev. Sci. Instrum., 65, 1858–1876, 1994. 28569

Whalley, L. K., Edwards, P. M., Furneaux, K. L., Goddard, A., Ingham, T., Evans, M. J., Stone, D., Hopkins, J. R., Jones, C. E., Karunaharan, A., Lee, J. D., Lewis, A. C., Monks, P. S., Moller, S. J., and Heard, D. E.: Quantifying the magnitude of a missing hydroxyl radical source in a tropical rainforest, Atmos. Chem. Phys., 11, 7223–7233, doi:10.5194/acp-11-7223-2011, 2011. 28565

Whalley, L. K., Blitz, M. A., Desservettaz, M., Seakins, P. W., and Heard, D. E.: Reporting the sensitivity of Laser Induced Fluorescence instruments used for HO<sub>2</sub> detection to an interference from RO<sub>2</sub> radicals and introducing a novel approach that enables HO<sub>2</sub> and certain RO<sub>2</sub> types to be selectively measured, Atmos. Meas. Tech. Discuss., 6, 6249–6292, doi:10.5194/amtd-6-6249-2013, 2013. 28573, 28584

White, J.: Long optical paths of large aperture, J. Opt. Soc. Am., 32, 285–288, 1942. 28569

Williams, J.: Organic trace gases in the atmosphere: an overview, Environ. Chem., 1, 125–136, doi:10.1071/EN04057, 2004. 28564

Williams, J., Pöschl, U., Crutzen, P., Hansel, A., Holzinger, R., Warneke, C., Lindinger, W., and Lelieveld, J.: An atmospheric chemistry interpretation of mass scans obtained from a proton transfer mass spectrometer flown over the Tropical Rainforest of Surinam, J. Atmos. Chem., 38, 133–166, doi:10.1023/A:1006322701523, 2001. 28568

Williams, J., Crowley, J., Fischer, H., Harder, H., Martinez, M., Petäjä, T., Rinne, J., Bäck, J., Boy, M., Dal Maso, M., Hakala, J., Kajos, M., Keronen, P., Rantala, P., Aalto, J., Aaltonen, H., Paatero, J., Vesala, T., Hakola, H., Levula, J., Pohja, T., Herrmann, F., Auld, J., Mesarchaki, E., Song, W., Yassaa, N., Nölscher, A., Johnson, A. M., Custer, T., Sinha, V., Thieser, J., Pouvesle, N., Taraborrelli, D., Tang, M. J., Bozem, H., Hosaynali-Beygi, Z., Axinte, R., Oswald, R., Novelli, A., Kubistin, D., Hens, K., Javed, U., Trawny, K., Breitenberger, C., Hidalgo, P. J., Ebben, C. J., Geiger, F. M., Corrigan, A. L., Russell, L. M., Ouwersloot, H. G., Vilà-Guerau de Arellano, J., Ganzeveld, L., Vogel, A., Beck, M., Baylerle, A., Kampf, C. J., Bertelmann, M., Köllner, F., Hoffmann, T., Valverde, J., González, D., Riekkola, M.-L., Kulmala, M., and Lelieveld, J.: The summertime Boreal forest field measurement intensive (HUMPPA-COPEC-2010): an overview of meteorological and chemical influences, Atmos. Chem. Phys., 11, 10599–10618, doi:10.5194/acp-11-10599-2011, 2011. 28566, 28567

**HO<sub>x</sub> photochemistry  
in a boreal forest**

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Yassaa, N., Song, W., Lelieveld, J., Vanhatalo, A., Bäck, J., and Williams, J.: Diel cycles of isoprenoids in the emissions of Norway spruce, four Scots pine chemotypes, and in Boreal forest ambient air during HUMPPA-COPEC-2010, *Atmos. Chem. Phys.*, 12, 7215–7229, doi:10.5194/acp-12-7215-2012, 2012. 28568
- 5 York, D., Evensen, N. M., Martínez, M. L., and Delgado, J. D. B.: Unified equations for the slope, intercept, and standard errors of the best straight line, *Am. J. Phys.*, 72, 367–375, doi:10.1119/1.1632486, 2004. 28619

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

**Table 1.** Instrumentation applied for above canopy observations during HUMPPA-COPEC2010.

Species/Quantity	Time resolution	Accuracy (1 $\sigma$ )	Precision (1 $\sigma$ )	Lower limit of detection	Technique <sup>b</sup>
OH	~ 4 min	30% (2 $\sigma$ )	$\sim 5 \times 10^5$ molec cm <sup>-3</sup>	$\sim 9 \times 10^5$ molec cm <sup>-3</sup>	IPI-LIF-FAGE
HO <sub>2</sub>	~ 15 s	30% (2 $\sigma$ )	< 0.8 pptV	0.4 pptV	LIF-FAGE
OH	30 s	32%	—	$\sim 5 \times 10^4$ molec cm <sup>-3</sup>	CIMS
O <sub>3</sub>	3 s	1%	—	~ 1 ppbV	UV
NO, NO <sub>2</sub>	1 s	~ 5%	~ 14 pptV	5 pptV	CLD (+Bluelight converter)
NO <sub>3</sub>	5 s	—	—	2–4 pptV	CRD
N <sub>2</sub> O <sub>5</sub>	5 s	—	—	5–10 pptV	CRD
HONO	30 s	10%	~ 1–2%	< 5 pptV	LOPAP
H <sub>2</sub> O <sub>2</sub> and ROOH	5 min	25–30%	8 pptV (3 $\sigma$ )	15 pptV	Dual enzyme
CO	1 s	< 10%	< 10%	~ 1 ppbV	UV
HCHO	5 min	19%	~ 700 pptV (3 $\sigma$ )	9 pptV	Hantsch
Alkanes, alkenes, isoprene, monoterpenes	60 min	—	—	1 pptV	GC-MS
Methanol, acetone, acetonitrile, total terpenes, benzol, toluene isoprene <sup>a</sup>	6 min	—	—	~ 50 pptV	PTR-MS
Total OH reactivity	1 min	16%	3–4 s <sup>-1</sup>	3 s <sup>-1</sup>	CRM
J <sub>O(1D)</sub> , J <sub>NO<sub>2</sub></sub>	1 s	~ 15%	1%	—	Filterradiometer

<sup>a</sup> Isoprene measurements might be affected by 2-methyl-3-buten-2-ol (MBO) detected at the same mass to charge ratio.

<sup>b</sup> IPI-LIF-FAGE = InletPreinjector-Laser Induced Fluorescence-FAst Gas Expansion; CIMS = Chemical Ionization Mass Spectrometry; UV = UltraViolet absorption/fluorescence; CLD=Chemiluminence Detector; CRD= Cavity Ring Down; LOPAP = Long Path Absorption Photometer; GC-MS = Gas Chromatography-Mass Spectrometry; PTR-MS = Proton Transfer-Reaction Mass Spectrometer; CRM = Comparative Reactivity Method.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Deposition rates/velocities used in the model.


Species	$v_D$ ( $\text{cm s}^{-1}$ )	Deposition rate ( $\text{s}^{-1}$ )
O <sub>3</sub>	0.56 <sup>a</sup>	–
NO	0.44 <sup>b</sup>	–
NO <sub>2</sub>	0.44 <sup>b</sup>	–
HCHO	0.50 <sup>b</sup>	–
HNO <sub>3</sub> , org. nitrates	3.50 <sup>b</sup>	–
PAN	0.29 <sup>b</sup>	–
H <sub>2</sub> O <sub>2</sub> , org. peroxides	–	$4 \times 10^{-5c}$

<sup>a</sup> mean value derived from measurements during HUMPPA-COPEC-2010,

<sup>b</sup> values taken from Evans et al. (2000),

<sup>c</sup> estimated by best fit of calculated to measured H<sub>2</sub>O<sub>2</sub> mixing ratio.

**Table 3.** HO<sub>x</sub> budget under different conditions of observed radiation (high, left; low, right panels) and total OH reactivity (moderate, upper; high, bottom panels). Radical production (green), recycling (blue), and loss (red) pathways are indicated by bold arrows. All rates are given in 10<sup>6</sup> molec cm<sup>-3</sup> s<sup>-1</sup>.



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

**Table 4.** Sensitivity analysis with respect to changes in observed, model constraining parameters. The values presented refer to a dataset around local solar noon (1 p.m. in UTC+2).

	OH (molec cm <sup>-3</sup> )	mod/mod <sub>reference</sub>	HO <sub>2</sub> (molec cm <sup>-3</sup> )	mod/mod <sub>reference</sub>
Observation	3.42 × 10 <sup>6</sup>		3.63 × 10 <sup>8</sup>	
Reference simulation	5.20 × 10 <sup>6</sup>	1	2.87 × 10 <sup>8</sup>	1
2× NO	6.47 × 10 <sup>6</sup>	1.24	2.22 × 10 <sup>8</sup>	0.77
2× O <sub>3</sub>	6.42 × 10 <sup>6</sup>	1.23	3.32 × 10 <sup>8</sup>	1.16
2× H <sub>2</sub> O	6.62 × 10 <sup>6</sup>	1.27	3.31 × 10 <sup>8</sup>	1.15
0.5× CO	5.63 × 10 <sup>6</sup>	1.08	2.65 × 10 <sup>8</sup>	0.92
2× CO	4.55 × 10 <sup>6</sup>	0.88	3.23 × 10 <sup>8</sup>	1.13
2× H <sub>2</sub> O <sub>2</sub>	5.27 × 10 <sup>6</sup>	1.01	2.92 × 10 <sup>8</sup>	1.02
2× HCHO	5.30 × 10 <sup>6</sup>	1.02	3.07 × 10 <sup>8</sup>	1.07
0.5× deposition	5.59 × 10 <sup>6</sup>	1.08	3.17 × 10 <sup>8</sup>	1.10
2× deposition	5.83 × 10 <sup>6</sup>	1.12	3.12 × 10 <sup>8</sup>	1.09
2× photolysis	7.11 × 10 <sup>6</sup>	1.37	3.63 × 10 <sup>8</sup>	1.26
10× photolysis	1.77 × 10 <sup>7</sup>	3.40	6.88 × 10 <sup>8</sup>	2.40

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 1.** Actinic flux density of the mercury lamp used for calibration of the HORUS instrument. The actinic flux density  $\Phi_0$  is derived from the photolysis of  $\text{N}_2\text{O}$  at different mixing ratios and the subsequent detection of the  $\text{NO}$  yields. The data shown here is the combined dataset from two actinometric measurements, one of them conducted before (red dots) and a second one after (black triangles) the field campaign, indicating a good long-term stability of the lamp actinic flux density. Errorbars indicate the propagated statistical variability of the calculated flux density.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 2.** Sensitivity change inside the second detection axis of HORUS as a function of the associated laser scatter signal. A linear correlation ( $R^2 = 0.96$ ) with a non-significant offset was observed during instrument calibrations. The functional dependency obtained from linear regression was applied to the measurements to correct for changes in instrument sensitivity by continuously monitoring the laser scatter signal. A similar correction function for the change in sensitivity of the first detection cell has also been determined empirically and applied.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.



**Fig. 3.** Comparison of the measured H<sub>2</sub>O<sub>2</sub> with simulation based on HO<sub>2</sub> measurements by LIF. The measured HO<sub>2</sub> is needed to explain the observed hydrogen peroxide.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 4.** Histogram of the Monte-Carlo simulation ( $N = 9999$ ) for the reaction rate coefficients on a single dataset in the early afternoon above canopy. The resulting distribution shows a slight skewness with a median of  $6.15 \times 10^6 \text{ molec cm}^{-3}$  and a mean of  $(6.21 \pm 0.86) \times 10^6 \text{ molec cm}^{-3}$ . The dark blue bars indicate the  $\pm 1\sigma$  range.


HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.



**Fig. 5.** Comparison of OH measurements by IPI-LIF-FAGE technique and by CIMS on ground (University of Helsinki, Petäjä et al., 2009). Nighttime OH was observed by both techniques.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 6.** Comparison of OH measurements by IPI-LIF-FAGE technique and by CIMS on ground (University of Helsinki, Petäjä et al., 2009) based on 30 min average data. Errorbars indicate the precision of the associated measurements. Linear regression following the method by York et al. (2004) yields a slope of  $1.20 \pm 0.14$  and an insignificant offset (offset:  $(2 \pm 93) \times 10^3 \text{ molec cm}^{-3}$ ).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 7.** Simultaneous OH measurements on the ground and above the canopy. Concentration maxima observed on the tower are up to a factor of 3 times higher than on the ground. Both instruments measured similar values during night.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 8.** Linear correlation of OH and  $J_{O(1D)}$  is found during HUMPPA-COPEC-2010. The comparison of the correlation on ground and above canopy shows that higher OH values on tower are driven by enhanced radiation. Similar slopes, mainly depending on the abundance of VOCs and NO<sub>x</sub>, do not suggest differences in the chemical regime. Small symbols show 5 min dataset, large symbols are mean of data binned in steps of  $3 \times 10^{-6} \text{ s}^{-1}$ .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 9.** Average diurnal OH production below (left panel, 27–30 July) and budget above (right panel, 1–8 August) the canopy. The whiskers indicate the variability of the total loss rate, calculated from total OH reactivity and hydroxyl radical measurement using LIF data.



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.



**Fig. 11.** Inclusion of the terpene mechanism MTM. HO<sub>2</sub> is still underpredicted in cases when the model reactivity does not match the measured reactivity. This indicates that the missing reactivity is an unaccounted source of HO<sub>2</sub>. The OH is reproduced reasonably well, suggesting that the missing source of OH due to the underpredicted HO<sub>2</sub> is compensated by the missing OH reactivity in the simulation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 12.** Tuning the model to the observed total OH reactivity by introducing an additional  $\alpha$ -pinene equivalent slightly improves the HO<sub>2</sub> agreement. The production of OH from the additional HO<sub>2</sub> is not sufficient to compensate for the enhanced reactivity in the model.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 13.** Uncertainties of HO<sub>x</sub> simulations (using the terpene mechanism) with respect to measurement uncertainties. The solid line represents the 1 : 1 ratio of observed and simulated HO<sub>x</sub>. Data points showing significant discrepancies within their uncertainties by more than a factor of  $2^{\pm 1}$  (dashed lines) are shown in red.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.



**Fig. 14.** Simulated contributions to the OH production during HUMPPA-COPEC-2010. Partitioning at different conditions of radiation and observed total OH reactivity with respect to the total OH production derived from measured OH concentrations and the total OH reactivity.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.




**Fig. 15.** Simulated contributions to the observed total OH reactivity during HUMPPA-COPEC-2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



HO<sub>x</sub> photochemistry  
in a boreal forest

K. Hens et al.



**Fig. 16.** Simulated production and loss rates of HO<sub>2</sub> during HUMPPA-COPEC-2010 under conditions of different radiation and total observed OH reactivity.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

