

# Supplement:

## VARIATIONS IN TROPOSPHERIC SUBMICRON PARTICLE SIZE DISTRIBUTIONS ACROSS THE EUROPEAN CONTINENT 2008-2009

**D. C. S. Beddows<sup>1</sup>, M. Dall'Osto<sup>2</sup>, Roy M. Harrison<sup>1\*†</sup>, M. Kulmala<sup>3</sup>,  
A. Asmi<sup>3</sup>, A. Wiedensohler<sup>4</sup>, P. Laj<sup>5</sup>, A.M. Fjaeraa<sup>6</sup>, K. Sellegri<sup>7</sup>,  
W. Birmili<sup>4</sup>, N. Bukowiecki<sup>8</sup>, E. Weingartner<sup>8</sup>, U. Baltensperger<sup>8</sup>,  
V. Zdimal<sup>9</sup>, N. Zikova<sup>9</sup>, J.-P. Putaud<sup>10</sup>, A. Marinoni<sup>11</sup>, P. Tunved<sup>12</sup>,  
H.-C. Hansson<sup>12</sup>, M. Fiebig<sup>6</sup>, N. Kivekäs<sup>13,14</sup>, E. Swietlicki<sup>13</sup>,  
H. Lihavainen<sup>14</sup>, E. Asmi<sup>14</sup>, V. Ulevicius<sup>15</sup>, P. P. Aalto<sup>3</sup>, N. Mihalopoulos<sup>16</sup>,  
N. Kalivitis<sup>16</sup>, I. Kalapov<sup>17</sup>, G. Kiss<sup>18</sup>, G. de Leeuw<sup>3, 14,19</sup>, B. Henzing<sup>19</sup>,  
C. O'Dowd<sup>20</sup>, S. G. Jennings<sup>20</sup>, H. Flentje<sup>21</sup>, F. Meinhardt<sup>22</sup>, L. Ries<sup>23</sup>,  
H.A.C. Denier van der Gon<sup>19</sup>, A.J.H. Visschedijk<sup>19</sup>**

<sup>1</sup> National Centre for Atmospheric Science, School of Geography, Earth and Environmental Sciences, University of Birmingham, B15 2TT, UK

<sup>2</sup> Institut de Ciències del Mar, CSIC, Pg Marítim de la Barceloneta 37-49 08003 Barcelona, Spain

<sup>3</sup> Department of Physics, University of Helsinki, P.O. Box 64, Helsinki, Finland

<sup>4</sup> Leibniz Institute for Tropospheric Research, Permoserstraße 15, 04318 Leipzig, Germany

<sup>5</sup> Laboratoire de Glaciologie et Géophysique de l'Environnement Université Joseph Fourier, Grenoble 1/CNRS, 38400 St. Martin d'Hères, France

<sup>6</sup> NILU - Norwegian Institute for Air Research Instituttveien 18, 2027 Kjeller, Norway

<sup>7</sup> Laboratoire de Météorologie Physique, UMR 6016, CNRS/University of Clermont-Ferrand, Clermont-Ferrand, France

<sup>8</sup> Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen PSI, Switzerland

<sup>9</sup> Laboratory of Aerosol Chemistry and Physics, Institute of Chemical Process Fundamentals of the AS CR, v.v.i., Rozvojova 135, 16502 Praha 6, Czech Republic

<sup>10</sup> European Commission, Joint Research Centre, Institute for Environment and Sustainability, 21027 (VA), Italy

<sup>11</sup> CNR-ISAC, Institute of Atmospheric Sciences and Climate, 40129, Bologna, Italy

<sup>12</sup> Department of Applied Environmental Science (ITM), Stockholm University, 10691 Stockholm, Sweden

<sup>13</sup> Department of Physics, Lund University, SE-22100, Lund, Sweden

<sup>14</sup> Finnish Meteorological Institute, Erik Palménin aukio 1, P.O. Box 503, FI-00101, Helsinki, Finland

<sup>15</sup> Center for Physical Sciences and Technology, Savanoriu 231, 02300 Vilnius, Lithuania

<sup>16</sup> Environmental Chemical Processes Laboratory, Department of Chemistry, University of Crete, Greece

<sup>17</sup> Institute of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Blvd. Tzarigradsko chaussee, 72, 1784 Sofia, Bulgaria

<sup>18</sup> MTA-PE Air Chemistry Research Group, P.O. Box 158, 8201 Veszprém, Hungary

<sup>19</sup> Netherlands Organisation for Applied Scientific Research TNO, Princetonlaan 6, 3508 TA Utrecht, The Netherlands

<sup>20</sup> National University of Ireland Galway, University Road, Galway, Ireland

<sup>21</sup> German Meteorological Service, Hohenpeissenberg Observatory, Albin-Schwaiger Weg 10, 82383 Hohenpeißenberg, Germany

<sup>22</sup> German Federal Environment Agency (UBA), Messnetzzentrale, Paul-Ehrlich-str. 29, 63225, Langen, Germany

<sup>23</sup> German Federal Environment Agency (UBA), Platform Zugspitze of GAW Global Station Zugspitze/Hohenpeissenberg, Zugspitze 5, 28475 Zugspitze, Germany

---

\* To whom correspondence should be addressed.

Tele: +44 121 414 3494; Fax: +44 121 414 3708; Email: r.m.harrison@bham.ac.uk

†

Also at: Department of Environmental Sciences / Center of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203, Jeddah, 21589, Saudi Arabia














Figure S1. Cluster Centres derived from the *k*-Means analysis of the hourly averaged spectra.






1






2



3



4



5

6  
7  
8  
9




Figure S2. Average clustered hourly particle size distributions (*cluster 1-9 left hand panels*) and the spatial distribution of each cluster (*right hand panels*). The solid black line shows the average spectrum and the dashed lines show the 10<sup>th</sup>, 25<sup>th</sup>, 75<sup>th</sup> and 90<sup>th</sup> percentile spectrum. The maximum and minimum spectra are traced out by the extremities of the shaded areas. When counting the spectrum types within the whole data set, the sites which collected above the 90<sup>th</sup>, 75<sup>th</sup> and 50<sup>th</sup> percentile were marked with a progressively lighter orange colour. Circles denote boundary layer sites and triangles sites of relatively high altitude. The right hand panel shows the colour maps plotted using the average day of hourly spectra for each of the clusters.





Figure S3. Cluster Centres derived from the *k*-Means analysis of the weekly averaged spectra.





1


2

3

4

5

Figure S4. Average clustered weekly particle size distributions (*cluster 1-5 left hand panels*) and the spatial distribution of each cluster (*right hand panels*). The solid black line shows the average spectrum and the dashed lines show the 10<sup>th</sup>, 25<sup>th</sup>, 75<sup>th</sup> and 90<sup>th</sup> percentile spectrum. The maximum and minimum spectra are traced out by the extremities of the shaded areas. When counting the spectrum types within the whole data set, the sites which collected above the 90<sup>th</sup>, 75<sup>th</sup> and 50<sup>th</sup> percentile were marked with a progressively lighter orange colour. Circles denote boundary layer sites and triangles sites of relatively high altitude. The right hand panel shows the colour maps plotted using the average day of hourly spectra for each of the clusters.




Cluster 1, 2 and 3 are Central European site dominant.

Clusters 5 & 4 are Atlantic-Arctic-Nordic site dominant.

Cluster 9 is a Central-Europe-Balkan site dominant.

Clusters 6, 7 & 8 are Nordic-Central European site dominant.

**Figure S5. Frequency percentage of the clusters measured at each of the sites: (left hand panel) sites ordered by latitude and (right hand panel) sites ordered by longitude. [Bar colours correspond to the different site types in Table 1: rural – green; mostly remote – grey; weakly influenced – blue and agglomeration – orange. Sites with blank bars have no type assignment.]**



**Figure S6. Seasonal percentages for the observed clusters.**







Figure S7. Using the volume distribution, the fitted modal-diameters of each cluster (1-9) is plotted against the Weighted Latitude, calculated for each cluster, using  $WL = \frac{\sum_i^{24} W_i \cdot X_i}{\sum_i^{24} W_i}$ , where  $X_i$  is the latitude/longitude of the sites where the cluster is detected and  $W_i$  corresponding population of the cluster across for the 24 sites.



(a) (b)  
Figure S8. Colour maps derived from the averaging of (a) clusters 1-4 and (b) 5-9, showing the diurnal trends of the spectra collected inside and outside of central Europe respectively.




Figure S9. Peak fitted clusters.







Figure S10. Plot showing how the modal diameters of the peaks (a) peak 1 and (b) peak 2, fitted to the average cluster spectra (shown in Figure S9) vary with the hour of their maximum occurrence. The red (1,2,3 and 4) and green (5,6,7,8 and 8) colours depict two aggregated trends observed in the data based on a South to North and West to East air mass movement. The fitted curves for figure (a) are  $MD = 1.5 \times HR + 24$  for clusters 1- 4 and  $MD = 1.3 \times HR + 9.7$  for clusters 5 - 9.




2008/12/20




2008/12/21



2008/12/22






2008/12/23




2008/12/24


**Figure S11. Synoptic chart available at midnight of the sequential days along the trajectory arriving at FKL on the 24<sup>th</sup> of December 2008 at midnight (Figure 8).**




**Figure S12. Meteorology along the trajectory arriving at FKL on the 24<sup>th</sup> of December 2008 at midnight (Figure 8).**




2009/04/12 2009/04/13 2009/04/14 2009/04/15 2009/04/16  
**Figure S13. Synoptic chart available at midnight of the sequential days along the trajectory arriving at MHD on the 16<sup>th</sup> of April 2009 at 18:00 (Figure 9).**




**Figure S14. Meteorology along the trajectory arriving at MHD on the 16<sup>th</sup> of April 2009 at 18:00 (Figure 9).**




2008/02/03




2008/02/04



2008/02/05







2008/02/06




2008/02/07

**Figure S15. Synoptic chart available at midnight of the sequential days along the trajectory arriving at ZEP on the 7<sup>th</sup> of February 2008 at 06:00 (Figure 10).**




**Figure S16. Meteorology along the trajectories arriving at ZEP on the 7<sup>th</sup> of February 2008 at 06:00 (Figure 10).**







2009/10/13




2009/10/14



2009/10/15






2009/10/16



2009/10/17

**Figure S17. Synoptic chart available at midnight of the sequential days along the trajectory arriving at CMN on the 17<sup>th</sup> of October 2009 at 18:00 (Figure 11).**



**Figure S18. Meteorology along the trajectory arriving at CMN on the 17<sup>th</sup> of October 2009 at 18:00 (Figure 11).**




Figure S19. Average clustered particle size distributions (*cluster 1-9 shown in left hand panels in Figure 3 but plotted on log scales*). The solid black line shows the average spectrum and the dashed lines show the 10<sup>th</sup>, 25<sup>th</sup>, 75<sup>th</sup> and 90<sup>th</sup> percentile spectrum.