

Linear trends in cloud top height from passive observations in the oxygen A-band

L. Lelli¹, A. A. Kokhanovsky^{*1}, V. V. Rozanov¹, M. Vountas¹, and J. P. Burrows¹

¹Institute of Environmental Physics and Remote Sensing, University of Bremen, Otto-Hahn-Allee 1, 28334 Bremen, Germany

^{*}now at EUMETSAT, Eumetsat Allee 1, D-64295 Darmstadt, Germany

Correspondence to: L. Lelli (luca@iup.physik.uni-bremen.de)

Abstract. Measurements by the hyperspectral spectrometers GOME, SCIAMACHY, and GOME-2 are used to determine the rate of linear change (and trends) in cloud top height (CTH) in the period between June 1996 and May 2012. The retrievals are obtained from Top-Of-Atmosphere (TOA) backscattered solar light in the oxygen A-band using the Semi-Analytical CloUd Retrieval Algorithm SACURA. The physical framework relies on the asymptotic equations of radiative transfer, valid for optically thick clouds. Using linear least-squares techniques, a global trend of -1.78 ± 2.14 m yr⁻¹ in deseasonalised CTH has been found, in the latitude belt within $\pm 60^\circ$, with diverging tendencies over land ($+0.27 \pm 3.2$ m yr⁻¹) and ocean (-2.51 ± 2.8 m yr⁻¹). The El Niño-Southern Oscillation (ENSO), strongly coupled to CTH, forces clouds to lower altitudes. The global ENSO-corrected trend in CTH amounts to -0.49 ± 2.22 m yr⁻¹. At a global scale, no explicit regional pattern of statistically significant trends (at 95% confidence level, estimated with bootstrap technique) have been found, which would be representative of typical natural synoptical features. One exception is North Africa, which exhibits the strongest upward trend in CTH sustained by an increasing trend in water vapor.

15 1 Introduction

Tropospheric clouds are main players in the Earth's climate system. Characterization of long-term global cloud properties (top and bottom height, optical thickness and albedo, effective radius of droplets) aims to support radiation budget assessments as well as trace gas retrievals, and analysis of cloud interactions with aerosol particles and gases in the atmosphere. The altitude of a cloud plays a cardinal role because clouds at different heights (i.e., temperatures) exert different feedbacks. High-altitude clouds absorb infrared radiation that comes from the lower atmosphere and radiate like

blackbodies. Since the temperature contrast between elevated clouds and lower atmosphere is high, the clouds have a positive contribution on the local net energy balance. At the same time, their albedo is small because they are, in most cases, comparably thin. Thus, they can warm the atmosphere more than they cool it, exerting a positive feedback.

Conversely, low-altitude clouds are strongly reflecting objects owing to their high optical density but they ineffectually shield infrared radiation emitted by atmospheric gases to the outer space. The reason is that the temperature of low clouds is closer to the temperature of the ground and the local net energy balance is close to zero. These clouds in turn can cool the climate system more than they warm it and thus exert a negative feedback.

Loeb et al. (2012) observed this mechanism in a study that focused on clouds in the tropical belt. However, this situation may change when considering the Pacific Northeast over a decadal time window. Evidence of a positive feedback by low-level clouds has already been demonstrated (Clement et al., 2009). It is therefore likely that no general description is possible on a global scale and regional studies should be conducted instead.

Moreover, a critical aspect of any trend analysis is that, due to the limitation of a relatively short derived CTH time serie, no statistical significance may be found. Previous works do not provide clear indications. The analysis of the 7-year GOME record (June 1996 – May 2003) by Loyola et al. (2010) shows a change of -4.8 m yr^{-1} for the latitudinal belt within $\pm 60^\circ$. Davies and Molloy (2012) reported a decreasing global trend of $-40 \text{ m decade}^{-1}$, as seen by MISR on the Terra platform. Evan and Norris (2012) showed that the previous results were affected by instrumental artifacts. Corrected MISR time series exhibit opposite tendency, that is an increase of $+54 \text{ m decade}^{-1}$, supported by a MODIS (on Terra) upward trend of $+60.9 \text{ m decade}^{-1}$. The considered record length amounts to 8 years, from March 2003 throughout February 2010.

In the present study it has been attempted to extend the cloud records including measurements of the three sensors GOME, SCIAMACHY and GOME-2 (hereinafter termed GSG). Since the launch of GOME, back in 1995, more than 17 years of data are now available and they form the observational basis of the present analysis (June 1996 – May 2012).

The manuscript has the following structure. Section 2 describes how the monthly-sampled cloud record is calculated from measured radiances. Section 3 introduces the model used to derive the rate of linear change in cloud top height. Section 4 is devoted to the main results of the work. First, an analysis of ENSO as reflected in cloud properties is presented and trends in cloud top height are assessed. In the last section, the findings are summarized also giving conclusions and outlook for future work.

55 2 Data and Methods

The spaceborne nadir-viewing spectrometers Global Ozone Monitoring Experiment (GOME, Burrows et al. (1999)), Scanning Imaging Absorption spectrometer for Atmospheric CHartographY (SCIAMACHY, Bovensmann et al. (1999)) and GOME-2 (Callies et al., 2000) measure solar backscattered light at a spectral resolution of 0.24–1.48 nm (see Table 1 for their characteristics). At the time
60 of processing, GOME L1b data were in their version 4.00 and SCIAMACHY L1b version 7.03 with consolidation degree U. GOME-2 L1b data for time window 01/2007–12/2009 were in reprocessed stage, while near-real time data have been used for 01/2010–05/2012. Cloud fraction is obtained from broadband Polarization Measuring Devices (PMD) measurements with the Optical Cloud Recognition Algorithm (OCRA) (Loyola and Ruppert, 1998). For GOME, cloud fraction is
65 delivered in bundle with L1b data, while cloud fraction for GOME-2 is taken from the off-line L2 reprocessed dataset made available by DLR in the framework of EUMETSAT’s O3M Satellite Application Facility. Cloud fraction for SCIAMACHY has been calculated at University of Bremen with the in-house OCRA implementation.

In particular, the sensors cover the wavelength range 758–772 nm (at ≈ 0.4 nm resolution) where
70 is situated one of the absorption bands of molecular oxygen, the O₂ A-band. The A-band is exploited by the Semi-Analytical CloUd Retrieval Algorithm (SACURA) (Rozanov and Kokhanovsky, 2004) to generate cloud top height (CTH) records in the following way: given the local fractional cloud cover (f), the sun-normalised Earthshine reflectances are weighted with the Independent Pixel Approximation (IPA) (Marshak et al., 1995). In this way, the measured scene reflectance R_{mes} is scaled
75 to the cloud reflectance R_{cl} generated by a full cloudy ($f = 1$) and a cloud-free ($f = 0$) pixel with

$$R_{mes} = f R_{cl} + (1 - f) R_s. \quad (1)$$

The clear-sky reflectance R_s is substituted by the Minimum Lambert-Equivalent Reflectivity (MLER) taken from the global database Tropospheric Emission Monitoring Internet Service (TEMIS) (Koelemeijer et al., 2003).

80 It has been demonstrated (Kokhanovsky et al., 2007; Lelli et al., 2012) that cloud top height is not substantially affected by the scaling of the cloud reflection function with different values of fractional cloud cover as long as f is known from an independent source. This statement is true regardless of cloud fraction and of instrumental spatial resolution, because the algorithm makes use of spectral ratios. From a physical point of view, it is also clear that, on the coarse footprint scales
85 of the sensors used in this work, the cloud reflectance is dominated by the contribution of photons scattered directly back to the platform and not by horizontal photon transport. This is not true for spatially better resolved instruments for which a cloud volume may have side lengths comparable with the footprint size.

The cloud optical thickness (COT) is retrieved in the continuum at 758 nm and used for the
90 computation of the forward TOA cloud reflectance with asymptotic approximations of radiative

transfer (Kokhanovsky and Rozanov, 2004). Then, the minimal difference between R_{cl} and the forward spectrum (modeled with the analytical equations of radiative transfer) is iteratively looked for along the whole O₂ A-band, at the nominal spectral sampling of the instruments. The distinctive feature of SACURA is the modeling of clouds as scattering layers instead of Lambertian reflectors
95 (Koelemeijer et al., 2001), therefore radiative transfer throughout, above and below the clouds is taken into account. A comprehensive description of the algorithm can be found in Lelli et al. (2012, and references therein), where also accuracy, validation and selection criteria of CTH retrievals are given.

Cloud top height retrievals (b) are firstly binned in 13 height layers (k) (see Table 2 for the
100 grid definition), projected onto a equidistant rectangular grid of 0.5° sided-cells at latitude (i) and longitude (j), counted-averaged over all altitudes (being c_k the retrieval counts at height bin k) at month t and the CTH means h_t are computed with

$$h_t(i, j) = \frac{\sum_k c_{kt}(i, j) b_{kt}(i, j)}{\sum_k c_{kt}(i, j)}. \quad (2)$$

This formulation has been chosen for consistency with Evan and Norris (2012, Eq. (1)). The values
105 for h_t are plotted in Fig. 1 for global (top) and tropical (bottom plot) region. In both cases, the time series of the individual instruments exhibit in-phase seasonality for the respective overlapping months, whereas the relative offsets (top plot) cannot be explained by the mere diurnal cycle clouds have, owing to the different overpass time of the instruments which fly 1/2 hour ahead of each other, GOME-2 crossing first the equator at 9:30 AM. In case of multi-layered cloud systems it is expected
110 that sensors with coarser footprints, such as GOME (320×40 km²), will observe more high clouds at the expenses of low clouds as compared to finer resolved sensors, such as SCIAMACHY (60×40 km²) and GOME-2 (80×60 km²). On a individual pixel scale, this effect can be seen in Lelli et al. (2012, Fig. 8, p. 1559), in which a GOME-derived CTH was co-located with a ground-based radar-derived CTH for a three-layer cloudy scene. GOME retrieved the height of the uppermost
115 cloud layer. On larger time and spatial scales, this situation is depicted in Fig. 2, which shows retrieval counts c_k (normalised to the total number of counts) as function of height bin k , for different instruments and for global (left) and tropical (mid plot) region. The right plot of Fig. 2 shows the count difference between GOME and SCIAMACHY in both global and tropical region. Especially in the tropics, where heterogeneity is frequent, GOME counts increase toward higher layers of the
120 troposphere. These values can be devised as correction factors for the harmonisation of time series among the instruments, given that the same scene is sensed and no time lag occurs between different measurements. Therefore, in the current analysis, it is preferred to compute the CTH anomalies of Fig. 3 subtracting the climatological seasonal cycle for each instrument separately, in order to avoid the introduction of *post-hoc* artifacts in the record.

125 While the CTH anomalies from GOME and SCIAMACHY are in almost perfect agreement, the transition between SCIAMACHY and GOME-2 records is not smooth. The reason for discrepancies may be manifold. On one hand, a change in the PMD pixel definitions of GOME-2 was devised in

April 2008 (EUMETSAT, 2010), which may have impacted cloud fraction f and the cloud reflection function R_{cl} of Eq. 1. On the other hand, radiometric calibration issues may also influence retrievals
 130 inferred from different instruments. For instance, van Diedenhoven et al. (2005) have shown that a difference of 20 hPa in surface pressure retrieved in the oxygen A-band by GOME and SCIAMACHY could be corrected adding an offset of 0.86% to the TOA reflectance in the continuum at 756 nm. However, tests have shown that radiometric uncertainties have almost no impact in the retrieved CTH with SACURA. This is a feature of the algorithm, which is based on spectral ratios
 135 and on the concurrent fit of CTH and CBH along the whole band.

For the ensuing analysis, the time series of GOME and SCIAMACHY have been joined in June 2003 because the on-board tape recorder of GOME broke down and global coverage wasn't provided after May 2003. Time series of SCIAMACHY and GOME-2 have been joined in May 2008. Given that SCIAMACHY is a thoroughly tested instrument and GOME-2 PMD definitions were updated
 140 in April 2008, we used SCIAMACHY anomalies prior May 2008, even though GOME-2 data were available. For the rest of the time serie, GOME-2 anomalies were used, as both instruments converge and exhibit good overlap. In fact, both instruments sense the strong negative CTH anomaly in year 2011. This feature dominates the whole time serie and it gives us reasonable confidence that the order of magnitude as well as the sign of the CTH trend aren't substantially affected, when using
 145 SCIAMACHY data.

3 Trend model

The basic requirement for the application of linear least-square regression is that the variable of interest is not persistent (Wilks, 2011). Persistence is the tendency for a variable to remain similar in time and is measured as the degree of temporal autocorrelation r_p . The dataset of this study is
 150 sampled on a monthly basis, thus autocorrelation at lag $p=1$ reveals the degree of persistence that CTH has after one month. The upper plot of Fig. 4 shows the autocorrelation r_p of CTH anomalies up to 4 months lag, for global and tropical region. It can be seen that in both cases the altitude of clouds does not exhibit almost any persistence. That is, the actual anomaly is not affected by values leading by one month. Since characteristic time scales of clouds range from minutes for individual
 155 shallow cloud cumulus to many hours for large stratiform cloud systems, over a month their features are randomised by intervening winds and persistence is not found. Therefore methods for normally distributed (gaussian) variables can be applied, given that the dataset has been deseasonalised. In the lower plot of Fig. 4 the mapped CTH autocorrelation signal (not exceeding $r_p=0.13$ over the Pacific Ocean) resembles ENSO-related patterns, stretching across the globe over the Atlantic and
 160 the Indian Oceans.

The trend magnitude β is calculated, at latitude and longitude (i, j) , with

$$Y_t(i, j) = \alpha(i, j) + \beta(i, j) \cdot X_t + \epsilon_t(i, j) \quad t = 1, \dots, 192 \quad (3)$$

where Y_t are the CTH anomalies, X is the temporal variable sampled at monthly step t , α and ϵ_t the offset and the random noise, respectively. We emphasize that the time series of absolute values
165 have been neither corrected for the impact of spatial resolution nor fitted directly. This is because the time series of Fig. 1 are shifted and the three instruments have different sensing local times. The chosen strategy has been to regress time series of anomalies, instead. This approach is analogous to the customary technique described in Mieruch et al. (2008, Eq. 1, p. 495, and references therein) and is reported below for convenience.

170 Being Y_t the monthly mean of the variable of interest at time t (for each geolocation point on the map), μC_t the offsets of the regression line, S_t the seasonal component, ω the desired change rate of the variable at time step X_t and N_t the noise, in the r.h.s. of the following equation

$$Y_t = \mu C_t + S_t + \omega X_t + \delta U_t + N_t \quad (4)$$

the term δU_t describes the level shift δ allowed when concatenating time series from different in-
175 struments at time T_0 , with a step function U_t defined as

$$U_t = \begin{cases} 0, & t < T_0 \\ 1, & t \geq T_0 \end{cases} \quad (5)$$

The removal of the sample mean for each respective month from the time serie of absolute values allows the seasonality to be accounted for. The term δU_t is incorporated by performing this step separately for each instrument. This is because the sample mean of anomalies is, by definition,
180 centered about zero (Wilks, 2011) and the constant μ can be neglected (being μ the mean value of Y_t at time $t=0$). Eventually, Eq. 4 reduces to Eq. 3 and potential autocorrelative effects are embedded in the noise term.

Among the techniques for the estimation of confidence intervals of the trend magnitude β , the *bootstrap resampling* (Efron and Tibshirani, 1993) enables the treatment of potentially non-normal
185 data without any assumption on the underlying probability distribution. It belongs to the group of nonparametric methods and the knowledge of the analytical form of the statistics is not required (Mudelsee, 2010). The guiding idea is the “plug-in” principle: a new scrambled dataset is drawn (resampling with replacement) from the original dataset and a new β' is calculated with Eq. (3). This procedure is repeated n times and an empirical sample distribution for β is estimated. This
190 empirical distribution can be regarded as the approximated estimate to the unknown β distribution and σ_β can be associated with the random effects reflected in the data. The required confidence level is customary chosen as 95%, implying a 2.5% cutting-off probability level on either sides for a two-tailed distribution. When the probability density function of the unknown distribution is related to its standardised Gaussian approximation, the cutting-off level amounts to the mean
195 value ≈ 1.96 standard deviations σ_β (Wilks, 2011, p.140). Therefore, at locations (i, j) where the ratio $|\beta/\sigma_\beta| > 2$, the trend magnitude β exceeds natural variability and is considered statistically significant.

Fig. 5 shows the normalised probability density function (PDF) and the cumulative density function (CDF) of β , calculated with 10^4 bootstrap resamples, for global (left), tropical (mid) regions
200 and the Central East Pacific (right). The red tails of the PDF portray the confidence level calculated with the percentile method at 2.5% and 97.5% quantile and provide the estimate of the uncertainty associated with the trend value β . We note that the condition of normality is reasonably satisfied even in the region of highest autocorrelation (i.e., Central East Pacific, 170° W- 120° W, 5° N- 5° S), as can be seen in the right plot of Fig. 5. Moreover, Fig. 6 displays the theoretical residual quantiles,
205 estimated after regression with the linear model and application of the parameter estimates $\hat{\alpha}$ and $\hat{\beta}$ of Eq. 3 against the sample distribution of global CTH anomalies. Since the majority of points cluster about the straight line, the linear model seems to be a reasonable assumption, given that the autocorrelation functions of Fig. 4 drops almost to zero after one month.

Therefore, the major process that still can influence the time serie is the quasi-stationary ENSO.
210 Laken et al. (2012) indirectly came to similar conclusions, while analysing the effect of solar activity on cloud altitudes with MODIS Terra and Aqua measurements. Norris (2005) comes to the same conclusion as well. The ENSO periodicity is still matter of on-going research (Solomon and Newman, 2012) and it is not the focus of this work.

4 Results

215 4.1 Cloud property changes coupling to ENSO

The El-Niño-Southern Oscillation (ENSO) has been recognized as the largest natural fluctuation that can establish teleconnections and modulate cloud properties. ENSO has to be understood as the tie between two different processes (Trenberth, 1997): the term *El Niño* is assigned to a yearly warming of the ocean current which streams along the coasts of Peru and Ecuador in late December. Regularly,
220 but not every year, a more pronounced warming of the wide tropical-east pacific pool occurs, driven by the magnitude of trade wind flows. The atmospheric component of this perturbation is termed *Southern Oscillation*. The opposite phase, namely *La Niña*, corresponds to a cooling of the pacific basin. Quantitatively, ENSO is identified when SST gradient changes over the Central East Pacific (170° W - 120° W, 5° N - 5° S) exceed $\pm 0.5^\circ$ C, resulting in the standard Niño 3.4 climate index
225 (U. S. National Centers for Environmental Prediction (NCEP), 1993).

In Fig. 7 the Niño 3.4 index has been plotted together with the unsmoothed CTH and CF anomalies for the 17 years of the GSG dataset and high ($R = +0.77$) and moderate ($R = +0.31$) correlations are observed for CTH and CF, respectively. It can also be noted that the maxima of the CF time serie are not in phase with the warm ENSO phases and are delayed by approximately 2 months.
230 The CTH and CF anomaly time series have been partitioned for high clouds (defined as clouds with altitude $h > 7$ km), middle clouds ($3.2 \text{ km} < h < 7 \text{ km}$) and low clouds ($h < 3.2 \text{ km}$), smoothed with a 6-month running mean and plotted in Fig. 8. The correlation coefficients (calculated with

unsmoothed data) reported on the right clearly indicate that a warm (cold) ENSO phase corresponds to a decrease (increase) in CF for low ($R = -0.55$) and mid ($R = -0.31$) clouds, while high clouds increase (decrease) their coverage ($R = +0.53$). In Fig. 8 it is also shown that the lagged extrema in column-averaged CF of Fig. 7 originate in the lowest layers of the troposphere, because CF for low- and mid-level clouds always anticipate the extrema for high clouds by a constant lag of 2 months.

More precisely, the coupling of CTH and CF to El Niño can be understood inspecting Fig. 9, where R for either cloud parameters has been plotted as function of altitude bin h . Starting from the Planet Boundary Layer (PBL) ($h < 2$ km), a warmer ocean surface produces evaporation and CF increases ($R = +0.39$), while CTH does not exhibit almost any correlation ($R = -0.11$). Departing from the PBL ($2 \text{ km} \leq h < 8$ km), the cloud field is compressed by winds (CF decreases in response to warming, $-0.30 \leq R \leq -0.41$) and convection lifts clouds higher up (CTH increases in response to warming, $+0.16 \leq R \leq +0.38$). Within the upper layer of the troposphere ($h \geq 8$ km), clouds rise until the tropopause ($h \approx 16\text{--}17$ km) stops their ascent ($-0.11 \leq R \leq 0$) and, concurrently, extend (CF increases, $+0.19 \leq R \leq +0.44$), following the divergent air flow toward zones of lower pressure. This is consistent with the picture of a reinforced meridional circulation which squeezes clouds at low altitudes, while high clouds spread out in the horizontal direction and air masses close the Hadley cell. In fact, positive correlations between CF, SST and surface convergence are found by Norris (2005) for the International Satellite Cloud Climatology Project (ISCCP) high cloud amount and the ground-based Extended Edited Cloud Report Archive (EECRA) high-cloud record, calculated over almost the same region. Conversely, the warm pool situated in the western tropical Pacific ($30^\circ \text{N}\text{--}30^\circ \text{S}$ and $100^\circ \text{E}\text{--}160^\circ \text{E}$) is characterized by the reversed situation: deep convective clouds occur during cold La Niña periods. Hence negative correlations between CF and SST are found (Norris, 2005; Marchand, 2013). This dipole-like ENSO influence on cloud properties can be seen in Fig. 10, where diagrams of COT-CTH are drawn for the strongest ENSO within the GSG record (i.e. El Niño April 1997 - May 1998 and the ensuing La Niña June 1998 - June 2001) over the Niño 3.4 index box and the tropical warm pool.

Bjerknes (1969) already pointed out how trade winds and sea surface temperature (SST) mutually strengthen for both ENSO phases. Interestingly, he also argued that the ocean response to a change of the wind circulation patterns during El Niño has a characteristic lag of 2 months (this mechanism is known as *Bjerknes feedback*), such as the lag seen in the CF anomaly time series of Fig. 7 and Fig. 8. It is suggestive to regard this similarity as the manifestation of a two-way coupling between cloud properties and sea surface temperature. However, the dynamical interpretation rather than the energetic interpretation is here preferred for two reasons: not only the curves and the correlations of Fig. 8 have been calculated for the narrow Niño 3.4 climate index box, located in the Central East Pacific, where the effect of the weakening of the longitudinal Walker circulation (corresponding to the onset of teleconnections) is at its maximum, but also Dessler (2011) showed that cloud changes have a negligible impact on SST, due to heat storage capacity and heat transport of the ocean, which

270 dominates the ENSO climate variability.

The relationship between CTH, CF and ENSO has been investigated by Erlykin and Wolfendale (2010). Through the analysis of the ISCCP cloud cover and top pressure, they assert that both ENSO phases and global Earth's low troposphere temperature are negatively correlated with low-cloud amount. Thus, in the periods influenced by warm El Niño, low clouds are lifted higher up and are 275 reclassified as mid-level clouds. The opposite holds true for periods driven by La Niña. Since the last decade is showing more frequently cold episodes (59% versus 41% occurrence of warm phases, see also Loeb et al. (2012)), it is reasonable to suppose that the ENSO pulls clouds down.

4.2 Trends in cloud top height

Fig. 11 supports the argument introduced in the previous section: the removal of ENSO (excluding 280 any data within longitudes of 170° W and 120° W and latitudes of 5° N and 5° S) shifts the overall trend, calculated in the latitudinal belt $\pm 60^\circ$ (in order to lessen the influence of ice-capped regions), from -30.26 m to -8.33 m over 17 years.

The time series has been broken up for underlying surface and a trend of -2.51 m yr⁻¹ over ocean and $+0.27$ m yr⁻¹ over land are identified (lower plot of Fig. 11). No exhaustive explanation 285 can be found for this divergent land/water trend on a global scale, due to the complexity and variety of tropospheric processes influencing the clouds. On a global scale, such effects might average out and give rise to well-defined cloud changes. Even so, it was noted (Wagner et al., 2008) that CTH reacts in response to a change in near-surface temperature (ST). The authors analyzed 7.5 years of GOME data and showed not only a strong positive correlation between CTH and ST, but also that 290 ST changes are highest over land. Thus, at the simplest level, global patterns of ST might explain the information concealed in the diverging land/ocean CTH trends.

The findings are summarised in Table 3 together with previous results from independent datasets. The reported values show discrepancies. Indeed, the length of datasets as well as the instrumental spatial resolution can alter the rate of linear change in CTH, because different cloud features are 295 sensed, both in space and time. It must be also noted that retrieved cloud properties and, therefore, trends are affected by the choice of the spectral window used in the algorithms. IR-based algorithms are sensitive to cirrus clouds, whereas NIR-based algorithms, such as SACURA, are valid for thick clouds. Moreover, SACURA takes into account O₂ absorption inside the clouds and the retrieved cloud height cannot be regarded as *effective*, since it will be closer to its geometrical top in the case 300 of a single-layer cloud.

Fig. 12 shows global maps of trend magnitude β and standard deviation σ_β , degraded to 5°-sided cell mesh for computational efficiency and smoothed with the 8 neighboring cells. The maps indicate that trend magnitudes are zonally partitioned. The strongest trends are found over North Africa and the Arabian subcontinent ($> +45$ m yr⁻¹), Central Eastern Pacific (-30 m yr⁻¹) and Indian Ocean 305 (-15 m yr⁻¹). Conversely, the mid-latitudes (60°N-30°N and 30°S-60°S) are homogeneously char-

acterised by a slightly decreasing trend (-2 m yr^{-1} on average). Fig. 12 shows also that the trend variability follows the oceanic contours, being over water almost always greater than 20 m yr^{-1} in the latitude belt $\pm 30^\circ$ and smaller than 15 m yr^{-1} otherwise. Table 4 summarizes zonal values of CTH trend for the tropics and the mid-latitudes, over land and water masses.

310 Analogous conclusions are indirectly drawn by Marchand (2013), who analysed MISR (on board the TERRA platform) cloud amount as function of cloud (top) height. Specifically, linear trends of cloud amount over oceanic regions are shown in Marchand (2013, Fig. 6, p. 1946). It can be seen that where high cloud amount decreases low clouds are more frequent. This effect can be seen over the North-East Pacific and over the Central East Pacific. Hence the author suggests a net decrease in
315 mean cloud top height. Notwithstanding the different length of the time series (MISR covering the period 2001 to 2011), we observe a similar tendency in CTH over the same regions, which is evident in the left plot of Fig. 12. The opposite also holds true: regions showing increasing occurrence of high clouds exhibit a decrease in low cloud amount. Consequently, the mean cloud top height is expected to increase, as seen over Indonesia and, to a lesser extent, over the Central Atlantic.

320 Statistically significant (at 95% confidence level) trends are plotted in Fig. 13. No patterns are discernible that can be easily classifiable in the context of natural synoptic cloud variability, except for the decrease over Central East Pacific and the increase over North Africa. The main difference between Fig. 12 and Fig. 13 is that the CTH increase over the Arabian peninsula is not significant. To investigate this effect, the two regions are magnified on a continental scale, labeled R1 (North
325 Africa) and R2 (Arabian Peninsula), and their trends in CTH are plotted in Fig. 14 (top row). Both regions are naturally arid areas, which do not exhibit any notable evaporation. Any cloud field over R1 and R2 is likely the result of moist air transport from neighboring water basins. For this reason, a water vapor (H_2O) dataset has been added and analysed in Fig. 14. The dataset has been derived from the same instruments used for generation of the cloud data records and is described by Noël
330 et al. (2004, 2005) and Mieruch et al. (2008).

It can be seen that the significant CTH trend over R1 is sustained by an increasing trend in H_2O . Conversely, the decreasing trend in H_2O over R2, which has been reported to be statistically significant (Mieruch et al., 2008), is anticorrelated with CTH. An explanation may be found in large-scale outflows of fine-mode soot particles, that are produced in the Indian subcontinent during biomass
335 burning seasons and transported over R2 (but not over R1) by easterly winds during the winter phase of the Indian monsoon. Indeed, most of the decreasing trend in H_2O is seen during winter seasons (Fig. 14, mid right), while almost no trend is seen in summer months (Fig. 14, mid left). The absorption of solar radiation by soot gives rise to two competing effects. On one hand, soot warms the atmospheric column (Feingold, 2005) and may cause clouds to dry out (Ackerman et al., 2000).
340 On the other hand, soot depletes the amount of radiation reaching the surface, exerting a negative radiative forcing (Nakajima and Schulz, 2009), and the amount of energy at disposal for evaporation is diminished. Further arguments supporting a decrease in surface insulation of the Indian Ocean

have been reviewed, among others, by Turner and Annamalai (2012).

In general, changes in column-averaged cloud top height might be explained by changes in cloud
345 cover in different altitude layers. When looking at the trend in annual cloud fraction anomalies (not
shown here) a constant decreasing tendency is suggested. This is consistent with long-term changes
derived by other instruments (Stubenrauch et al., 2013) for the same region. In the bottom row of
Fig. 14 the breakups in summer (JJA) and winter (DJF) months are portrayed. Cloud fraction trends
don't exhibit seasonality and are not correlated with the seasonal Indian monsoon because the trends
350 over R1 and R2 are commensurate and have equal sign, which wouldn't be otherwise. This finding
also complies with Norris (2001).

Moreover, the question whether a change in cloud parameters (fraction, optical thickness and top
height) translates into a change in columnar water vapor (due to the assumption in the algorithm
of ghost column under the cloud) has been recently addressed by du Piesanie et al. (2013). They
355 analysed SCIAMACHY water vapor columns and cloud products, generated with the same algo-
rithms of the datasets used in this work (AMC-DOAS, Noël et al. (2004, 2005); SACURA, Rozanov
and Kokhanovsky (2004)). The authors showed that water vapor columns are influenced neither by
changes in cloud fraction nor in cloud optical thickness. The dependence of H₂O total column on
changes in cloud top height has been explored for cases with CF \geq 0.9. Even for these very cloudy
360 scenes, a change in CTH of \approx 450 m over 17 years can't explain the decrease of \approx -0.85 g/cm⁻²
above the Arabian Peninsula for the same time span. This argument rules out algorithmic artifacts
due to the shielding of water vapor by clouds and points to a real process, that presumably takes
place over the northern Indian Ocean.

5 Conclusions

365 For the first time, spaceborne passive measurements of the O₂ A-band have been used to identify
the rate of linear change (trend) in cloud top height anomalies for the period June 1996–May 2012.
The instruments used in this work are hyperspectrally resolved spectrometers (i.e. GOME, SCIA-
MACHY and GOME-2), payloads of polar orbiting platforms which cross the Equator at mid morn-
ing (local time). In particular, it has been seen that the instrumental spatial resolution impacts the
370 calculation of mean values of apparent cloud top height at a monthly sampling. Owing to the sens-
ing time lag among the instruments, the anomaly time series have been calculated for each sensor
separately, once the climatological mean state of the cloud property of interest has been subtracted.
Given that cloud top height, at monthly sampling, does not exhibit any significant persistence in
time, the rate of linear change has been calculated with linear regression methods. The estimation
375 of the distribution of cloud top height trends with bootstrap technique enabled the computation of
confidence intervals and, consequently, the assessment of statistical significance.

First, cloud top height and, to a minor extent, cloud cover have been found to be coupled to ENSO

in a dipole-like fashion across the Pacific Ocean. Patterns of cloud top height autocorrelation of magnitude 0.12–0.13 (vanishing after approximately 6 months) are seen over the Central East Pacific, Atlantic and Indian Oceans, resembling the large scale fluctuations induced by ENSO. Moreover, the hallmark of the cold tongue in the central eastern Pacific is correlated with the strongest decreasing trend in cloud top height. The anomaly time serie of Fig. 3 clearly exhibits two dips, centered at the beginning of year 2000 and 2011. They resemble La Niña’s periodicity, pointing to the ENSO’s role played in the overall decreasing trend in cloud top height.

Second, except for trends derived from the same instrument (i.e. GOME), the comparison with trends previously published in the literature displays differences, which cannot be easily understood within a single explanation. In the case of MODIS and MISR trends (Evan and Norris, 2012), one notes that those instruments not only differ in spatial sampling but also are sensitive to thin clouds, in both solid and liquid phase. These clouds cannot be detected with the hyperspectral sensors in the O₂ A-band used in this work. Thus, the biases shown in Table 3 might point to a recent increasing trend of thin, typically cirrus, clouds. This argument could be also substantiated by the moderate decreasing CTH trend seen in both hemispherical mid-latitudes: since there is evidence on the widening of the Hadley cell (Allen et al., 2012, and references therein), the poleward displacement of cloudiness (Eastman and Warren, 2013) and the concurrent upward shift of high clouds on a synoptic scale (inferred from modeling studies, Singh and O’Gorman (2012)), this picture would imply a systematic thermodynamic phase change in high clouds. As a consequence, a lower occurrence of warm (liquid) clouds in the upper atmospheric layers suggests a higher occurrence of cirrus clouds (undetectable with the O₂ A-band), with the net effect of lowering the mean columnar cloud top height, as seen in the current dataset.

Preliminary analysis of the trends seen over North Africa and Arabian Peninsula suggests a semi-direct aerosol effect on clouds. Natural and anthropogenic soot particless have an impact on cloud top height, modulating insulation and water evaporation over the Indian Ocean. Subsequently, during the winter phase of the Indian Monsoon, transport of moist air masses over the arid Arabian Peninsula exerts a negative feedback and decreases statistical significance of clouds’ vertical displacement. In fact, this effect has been already seen over Europe during the economic, industrial and infrastructural adjustments following the fall of the East Bloc (Devasthale et al., 2005). However, this topic deserves in-depth investigation, which we defer to a later publication.

In general, the representativeness of every trend is affected by the record length under study. 17 years of data still might not be enough to provide any decisive answer to current open questions involving clouds. Therefore, future planned Earth’s observing missions, such as Metop-C and Sentinel-5 precursor with their respective payloads GOME-2 and TROPOMI (Veefkind et al., 2012), will enable the extension of the existing cloud records until and beyond 2022, supporting attribution studies of cloud property changes to natural or human causes and assessments of the cloud feedback sign within the climate system.

415 *Acknowledgements.* The authors are grateful to D. Loyola and W. von Hoyningen-Huene for discussions related
to cloud and aerosol remote sensing. We also acknowledge S. Noël for providing the water vapor dataset, H.
Bovensmann for his expertise in instrumental radiometry and R. Hommel for suggestions on a draft version
of the manuscript. The two anonymous reviewers and B. van Diedenhoven are credited for improving the
manuscript as well. A special thank goes to W. Lotz for sharing his serious programming knowledge, which
420 enabled accurate number crunching. SCIAMACHY and GOME-2 radiances have been made available by the
European Space Agency (ESA) and the European Organisation for the Exploitation of Meteorological Satellites
(EUMETSAT), respectively. GOME radiances and cloud fraction data (for GOME and GOME-2) have been
made available by the German Aerospace Center (DLR) and generated under the auspices of the O3MSAF
project funded by EUMETSAT and national contributions. Generic Mapping Tools (GMT, Wessel and Smith
425 (1998)) and Climate Data Operators (CDO, Schulzweida et al. (2012)) were used for this work. L. Lelli was
supported by the DLR Sentinel-5 precursor project (grant nr. 50EE1247).

References

- Ackerman, A. S., Toon, O. B., Stevens, D. E., Heymsfield, A. J., Ramanathan, V., and Welton, E. J.: Reduction of Tropical Cloudiness by Soot, *Science*, 288, 1042–1047, doi:10.1126/science.288.5468.1042, 2000.
- 430 Allen, R. J., Sherwood, S. C., Norris, J. R., and Zender, C. S.: Recent Northern Hemisphere tropical expansion primarily driven by black carbon and tropospheric ozone, *Nature*, 485, 350–354, doi:10.1038/nature11097, 2012.
- Bjerknes, J.: Atmospheric teleconnections from the Equatorial Pacific, *Monthly Weather Review*, 97, 163–172, doi:10.1175/1520-0493(1969)097<0163:ATFTEP>2.3.CO;2, 1969.
- 435 Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission objectives and measurement modes, *J. Atmos. Sci.*, 56, 127–150, doi:doi:10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2, 1999.
- Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V. V., Ladstätter Weissenmayer, A., Richter, A., DeBeek, R., Hoogen, R., Bramstedt, K., Eichmann, K. U., Eisinger, M., and Perner, D.: The Global Ozone Monitoring Experiment (GOME): Mission Concept and First Scientific Results, *J. Atmos. Sci.*, 56, 151–175, doi:10.1175/1520-0469, 1999.
- 440 Callies, J., Corpaccioli, E., Eisinger, M., Hahne, A., and Lefebvre, A.: GOME-2- Metop’s second-generation sensor for operational ozone monitoring, *ESA bulletin*, 102, 28–36, 2000.
- Clement, A. C., Burgman, R., and Norris, J. R.: Observational and Model Evidence for Positive Low-Level Cloud Feedback, *Science*, 325, 460–464, doi:10.1126/science.1171255, 2009.
- 445 Davies, R. and Molloy, M.: Global cloud height fluctuations measured by MISR on Terra from 2000 to 2010, *Geophys. Res. Lett.*, 39, L03 701, doi:10.1029/2011GL050506, 2012.
- Dessler, A. E.: Cloud variations and the Earth’s energy budget, *Geophys. Res. Lett.*, 38, L19 701, doi:10.1029/2011GL049236, 2011.
- 450 Devasthale, A., Kruger, O., and Graßl, H.: Change in Cloud-Top Temperatures Over Europe, *IEEE Geo. Rem. Sens. Lett.*, 2, 333–336, 2005.
- du Piesanie, A., PETERS, A. J. M., Aben, I., Schrijver, H., Wang, P., and Noël, S.: Validation of two independent retrievals of SCIAMACHY water vapour columns using radiosonde data, *Atmos. Meas. Tech.*, 6, 2925–2940, doi:10.5194/amt-6-2925-2013, 2013.
- 455 Eastman, R. and Warren, S. G.: A 39-Yr Survey of Cloud Changes from Land Stations Worldwide 1971–2009: Long-Term Trends, Relation to Aerosols, and Expansion of the Tropical Belt, *J. Clim.*, 26, 1286–1303, doi:10.1175/JCLI-D-12-00280.1, 2013.
- Efron, B. and Tibshirani, R. J.: *An Introduction to the Bootstrap*, Chapman & Hall, New York, 1993.
- Erlykin, A. D. and Wolfendale, A. W.: Global Cloud Cover and the Earth’s Mean Surface Temperature, *Surveys in Geophysics*, 31, 399–408, doi:10.1007/s10712-010-9098-7, 2010.
- 460 EUMETSAT: GOME-2 PMD Band Definitions 3.0 and PMD Calibration, Tech. Rep. EUM/OPS-EPS/DOC/07/0601, EUMETSAT, Darmstadt, Germany, 2010.
- Evan, A. T. and Norris, J. R.: On global changes in effective cloud height, *Geophys. Res. Lett.*, 39, L19 710, doi:10.1029/2012GL053171, 2012.
- 465 Feingold, G.: On smoke suppression of clouds in Amazonia, *Geophys. Res. Lett.*, 32, L02 804, doi:10.1029/2004GL021369, 2005.

- Koelemeijer, R. B. A., Stammes, P., Hovenier, J. W., and de Haan, J. F.: A fast method for retrieval of cloud parameters using oxygen A band measurements from the Global Ozone Monitoring Experiment, *J. Geophys. Res.*, 106, 3475–3490, doi:10.1029/2000JD900657, 2001.
- 470 Koelemeijer, R. B. A., de Haan, J. F., and Stammes, P.: A database of spectral surface reflectivity in the range 335–772 nm derived from 5.5 years of 0.5 observations, *J. Geophys. Res.*, 108, 4070, doi:10.1029/2002JD002429, <http://www.temis.nl/data/ler.html>, 2003.
- Kokhanovsky, A. A. and Rozanov, V. V.: The physical parameterization of the top-of-atmosphere reflection function for a cloudy atmosphere–underlying surface system: the oxygen A-band case study, *J. Quant. Spectrosc. Ra.*, 85, 35–55, doi:10.1016/S0022-4073(03)00193-6, 2004.
- 475 Kokhanovsky, A. A., Mayer, B., Rozanov, V. V., Wapler, K., Burrows, J. P., and Schumann, U.: The influence of broken cloudiness on cloud top height retrievals using nadir observations of backscattered solar radiation in the oxygen A-band, *J. Quant. Spectrosc. Ra.*, 103, 460 – 477, doi:10.1016/j.jqsrt.2006.06.003, 2007.
- Laken, B., Palle, E., and Miyahara, H.: A Decade of the Moderate Resolution Imaging Spectroradiometer: Is a
480 Solar–Cloud Link Detectable?, *J. Clim.*, 25, 4430–4440, doi:10.1175/JCLI-D-11-00306.1, 2012.
- Lelli, L., Kokhanovsky, A. A., Rozanov, V. V., Vountas, M., Sayer, A. M., and Burrows, J. P.: Seven years of global retrieval of cloud properties using space-borne data of GOME, *Atmos. Meas. Tech.*, 5, 1551–1570, doi:10.5194/amt-5-1551-2012, 2012.
- Loeb, N. G., Kato, S., Su, W., Wong, T., Rose, F. G., Doelling, D. R., Norris, J. R., and Huang, X.: Ad-
485 vances in Understanding Top-of-Atmosphere Radiation Variability from Satellite Observations, *Surveys in Geophysics*, 33, 359–385, doi:10.1007/s10712-012-9175-1, 2012.
- Loyola, D. G. and Ruppert, T.: A new PMD cloud-recognition algorithm for GOME, *ESA Earth Observation Quarterly*, 58, 45–47, 1998.
- Loyola, D. G., Thomas, W., Spurr, R., and Mayer, B.: Global patterns in daytime cloud proper-
490 ties derived from GOME backscatter UV-VIS measurements, *Int. J. Remote Sens.*, 31, 4295–4318, doi:10.1080/01431160903246741, 2010.
- Marchand, R.: Trends in ISCCP, MISR, and MODIS cloud-top-height and optical-depth histograms, *J. Geophys. Res. Atmos.*, 118, 1941–1949, doi:10.1002/jgrd.50207, 2013.
- Marshak, A., Davis, A., Wiscombe, W., and Titov, G.: The verisimilitude of the independent pixel approxi-
495 mation used in cloud remote sensing, *Rem. Sens. Environ.*, 52, 72–78, doi:10.1016/0034-4257(95)00016-T, 1995.
- Mieruch, S., Noël, S., Bovensmann, H., and Burrows, J. P.: Analysis of global water vapour trends from satellite measurements in the visible spectral range, *Atmos. Chem. Phys.*, 8, 491–504, doi:10.5194/acp-8-491-2008, 2008.
- 500 Mudelsee, M.: *Climate Time Series Analysis: Classical Statistical and Bootstrap Methods*, Atmospheric and Oceanographic Sciences Library, Vol. 42, Springer, Dordrecht Heidelberg London New York, doi:10.1007/978-90-481-9482-7, 2010.
- Nakajima, T. and Schulz, M.: What Do We Know about Large-scale Changes of Aerosols, Clouds, and the Radiation Budget?, in: *Clouds in the Perturbed Climate System*, edited by Heintzenberg, J. and Charlson, R. J., Strüngmann Forum Reports, pp. 401–432, MIT Press, Cambridge, Massachusetts, 2009.
- 505 Noël, S., Buchwitz, M., and Burrows, J. P.: First retrieval of global water vapour column amounts from SCIA-

- MACHY measurements, *Atmos. Chem. Phys.*, 4, 111–125, doi:10.5194/acp-4-111-2004, 2004.
- Noël, S., Buchwitz, M., Bovensmann, H., and Burrows, J. P.: Validation of SCIAMACHY AMC-DOAS water vapour columns, *Atmos. Chem. Phys.*, 5, 1835–1841, doi:10.5194/acp-5-1835-2005, 2005.
- 510 Norris, J. R.: Has northern Indian Ocean cloud cover changed due to increasing anthropogenic aerosol?, *Geophys. Res. Lett.*, 28, 3271–3274, doi:10.1029/2001GL013547, 2001.
- Norris, J. R.: Trends in upper-level cloud cover and surface divergence over the tropical Indo-Pacific Ocean between 1952 and 1997, *J. Geophys. Res.*, 110, D21 110, doi:10.1029/2005JD006183, 2005.
- Rozanov, V. V. and Kokhanovsky, A. A.: Semianalytical cloud retrieval algorithm as applied to the cloud top altitude and the cloud geometrical thickness determination from top-of-atmosphere reflectance measurements in the oxygen A band, *J. Geophys. Res.*, 109, 4070, doi:10.1029/2003JD004104, 2004.
- 515 Schulzweida, U., Müller, R., Kornblüh, L., Ansoerge, C., and Quast, R.: CDO - Climate Data Operators - v1.5.9, Center for Marine and Atmospheric Sciences (ZMAW), Max-Planck Institute for Meteorology, University of Hamburg, <https://code.zmaw.de/projects/cdo>, 2012.
- 520 Singh, M. S. and O’Gorman, P. A.: Upward Shift of the Atmospheric General Circulation under Global Warming: Theory and Simulations, *J. Clim.*, 25, 8259–8276, doi:10.1175/JCLI-D-11-00699.1, 2012.
- Solomon, A. and Newman, M.: Reconciling disparate twentieth-century Indo-Pacific ocean temperature trends in the instrumental record, *Nature Climate Change*, 2, 691–699, doi:10.1038/NCLIMATE1591, 2012.
- Stubenrauch, C. J., Rossow, W. B., Kinne, S., Ackerman, S., Cesana, G., Chepfer, H., Getzewich, B., Di Girolamo, L., Guignard, A., Heidinger, A., Maddux, B., Menzel, P., Minnis, P., Pearl, C., Platnick, S., Riedi, J., Sun-Mack, S., Walther, A., Winker, D., Zeng, S., and Zhao, G.: Assessment of Global Cloud Datasets from Satellites: Project and Database Initiated by the GEWEX Radiation Panel, *Bull. Amer. Meteorol. Soc.*, pp. 0003–0007, doi:10.1175/BAMS-D-12-00117, 2013.
- Trenberth, K. E.: The definition of El Niño, *Bull. Amer. Meteorol. Soc.*, 78, 2771–2777, doi:10.1175/1520-0477(1997)078<2771:TDOENO>2.0.CO;2, 1997.
- 530 Turner, A. and Annamalai, H.: Climate change and the South Asian summer monsoon, *Nature Climate Change*, 2, 587–595, doi:10.1038/nclimate1495, 2012.
- U. S. National Centers for Environmental Prediction (NCEP): Oceanic Niño Index (ONI), National Oceanic and Atmospheric Administration (NOAA), <http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/enso.shtml>, (last access: January 2013), 1993.
- 535 van Diedenhoven, B., Hasekamp, O. P., and Aben, I.: Surface pressure retrieval from SCIAMACHY measurements in the O₂ A Band: validation of the measurements and sensitivity on aerosols, *Atmos. Chem. Phys.*, 5, 2109–2120, doi:10.5194/acp-5-2109-2005, 2005.
- Veeffkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., Van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Rem. Sens. Environ.*, 120, 70–83, doi:10.1016/j.rse.2011.09.027, 2012.
- 540 Wagner, T., Beirle, S., Deutschmann, T., Grzegorski, M., and Platt, U.: Dependence of cloud properties derived from spectrally resolved visible satellite observations on surface temperature, *Atmos. Chem. Phys.*, 8, 2299–2312, doi:10.5194/acp-8-2299-2008, 2008.
- 545

Wessel, P. and Smith, W. H. F.: New, improved version of Generic Mapping Tools (GMT) released, Eos Trans. AGU, 79, 579–579, doi:10.1029/98EO00426, 1998.

550 Wilks, D. S.: Statistical Methods in the Atmospheric Sciences, Volume 100, Third Edition (International Geophysics), Academic Press, 2011.

Table 1. Technical specifications of the instruments used in this work. The bottom part of the table describes the Polarization Measuring Devices (PMD).

	GOME	SCIAMACHY	GOME-2
Data availability	1996 – 2011 ^{a,b}	2002 – 2012 ^c	2007 – 2022 ^d
Equator crossing (LT)	10:30 AM	10:00 AM	9:30 AM
Global coverage [days]	3	6	1.5
Spectral coverage [nm]	237 – 794	240 – 2400	237 – 794
Spectral resolution [nm]	0.38	0.44	0.48
Viewing geometry	nadir	nadir, limb, occultation	nadir
Ground pixel size [km ²]	320 × 40	60 × 40	80 × 60
Swath width [km]	≈ 960	≈ 1000	≈ 1920
Polarization	3p	6p	12 (s,p)
Spectral resolution [nm]	100 – 160	55 – 120	2.8 – 40
Spatial resolution [km ²]	40 × 20	30 × 7.5	40 × 5

^aGlobal coverage lost in May 2003. ^bPayload has been switched-off since July 2011. ^cLost contact on April 8, 2012. ^dForeseen extension of GOME-2 records aboard Metop-B/C.

Table 2. Description of the height grid for calculation and analysis of cloud top height time series.

Cloud type	Height range [km]	Bins
Low	0 – 3	1 – 1.5 km, 1.5 – 2 km, 2 – 2.5 km, 2.5 – 3 km
Middle	3 – 7	3 – 5 km, 5 – 7 km
High	7 – 17	7 – 9, 9 – 11 km, 11 – 13 km, 13 – 15 km, 15 – 17 km

Table 3. Overview of trends in cloud height [m decade⁻¹]^a. Trends derived from the GSG dataset are given with (left column) and without (right column) ENSO, that has been filtered masking any data within the Niño 3.4 climate index box 170° W - 120° W, 5° N - 5° S

Base period [year/month]	MISR	MISR corr.	MODIS	GOME	GSG	
03/03 – 11/02	-40.1 ¹	+54.3 ²	+60.9 ²	-	-10.7	+8.4
96/06 – 03/05	-	-	-	-47.9 ^{*,3}	-67.2 [*]	+25.2 [*]
96/06 – 12/05	-	-	-	-	-17.8	-4.9

^aThe GSG trends are recalculated for the length of the referenced dataset and latitude belt $\pm 60^\circ$.

¹Davies and Molloy (2012). ²Evan and Norris (2012). ³Loyola et al. (2010). *Extrapolated to decade.

Table 4. Overview of zonal trends in cloud top height [m yr⁻¹]. ENSO excluded masking any data within the box 170° W - 120° W, 5° N - 5° S. Bootstrap resamples $n = 10^3$. The zonal values are not weighted by the respective land and water abundances.

	Belt		Land + Water	Land	Water
With ENSO	Tropics	5° N - 5° S	-4.34±5.65	-1.56±4.02	-5.15±8.21
		20° N - 20° S	-2.16±2.97	1.83±4.40	-3.39±5.32
	Mid latitude	30° N - 60° N	-2.17±1.52	-2.85±4.23	-1.52±3.68
		30° S - 60° S	-2.71±2.59	-2.70±9.25	-2.71±2.47
Without ENSO	Tropics	5° N - 5° S	-1.80±6.00	-1.43±5.05	-1.99±8.52
		20° N - 20° S	0.53±3.53	5.93±5.33	-1.74±4.36
	Mid latitude	30° N - 60° N	-2.11±3.09	-2.72±4.60	-1.53±3.70
		30° S - 60° S	-2.78±2.54	-3.24±8.77	-2.75±2.35




Fig. 1. Time series of monthly means of cloud top height for global (top) and tropical (bottom) region.




Fig. 2. Relative pixel count as function of height for different instruments for global (left) and tropical (mid) region. (Right) Count difference between GOME and SCIAMACHY. The coarse GOME footprint favors the detection of high clouds over low clouds.




Fig. 3. Monthly means of cloud top height anomaly. The dashed gray vertical lines show the months at which the time series have been merged (June 2003 and May 2008).




Fig. 4. (Upper plot) Autocorrelation function of cloud top height anomalies. Unlagged data ($p = 0$) correlate perfectly with themselves ($r_0 = 1$). r_p decays to almost 0 already at $p = 1$. (Lower plot) Maps of CTH autocorrelation for the first 4 months. ENSO-related patterns are seen over Pacific, Indian and Atlantic Oceans.




Fig. 5. Normalised bootstrap (resamples $n = 10^4$) distributions of trend β for global (left), tropical (mid) regions and Central East Pacific (170° W- 120° W, 5° N- 5° S). The red tails indicate the confidence interval (CI) at 95%. A Gaussian distribution and a cumulative distribution function are plotted in red and green, respectively.




Fig. 6. Normal quantile-quantile plot for global cloud top height anomalies versus the theoretical residual quantiles estimated after regression with Eq. 3 and application of the parameter estimates $\hat{\alpha}$ and $\hat{\beta}$.




Fig. 7. Time series of cloud top height (top) and cloud fraction (bottom) anomaly over the Central East Pacific with the El Niño 3.4 index.




Fig. 8. (Top plot) Time series of cloud top height (black curve) and cloud fraction (purple) anomaly over the Central East Pacific with the El Niño 3.4 index and Pearson's correlation coefficient. (Mid plot) CTH and (bottom plot) CF anomalies subset for low- (blue), mid- (orange), and high-level (red) clouds. Time series smoothed with a 6-month running mean. Correlation coefficients calculated with unsmoothed data.




Fig. 9. Correlation coefficient of CTH and CF with the El Niño 3.4 index as function of height for the 17 years of the GSG dataset. Region considered: Central East Pacific (170° W-120° W, 5° N-5° S).




Fig. 10. ISCCP-like diagrams of cloud optical thickness vs. cloud top height for a warm (left column) and cold (right column) ENSO phase between April 1997 and June 2011 over the Niño 3.4 index box (top row) and the western tropical warm pool (bottom row).




Fig. 11. (Upper plot) Global trend β in CTH anomalies in the latitude belt $\pm 60^\circ$, with (red line) and without (blue line) ENSO region. (Lower plot) The time series are broken up for underlying surface. Abundances of land and water masses between 60°N and 60°S are 0.264 and 0.736, respectively.




Fig. 12. Global map of linear trend β in CTH (left) and standard deviation σ_β (right).




Fig. 13. Global trend β in CTH anomaly, statistically significant at 95% confidence level. Data are gridded onto a mesh of 2° -sided cells.




Fig. 14. Maps of linear trends over North Africa (R1) and Arabian Peninsula (R2) for (top left) CTH anomalies and (top right) columnar water vapor anomalies [g/m²/yr]. The latter is subset for (mid left) summer (JJA) and (mid right) winter (DJF) months. Linear trends for CF anomalies are plotted in the bottom row, for summer (left) and winter (right) months.