



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

On the abundance and source contributions of dicarboxylic acids in size-resolved aerosol particles at continental sites in Central Europe

D. van Pinxteren¹, C. Neusüß², and H. Herrmann¹

¹Leibniz-Institut für Troposphärenforschung (TROPOS), Permoserstr. 15, 04318 Leipzig, Germany

²Hochschule Aalen, Beethovenstr. 1, 73430 Aalen, Germany

Received: 28 November 2013 – Accepted: 29 November 2013 – Published: 9 December 2013

Correspondence to: H. Herrmann (herrmann@tropos.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Dicarboxylic acids (DCAs) are among the most abundant organic compounds observed in atmospheric aerosol particles and have been extensively studied at many places around the world. The importance of the various primary sources and secondary formation pathways discussed in the literature is often difficult to assess from field studies, though. In the present study, a large dataset of size-resolved DCA concentrations from several inland sites in Germany is combined with results from a recently developed approach of statistical back-trajectory analysis and additional data. Principal component analysis is then used to reveal the most important factors governing the abundance of DCAs in different particle size ranges. The two most important sources revealed are (i) photochemical formation in polluted air masses, likely occurring in the gas phase on short timescales (gasSOA), and (ii) secondary reactions in anthropogenically influenced air masses, likely occurring in the aqueous phase on longer timescales (aqSOA). While the first source strongly impacts DCA concentrations mainly in small and large particles, the second one enhances accumulation mode DCAs and is responsible for the bulk of the observed concentrations. Primary sources were found to be minor (sea salt, soil resuspension) or non-existent (biomass burning, traffic). The results can be regarded representative for typical central-european continental conditions.

1 Introduction

Low-molecular weight dicarboxylic acids (LMW-DCAs) are arguably the most abundant compound group typically identified in atmospheric organic aerosol particles. As such they have a strong impact on the properties of organic particles and can play a role in the hygroscopic growth of particles (Choi and Chan, 2002; Prenni et al., 2003), the activation of cloud condensation nuclei (Pradeep Kumar et al., 2003; Booth et al., 2009), and ice formation (Zobrist et al., 2006). Furthermore, they often serve as tracers for the

ACPD

13, 32093–32131, 2013

Source contributions of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



oxidative processing of water-soluble organic aerosol in the atmosphere (Sorooshian et al., 2010; Ervens et al., 2011).

A variety of both primary and secondary sources of LMW-DCAs have been proposed in the literature. Among the primary ones are the rather minor sources cigarette smoke (Rogge et al., 1994) and meat charbroiling emissions (Rogge et al., 1991), as well as the potentially more relevant one fossil fuel combustion (Fraser et al., 1998; Kawamura and Kaplan, 1987). Evidence for the latter one is conflicting, though, with more recent studies neglecting a contribution of traffic to LMW-DCA concentrations in the atmosphere (Rogge et al., 1993a; Huang and Yu, 2007; Stone et al., 2010). DCAs were observed in high concentrations in biomass burning plumes (Jaffrezo et al., 1998; Narukawa et al., 1999; Graham et al., 2002; Gao et al., 2003; Allen et al., 2004; Falkovich et al., 2005; Kundu et al., 2010), although it is unclear to what extent they are directly emitted (Jaffrezo et al., 1998) and/or formed during the processing of the plume (Allen et al., 2004; Gao et al., 2003). In marine influenced air masses, direct emission of DCAs associated with sea salt particles can be a primary source (Matsumoto et al., 1998), as sea water has been shown to contain non-negligible concentrations of DCAs (Tedetti et al., 2006).

In general, secondary formation from precursor gases is believed to be a more important source of LMW-DCAs in the atmosphere (Rogge et al., 1993b; Yao et al., 2002; Kleefeld et al., 2002; Röhrli and Lammel, 2000; Wang and Shooter, 2004; Huang et al., 2005). In field studies, increasing concentrations of organic acids are often observed with increasing age of anthropogenic emission plumes (Veres et al., 2011; van Pinxteren et al., 2009; Sorooshian et al., 2006; Satsumabayashi et al., 1990). In smog chamber studies, LMW-DCAs have been observed as reaction products from anthropogenic precursors such as aromatic compounds (Borras and Tortajada-Genaro, 2012; Sorooshian et al., 2007; Sato et al., 2007; Fisseha et al., 2004) or cyclic alkenes (Hamilton et al., 2006; Gao et al., 2004; Kalberer et al., 2000; Hatakeyama et al., 1985; Hatakeyama et al., 1987) as well as from complex compound mixtures such as diesel exhaust (Samy and Zielinska, 2010). Biogenic precursors have been postu-

Source contributions of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lated to significantly contribute to LMW-DCA concentrations as well (Lim et al., 2005; Carlton et al., 2006; Ervens et al., 2008). In fact, a recent global 3-D modelling study suggested isoprene to be by far the most important precursor of oxalic acid in the atmosphere (70%), with anthropogenic precursors contributing only by about 21% on a global scale (Myriokefalitakis et al., 2011). It was also shown, however, that the model significantly underpredicted observed oxalic acid concentrations especially in urban areas. In the marine atmosphere, LMW-DCA were proposed to be related to the degradation of longer-chain DCA, produced from the oxidation of fatty acids (Kawamura et al., 1996b; Miyazaki et al., 2010; Legrand et al., 2007; Turekian et al., 2003), heterogeneous bromine chemistry (Kawamura et al., 2005; Narukawa et al., 2003a), acetylene and ethene oxidation (Warneck, 2003), and the oxidation of glyoxal and/or glyoxylic acid (Rinaldi et al., 2011; Miyazaki et al., 2010; Narukawa et al., 2003a). In comparison to continental source strengths, however, evidence from field studies suggests such marine sources to be less important (Kerminen et al., 1999; Mochida et al., 2003a; Bardouki et al., 2003; Kawamura and Usukura, 1993; Kleefeld et al., 2002; Matsumoto et al., 1998; Narukawa et al., 2003b; Röhrl and Lammel, 2001; Agarwal et al., 2010; Neusüß et al., 2000).

The large variety of secondary sources in diverse atmospheric regimes is a direct result of LMW-DCAs being late-generation products in the oxidation pathways of primary gaseous precursors. In field studies, it is therefore often difficult to judge on the contribution of the different sources and formation pathways to the observed concentrations. The aim of the present study was to elucidate the factors governing the abundance of LMW-DCAs at typical European continental sites. Size-resolved concentration data from several campaigns were compiled to ensure robust results from a statistical source apportionment approach (principal component analysis, PCA). An attempt was made to discriminate impacts from natural vegetation vs. anthropogenic emissions on the formation of LMW-DCAs by including the results of advanced back-trajectory analysis and further data into the PCA.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Some of the data used in this study have already been published within the context of the respective campaigns (Neusüß et al., 2002; Plewka et al., 2004; Gnauk et al., 2005; Müller et al., 2005; Brüggemann et al., 2009). This refers mainly to inorganic ions and OC/EC concentrations, which are used as indicators for certain sources in the source apportionment. The DCA data used in this study have not been published before, with the exception of 8 out of 20 samples from Melpitz (Plewka et al., 2004) and 3 out of 10 samples from Goldlauter (van Pinxteren et al., 2005; Müller et al., 2005).

2 Materials and methods

2.1 Sampling

Particle sampling was performed within several field campaigns at different locations in Germany between 1997 and 2005. They include the rural sites Melpitz, 45 km north-east of Leipzig (51°32′ N, 12°55′ E, TROPOS research site, 20 samples in October–November 1997, sampling time 8–27 h sample⁻¹) and Falkenberg, 50 km southeast of Berlin (52°10′ N, 14°7′ E, research site of German weather service DWD, 23 samples in July–August 1998, 7–23 h sample⁻¹), the rural site Goldlauter in the Thuringian forest area close to Mt. Schmücke (50°38′ N, 10°45′ E, 10 samples in October 2001 and October 2002, 5–16 h sample⁻¹), an urban background site in Leipzig (51°21′ N, 12°26′ E, TROPOS institute, 30 samples in July 2003–August 2005, 24 h sample⁻¹) and a traffic-impacted site in Dresden (51°4′ N, 13°44′ E, close to Dresden-Neustadt train station, 12 samples in September 2003–August 2004, 24 h sample⁻¹).

Size-resolved particle sampling was done at all sites using 5-stage Berner impactors (Hauke, Austria) with 50 % cut-offs at 0.05, 0.14, 0.42, 1.2, 3.5, and 10 µm aerodynamic diameter (D_p) and a flow rate of 75 L min⁻¹. The impactors were equipped with aluminum foils as impaction substrates, which were heated at 350 °C for at least 2 h (aluminum) to reduce blank levels prior to sampling. During some campaigns, a second impactor was run in parallel and equipped with Tedlar[®] foils which were cleaned

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with 5 % aqueous H_2O_2 . Sampling durations varied between a few hours and 1 day (typically 12 or 24 h), depending on the aims of the respective campaign. Directly after sampling, the impaction substrates were stored at -20°C until analysis, which was usually performed within about a year from the time of sampling.

5 Low pressure impactors can suffer from sampling artifacts such as evaporation of semi-volatiles from the lowest stages or particle bounce-off. To reduce particle bounce, d RH-controlled sampling was performed during some of the campaigns (Melpitz, Falkenberg, Goldlauter) by installing the impactor downstream a bundle of 7 3/4 in. stainless steel tubes (1.5 m length), which were temperature controlled to keep the RH of the
10 the sampling air at approx. $60 \pm 10\%$. Inlet heights were between approx. 4 and 20 m above ground for the different campaigns.

2.2 Measurements

Total particle mass was determined by weighing the impaction substrates before and after sampling on a UMT-2 microbalance (Mettler-Toledo, Switzerland) with a reading
15 precision of $0.1\ \mu\text{g}$ and a standard deviation of ca. 1 %. For RH equilibration, the substrates were stored in a closed box at constant temperature ($20 \pm 2^\circ\text{C}$) and RH ($55 \pm 5\%$) for usually 48 h (at least 12 h). A fraction of the Tedlar[®] foil substrate (or aluminum substrate where only one impactor was run) was extracted in 1–2 mL deionized water by ultrasonication and shaking, while a fraction of the aluminum substrate was
20 used for the determination of organic and elemental carbon (OC/EC).

The aqueous particle extracts were analyzed for inorganic cations by capillary electrophoresis (CE) or ion chromatography (IC), while inorganic anions as well as the DCA were always determined by CE. Details of the respective methods can be found elsewhere (Neusüß et al., 2000; Brüggemann and Rolle, 1998). The applied extraction
25 and analysis method determines the sum of both the dissociated and free acid form of a respective DCA in the particles. To differentiate between the two forms, parallel extractions in organic solvents together with an appropriate analysis method would be required (Yang and Yu, 2008).

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



OC/EC determination was done by a two-step thermographic method using a C-mat 5500 carbon analyzer (Ströhlein, Germany). The temperature for OC volatilization under nitrogen was 590 °C during the early campaigns (Melpitz and Falkenberg), while for all other samples 650 °C was used. EC was determined at 650 °C under oxygen.

5 More details of the method are given elsewhere (Neusüß et al., 2000; Gnauk et al., 2008).

Mixing ratios of the trace gases O₃, NO, NO₂, SO₂, and CO were measured at all sites except Falkenberg using standard trace gas monitors. The data were averaged within the sampling intervals and used in the source apportionment approach to reflect local pollution conditions.

2.3 Back trajectory analysis

In this study a method of back trajectory calculation and GIS analysis to obtain proxy parameters for the impacts of several land cover classes on the sampled air masses has been applied to the dataset of samples from the various campaigns. The method is described in full detail by van Pinxteren et al. (2010) and will only briefly be outlined here. 96 h back trajectories were calculated using HYSPLIT v4.9 (Draxler and Hess, 1998) with input fields obtained from the FNL archive at NOAA's Air Resource Laboratory (<http://ready.arl.noaa.gov/archives.php>). The model was run in the ensemble mode with a starting height of 500 m a.g.l. and offsets of one meteorological grid point in the horizontal dimension and 0.01 sigma units (ca. 250 m) in the vertical dimension, yielding an ensemble of 27 trajectories for all possible offsets in the *x*-, *y*-, and *z*-dimension for a given starting time. Trajectory ensembles were calculated for every full hour during a sampling interval and intersected with land cover data from the Global Land Cover 2000 project of the European Commission Joint Research Centre (GLC2000 database, <http://bioval.jrc.ec.europa.eu/products/glc2000/glc2000.php>) within a database system with GIS functionality (PostgreSQL 9.1 + PostGIS 2.0). The outcome is a dimensionless residence time index (RTI), which – on a relative scale between 0 and 1 – reflects the (weighted) time the sampled air masses have resided above certain land cover

**Source contributions
of dicarboxylic acids**

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



categories (water, natural vegetation, agricultural lands, bare areas, and urban areas). The land cover classes were obtained by reclassifying the original 22 classes of the GLC2000 dataset into the broader categories as given by van Pinxteren et al. (2010). Given that emissions from a given area will be strongly influenced by its land use (e.g. biogenic emissions from areas with natural vegetation, anthropogenic emissions from urban areas), the index can be regarded a measure for potential impacts of such typical emissions from the different land cover categories to the sampled air mass during their transport to the receptor site. The residence time index is calculated in a time-weighted manner to account for a decreasing impact of areas far away from the receptor site (i.e. backward in trajectory time).

In a similar way, the meteorological output of HYSPLIT can be averaged in a weighted manner “along the trajectories” to yield a proxy for the solar flux during the past 4 days of the sampled air mass. Additionally, the HYSPLIT solar flux as well as the HYSPLIT mixing layer depth (modelled from the input meteorological fields) can be averaged at the location of the receptor site over a given sampling interval and the mean length of the trajectories during a sampling interval can be calculated. These parameters proved helpful in a previous application of the method (van Pinxteren et al., 2010) and are used in the present source apportionment study as well.

2.4 Principal component analysis

PCA was performed on the dataset of particulate concentrations, trace gas mixing ratios, and trajectory parameters using the open source statistical software package R (<http://www.r-project.org>). All concentration data (both particulate and gas data) as well as the two solar flux and the mixing height variables were log₁₀-transformed to reduce the skewness in the variables. The skewness results from the approximate lognormal distribution of these variables and can potentially distort the results of a PCA. After log-transformation, the data was mean-centered, scaled to unit variance and the correlation matrix was used for calculation of the principal components, which is necessary if variables on very different scales are to be used. As PCA requires a complete set of

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data, concentrations below the detection limit were set to half the detection limit, while missing data (e.g. due to instrumental problems or samples not analyzed for the full suite of compounds) was replaced by the mean value of the respective parameter. It has to be noted that during the Melpitz campaign only a fraction of the samples (6 out of 20) was analyzed for the full suite of DCAs, while oxalate was analyzed from all of these samples. By replacing the missing data with their mean values they are set to the non-informative point at the origin of the mean-centered data and do thus not influence the outcome of the PCA. The number of factors to extract was defined by examining the scree plot of eigenvalues vs. number of PCs (Cattell, 1966) and by comparing the magnitudes of the obtained eigenvalues with those obtained from a random data matrix of the same size as the original (so-called parallel analysis; Horn, 1965). Varimax rotation was applied to the extracted principal components to result in rotated components with easier-to-interpret component loadings.

3 Results and discussion

3.1 PM₁₀ concentrations and size distributions of DCAs

In Table 1 the concentrations of dicarboxylic acids obtained in this work are summarized. Literature data from comparable European rural and urban sites are given for comparison in Table 1. Generally, the mean DCA concentrations obtained in this study do not differ more than a factor of about 2 between the 5 different sites. The urban sites do not necessarily show higher concentrations than the rural sites. Highest concentrations are always observed for oxalic followed by malonic acid. Succinic, glutaric (incl. 2 unresolved isomers; Neusüß et al., 2000), and malic acid often show comparable concentration ranges, with the exception of the summer-time Falkenberg data, where glutaric and malic acid were considerably higher than succinic acid. Tartronic and tartaric acid are observed at rather low concentrations of only a few ng m⁻³, if at all.

A notable exception is again the Falkenberg data, where tartronic acid was observed at rather high concentrations.

Compared to the literature data, this work's DCA concentrations at urban sites usually agree within a factor of 2 with similar sites in Europe. Oxalic acid determined by Röhrl and Lammel (2001) during summer in Leipzig and Eichstädt somewhat exceeds the concentration observed at the other sites, which might be due to the sampling of total suspended matter (TSP) and/or – as indicated by the authors – possible positive sampling artefacts for oxalic acid. Concentrations at rural sites agree to a similar extent, with the exceptions of rather high TSP oxalic and malonic acid in Falkenberg (Röhrl and Lammel, 2001) and PM_{2.5} oxalic acid at K-pusztá in the central plain of Hungaria (Legrand et al., 2007). Unusually low concentrations of C3-C5 DCAs were recently reported for a rural site in Sweden (Hyder et al., 2012), possibly related to the choice of extraction solvent and/or derivatization agent for GC/MS analysis applied in this study. Yang and Yu (2008) demonstrated that the combination of solvent extraction and trimethylsilylation GC/MS recovers only a fraction of total malonic acid (including malonate) concentrations in PM_{2.5} samples, while – as reported by Hyder et al. (2012) as well – oxalic acid was not detected at all.

The size distributions of DCAs are given in Fig. 1 as averages over all sites. Highest concentrations for all DCAs were observed in accumulation mode particles ($D_p = 0.14$ – $1.2 \mu\text{m}$) with significantly lower concentrations in ultrafine ($D_p = 0.05$ – $0.14 \mu\text{m}$) and coarse ($D_p = 1.2$ – $10 \mu\text{m}$) particles. Such size distributions are typical for European continental DCA measurements and have been observed in similar ways before (Kerminen et al., 2000; Meszaros et al., 1997; Müller et al., 2005; Legrand et al., 2007).

3.2 Back-trajectory analysis

The residence time indices as main results from the statistical back-trajectory analysis are shown in Fig. 2. As expected from the location of the sites (in-land continental), the sampled air masses had higher (time-weighted) residence times from the two main continental land cover classes agriculture and natural vegetation as compared to the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



residence time above marine areas (water). Urban areas make up only a tiny fraction of continental surface area, which is why the values of this index are much smaller. The indices show a large variability, indicating that very different air masses were sampled in the various campaigns. Further output parameters from the back-trajectory analysis as described above are given in Fig. 2 as well. In agreement with the campaign season, solar flux and mixing depth were quite different at the different sites with higher values for the Falkenberg summer measurements, lower ones for the fall campaigns in Melpitz and Goldlauter and highly variable values for the year-round measurements in Dresden and Leipzig.

3.3 Source apportionment

To assess the latent factors controlling the abundance of DCAs in different particle sizes, PCA was performed on the dataset of size-resolved DCA concentrations. The following further particle constituents were included in the analysis in order to aid the interpretation of the resolved principal components: EC (primary, anthropogenic combustion or biomass burning), OC (both primary and secondary, both anthropogenic and biogenic), sulfate (mainly secondary, anthropogenic combustion), potassium (primary, biomass burning), sodium (primary, sea salt and/or soil), and calcium (primary, dust/soil). Due to missing concentration data for sodium and calcium on the smallest particles (impactor stages 1 and 2, all data below detection limit), concentrations from impactor stages 3–5 only were included for these species. Additionally, trace gas mixing ratios of NO_x , CO, and SO_2 (anthropogenic combustion) and O_3 (photochemically formed) were included. All the back-trajectory parameters were included as well with the exception of the residence time indices for bare areas (negligible values for sampled air masses) and water (simply anti-correlated to the continental indices).

In PCA, the number of PCs to extract is a critical (and to some extent subjective) decision which can strongly affect the interpretability of the result. Many different procedures exist to aid this decision (Jolliffe, 2002). One of the most common ones is the scree plot of successive eigenvalues for all calculated PCs, which is shown in Fig. 3.

Source contributions of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As can be seen, the rate of decline is fast for the first 7 PCs and then clearly levels off. The “elbow” of the plot (i.e. PC 7) is often considered to indicate the number of components to extract (Jolliffe, 2002). On the other hand, the eigenvalue of PC 7 is already significantly lower than that obtained from a random matrix of univariate normal data (“simulated data” in Fig. 3) and from random samples (randomized across rows) of the original data (“resampled data” in Fig. 3). Thus, in the final solution the first 6 PCs were chosen to be subjected to Varimax rotation. The “correctness” of the 6 factor solution was, however, verified by varying the number of extracted PCs between 6 and 7, as will be discussed below. In Table 2 the loadings of the variables on the 6 rotated components are shown, which basically describe the correlation of the variable with the component. Note that for better readability loadings with $|x| < 0.2$ were considered insignificant and removed, while loadings with $|x| \geq 0.6$ were considered “high” and printed bold.

3.3.1 PC 1: anthropogenically influenced gasSOA

The first rotated component (PC 1) explains the largest part of the total variance in the dataset (19%) and has significant loadings for all DCAs in basically all particle size ranges. Highest loadings are observed for DCAs in smallest ($D_p = 0.05\text{--}0.14\ \mu\text{m}$) and largest ($D_p = 3.5\text{--}10\ \mu\text{m}$) particles, while they tend to decrease towards accumulation mode particles ($D_p = 0.42\text{--}1.2\ \mu\text{m}$) (even non-significant for succinic acid). The hydroxy-DCA malic acid shows a somewhat different behavior with continuously decreasing loadings from smallest towards largest particle sizes. High loadings are also observed for the two solar flux variables and for O_3 . PC 1 likely represents photochemical formation of DCAs in the gas phase (gasSOA) during days with high intensity of solar radiation. The gas phase formation is inferred (i) from the typically low ambient relative humidity (RH) and cloud cover during such days (RH during sampling is strongly anti-correlated to the solar flux parameters in our dataset), leading to low amounts of liquid water in the atmosphere and (ii) from the strongly increasing loadings towards smaller particle sizes, reflecting the higher surface area available for condensation. The

**Source contributions
of dicarboxylic acids**

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high loading of sulfate in very small particles is consistent with PC1 representing a gas-phase source, as gaseous formation and subsequent condensation of sulfuric acid is a well-known process under the influence of strong solar radiation (Hamed et al., 2010). The high loadings of DCAs in coarse particle size ranges might be explained by their repartitioning onto crustal material and/or mineral dust particles. Calcium – a typical crust component – has a quite high loading to PC 1 for the largest particle size range as well. Due to their less acidic character (Scheinhardt et al., 2013), the diacids might strongly condense to such particles despite the lower surface area they provide. Such a preferential partitioning of DCA has already been suggested previously (Kerminen et al., 2001). The different trend of malic acid loadings with respect to coarse particles is consistent with the lower vapor pressures of hydroxyl-DCAs and thus less favorable evaporation from smaller particles. Heterogeneous photochemical reactions on dust could be another plausible explanation for the observed coarse mode loadings. Regarding the timescale of the photochemical production of DCA in PC 1, the very similar magnitude of the two solar flux loadings indicate a rather local and thus quick formation. If major fractions of particulate DCAs would have been formed during previous days already and just been transported to the site, one could expect the solar flux along the back trajectory loading to be higher than the local solar flux loading. Even though this might not be a very strong indication, it is consistent with a study on highly time-resolved oxalate concentrations, showing a strong diurnal profile during summer and thus a rather fast formation (Fisseha et al., 2006). Among the residence time indices, urban areas have the highest loading, followed by a minor loading of agricultural areas, and a non-significant loading of natural vegetation, indicating a significant influence of anthropogenic emissions to this factor. Overall, PC 1 likely represents secondary organic aerosol formed in the gas phase of polluted air masses (gasSOA) as an important source of DCAs.

3.3.2 PC 2: anthropogenically influenced aqSOA

The second rotated component (PC 2) explains another 15% of the total variance. Moderate to high loadings on PC 2 were found for all DCAs in the particle size range $D_p = 0.14\text{--}3.5\ \mu\text{m}$ (impactor stages 2 to 4) with maximum loadings in accumulation mode particles with $D_p = 0.42\text{--}1.2\ \mu\text{m}$. Similar loading patterns with highest values in medium-sized particles were observed for OC, EC, sulfate, and potassium. Among the residence time indices, agricultural areas, urban areas, and natural vegetation show high, moderate and small loadings, respectively. The mean trajectory length is anticorrelated to the component. PC 2 likely represents aged air masses with long residence times above continental areas and a strong input of anthropogenic emissions including biomass burning. The high loading of agricultural areas might reflect the impact of sources located outside urban areas, such as industrial facilities, roads, and power plants. The moderate loading of SO_2 , whose main source in Europe is energy production (European Environment Agency, 2010), seems to support especially the latter suggestion. The solar flux parameters do not show significant loadings on PC 2, indicating that the formation of DCAs in this factor is not primarily driven by strong radiation. Together with the regional character of the source (highest loadings in accumulation mode particles), it can be assumed that DCA production takes place on a slower timescale as compared to PC 1. In principal, contributions from primary sources to PC 2 cannot be excluded, as EC and – to a lesser extent – potassium correlate with the factor as well. However, the two primary sources which are most often discussed in the context of DCA emissions (traffic and biomass burning) are resolved by the model in PCs 3 and 4. PC 2 might thus represent the SOA from traffic and/or biomass burning, while PCs 3 and 4 represent their primary components. The similarity between sulfate and DCA loadings on PC 2 implies a high correlation of their concentrations in this source. Good correlation of sulfate with oxalate (and sometimes other DCAs) has often been reported from field studies (Sorooshian et al., 2006; Hsieh et al., 2007; Yu et al., 2005; Miyazaki et al., 2009; Hsieh et al., 2008) and can be taken as an indication of aqueous

Source contributions of dicarboxylic acids

D. van Pinxteren et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



phase formation of DCAs (Sorooshian et al., 2006; Yu et al., 2005) due to aqueous SO_2 oxidation being the dominant source of sulfate in the atmosphere (Yu et al., 2005). From size-resolved DCA measurements, a dominant “droplet mode” with D_p ca. 0.7–0.9 μm as opposed to a less abundant “condensation mode” with D_p ca. 0.2–0.4 μm can be taken as evidence for aqueous phase formation as well (Lan et al., 2011; Hsieh et al., 2009; Yao et al., 2003), in analogy to earlier discussions on the abundance of sulfate in these two particle modes (Hering and Friedlander, 1982; John et al., 1990; Meng and Seinfeld, 1994). Even though more direct evidence of aqueous phase formation would be desirable (e.g. the concentrations of aqueous phase oxidants or the amount of available particle and/or cloud water), it can be concluded that aqueous phase formation of DCAs likely plays an important role in the source resolved by PC 2 and it can thus be attributed to aqueous phase SOA (aqASOA) with a significant anthropogenic influence on either the precursors and/or the oxidant chemistry.

3.3.3 PC 3: traffic

The third rotated component (PC 3) explains 10 % of the variance and has high positive loadings for NO_x and CO. EC shows a high loading in ultrafine particles ($D_p = 0.05$ –0.14) and moderate and continuously decreasing loadings in larger particle size fractions, while the solar flux parameters and the mixing depth at the receptor site have negative loadings. PC 3 can clearly be attributed to a traffic source, presumable in winter time conditions, where pollution can build up quickly due to less efficient boundary layer mixing processes. All significant DCA loadings on PC 3 are negative, thus strongly implying that primary traffic emissions are not a significant source of DCAs in this dataset.

3.3.4 PC 4: biomass burning

The fourth rotated component (PC 4) explains another 9 % of the total data variance and contains mainly potassium with high loadings in all particle size ranges. In addi-

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion, sulfate, sodium, and calcium show significant loadings within some size ranges and SO_2 has a rather strong loading as well. As fine particle potassium is a classical tracer of biomass burning (Khalil and Rasmussen, 2003), PC 4 is assumed to represent a primary biomass burning source. No DCA variable loads significantly on this factor, which is why biomass burning is unlikely to be a primary source of DCAs in the present study.

3.3.5 PC 5: sea salt

The fifth rotated component (PC 5) explains 8% of total variance. High loadings of coarse mode sodium together with moderate loadings of coarse mode sulfate are observed. The mean trajectory length shows a moderate positive loading, while the residence time indices for natural vegetation and agriculture are anticorrelated to the factor. PC 5 can be attributed to (processed) sea salt. Among the DCAs, only oxalate in coarse mode particles shows small loadings, while other DCAs have either insignificant or negative loadings. Sea salt might thus contribute to some small extent to coarse mode oxalate concentrations at continental sites in Germany. A sea salt mode of DCA is usually observed in marine regions (Neusüß et al., 2000; Kerminen et al., 1999, 2000; Mochida et al., 2003b). Whether the oxalate measured at the inland sites of this study indeed originates from sea salt or from the processing of these particles on their way to the receptor sites (e.g. by condensation to less acidic sea salt particles) cannot be concluded from the data, though.

3.3.6 PC 6: soil

The sixth rotated component (PC 6) explains 7% of the data variance and contains moderate to high loadings of OC in all particle size ranges and of sulfate and calcium in coarse mode particles. EC shows significant loadings in small and large particles, while minor loadings are observed for a few other variables. Based on this loadings pattern the attribution of a source to this factor is less evident than for the previous

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



components. Mainly due to the loadings of calcium, sulfate, and OC, PC 6 is tentatively interpreted as a primary soil source. The EC loadings might then result from complex high-molecular-weight soil organic matter (humic substances), which does likely not fully evaporate at 650 °C and is thus incorrectly attributed to the EC fraction of total particulate carbon in our method of OC/EC determination. Among the DCAs, minor loadings of coarse mode oxalate and malonate are observed, while coarse mode succinic acid shows moderate loadings to this factor. Organic acids, including the DCAs of this study, are known to be important constituents of soil organic matter (Jones, 1998), thus resuspended soil in coarse mode particles can be assumed to contribute to coarse mode DCA concentrations.

As discussed above, a 7 factor solution for the PCA could in principle be justified from the scree plot as well. The main difference to the 6 factor solution is that the loadings of the original PC 6 are then distributed between the new PC 6 and PC 7, i.e. PC 6 seems to be split into two factors. One factor tends to hold the loadings on impactor stage 4 ($D_p = 1.2\text{--}3.5\ \mu\text{m}$), while the other tends to take over the stage 5 ($D_p = 3.5\text{--}10\ \mu\text{m}$) loadings to some extent. The variables for which significant loadings are observed for the new PC 6 and PC 7 are the same as for the original PC 6 discussed here. The interpretability of the new PC 6 and PC 7 is even more ambiguous than for the original PC 6. This is why the 7 factor solution has been considered to resolve two artificial sources instead of one real one and has been discarded.

3.4 Discussion of main DCA sources

The results of the PCA show that the major part of the variance of DCAs in different particle size ranges can be attributed to two main sources: (1) Fast photochemical formation during intense radiation days, likely due to gas phase oxidation of precursors in anthropogenically influenced air masses (gasSOA), and (2) slower, but presumably more continuous secondary formation in anthropogenically polluted air masses, possibly with an important contribution of aqueous phase oxidation processes (aqSOA).

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Much smaller contributions to the abundance of LMW-DCAs can be expected from primary biomass burning emissions, (processed) sea salt, and primary soil emissions.

While the importance of secondary processes in the formation of LMW-DCAs is well-established, they are typically assumed to be products of aqueous phase reactions (Ervens et al., 2011; Kanakidou et al., 2005). From the results of this study, however, a significant contribution from gas-phase reactions seems possible (PC 1) under certain conditions. For the 10 % of samples which have the highest scores on PC 1 (i.e. the days with presumably the highest impact of gasSOA) the sum of impactor stage 1 and 2 ($D_p = 0.05\text{--}0.42\ \mu\text{m}$) concentrations for a given DCA typically makes up 45–55 % of its total PM_1 and 30–55 % of its total PM_{10} concentration. Making the (very crude) assumption that stage 1 and stage 2 concentrations represent DCAs formed solely from gas phase oxidation processes, these results would indicate that roughly up to one third to one half of total DCA concentrations can be attributed to gas phase sources under appropriate conditions. As condensed phase reactions can contribute to DCA concentrations in these particle sizes as well, the contribution of gas phase sources will likely be lower, though. To the best of the authors knowledge, the formation of LMW-DCAs is at present not included in comprehensive atmospheric gas phase chemistry models. For example, none of the DCAs determined in this study is incorporated in the Master Chemical Mechanism v3.2 (<http://mcm.leeds.ac.uk/MCM>). Two possibilities of gas-phase formation of organic acids are well-known, though: (i) ozonolysis of unsaturated compounds (see Orzechowska and Paulson (2005) and references therein), and (ii) the reactions of acylperoxy radicals with HO_2 radicals (see Calvert et al., 2011, and references therein). Ozonolysis of semi-volatile unsaturated fatty acids has since long been suggested as a possible source of LMW-DCAs in the atmosphere (Kawamura et al., 1996a) and DCAs have frequently been observed in the aerosol phase in chamber studies of unsaturated hydrocarbons (Borras and Tortajada-Genaro, 2012; Sorooshian et al., 2007; Sato et al., 2007; Fisseha et al., 2004; Hamilton et al., 2006; Gao et al., 2004; Kalberer et al., 2000; Hatakeyama et al., 1985; Hatakeyama et al., 1987). Depending on the position of the C-C double bond in the monocarboxylic acid

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



precursor, the C2 to C5 DCAs observed in this study could easily result from such reactions. Ozone has recently been suggested to be an important driver in the formation of carboxylic acid containing SOA (Liu et al., 2011) and it is correlated with PC 1 in our study as well. The oxidation of ω -oxo-monocarboxylic acids to DCAs has also been proposed in the past (Kawamura et al., 1996a). This reaction would proceed via abstraction of the aldehydic H-Atom (e.g. by OH), addition of molecular oxygen and reaction of the formed acylperoxy radical with HO₂ to result in the corresponding DCA. The relevance of these qualitatively feasible formation pathways, however, depends on their competition with the manifold other reaction channels in the case of ozonolysis (Orzechowska and Paulson, 2005) and – for continental conditions – mainly with the reaction of acylperoxy radicals with NO, leading to chain-shortened aldehydes (Calvert et al., 2011). In general, these pathways can be expected to be of minor importance only, although kinetic data product studies of relevant precursors (unsaturated or oxygenated acids) is very limited (Calvert et al., 2011). If the concentrations of precursors (and oxidants) are high enough, however, they might significantly contribute to the observed LMW-DCA concentrations even in low yields.

Another possible gas-phase formation pathway of carboxylic acids could be the reaction of enols with OH. Recently, the common assumption of keto-enol tautomerization of gas-phase aldehydes and ketones being negligible in the atmosphere has been challenged by both theoretical and experimental studies (Andrews et al., 2012; Clubb et al., 2012; da Silva, 2010). It was suggested that a significant proportion of carbonyl compounds might exist in their (highly reactive) enol form. Addition of OH to the C-C double bond then results in a carboxylic acid to be formed, which could account for the discrepancies between typical modeled vs. measured budgets of organic acids in the atmosphere (Archibald et al., 2007). This could hypothetically be another pathway for the gas-phase oxidation of keto-acids to LMW-DCAs. Naturally, much more thermodynamic and kinetic data and its implementation into models is needed to judge on the relevance of such a formation pathway.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Despite a potential relevance of gas-phase processes (PC 1), in terms of total DCA concentrations, PC 2 likely represents the most important source, as highest concentrations of DCAs are typically observed on impactor stages 2 and 3 ($D_p = 0.14\text{--}1.2\ \mu\text{m}$, see Fig. 1). This is in-line with the acknowledged importance of aqueous phase oxidation processes on the formation of LMW-DCAs in the atmosphere (Ervens et al., 2011; Tilgner and Herrmann, 2010). Water-soluble products of VOC oxidation can readily be taken up by deliquesced particles and cloud droplets. Especially by the hydration of aldehyde groups to gem-diols, this uptake opens a reaction channel towards carboxylic acids which is not available in the gas phase.

As outlined above (introduction), the precursors of DCAs are manifold and can originate from both continental and marine as well as from anthropogenic and biogenic sources. For the sampling sites of the present study, the results of back trajectory analysis suggest that continental sources dominate (negative correlation with trajectory length) and indicate an influence of anthropogenic activities on the DCA concentrations for both PC 1 (correlation with urban areas RTI) and PC 2 (correlation with agriculture and urban areas RTI). It has to be noted that correlations with primary anthropogenic trace gases are weak or even absent in accordance with the secondary nature of the two main sources, which implies highest concentrations away from direct sources (dispersion/dilution of trace gases). These results might indicate the importance of anthropogenic precursors on DCA concentrations. There is, however, evidence from several field studies on SOA formation that despite a strong anthropogenic influence to the studied air masses (typically deduced from correlations with anthropogenic tracers), a dominant fraction of carbonaceous material is composed of modern carbon (in contrast to fossil carbon) and thus originates from biogenic sources (El Haddad et al., 2013; Worton et al., 2011; Weber et al., 2007). The coupling of an anthropogenic and biogenic component in sampled air masses has been suggested to potentially enhance biogenic SOA formation, e.g. through higher oxidant concentrations due to higher NO_x and/or higher partitioning of biogenic SOA due to higher concentrations of (primary or

secondary) anthropogenic organic aerosol (Carlton et al., 2010). Thus, in the absence of additional data, firm conclusions on the nature of their VOCs cannot easily be drawn.

4 Conclusions

A relatively large set of size-resolved samples of aerosol particles has been used to study the abundance of LMW-DCA in different particle size ranges at several continental sites in Germany. PCA was used to qualitatively distinguish between different sources and formation pathways. Together with additional data from back trajectory analysis and trace gas measurements, two main sources of DCA could be identified: photochemical formation in the gas phase, strongly affecting concentrations of small and large particles, and secondary formation via aqueous phase processes, leading to increased concentrations in accumulation mode particles mainly. For both sources, an anthropogenic influence is apparent from the back trajectory analysis, either by providing the precursors for DCA formation, or by triggering enhanced biogenic SOA formation e.g. through increased oxidant levels. Compared to the secondary sources, primary emissions of DCAs from traffic, biomass burning, sea salt, and soil are considered to contribute much less or even nothing at all to the concentrations of DCAs at the sites studied. The results of this study suggest that beyond the well-known importance of aqueous phase formation, gas-phase formation of LMW-DCAs can be important under conditions of high photochemistry in anthropogenically influenced air masses as well.

Acknowledgements. Financial support for the campaigns was granted by the German Federal Ministry of Education and Research (BMBF) as part of the projects 07AF102A/1 (MINT, Melpitz) and 07AF302/6 (LACE, Falkenberg) within the German AFS framework and as part of the project 07ATF01 (FEBUKO, Goldlauter) within the AFO2000 framework. Financial support for the measurements in Dresden and Leipzig was received from the Saxon State Ministry of the Environment and Agriculture and the Forschungsvereinigung Automobiltechnik e.V., respectively.

Source contributions of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Agarwal, S., Aggarwal, S. G., Okuzawa, K., and Kawamura, K.: Size distributions of dicarboxylic acids, ketoacids, α -dicarbonyls, sugars, WSOC, OC, EC and inorganic ions in atmospheric particles over Northern Japan: implication for long-range transport of Siberian biomass burning and East Asian polluted aerosols, *Atmos. Chem. Phys.*, 10, 5839–5858, doi:10.5194/acp-10-5839-2010, 2010.
- Allen, A. G., Cardoso, A. A., and da Rocha, G. O.: Influence of sugar cane burning on aerosol soluble ion composition in Southeastern Brazil, *Atmos. Environ.*, 38, 5025–5038, 2004.
- Andrews, D. U., Heazlewood, B. R., Maccarone, A. T., Conroy, T., Payne, R. J., Jordan, M. J. T., and Kable, S. H.: Photo-tautomerization of acetaldehyde to vinyl alcohol: a potential route to tropospheric acids, *Science*, 337, 1203–1206, doi:10.1126/science.1220712, 2012.
- Archibald, A. T., McGillen, M. R., Taatjes, C. A., Percival, C. J., and Shallcross, D. E.: Atmospheric transformation of enols: a potential secondary source of carboxylic acids in the urban troposphere, *Geophys. Res. Lett.*, 34, L21801, doi:10.1029/2007gl031032, 2007.
- Bardouki, H., Liakakou, H., Economou, C., Sciare, J., Smolík, J., Zdímal, V., Eleftheriadis, K., Lazaridis, M., Dye, C., and Mihalopoulos, N.: Chemical composition of size-resolved atmospheric aerosols in the eastern Mediterranean during summer and winter, *Atmos. Environ.*, 37, 195–208, 2003.
- Booth, A. M., Topping, D. O., McFiggans, G., and Percival, C. J.: Surface tension of mixed inorganic and dicarboxylic acid aqueous solutions at 298.15 K and their importance for cloud activation predictions, *Phys. Chem. Chem. Phys.*, 11, 8021–8028, doi:10.1039/B906849j, 2009.
- Borras, E. and Tortajada-Genaro, L. A.: Secondary organic aerosol formation from the photo-oxidation of benzene, *Atmos. Environ.*, 47, 154–163, doi:10.1016/j.atmosenv.2011.11.020, 2012.
- Brüggemann, E. and Rolle, W.: Changes of some components of precipitation in East Germany after the unification, *Water Air Soil Poll.*, 107, 1–23, 1998.
- Brüggemann, E., Gerwig, H., Gnauk, T., Müller, K., and Herrmann, H.: Influence of seasons, air mass origin and day of the week on size-segregated chemical composition of aerosol particles at a kerbside, *Atmos. Environ.*, 43, 2456–2463, doi:10.1016/j.atmosenv.2009.01.054, 2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Calvert, J. G., Mellouki, A., Orlando, J. J., Pilling, M. J., and Wallington, T. J.: The Mechanisms of Atmospheric Oxidation of the Oxygenates, Oxford University Press, New York, 2011.
- Carlton, A. G., Turpin, B. J., Lim, H. J., Altieri, K. E., and Seitzinger, S.: Link between isoprene and secondary organic aerosol (SOA): pyruvic acid oxidation yields low volatility organic acids in clouds, *Geophys. Res. Lett.*, 33, L06822, doi:10.1029/2005gl025374, 2006.
- Carlton, A. G., Pinder, R. W., Bhave, P. V., and Pouliot, G. A.: To what extent can biogenic SOA be controlled?, *Environ. Sci. Technol.*, 44, 3376–3380, doi:10.1021/Es903506b, 2010.
- Cattell, R. B.: The scree test for the number of factors, *Multivar. Behav. Res.*, 1, 245–276, 1966.
- Choi, M. Y. and Chan, C. K.: The effects of organic species on the hygroscopic behaviors of inorganic aerosols, *Environ. Sci. Technol.*, 36, 2422–2428, 2002.
- Clubb, A. E., Jordan, M. J. T., Kable, S. H., and Osborn, D. L.: Phototautomerization of acetaldehyde to vinyl alcohol: a primary process in UV-irradiated acetaldehyde from 295 to 335 nm, *J. Phys. Chem. Lett.*, 3, 3522–3526, doi:10.1021/jz301701x, 2012.
- da Silva, G.: Carboxylic acid catalyzed keto-enol tautomerizations in the gas phase, *Angew. Chem. Int. Ed.*, 49, 7523–7525, doi:10.1002/anie.201003530, 2010.
- Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT_4 modeling system of trajectories, dispersion, and deposition, *Aust. Meteorol. Mag.*, 47, 295–308, 1998.
- El Haddad, I., D'Anna, B., Temime-Roussel, B., Nicolas, M., Boreave, A., Favez, O., Voisin, D., Sciare, J., George, C., Jaffrezo, J.-L., Wortham, H., and Marchand, N.: Towards a better understanding of the origins, chemical composition and aging of oxygenated organic aerosols: case study of a Mediterranean industrialized environment, Marseille, *Atmos. Chem. Phys.*, 13, 7875–7894, doi:10.5194/acp-13-7875-2013, 2013.
- Ervens, B., Carlton, A. G., Turpin, B. J., Altieri, K. E., Kreidenweis, S. M., and Feingold, G.: Secondary organic aerosol yields from cloud-processing of isoprene oxidation products, *Geophys. Res. Lett.*, 35, L02816, doi:10.1029/2007gl031828, 2008.
- Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069–11102, doi:10.5194/acp-11-11069-2011, 2011.
- European Environment Agency: Sector share of sulphur oxides emissions – 2010, available at: http://www.eea.europa.eu/data-and-maps/figures/ds_resolveuid/4b414b0cfe27461b8ab75b8a455c1e8e (last access: 10 October 2013), 2010.
- Falkovich, A. H., Graber, E. R., Schkolnik, G., Rudich, Y., Maenhaut, W., and Artaxo, P.: Low molecular weight organic acids in aerosol particles from Rondônia, Brazil, dur-

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing the biomass-burning, transition and wet periods, *Atmos. Chem. Phys.*, 5, 781–797, doi:10.5194/acp-5-781-2005, 2005.

Fisseha, R., Dommen, J., Sax, M., Paulsen, D., Kalberer, M., Maurer, R., Hofler, F., Weingartner, E., and Baltensperger, U.: Identification of organic organic aerosol and the acids in secondary corresponding gas phase from chamber experiments, *Anal. Chem.*, 76, 6535–6540, 2004.

Fisseha, R., Dommen, J., Gaeggeler, K., Weingartner, E., Samburova, V., Kalberer, M., and Baltensperger, U.: Online gas and aerosol measurement of water soluble carboxylic acids in Zurich, *J. Geophys. Res.-Atmos.*, 111, D12316, doi:10.1029/2005jd006782, 2006.

Fraser, M. P., Cass, G. R., and Simoneit, B. R. T.: Gas-phase and particle-phase organic compounds emitted from motor vehicle traffic in a Los Angeles roadway tunnel, *Environ. Sci. Technol.*, 32, 2051–2060, 1998.

Gao, S., Hegg, D. A., Hobbs, P. V., Kirchstetter, T. W., Magi, B. I., and Sadilek, M.: Water-soluble organic components in aerosols associated with savanna fires in southern Africa: identification, evolution, and distribution, *J. Geophys. Res.-Atmos.*, 108, 8491, doi:10.1029/2002jd002324, 2003.

Gao, S., Keywood, M., Ng, N. L., Surratt, J., Varutbangkul, V., Bahreini, R., Flagan, R. C., and Seinfeld, J. H.: Low-molecular-weight and oligomeric components in secondary organic aerosol from the ozonolysis of cycloalkenes and alpha-pinene, *J. Phys. Chem. A*, 108, 10147–10164, 2004.

Gnauk, T., Brüggemann, E., Müller, K., Chemnitzer, R., Rüd, C., Galgon, D., Wiedensohler, A., Acker, K., Auel, R., Wieprecht, W., Möller, D., Jaeschke, W., and Herrmann, H.: Aerosol characterisation at the FEBUKO upwind station Goldlauter (I): particle mass, main ionic components, OCEC, and mass closure, *Atmos. Environ.*, 39, 4209–4218, 2005.

Gnauk, T., Muller, K., van Pinxteren, D., He, L. Y., Niu, Y. W., Hu, M., and Herrmann, H.: Size-segregated particulate chemical composition in Xinken, Pearl River Delta, China: OC/EC and organic compounds, *Atmos. Environ.*, 42, 6296–6309, doi:10.1016/j.atmosenv.2008.05.001, 2008.

Gómez-González, Y., Wang, W., Vermeylen, R., Chi, X., Neiryneck, J., Janssens, I. A., Maenhaut, W., and Claeys, M.: Chemical characterisation of atmospheric aerosols during a 2007 summer field campaign at Brasschaat, Belgium: sources and source processes of biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 125–138, doi:10.5194/acp-12-125-2012, 2012.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Graham, B., Mayol-Bracero, O. L., Guyon, P., Roberts, G. C., Decesari, S., Facchini, M. C., Artaxo, P., Maenhaut, W., Koll, P., and Andreae, M. O.: Water-soluble organic compounds in biomass burning aerosols over Amazonia – 1. Characterization by NMR and GC-MS, *J. Geophys. Res.-Atmos.*, 107, 8047, doi:10.1029/2001jd000336, 2002.
- 5 Hamed, A., Birmili, W., Joutsensaari, J., Mikkonen, S., Asmi, A., Wehner, B., Spindler, G., Jaatinen, A., Wiedensohler, A., Korhonen, H., Lehtinen, K. E. J., and Laaksonen, A.: Changes in the production rate of secondary aerosol particles in Central Europe in view of decreasing SO₂ emissions between 1996 and 2006, *Atmos. Chem. Phys.*, 10, 1071–1091, doi:10.5194/acp-10-1071-2010, 2010.
- 10 Hamilton, J. F., Lewis, A. C., Reynolds, J. C., Carpenter, L. J., and Lubben, A.: Investigating the composition of organic aerosol resulting from cyclohexene ozonolysis: low molecular weight and heterogeneous reaction products, *Atmos. Chem. Phys.*, 6, 4973–4984, doi:10.5194/acp-6-4973-2006, 2006.
- Hatakeyama, S., Tanonaka, T., Weng, J. H., Bandow, H., Takagi, H., and Akimoto, H.: Ozone cyclohexene reaction in air: Quantitative analysis of particulate products and the reaction mechanism, *Environ. Sci. Technol.*, 19, 935–942, 1985.
- 15 Hatakeyama, S., Ohno, M., Weng, J. H., Takagi, H., and Akimoto, H.: Mechanism for the formation of gaseous and particulate products from ozone-cycloalkene reactions in air, *Environ. Sci. Technol.*, 21, 52–57, 1987.
- 20 Hering, S. V. and Friedlander, S. K.: Origins of aerosol sulfur size distributions in the Los-Angeles Basin, *Atmos. Environ.*, 16, 2647–2656, 1982.
- Horn, J.: A rationale and test for the number of factors in factor analysis, *Psychometrika*, 30, 179–185, 1965.
- Hsieh, L. Y., Kuo, S. C., Chen, C. L., and Tsai, Y. I.: Origin of low-molecular-weight dicarboxylic acids and their concentration and size distribution variation in suburban aerosol, *Atmos. Environ.*, 41, 6648–6661, doi:10.1016/j.atmosenv.2007.04.014, 2007.
- 25 Hsieh, L. Y., Chen, C. L., Wan, M. W., Tsai, C. H., and Tsai, Y. I.: Speciation and temporal characterization of dicarboxylic acids in PM_{2.5} during a PM episode and a period of non-episodic pollution, *Atmos. Environ.*, 42, 6836–6850, doi:10.1016/j.atmosenv.2008.05.021, 2008.
- 30 Hsieh, L. Y., Kuo, S. C., Chen, C. L., and Tsai, Y. I.: Size distributions of nano/micron dicarboxylic acids and inorganic ions in suburban PM episode and non-episodic aerosol, *Atmos. Environ.*, 43, 4396–4406, doi:10.1016/j.atmosenv.2009.04.034, 2009.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Huang, X. F. and Yu, J. Z.: Is vehicle exhaust a significant primary source of oxalic acid in ambient aerosols?, *Geophys. Res. Lett.*, 34, L02808, doi:10.1029/2006gl028457, 2007.
- Huang, X. F., Hu, M., He, L. Y., and Tang, X. Y.: Chemical characterization of water-soluble organic acids in PM_{2.5} in Beijing, China, *Atmos. Environ.*, 39, 2819–2827, 2005.
- Hyder, M., Genberg, J., Sandahl, M., Swietlicki, E., and Jonsson, J. A.: Yearly trend of dicarboxylic acids in organic aerosols from south of Sweden and source attribution, *Atmos. Environ.*, 57, 197–204, doi:10.1016/j.atmosenv.2012.04.027, 2012.
- Ion, A. C., Vermeylen, R., Kourtchev, I., Cafmeyer, J., Chi, X., Gelencsér, A., Maenhaut, W., and Claeys, M.: Polar organic compounds in rural PM_{2.5} aerosols from K-pusztá, Hungary, during a 2003 summer field campaign: Sources and diel variations, *Atmos. Chem. Phys.*, 5, 1805–1814, doi:10.5194/acp-5-1805-2005, 2005.
- Jaffrezo, J. L., Davidson, C. I., Kuhns, H. D., Bergin, M. H., Hillamo, R., Maenhaut, W., Kahl, J. W., and Harris, J. M.: Biomass burning signatures in the atmosphere of central Greenland, *J. Geophys. Res.-Atmos.*, 103, 31067–31078, 1998.
- John, W., Wall, S. M., Ondo, J. L., and Winklmayr, W.: Modes in the size distributions of atmospheric inorganic aerosol, *Atmos. Environ.*, 24, 2349–2359, 1990.
- Jolliffe, I. T.: *Principal Component Analysis*, 2nd edn., Springer Series in Statistics, Springer, New York, 2002.
- Jones, D. L.: Organic acids in the rhizosphere - a critical review, *Plant Soil*, 205, 25–44, 1998.
- Kalberer, M., Yu, J., Cocker, D. R., Flagan, R. C., and Seinfeld, J. H.: Aerosol formation in the cyclohexene-ozone system, *Environ. Sci. Technol.*, 34, 4894–4901, doi:10.1021/es001180f, 2000.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- Kawamura, K. and Kaplan, I. R.: Motor exhaust emissions as a primary source for dicarboxylic acids in Los Angeles ambient air, *Environ. Sci. Technol.*, 21, 105–110, 1987.
- Kawamura, K. and Usukura, K.: Distributions of low molecular weight dicarboxylic acids in the North Pacific aerosol samples, *J. Oceanogr.*, 49, 271–283, 1993.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kawamura, K., Kasukabe, H., and Barrie, L. A.: Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in arctic aerosols: one year of observations, *Atmos. Environ.*, **30**, 1709–1722, 1996a.
- 5 Kawamura, K., Sempere, R., Imai, Y., Fujii, Y., and Hayashi, M.: Water soluble dicarboxylic acids and related compounds in Antarctic aerosols, *J. Geophys. Res.-Atmos.*, **101**, 18721–18728, 1996b.
- Kawamura, K., Imai, Y., and Barrie, L. A.: Photochemical production and loss of organic acids in high Arctic aerosols during long-range transport and polar sunrise ozone depletion events, *Atmos. Environ.*, **39**, 599–614, 2005.
- 10 Kerminen, V. M., Teinila, K., Hillamo, R., and Makela, T.: Size-segregated chemistry of particulate dicarboxylic acids in the Arctic atmosphere, *Atmos. Environ.*, **33**, 2089–2100, 1999.
- Kerminen, V. M., Ojanen, C., Pakkanen, T., Hillamo, R., Aurela, M., and Merilainen, J.: Low-molecular-weight dicarboxylic acids in an urban and rural atmosphere, *J. Aerosol Sci.*, **31**, 349–362, 2000.
- 15 Kerminen, V. M., Hillamo, R., Teinila, K., Pakkanen, T., Allegrini, I., and Sparapani, R.: Ion balances of size-resolved tropospheric aerosol samples: implications for the acidity and atmospheric processing of aerosols, *Atmos. Environ.*, **35**, 5255–5265, 2001.
- Khalil, M. A. K. and Rasmussen, R. A.: Tracers of wood smoke, *Atmos. Environ.*, **37**, 1211–1222, doi:10.1016/s1352-2310(02)01014-2, 2003.
- 20 Kleefeld, S., Hoffer, A., Krivacsy, Z., and Jennings, S. G.: Importance of organic and black carbon in atmospheric aerosols at Mace Head, on the West Coast of Ireland (53° 19' N, 9° 54' W), *Atmos. Environ.*, **36**, 4479–4490, 2002.
- Kourtchev, I., Warnke, J., Maenhaut, W., Hoffmann, T., and Claeys, M.: Polar organic marker compounds in PM_{2.5} aerosol from a mixed forest site in western Germany, *Chemosphere*, **73**, 1308–1314, doi:10.1016/j.chemosphere.2008.07.011, 2008.
- 25 Kourtchev, I., Copolovici, L., Claeys, M., and Maenhaut, W.: Characterization of atmospheric aerosols at a forested site in Central Europe, *Environ. Sci. Technol.*, **43**, 4665–4671, doi:10.1021/es803055w, 2009.
- Kundu, S., Kawamura, K., Andreae, T. W., Hoffer, A., and Andreae, M. O.: Molecular distributions of dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls in biomass burning aerosols: implications for photochemical production and degradation in smoke layers, *Atmos. Chem. Phys.*, **10**, 2209–2225, doi:10.5194/acp-10-2209-2010, 2010.
- 30

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lan, Z.-J., Chen, D.-L., Li, X., Huang, X.-F., He, L.-Y., Deng, Y.-G., Feng, N., and Hu, M.: Modal characteristics of carbonaceous aerosol size distribution in an urban atmosphere of South China, *Atmos. Res.*, 100, 51–60, doi:10.1016/j.atmosres.2010.12.022, 2011.
- Legrand, M., Preunkert, S., Oliveira, T., Pio, C. A., Hammer, S., Gelencser, A., Kasper-Giebl, A., and Laj, P.: Origin of C-2-C-5 dicarboxylic acids in the European atmosphere inferred from year-round aerosol study conducted at a west–east transect, *J. Geophys. Res.-Atmos.*, 112, D23s07, doi:10.1029/2006jd008019, 2007.
- Lim, H. J., Carlton, A. G., and Turpin, B. J.: Isoprene forms secondary organic aerosol through cloud processing: Model simulations, *Environ. Sci. Technol.*, 39, 4441–4446, doi:10.1021/Es048039h, 2005.
- Limbeck, A., Kraxner, Y., and Puxbaum, H.: Gas to particle distribution of low molecular weight dicarboxylic acids at two different sites in central Europe (Austria), *J. Aerosol Sci.*, 36, 991–1005, 2005.
- Liu, S., Day, D. A., Shields, J. E., and Russell, L. M.: Ozone-driven daytime formation of secondary organic aerosol containing carboxylic acid groups and alkane groups, *Atmos. Chem. Phys.*, 11, 8321–8341, doi:10.5194/acp-11-8321-2011, 2011.
- Maenhaut, W., Wang, W., and Chi, X. G.: Semivolatile behaviour and filter sampling artifacts for dicarboxylic acids during summer campaigns at three forested sites in Europe, *Boreal Environ. Res.*, 16, 273–287, 2011.
- Matsumoto, K., Nagao, I., Tanaka, H., Miyaji, H., Iida, T., and Ikebe, Y.: Seasonal characteristics of organic and inorganic species and their size distributions in atmospheric aerosols over the northwest Pacific Ocean, *Atmos. Environ.*, 32, 1931–1946, 1998.
- Meng, Z. Y. and Seinfeld, J. H.: On the source of the submicrometer droplet mode of urban and regional aerosols, *Aerosol Sci. Tech.*, 20, 253–265, 1994.
- Meszaros, E., Barcza, T., Gelencser, A., Hlavay, J., Kiss, G., Krivacsy, Z., Molnar, A., and Polyak, K.: Size distributions of inorganic and organic species in the atmospheric aerosol in Hungary, *J. Aerosol Sci.*, 28, 1163–1175, 1997.
- Miyazaki, Y., Aggarwal, S. G., Singh, K., Gupta, P. K., and Kawamura, K.: Dicarboxylic acids and water-soluble organic carbon in aerosols in New Delhi, India, in winter: characteristics and formation processes, *J. Geophys. Res.-Atmos.*, 114, D19206, doi:10.1029/2009jd011790, 2009.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Miyazaki, Y., Kawamura, K., and Sawano, M.: Size distributions and chemical characterization of water-soluble organic aerosols over the western North Pacific in summer, *J. Geophys. Res.-Atmos.*, 115, D23210, doi:10.1029/2010jd014439, 2010.
- Mochida, M., Kawabata, A., Kawamura, K., Hatsushika, H., and Yamazaki, K.: Seasonal variation and origins of dicarboxylic acids in the marine atmosphere over the western North Pacific, *J. Geophys. Res.-Atmos.*, 108, 4193, doi:10.1029/2002jd002355, 2003a.
- Mochida, M., Umemoto, N., Kawamura, K., and Uematsu, M.: Bimodal size distribution of C-2-C-4 dicarboxylic acids in the marine aerosols, *Geophys. Res. Lett.*, 30, 1672, doi:10.1029/2003gl017451, 2003b.
- Müller, K., van Pinxteren, D., Plewka, A., Svrčina, B., Kramberger, H., Hofmann, D., Bächmann, K., and Herrmann, H.: Aerosol characterisation at the FEBUKO upwind station Goldlauter (II): detailed organic chemical characterisation, *Atmos. Environ.*, 39, 4219–4231, 2005.
- Myriokefalitakis, S., Tsigaridis, K., Mihalopoulos, N., Sciare, J., Nenes, A., Kawamura, K., Segers, A., and Kanakidou, M.: In-cloud oxalate formation in the global troposphere: a 3-D modeling study, *Atmos. Chem. Phys.*, 11, 5761–5782, doi:10.5194/acp-11-5761-2011, 2011.
- Narukawa, M., Kawamura, K., Takeuchi, N., and Nakajima, T.: Distribution of dicarboxylic acids and carbon isotopic compositions in aerosols from 1997 Indonesian forest fires, *Geophys. Res. Lett.*, 26, 3101–3104, 1999.
- Narukawa, M., Kawamura, K., Anlauf, K. G., and Barrie, L. A.: Fine and coarse modes of dicarboxylic acids in the Arctic aerosols collected during the Polar Sunrise Experiment 1997, *J. Geophys. Res.-Atmos.*, 108, 4575, doi:10.1029/2003jd003646, 2003a.
- Narukawa, M., Kawamura, K., Okada, K., Zaizen, Y., and Makino, Y.: Aircraft measurement of dicarboxylic acids in the free tropospheric aerosols over the western to central North Pacific, *Tellus B*, 55, 777–786, 2003b.
- Neusüß, C., Pelzing, M., Plewka, A., and Herrmann, H.: A new analytical approach for size-resolved speciation of organic compounds in atmospheric aerosol particles: methods and first results, *J. Geophys. Res.-Atmos.*, 105, 4513–4527, 2000.
- Neusüß, C., Wex, H., Birmili, W., Wiedensohler, A., Koziar, C., Busch, B., Brüggemann, E., Gnauk, T., Ebert, M., and Covert, D. S.: Characterization and parameterization of atmospheric particle number-, mass-, and chemical-size distributions in central Europe during LACE 98 and MINT, *J. Geophys. Res.-Atmos.*, 107, 8127, doi:10.1029/2001JD000514, 2002.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oliveira, C., Pio, C., Alves, C., Evtugina, M., Santos, P., Gonçalves, V., Nunes, T., Silvestre, A. J. D., Palmgren, F., Wåhlin, P., and Harrad, S.: Seasonal distribution of polar organic compounds in the urban atmosphere of two large cities from the North and South of Europe, *Atmos. Environ.*, 41, 5555–5570, doi:10.1016/j.atmosenv.2007.03.001, 2007.

Orzechowska, G. E. and Paulson, S. E.: Photochemical sources of organic acids, 1. Reaction of ozone with isoprene, propene, and 2-butenes under dry and humid conditions using SPME, *J. Phys. Chem. A*, 109, 5358–5365, 2005.

Plewka, A., Gnauk, T., Brüggemann, E., Neusüß, C., and Herrmann, H.: Size-resolved aerosol characterization for a polluted episode at the IfT research station Melpitz in autumn 1997, *J. Atmos. Chem.*, 48, 131–156, 2004.

Pradeep Kumar, P., Broekhuizen, K., and Abbatt, J. P. D.: Organic acids as cloud condensation nuclei: Laboratory studies of highly soluble and insoluble species, *Atmos. Chem. Phys.*, 3, 509–520, doi:10.5194/acp-3-509-2003, 2003.

Prenni, A. J., De Mott, P. J., and Kreidenweis, S. M.: Water uptake of internally mixed particles containing ammonium sulfate and dicarboxylic acids, *Atmos. Environ.*, 37, 4243–4251, 2003.

Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Fuzzi, S., Ceburnis, D., O'Dowd, C. D., Sciare, J., Burrows, J. P., Vrekoussis, M., Ervens, B., Tsigaridis, K., and Facchini, M. C.: Evidence of a natural marine source of oxalic acid and a possible link to glyoxal, *J. Geophys. Res.-Atmos.*, 116, D16204, doi:10.1029/2011jd015659, 2011.

Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of fine organic aerosol, 1. Charbroilers and meat cooking operations, *Environ. Sci. Technol.*, 25, 1112–1125, 1991.

Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of fine organic aerosol, 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks, *Environ. Sci. Technol.*, 27, 636–651, 1993a.

Rogge, W. F., Mazurek, M. A., Hildemann, L. M., Cass, G. R., and Simoneit, B. R. T.: Quantification of urban organic aerosols at a molecular level: identification, abundance and seasonal variation, *Atmos. Environ.*, 27, 1309–1330, 1993b.

Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of fine organic aerosol, 6. Cigarette smoke in the urban atmosphere, *Environ. Sci. Technol.*, 28, 1375–1388, 1994.

Röhl, A. and Lammel, G.: Dicarboxylic acids in atmospheric aerosols: local or regional formation?, *J. Aerosol Sci.*, 31, S348–S349, 2000.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Röhl, A. and Lammel, G.: Low molecular weight dicarboxylic acids and glyoxylic acid: seasonal and air mass characteristics, *Environ. Sci. Technol.*, 35, 95–101, 2001.
- Röhl, A. and Lammel, G.: Determination of malic acid and other C-4 dicarboxylic acids in atmospheric aerosol samples, *Chemosphere*, 46, 1195–1199, 2002.
- 5 Saarikoski, S., Carbone, S., Decesari, S., Giulianelli, L., Angelini, F., Canagaratna, M., Ng, N. L., Trimborn, A., Facchini, M. C., Fuzzi, S., Hillamo, R., and Worsnop, D.: Chemical characterization of springtime submicrometer aerosol in Po Valley, Italy, *Atmos. Chem. Phys.*, 12, 8401–8421, doi:10.5194/acp-12-8401-2012, 2012.
- 10 Samy, S. and Zielinska, B.: Secondary organic aerosol production from modern diesel engine emissions, *Atmos. Chem. Phys.*, 10, 609–625, doi:10.5194/acp-10-609-2010, 2010.
- Sato, K., Hatakeyama, S., and Imamura, T.: Secondary organic aerosol formation during the photooxidation of toluene: NO_x dependence of chemical composition, *J. Phys. Chem. A*, 111, 9796–9808, doi:10.1021/Jp071419f, 2007.
- 15 Satsumabayashi, H., Kurita, H., Yokouchi, Y., and Ueda, H.: Photochemical formation of particulate dicarboxylic acids under long-range transport in central Japan, *Atmos. Environ.*, 24, 1443–1450, 1990.
- Scheinhardt, S., Müller, K., Spindler, G., and Herrmann, H.: Complexation of trace metals in size-segregated aerosol particles at nine sites in Germany, *Atmos. Environ.*, 74, 102–109, doi:10.1016/j.atmosenv.2013.03.023, 2013.
- 20 Sorooshian, A., Varutbangkul, V., Brechtel, F. J., Ervens, B., Feingold, G., Bahreini, R., Murphy, S. M., Holloway, J. S., Atlas, E. L., Buzorius, G., Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Oxalic acid in clear and cloudy atmospheres: analysis of data from International Consortium for Atmospheric Research on Transport and Transformation 2004, *J. Geophys. Res.-Atmos.*, 111, D23s45, doi:10.1029/2005jd006880, 2006.
- 25 Sorooshian, A., Ng, N. L., Chan, A. W. H., Feingold, G., Flagan, R. C., and Seinfeld, J. H.: Particulate organic acids and overall water-soluble aerosol composition measurements from the 2006 Gulf of Mexico Atmospheric Composition and Climate Study (GoMACCS), *J. Geophys. Res.-Atmos.*, 112, D13201, doi:10.1029/2007jd008537, 2007.
- 30 Sorooshian, A., Murphy, S. M., Hersey, S., Bahreini, R., Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Constraining the contribution of organic acids and AMS *m/z* 44 to the organic aerosol budget: on the importance of meteorology, aerosol hygroscopicity, and region, *Geophys. Res. Lett.*, 37, L21807, doi:10.1029/2010gl044951, 2010.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Stone, E. A., Hedman, C. J., Zhou, J. B., Mieritz, M., and Schauer, J. J.: Insights into the nature of secondary organic aerosol in Mexico City during the MILAGRO experiment 2006, *Atmos. Environ.*, 44, 312–319, doi:10.1016/j.atmosenv.2009.10.036, 2010.
- 5 Tedetti, M., Kawamura, K., Charriere, B., Chevalier, N., and Sempere, R.: Determination of low molecular weight dicarboxylic and ketocarboxylic acids in seawater samples, *Anal. Chem.*, 78, 6012–6018, doi:10.1021/Ac052226w, 2006.
- Tilgner, A. and Herrmann, H.: Radical-driven carbonyl-to-acid conversion and acid degradation in tropospheric aqueous systems studied by CAPRAM, *Atmos. Environ.*, 44, 5415–5422, doi:10.1016/j.atmosenv.2010.07.050, 2010.
- 10 Turekian, V. C., Macko, S. A., and Keene, W. C.: Concentrations, isotopic compositions, and sources of size-resolved, particulate organic carbon and oxalate in near-surface marine air at Bermuda during spring, *J. Geophys. Res.-Atmos.*, 108, 4157, doi:10.1029/2002jd002053, 2003.
- van Pinxteren, D., Plewka, A., Hofmann, D., Müller, K., Kramberger, H., Svrčina, B., Bächmann, K., Jaeschke, W., Mertes, S., Collett, J. L., and Herrmann, H.: Schmücke hill cap cloud and valley stations aerosol characterisation during FEBUKO (II): organic compounds, *Atmos. Environ.*, 39, 4305–4320, 2005.
- 15 van Pinxteren, D., Brüggemann, E., Gnauk, T., Iinuma, Y., Müller, K., Nowak, A., Achtert, P., Wiedensohler, A., and Herrmann, H.: Size- and time-resolved chemical particle characterization during CAREBeijing-2006: Different pollution regimes and diurnal profiles, *J. Geophys. Res.-Atmos.*, 114, D00g09, doi:10.1029/2008jd010890, 2009.
- 20 van Pinxteren, D., Brüggemann, E., Gnauk, T., Müller, K., Thiel, C., and Herrmann, H.: A GIS based approach to back trajectory analysis for the source apportionment of aerosol constituents and its first application, *J. Atmos. Chem.*, 67, 1–28, doi:10.1007/s10874-011-9199-9, 2010.
- 25 Veres, P. R., Roberts, J. M., Cochran, A. K., Gilman, J. B., Kuster, W. C., Holloway, J. S., Graus, M., Flynn, J., Lefer, B., Warneke, C., and de Gouw, J.: Evidence of rapid production of organic acids in an urban air mass, *Geophys. Res. Lett.*, 38, L17807, doi:10.1029/2011gl048420, 2011.
- 30 Wang, H. B. and Shooter, D.: Low molecular weight dicarboxylic acids in PM₁₀ a city with intensive solid fuel burning, *Chemosphere*, 56, 725–733, 2004.
- Warneck, P.: In-cloud chemistry opens pathway to the formation of oxalic acid in the marine atmosphere, *Atmos. Environ.*, 37, 2423–2427, 2003.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Weber, R. J., Sullivan, A. P., Peltier, R. E., Russell, A., Yan, B., Zheng, M., de Gouw, J., Warneke, C., Brock, C., Holloway, J. S., Atlas, E. L., and Edgerton, E.: A study of secondary organic aerosol formation in the anthropogenic-influenced southeastern United States, *J. Geophys. Res.-Atmos.*, 112, D13302, doi:10.1029/2007jd008408, 2007.
- 5 Worton, D. R., Goldstein, A. H., Farmer, D. K., Docherty, K. S., Jimenez, J. L., Gilman, J. B., Kuster, W. C., de Gouw, J., Williams, B. J., Kreisberg, N. M., Hering, S. V., Bench, G., McKay, M., Kristensen, K., Glasius, M., Surratt, J. D., and Seinfeld, J. H.: Origins and composition of fine atmospheric carbonaceous aerosol in the Sierra Nevada Mountains, California, *Atmos. Chem. Phys.*, 11, 10219–10241, doi:10.5194/acp-11-10219-2011, 2011.
- 10 Yang, L. M. and Yu, L. E.: Measurements of oxalic acid, oxalates, malonic acid, and malonates in atmospheric particulates, *Environ. Sci. Technol.*, 42, 9268–9275, doi:10.1021/Es801820z, 2008.
- Yao, X. H., Chan, C. K., Fang, M., Cadle, S., Chan, T., Mulawa, P., He, K. B., and Ye, B. M.: The water-soluble ionic composition of PM_{2.5} in Shanghai and Beijing, China, *Atmos. Environ.*, 15 36, 4223–4234, 2002.
- Yao, X. H., Lau, A. P. S., Fang, M., Chan, C. K., and Hu, M.: Size distributions and formation of ionic species in atmospheric particulate pollutants in Beijing, China: 2 – dicarboxylic acids, *Atmos. Environ.*, 37, 3001–3007, 2003.
- Yu, J. Z., Huang, X. F., Xu, J. H., and Hu, M.: When aerosol sulfate goes up, so does oxalate: Implication for the formation mechanisms of oxalate, *Environ. Sci. Technol.*, 39, 128–133, 20 2005.
- Zobrist, B., Marcolli, C., Koop, T., Luo, B. P., Murphy, D. M., Lohmann, U., Zardini, A. A., Krieger, U. K., Corti, T., Cziczo, D. J., Fueglistaler, S., Hudson, P. K., Thomson, D. S., and Peter, T.: Oxalic acid as a heterogeneous ice nucleus in the upper troposphere and its indirect aerosol effect, *Atmos. Chem. Phys.*, 6, 3115–3129, doi:10.5194/acp-6-3115-2006, 2006.
- 25

Table 1. Concentrations of dicarboxylic acids determined in this work and literature data from comparable European rural and urban sites. All data given in ng m^{-3} as mean \pm standard deviation (min–max). For Melpitz, Falkenberg, and Goldlauter data mean and standard deviation were calculated in a weighted manner to account for the variable sampling times applied at these sites.

Site	Season	#	Size	Oxalate	Malonate	Succinate	Glutarate	Tartronate	Malate	Tartrate	Ref.
Urban sites											
Leipzig	Summer/Winter	30	PM ₁₀	82 \pm 52 (10–180)	31 \pm 23 (bdl-72)	16 \pm 12 (bdl-60)	21 \pm 20 ^a (bdl-86)	4 \pm 5 (bdl-22)	17 \pm 19 (bdl-85)	4 \pm 7 (bdl-35)	This work
Dresden	Year-round	12	PM ₁₀	58 \pm 40 (11–135)	19 \pm 16 (1–46)	13 \pm 10 (2–36)	12 \pm 11 ^a (bdl-29)	3 \pm 3 (bdl-9)	10 \pm 12 (bdl-40)	1 \pm 1 (bdl-3)	This work
Veszprém, HU	Summer/Winter	17	PM ₁₆	105/129 ^b	25/30 ^b	27/33 ^b					Meszaros et al. (1997)
Helsinki, FI	Year-round	50	PM _{2.5}	100 ^c	10 ^c	12 ^c					Kerminen et al. (2000)
Eichstädt, DE	Summer	22	TSP	251	56	18	10				Röhrl and Lammel (2001)
Leipzig, DE	Summer	24	TSP	229	66	35	30				Röhrl and Lammel (2001)
Eichstädt, Leipzig, DE	Summer	na	TSP			30			54	6.7	Röhrl and Lammel (2002)
Vienna, AT	Winter	5	TSP	86 ^d	26 ^d	42 ^d	9 ^d				Limbeck et al. (2005)
Zurich, CH	Summer/Winter	na	PM ₁₀	135/104 ^b	29/13 ^b	10/6 ^b			21/11 ^b	2/1 ^b	Fisseha et al. (2006)
Oporto, PT	Summer/Winter	na	PM ₁₀		na/4 ^c	11/15 ^c	3/10 ^c				Oliveira et al. (2007)
Copenhagen, DK	Summer/Winter	na	PM ₁₀		0.5/1 ^c	18/12 ^c	8/12 ^c				Oliveira et al. (2007)
Jülich, DE	Summer	6	PM _{2.5}			8.1			39		Kourchev et al. (2008)
Brasschaat, BE	Summer	71	PM _{2.5}	75 ^e	49 ^e	11 ^e	4.7 ^e				Gomez-Gonzalez et al. (2012)
Rural sites											
Melpitz	Fall	20	PM ₁₀	52 \pm 26 (18–138)	22 \pm 16 (3–46)	19 \pm 6 (8–27)	16 \pm 11 ^a (5–31)	4 \pm 5 (0–11)	12 \pm 8 (6–26)	bdl	This work
Falkenberg	Summer	23	PM ₁₀	80 \pm 41 (15–209)	39 \pm 20 (7–64)	11 \pm 8 (1–27)	28 \pm 20 ^a (3–63)	14 \pm 13 (1–38)	31 \pm 22 (4–70)	bdl	This work
Goldlauter	Fall	10	PM ₁₀	36 \pm 23 (16–114)	18 \pm 16 (5–67)	12 \pm 11 (3–45)	10 \pm 11 ^a (bdl-35)	2 \pm 2 (3–32)	10 \pm 7 (3–32)	2 \pm 2 (bdl-6)	This work
Merseburg, DE	Winter	22	TSP	57	4.8	4.2					Röhrl and Lammel (2001)
Falkenberg, DE	Summer	17	TSP	343	64	25	21				Röhrl and Lammel (2001)
3 rural sites, DE	Summer/Winter	na	TSP			14			34	4	Röhrl and Lammel (2002)
K-pusztza, HU	Summer	63	PM ₂						38 ^e		Ion et al. (2005)
K-pusztza, HU	Summer/Winter	na	PM _{2.5}	264/300 ^b	29/28 ^b	21/37 ^b	19/21 ^b		37/19 ^b	6/4 ^b	Legrand et al. (2007)
K-pusztza, HU	Summer	63	PM _{2.5}	189		15	10		40		Kourchev et al. (2009)
Hyytiälä, FI	Summer	51	PM _{2.5}	104 ^e	20 ^e	13 ^e	7 ^e				Maenhaut et al. (2011)
Po Valley, IT	Spring	23	PM ₁	100 ^c	15 ^c	8 ^c			12 ^c		Saarikoski et al. (2012)
Vavihill, SE	Year-round	35	PM ₁₀		1.6 \pm 0.5	3.5 \pm 2.4	1.8 \pm 0.8				Hyder et al. (2012)

Abbr.: na: not available, bdl: below detection limit.

^a Sum of 3 C5 DCA isomers.

^b Mean conc. for summer and winter, respectively.

^c Estimated from graphical data representation.

^d Sum of front- and backup filters.

^e Median conc.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Loadings of PCA after Varimax rotation. Loadings with $|x| < 0.2$ are considered insignificant and removed, while loadings with $|x| \geq 0.6$ are considered as “high” and printed bold.

Variable Assigned source	PC1 gasSOA	PC2 aqSOA	PC3 traffic	PC4 BB	PC5 sea salt	PC6 soil
Oxalate 0.05–0.14 μm	0.81					
Oxalate 0.14–0.42 μm	0.70	0.42	–0.29			
Oxalate 0.42–1.2 μm	0.32	0.81				
Oxalate 1.2–3.5 μm	0.53	0.29	–0.42		0.26	0.23
Oxalate 3.5–10 μm	0.69				0.33	0.32
Malonate 0.05–0.14 μm	0.81				–0.22	
Malonate 0.14–0.42 μm	0.55	0.68	–0.25		–0.29	
Malonate 0.42–1.2 μm	0.38	0.77				
Malonate 1.2–3.5 μm	0.54	0.34	–0.61			0.26
Malonate 3.5–10 μm	0.68		–0.58			
Succinate 0.05–0.14 μm	0.71					
Succinate 0.14–0.42 μm	0.35	0.80			–0.34	
Succinate 0.42–1.2 μm		0.88			–0.23	
Succinate 1.2–3.5 μm	0.31	0.48	–0.32			0.48
Succinate 3.5–10 μm	0.64			–0.31		0.42
Malate 0.05–0.14 μm	0.79		–0.20		–0.26	–0.20
Malate 0.14–0.42 μm	0.70	0.40	–0.47			
Malate 0.42–1.2 μm	0.43	0.72	–0.44			
Malate 1.2–3.5 μm	0.32	0.28	–0.70			
Malate 3.5–10 μm	0.23		–0.75			
OC 0.05–0.14 μm			0.33		–0.53	0.46
OC 0.14–0.42 μm		0.25			–0.38	0.43
OC 0.42–1.2 μm		0.56			–0.44	0.31
OC 1.2–3.5 μm		0.28				0.61
OC 3.5–10 μm	0.49					0.36
EC 0.05–0.14 μm	0.20		0.59			0.40
EC 0.14–0.42 μm		0.55	0.40			0.28
EC 0.42–1.2 μm		0.61	0.36			
EC 1.2–3.5 μm		0.39	0.31			0.51
EC 3.5–10 μm	0.53					0.45

Source contributions of dicarboxylic acids

D. van Pinxteren et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Continued.

Variable Assigned source	PC1 gasSOA	PC2 aqSOA	PC3 traffic	PC4 BB	PC5 sea salt	PC6 soil
Sulfate 0.05–0.14 μm	0.62	0.21		0.40		
Sulfate 0.14–0.42 μm	0.20	0.63		0.49		
Sulfate 0.42–1.2 μm		0.90		0.22		
Sulfate 1.2–3.5 μm		0.50			0.23	0.40
Sulfate 3.5–10 μm				0.41	0.56	0.33
Potassium 0.05–0.14 μm				0.69		
Potassium 0.14–0.42 μm		0.32	0.29	0.71		
Potassium 0.42–1.2 μm	–0.24	0.37	0.40	0.62		
Potassium 1.2–3.5 μm				0.81		
Potassium 3.5–10 μm	0.20			0.65	0.38	–0.22
Sodium 0.42–1.2 μm				0.62	0.29	
Sodium 1.2–3.5 μm		–0.20			0.79	
Sodium 3.5–10 μm		–0.26			0.83	
Calcium 0.42–1.2 μm				0.72	–0.20	
Calcium 1.2–3.5 μm	0.33			0.33		0.54
Calcium 3.5–10 μm	0.58			0.28		0.35
O ₃	0.46			–0.54		–0.48
NO _x	0.26		0.82			0.29
CO			0.58			
SO ₂	–0.21	0.34	0.37	0.57		0.22
RTI _{Natural vegetation}		0.29			–0.71	
RTI _{Agriculture}	0.25	0.68			–0.40	
RTI _{Urban areas}	0.44	0.37				
Mean trajectory length	–0.46	–0.41			0.39	
Solar flux along trajectory	0.79		–0.42			
Solar flux at receptor	0.78		–0.26			–0.27
Mixing depth at receptor	0.52		–0.30		0.39	–0.44
Explained variance (%)	18	15	10	9	8	7
Cummulative variance (%)	18	34	44	53	61	68

**Source contributions
of dicarboxylic acids**

D. van Pinxteren et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Source contributions
of dicarboxylic acids

D. van Pinxteren et al.




Fig. 1. Mean size distributions of DCAs at all sites. Error bars indicate the 15th and 85th percentiles as a measure of data variability.

Source contributions
of dicarboxylic acids

D. van Pinxteren et al.




Fig. 2. Box-plots of residence time indices and further parameters of back trajectory analysis at different sampling sites. Bold line represents median, boxes lower and upper quartiles, whiskers 1.5 times the interquartile range, and dots indicate values outside this range.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Screen plot of PCA including results from parallel analysis.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

