



**Photosensitised
kinetics of biomass
burning aerosol**

S. M. Forrester and
D. A. Knopf

Photosensitised heterogeneous oxidation kinetics of biomass burning aerosol surrogates by ozone using an irradiated rectangular channel flow reactor

S. M. Forrester and D. A. Knopf

Stony Brook University, Institute for Terrestrial and Planetary Atmospheres/School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, New York, USA

Received: 6 March 2013 – Accepted: 7 March 2013 – Published: 20 March 2013

Correspondence to: D. A. Knopf (daniel.knopf@stonybrook.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Heterogeneous reaction kinetics involving organic aerosol and atmospheric oxidants such as ozone can be enhanced under visible or UV irradiation in the presence of a photosensitiser, with subsequent implications for the climate, cloud radiative properties, air quality, and source appointment. In this study we report the steady state reactive uptake coefficient, γ , of O_3 by levoglucosan and 5-nitroguaiacol acting as surrogates for biomass burning aerosol particles, with and without the presence of Pahokee peat acting as a photosensitiser. The reactive uptake has been determined in the dark and as a function of visible and UV-A irradiation and ozone concentration. In addition, γ was determined for 1:1, 1:10, and 1:100 by mass mixtures of Pahokee peat and 5-nitroguaiacol, and for a 10:1:3 mixture of levoglucosan, Pahokee peat, and 5-nitroguaiacol. We developed a novel irradiated rectangular channel flow reactor (I-RCFR) that was operated under low pressures of about 2–4 hPa, and allowed for uniform irradiation of the organic substrates. The I-RCFR was coupled to a chemical ionisation mass spectrometer and has been successfully validated by measuring the kinetics between various organic species and oxidants. γ of O_3 and levoglucosan in the dark and under visible and UV-A irradiation was determined to be in the range of $(2\text{--}11) \times 10^{-6}$ and did not change in the presence of Pahokee peat. The determined γ of O_3 and 5-nitroguaiacol in the dark was 5.7×10^{-6} and was only enhanced under UV-A irradiation, yielding a value of 3.6×10^{-5} . γ of the 1:1 Pahokee peat/5-nitroguaiacol substrate was enhanced under visible and UV-A irradiation to 2.4×10^{-5} and 2.8×10^{-5} , respectively. Decreasing the amount of Pahokee peat in the 5-nitroguaiacol/Pahokee peat substrate resulted in lower values of γ under visible irradiation, however, γ was consistent under UV-A irradiation regardless of the amount of Pahokee peat. The 10:1:3 mixture by mass of levoglucosan, Pahokee peat, and 5-nitroguaiacol, under both visible and UV-A irradiation yielded γ values of 2.8×10^{-5} and 1.4×10^{-5} , respectively. γ was determined as a function of photon flux for O_3 with the 1:1 Pahokee peat/5-nitroguaiacol substrate, yielding a linear relationship under both visible and UV-A irra-

ACPD

13, 7553–7598, 2013

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

diation. γ of O_3 with the 1:1 Pahokee peat/5-nitroguaiacol substrate was determined as a function of ozone concentration and exhibited an inverse dependence of γ on ozone concentration, commonly interpreted as a Langmuir-Hinshelwood mechanism. The reactive uptake data have been represented by a Langmuir-type isotherm. From the O_3 uptake data under visible irradiation, the following fit parameters have been derived: $k_s = (5.5 \pm 2.7) \times 10^{-19} \text{ cm}^2 \text{ s}^{-1} \text{ molecule}^{-1}$ and $K_{O_3} = (2.3 \pm 2.0) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1}$ and under UV-A irradiation: $k_s = (8.1 \pm 2.0) \times 10^{-19} \text{ cm}^2 \text{ s}^{-1} \text{ molecule}^{-1}$ and $K_{O_3} = (1.7 \pm 0.7) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1}$. The oxidative power, or the product of γ and $[O_3]$, was determined for O_3 with the 1:1 Pahokee peat/5-nitroguaiacol substrate and was in the range of $(1.2\text{--}26) \times 10^6 \text{ molecule cm}^{-3}$. Atmospheric particle lifetimes were estimated for a $0.4 \mu\text{m}$ 5-nitroguaiacol particle as a function of visible and UV-A irradiation and ozone concentration.

1 Introduction

Condensed phase organic material is ubiquitous in the atmosphere and can constitute a significantly large portion (20–90 %) of the submicron atmospheric aerosol mass fraction (Kanakidou et al., 2005; Zhang et al., 2007; Hallquist et al., 2009). Biomass burning is a large source of trace gases and organic aerosol to the atmosphere, and has a source strength that is comparable to fossil fuel burning (Crutzen et al., 1979; Logan et al., 1981; Crutzen and Andreae, 1990). Biomass burning aerosol (BBA) can have significant effects on air quality and human health, and can directly affect the atmosphere through scattering and absorption of radiation, or indirectly through influencing cloud radiative properties by impacting underlying microphysics (Hobbs and Radke, 1969; Ramanathan et al., 2001; Pope et al., 2009; Levin et al., 2010; Dusek et al., 2011; Engelhart et al., 2012).

Although BBA represents a complex mixture of inorganic and water soluble/insoluble organic compounds, some species are common in BBA and can be applied

as biomolecular markers for source apportionment of biomass burning plumes (Hawthorne et al., 1988, 1989; Simoneit, 2002; Robinson et al., 2006b; Fu et al., 2012). Two such biomolecular markers are the hemicellulose decomposition product, levoglucosan (1,6-anhydro- β -D-glucopyranose, $C_6H_{10}O_5$) and the substituted phenol, 5-nitroguaiacol (2-methoxy-5-nitrophenol, $C_7H_7NO_4$) (Hawthorne et al., 1988, 1989; Simoneit, 2002; Robinson et al., 2006b; Fu et al., 2012). To assess the aerosol source strength, biomolecular markers employed in apportionment studies should be inert with respect to chemical reaction during atmospheric transport. In the atmosphere, however, BBA particles are exposed to gas phase oxidants such as O_3 , resulting in multiphase chemical reactions, which can cause chemical and physical modification of the particles, a process known as chemical ageing (Hennigan et al., 2010; Kessler et al., 2010; Hennigan et al., 2011; Knopf et al., 2006, 2011; Shiraiwa et al., 2012). It has been shown that O_3 can be photochemically created in BB plumes, with reported concentration levels as high as 100 ppb (Delany et al., 1985; Andreae et al., 1988; Crutzen and Andreae, 1990; Ziemke et al., 2009; Konovalov et al., 2011; Dupont et al., 2012; Martins et al., 2012; Akagi et al., 2013). Therefore, heterogeneous oxidation reactions inside a BB plume can cause chemical modification of particles and biomolecular marker species, potentially having subsequent effects on source apportionment (Robinson et al., 2006b; Lin et al., 2010). Chemical ageing may also enhance aerosol toxicity, as well as yield changes in aerosol hygroscopicity, affecting cloud formation processes (Finlayson-Pitts and Pitts, 2000; Maria et al., 2004; Rudich et al., 2007; Andreae, 2009; Jimenez et al., 2009; Hennigan et al., 2011; Wang and Knopf, 2011). It has been shown recently that there can be significant degradation and volatilisation of typical BBA species such as levoglucosan, abietic acid, and 5-nitroguaiacol through heterogeneous oxidation reactions (Hennigan et al., 2010; Kessler et al., 2010; Knopf et al., 2011; Shiraiwa et al., 2012). The reactive uptake coefficient, γ , or the probability that a gas to particle collision results in loss of that gas molecule due to reaction, of O_3 with levoglucosan and 5-nitroguaiacol can be on the order of 10^{-5} in the absence of irradiation (Knopf et al., 2011). The corresponding oxidative powers ($\gamma \cdot [O_3]$)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



can be on the order of 3.3×10^7 and 2.0×10^8 molecule cm^{-3} , respectively, assuming 100 ppb of O_3 in a BB plume (Knopf et al., 2011). Compared to the oxidative power of OH with levoglucosan, 5.9×10^6 assuming $\gamma_{\text{OH}} = 0.52$ and $[\text{OH}] = 0.46$ ppt (Yokelson et al., 2009; Slade and Knopf, 2013), heterogeneous reactions of O_3 with compounds typical of BBA and thus particle and species degradation could be significant, however volatilisation of condensed phase species is expected to be much less for reactions involving O_3 compared to OH (Slade and Knopf, 2013). In summary, it is highly likely that BBA particles undergo chemical ageing by O_3 during atmospheric transport, with subsequent implications for source apportionment calculations.

Chemical ageing of BBA by gas phase oxidants can be amplified due to the presence of a photosensitiser. A photosensitiser is a light-absorbing compound that can further enhance heterogeneous kinetics through photosensitive interactions with other species (George et al., 2005; Jammoul et al., 2008; Sosedova et al., 2011). A photosensitising compound can absorb radiation and transfer its excitation energy to a neighbouring molecule that might not have otherwise absorbed radiation (Vione et al., 2006; Hoffmann, 2008; Alvarez et al., 2012). This activated molecule can then undergo other processes such as oligomerisation, HONO formation, or chemical ageing (Alvarez et al., 2012). Humic-like substances (HULIS) are commonly present in BBA, and have been shown to possess photosensitive properties (Facchini et al., 2000; Mayol-Bracero et al., 2002; Gelencsér et al., 2003; Lukács et al., 2007; Schmidl et al., 2008a,b; Fors et al., 2010; Konovalov et al., 2012). The presence of HULIS in BBA may enhance degradation of biomolecular markers if they are able to absorb the excitation energy of the photosensitiser and subsequently react more efficiently with atmospheric oxidants. Concentrations of HULIS can be highly variable in BB smoke, i.e. 0.6–5.8 wt% of HULIS in wood combustion smoke compared to 1.6–21.2 wt% from burning leaves (Schmidl et al., 2008a,b). The effect of varying the amount of photosensitiser in a photosensitised reactive uptake measurement has not been explored in detail.

It has been shown experimentally that phenols can efficiently quench the excited triplet state of a photosensitiser, such as an aromatic carbonyl compound, in solu-

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

tion (Canonica et al., 1995; Anastasio et al., 1997; Canonica et al., 2000). However, it is expected that these types of reactions may be even more efficient at the surface of aerosol particles due to enhanced light intensity at the surface (Nissenson et al., 2006). Although phenolic and polyphenolic compounds are ubiquitous in the atmosphere, they can be significantly enhanced in BB plumes, and have been suggested as tracers for BBA sources (Rogge et al., 1998; Fine et al., 2001). Recent studies have shown that reactive uptake of gas phase oxidants by adsorbed phenolic and polyphenolic compounds such as catechols, gentistic acid, tannic acid, 4-phenoxyphenol, and phenol in the presence of a photosensitiser can be significantly enhanced when exposed to ultra-violet (UV) or visible (VIS) radiation compared to dark conditions (George et al., 2005; Jammoul et al., 2008; Net et al., 2010; Sosedova et al., 2011).

Previous studies have observed that γ can have a linear dependence on VIS and UV photon flux, but some measurements were limited in the VIS photon flux intensity compared to the sun's total VIS irradiance (Stemmler et al., 2006; Jammoul et al., 2008; Brigante et al., 2008; D'Anna et al., 2009; Baduel et al., 2011; Sosedova et al., 2011; Zelenay et al., 2011). Previous studies have also shown that the uptake of gas phase oxidants may follow a Langmuir–Hinshelwood (LH) type reaction mechanism, where the oxidant first adsorbs on the organic substrate and subsequently reacts yielding γ values that are reduced at high oxidant concentrations (Stemmler et al., 2007; Jammoul et al., 2008; Brigante et al., 2008; D'Anna et al., 2009; Net et al., 2010; Baduel et al., 2011; Sosedova et al., 2011; Zelenay et al., 2011). A recent modelling study by Konovalov et al. (2012) indicated that there may be a positive feedback between the O_3 loss rate and O_3 mixing ratio because of this inverse dependence of γ on O_3 concentration (Konovalov et al., 2012).

This study will explore the effect of using the humic acid Pahokee peat (PP) as a proxy for atmospheric HULIS acting as a photosensitiser on γ of O_3 with solid films of levoglucosan (LEV) and 5-nitroguaiacol (NG) when exposed to UV and VIS radiation. We employ a novel irradiated rectangular channel flow reactor (I-RCFR) coupled to a custom-built chemical ionisation mass spectrometer (CIMS) to determine photo-

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

sensitised heterogeneous kinetics. The advantage of this approach is the capability to uniformly irradiate the organic substrate and conduct reactive uptake measurements at atmospherically relevant O₃ concentrations and at low pressures, thereby decreasing diffusion limiting effects. The presented study will also explore the effect of varying the amount of photosensitiser by investigating 1:1, 1:10, and 1:100 mixtures by mass of PP and NG. We will also investigate a 10:1:3 mixture by mass of LEV, NG, and PP. 5-nitroguaiacol is expected to efficiently absorb excitation energy from humic acid due to its phenolic structure, and therefore have enhanced reactivity towards ozone. We also investigate the effect of irradiation intensity on γ . This study employs a photon flux similar to that received by the sun for both VIS ($\Phi_p = 1.0 \times 10^{17}$ photons cm⁻² s⁻¹) and UV-A ($\Phi_p = 5.6 \times 10^{15}$ photons cm⁻² s⁻¹) radiation (Gueymard et al., 2002). Lastly, we will also explore the effect of O₃ concentration on γ at atmospherically relevant oxidant concentrations.

2 Experimental

2.1 Apparatus

Uptake experiments were conducted using an I-RCFR with controlled irradiance coupled to CIMS based on previous setups (Fig. 1) (Knopf et al., 2007, 2011). The I-RCFR is rectangular in shape allowing us to study the reactivity of a substrate with dimensions of 9.0 × 14.9 cm². Either a quartz trough or a sand-blasted glass block can be inserted to study the uptake of gas phase oxidants by liquids or highly viscous/amorphous solid films, respectively. The bottom of the flow reactor is equipped with a cooling jacket to control the temperature of the uptake experiments. Temperatures reported in this study are of the substrate surface. This was determined with a thermocouple directly contacting the glass block while the flow reactor was under low pressure and gas flow was present. Temperatures of the gas were also recorded to determine the mean ther-

mal velocity of the gas molecules. The temperature measurements were performed in the dark and under VIS and UV irradiation.

The top cover of the flow reactor is made of quartz glass to allow maximum transmission (90 % for VIS and UV) of irradiance to the reactive substrate. The quartz is covered with black vinyl for the dark reactive uptake measurements. The distance between the reactive surface and the inside of the cover is 7 mm. The O₃ enters the flow reactor through a 1/8" outer diameter movable injector with a u-shaped end, resulting in two gas flow exits to enhance lateral mixing of the O₃ downstream of the flow reactor. Carrier gases were introduced through two inlets at the back of the flow reactor and allowed to thoroughly mix laterally by use of a mixing barrier.

2.2 Uptake measurements

The pseudo-first order loss of the oxidant to the organic substrate was determined using two different methods. The first method employs monitoring the loss of the gas phase species as the injector was instantaneously pulled back 10 or 12 cm over the reactive substrate. The change in the gas phase oxidant's concentration was then monitored as a function of exposure time (~ 10 min). The steady state O₃ signal is used for calculation of γ . In some instances, e.g. for validation of literature values, we used the initial drop in the O₃ signal to calculate γ . The second method monitors the loss of the gas phase species as the injector is pulled back over the organic substrate in increments of 3 cm until reaching 12 cm. These methods will be termed pullback and stepwise, respectively, for the remainder of the manuscript. At least 6 independent experiments, each using a fresh organic substrate, were conducted for each investigated species or mixture.

2.3 Irradiation source

The VIS irradiation source used is a 400 W GrowBright high pressure sodium (HPS) lamp, fit inside an air-coolable reflector (HTG Supply). The UV irradiation source used

ACPD

13, 7553–7598, 2013

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

is a 400 W Eliminator high pressure mercury vapor blacklight bulb, also fit inside an air-coolable reflector. The lamps are positioned above the flow reactor in such a way that they can be moved closer or farther away, allowing for a change in the photon flux reaching the reactive substrate. The spectral irradiance reaching the reactive substrate was determined using a USB 2000 spectrometer coupled to a 50 μm optical fiber (Ocean Optics). The spectrometer was calibrated using an HL-2000-CAL calibrated tungsten lamp (Ocean Optics). A UV-VIS neutral density filter with an optical density of 3.0 and a wavelength range of 200–700 nm was used between the optical fiber and the irradiation source to avoid saturation of the spectrometer detector. The measurements were performed with the quartz cover between the fiber optics and the irradiation source. For the UV irradiance measurements, a UV bandpass filter with a center wavelength of 360 nm was used to eliminate all wavelengths except the main peak at 365.7 nm. This filter was positioned on top of the quartz cover during UV uptake experiments to eliminate possible influences from VIS irradiation.

2.4 Oxidant generation, detection, and flow conditions

The gases N_2 and He were first passed through a carbon filter (Supelco, Supelcarb HC) and a Drierite trap cooled with liquid nitrogen to remove any possible organic contamination. O_2 was passed through a Drierite trap cooled by a liquid ethanol/dry ice mixture. O_3 was generated by passing a flow of O_2 over a UV source (Jelight, model #600) before entering the flow reactor. The flow of the O_2/O_3 mixture varied between 1 and 5 $\text{cm}^3 \text{min}^{-1}$ at standard temperature and pressure (STP). A He gas carrier flow of about 50 $\text{cm}^3 \text{min}^{-1}$ STP entered the flow reactor and was allowed to thoroughly mix laterally. Typical flow velocities inside the I-RCFR were 60–65 cm s^{-1} . This resulted in Reynolds numbers of about 0.04, indicating laminar flow conditions. O_3 was detected as O_3^- ($m/z = 48$) by the mass spectrometer after chemical ionisation by SF_6^- ($m/z = 146$). SF_6^- was generated by passing a trace amount of SF_6 in about 1 L min^{-1} STP N_2 through a ^{210}Po source (Knopf et al., 2005, 2011). O_3 concentrations used in

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

these reactive uptake measurements ranged from 1.7×10^{11} – 1.9×10^{12} molecule cm^{-3} , representing 7–75 ppb. Experiments were conducted at 2–3 hPa.

For validation purposes, the uptake of NO_3 radicals by linoleic acid (cis,cis-9,12-octadecadienoic acid) was measured. NO_3 was generated by thermal decomposition of N_2O_5 , which was generated through the gas-phase reaction of NO_2 with excess O_3 , as discussed in detail in our previous work (Knopf et al., 2006, 2011). NO_3 was detected as NO_3^- after chemical ionisation by SF_6^- or I^- . I^- was generated by passing a trace amount of CH_3I in about 1 L min^{-1} STP N_2 through a ^{210}Po source (Knopf et al., 2006, 2011).

2.5 Substrate preparation

Pure PP substrates were prepared using a 1 mg mL^{-1} aqueous solution. PP only partially dissolves in water, leaving some particulate matter suspended in the solution. Pure LEV substrates were prepared using a 1.00 wt% aqueous solution, and pure NG substrates were prepared using a 1.00 wt% solution in methanol. 1:1 by mass PP/LEV substrates were prepared as a 1.00 wt% aqueous solution and 1:1, 1:10, and 1:100 by mass PP/NG substrates were prepared as 1.00 wt% solutions in methanol. 10:1:3 by mass LEV/PP/NG substrates were prepared as a 1.00 or 2.00 wt% solutions in methanol. Organic solutions were deposited on the glass block in ~ 0.6 mL increments and solvents were allowed to completely evaporate in a particle free environment. The thickness of the organic film is estimated to be ~ 0.2 – 0.4 μm based on the amount of applied solution.

2.6 Absorbance measurements

Absorbance measurements were conducted using a Cary 1 UV-VIS spectrophotometer. A 1 mg mL^{-1} aqueous solution of PP was centrifuged to remove particulate matter suspended in the solution. The absorbance of a 1 wt% aqueous solution of LEV was measured, but the results are not reported due to a complete lack of absorbance in the

300–700 nm wavelength regime. The absorbance of NG was determined by dissolving in both a 0.01 M HCl or a 0.01 M NaOH solution to account for effects of solution pH on absorbance.

2.7 Chemicals

5 Listed below are the chemicals, the corresponding purities, and manufacturer used in our studies: He (UHP), N₂ (UHP), O₂ (UHP), SF₆ (99.998 %), all acquired from Prax-air; 2-methoxy-5-nitrophenol (97 %), 1,6-anhydro-β-D-glucopyranose (99 %), cis-9-octadecenoic acid (99 %), cis-9,cis-12-octadecadienoic acid (97 %), methanol (99.9 %), all acquired from VWR; canola oil (not determined), Pahokee Peat humic acid standard
10 (International Humic Substances Society). Millipore water (resistivity > 18.2 MΩ cm) was used for preparation of aqueous solutions.

3 Results and discussion

3.1 Light source characterisation

The spectral irradiance of the light sources, as well as a standard solar spectrum tilted 37° from the sun, as given by the American Society for Testing and Materials (Gueymard et al., 2002), are shown in Fig. 2. The total irradiances from the sun are 1.0×10^{17} photons cm⁻² s⁻¹ from 400–700 nm, and 5.6×10^{15} photons cm⁻² s⁻¹ from 300–400 nm. The total irradiance for the visible lamp integrated from 400–700 nm is 5.9×10^{17} photons cm⁻² s⁻¹ at a distance of 7.6 cm and the total irradiance for the UV
15 lamp integrated from 300–400 nm is 8.1×10^{15} photons cm⁻² s⁻¹ at a distance of 5.1 cm. As shown in Fig. 2, the bulk of the photon flux density for the visible lamp lies between 550 and 650 nm. However, previous studies use fluorescent lamps which have significant photon flux from 400 to 500 nm (Jammoul et al., 2008; D'Anna et al., 2009). Furthermore, the sun emits irradiation continually in the visible wavelength regime, indicating that our VIS light source may lack shorter wavelength irradiation compared to
25

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the sun. Also shown in Fig. 2, the photon flux from the UV irradiation source peaks from 362 to 372 nm. However, the sun continually emits irradiation in the UV-A wavelength regime, again indicating that our UV light source may lack shorter wavelength irradiation compared to the sun.

The total irradiance was also measured as a function of distance from the reactive substrate for each of the lamps, as shown in Fig. 3. Our setup allows for the variation of the total VIS and UV photon flux by at least an order of magnitude.

3.2 Derivation of reactive uptake coefficient

The reactive uptake coefficient, γ , is derived using the method developed by Knopf and coworkers (Knopf et al., 2007). As mentioned above, the loss of ozone is measured as the injector is pulled back over a known distance of the organic substrate. Assuming first-order kinetics, we can describe the relationship between distance and concentration as follows:

$$C(l) = C_0 \exp\left(-\frac{k_{\text{obs}} l}{v_{\text{avg}}}\right), \quad (1)$$

where $C(l)$ is the O_3 concentration at position l , l is the length of the reactive substrate exposed to O_3 , C_0 is the concentration at $l = 0$, v_{avg} is the average flow velocity, and k_{obs} is the observed first-order reaction rate constant. We determine k_{obs} as the slope of the logarithmic change in the normalised O_3 signal as a function of reaction time, $t = l/v_{\text{avg}}$, as shown in Fig. 4. Efficient reactive uptake of O_3 can result in concentration gradients close to the substrate surface and thus diffusion effects must be considered. The diffusion coefficient of O_3 in He is taken as $394 \text{ Torr cm}^2 \text{ s}^{-1}$ at 298 K (Moise and Rudich, 2000). Correction for concentration gradients in the vertical direction and in the direction of the bulk gas flow is performed by numerical analysis of the laboratory data following the method described by Knopf et al. (2007). The analysis uses iterative measures to obtain the first order reaction rate constant, $k_{\text{w}}^{1\text{st}}$ from the experimentally

determined k_{obs} . γ is derived according to the first-order wall loss rate constant (Hanson, 1998; Knopf et al., 2007):

$$\frac{1}{\gamma} = \frac{cA}{4k_{\text{w}}^{\text{1st}}V} + \frac{1}{2}, \quad (2)$$

where A is the geometric surface area of the organic substrate, V is the volume of the gas above the reactive substrate, $k_{\text{w}}^{\text{1st}}$ is the first order wall loss rate constant, and c is the thermal molecular speed of O_3 .

The γ values reported in this study are steady state γ values, meaning they were calculated from a steady O_3 signal. The initial γ was also determined for comparison with some literature values, whereby the maximum drop in the O_3 signal was used to calculate γ .

3.3 Validation

To validate the I-RCFR we measured γ of oleic acid (cis-9-octadecenoic acid, $\text{C}_{18}\text{H}_{34}\text{O}_2$), canola oil, levoglucosan, 5-nitroguaiacol, and PP (under dark and UV irradiation conditions) with O_3 and linoleic acid with NO_3 , as shown in Table 1. The uptake of O_3 by liquid and solid oleic acid is in agreement with previous studies within experimental uncertainty (Moise and Rudich, 2002; Zahardis and Petrucci, 2007). The uptake of NO_3 by linoleic acid is in agreement with a study by Zhao et al. (2011). The uptake of O_3 by canola oil is, within experimental uncertainty, in agreement with a previous study by Knopf et al. (2007) that utilised an RCFR. The preceding reactive uptake measurements were performed using the stepwise method. The uptake of O_3 by LEV is in agreement with our previous study using a rotating-wall flow-tube reactor (Knopf et al., 2011). The reactive uptake coefficient of O_3 with NG is very close to the study by Knopf et al. (2011) and the minor difference may be due to differences in the morphology of the organic substrate surfaces. It is important to note that the γ values of LEV and NG with O_3 were determined using the initial drop of the O_3 signal, as in the previous study. Steady state reactive uptake of O_3 by PP under dark conditions is

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in agreement with a study by D'Anna et al. (2009). Under UV irradiation, however, the steady state reactive uptake of O_3 is lower than previously determined by D'Anna et al. (2009). This can be attributed to a lower total irradiance reaching the substrate, i.e. $8.1 \times 10^{15} \text{ photons cm}^{-2} \text{ s}^{-1}$ in this study compared to $2.4 \times 10^{16} \text{ photons cm}^{-2} \text{ s}^{-1}$ applied by D'Anna et al. (2009). The authors have determined that decreasing the photon flux causes γ to decrease, thus a lower γ in our study can be expected.

3.4 Reactive uptake of ozone

3.4.1 Dark experiments

Steady state reactive uptake coefficients of O_3 were determined in complete darkness, under dry conditions, and with atmospherically relevant O_2 levels. The temperature of the organic substrate was 20.0°C . Under these conditions each substrate can be assumed to be in a highly viscous or amorphous solid state (Mikhailov et al., 2009; Knopf et al., 2011). All dark experiments were performed using the pullback method. As shown in Table 2, γ of the LEV substrate is $(3.6 \pm 2.7) \times 10^{-6}$. Such a low γ can be expected when considering the low reactivity of the gas phase reaction between O_3 and alcohols (Atkinson and Carter, 1984). Steady state derived γ of NG is $(5.7 \pm 2.5) \times 10^{-6}$, about a factor of 2 lower compared to γ determined by applying the initial drop in the O_3 signal. The PP substrate yielded a γ of $(3.5 \pm 2.1) \times 10^{-6}$, which is in agreement with the value reported by D'Anna et al. (2009). γ obtained from the 1:1 PP/LEV substrate is similar to γ determined from the individual LEV and PP substrates, within experimental error. γ derived from the 1:1, 1:10, and 1:100 PP/NG substrates are all similar to γ determined from the individual NG and PP substrates, within experimental error. γ of the 10:1:3 LEV/PP/NG substrate exhibited a higher uncertainty and the resulting γ values are similar to that of the individual substrates within the experimental error. The larger uncertainty may be caused by differences in surface composition of the organic substrate.

3.4.2 Visible irradiation experiments

Steady state γ values measured in the presence of VIS irradiation were conducted under dry conditions, with atmospherically relevant O_2 , and with substrate temperatures of 23 °C. Most experiments were conducted using the pullback method, however, some of the experiments using the 1:1 PP/NG substrate were conducted with the stepwise method for comparison. These two methods yielded similar γ values, within experimental error. As shown in Table 2, γ for the pure PP solution is enhanced by more than a factor of 2 when exposed to VIS irradiation. This is expected due to the photosensitive properties of humic acid, and is similar to the effects observed by D'Anna et al. (2009). However, the enhancement in γ determined here is smaller than that reported by D'Anna et al. (2009), which may be due to differences in the spectral distribution of the irradiation source used in this study. As shown in Fig. 5, PP absorbs radiation more strongly at lower visible wavelengths, but the visible irradiation source used in our study provides the strongest irradiation between 550–650 nm. The uptake of O_3 by the pure LEV and NG substrates yields no enhancement when exposed to VIS irradiation. As stated previously, LEV does not absorb radiation in the experimental wavelength regime. As shown in Fig. 5, NG absorbs only at wavelengths smaller than 400 nm under acidic conditions (pH = 2) and at wavelengths smaller than 500 nm under basic conditions (pH = 12). The sparsity of photon flux density of our VIS irradiation source at wavelengths between 400–550 nm may explain the insignificant enhancement in γ for pure NG when exposed to VIS irradiation. The addition of the PP photosensitiser to LEV does not yield an enhancement in O_3 uptake, within the experimental error. Although LEV should be able to act as an electron donor and react with the triplet excited state of PP (Alvarez et al., 2012), it is not more readily able to adsorb or react with O_3 . The addition of PP to NG yields enhanced uptake of O_3 by a factor of 3.6, as given in Table 2. The substituted phenolic structure of NG can efficiently absorb excitation energy from PP, and subsequently adsorb or react with O_3 . Decreasing the ratio of PP to NG causes a decrease in γ , as shown in Fig. 6. This is most likely due to the in-

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

creasing dominance of NG, which does not efficiently absorb VIS radiation. The 10:1:3 LEV/PP/NG substrate yields an enhanced γ under VIS irradiation that lies between the γ derived from the 1:1 and 1:10 PP/NG substrates.

3.4.3 UV irradiation experiments

5 Steady state γ values measured in the presence of UV irradiation were conducted under dry conditions, with atmospherically relevant O_2 , and with surface temperatures of $20^\circ C$. Similar to the VIS irradiation experiments, most of the experiments were performed using the pullback method, aside from a few experiments using the 1:1 PP/NG substrate that were performed using the stepwise method. These two methods also
10 yielded similar γ values, within the experimental error. The PP substrate yields a γ that is a factor of ~ 5 higher when exposed to UV radiation compared to dark values. As mentioned previously, this is slightly smaller than γ reported by D'Anna et al. (2009), most likely because the UV irradiation applied to the organic substrate was smaller in our study. The LEV substrate does not yield enhanced uptake of O_3 under UV irradiation within experimental error. However, the NG substrate yields significantly enhanced
15 γ values when exposed to UV irradiation. This is most likely because NG absorbs efficiently at these wavelengths, regardless of pH, as shown in Fig. 5. Previous studies have also shown that nitrated phenols absorb efficiently at wavelengths smaller than 450 nm (Jacobson, 1999). The addition of PP to the LEV substrate yields a slight enhancement in γ compared to dark values, but this enhancement is most likely due to PP, and not an energy transfer between PP and LEV. The addition of PP to NG also yields enhanced γ values compared to γ determined in the dark. However, there is little difference in γ when varying the mass ratios of PP to NG, as shown in Fig. 6. This is most likely due to the efficient absorption of UV radiation by NG. The 10:1:3
20 LEV/PP/NG substrate yields a slightly enhanced γ that is similar to that of the 1:1 PP/LEV substrate.

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5 Effect of photon flux on O₃ uptake

We measured γ of the 1:1 PP/NG substrate with O₃ as a function of irradiation intensity. These experiments were conducted in dry conditions with atmospherically relevant O₂, and [O₃] = $(4.60 \pm 0.53) \times 10^{11}$ molecule cm⁻³. We observe a linear dependence of γ on photon flux using both VIS and UV irradiation, as shown in Fig. 7. γ as a function of VIS photon flux can be described by a linear relationship in the form of $\gamma = 3.5 \times 10^{-21} \cdot \Phi + 1.3 \times 10^{-6}$, where Φ is the photon flux in photons cm⁻² s⁻¹. Similarly, γ as a function of UV photon flux can be expressed as $\gamma = 2.8 \times 10^{-21} \cdot \Phi + 3.8 \times 10^{-6}$.

3.6 Effect of O₃ concentration on O₃ uptake

Most of the experiments conducted in this study yielded an initial fast uptake of O₃ followed by a slower recovery of the O₃ signal, possibly indicating adsorption of gas molecules to the organic substrate. Hence, γ as a function of O₃ concentration is determined. For total VIS and UV irradiation fluxes of 5.9×10^{17} and 8.1×10^{15} photons cm⁻² s⁻¹, respectively, γ of the 1:1 PP/NG substrate with O₃ concentrations ranging from 1.74×10^{11} to 1.89×10^{12} molecule cm⁻³ have been determined. The inverse dependency of γ on O₃ concentration is presented in Fig. 8. Figure 8a shows that under VIS irradiation γ increases by a factor of ~ 6 when [O₃] decreases from 71.3 to 7.5 ppb. A similar dependency of γ on [O₃] is observed when the substrate is irradiated by UV as shown in Fig. 8b. γ increases by a factor of ~ 3 when [O₃] decreases from 73.7 to 7.7 ppb. These trends in γ with [O₃] are similar to the decreases in γ observed by D'Anna et al. (2009) using only PP as a substrate and UV-A irradiation.

A decrease in uptake when the concentration of the gas phase oxidant increases is commonly interpreted as a Langmuir-Hinshelwood type uptake mechanism, where oxidant adsorption is followed by reaction on the surface and potentially in the bulk (Pöschl et al., 2001; Ammann et al., 2003; Shiraiwa et al., 2012). The data shown in Fig. 8 are represented by a Langmuir-type isotherm of the form (Ammann et al., 2003):

$$\gamma = \frac{4k_s[\text{Org}]_s K_{\text{O}_3}}{c\sigma_{\text{O}_3}(1 + K_{\text{O}_3}[\text{O}_3]_g)}, \quad (3)$$

where k_s is the second order rate constant, $[\text{Org}]_s$ is the surface concentration of the organic species, K_{O_3} is the O_3 adsorption/desorption equilibrium constant, and σ_{O_3} is the surface area occupied by one adsorbed molecule of O_3 . The experimental data under VIS irradiation are represented by a fit of Eq. (3), as shown in Fig. 8a. Assuming $[\text{Org}]_s = 5.9 \times 10^{14} \text{ molecule cm}^{-2}$ and $\sigma_{\text{O}_3} = 1.7 \times 10^{-15} \text{ cm}^2 \text{ molecule}^{-1}$ (Pöschl et al., 2001; Shiraiwa et al., 2009; Springmann et al., 2009; Kaiser et al., 2011), the determined fit parameters are $k_s = (5.5 \pm 2.7) \times 10^{-19} \text{ cm}^2 \text{ s}^{-1} \text{ molecule}^{-1}$ and $K_{\text{O}_3} = (2.3 \pm 2.0) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1}$. Derived k_s is similar in magnitude to k_s derived in previous studies for O_3 uptake by solid PAHs (Pöschl et al., 2001; Shiraiwa et al., 2009). Derived K_{O_3} is larger than typical values reported in the literature (Pöschl et al., 2001; Kwamena et al., 2007; Shiraiwa et al., 2009). Pöschl et al. (2001) report $K_{\text{O}_3} = (2.8 \pm 0.2) \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1}$ for BaP coated soot with O_3 , however, most K_{O_3} values for uptake of O_3 by solid aerosol are on the order of 10^{-14} – $10^{-16} \text{ cm}^3 \text{ molecule}^{-1}$ (Shiraiwa et al., 2009). Kwamena et al. (2007) suggest that O_3 preferentially partitions to nonpolar surfaces. This could explain the large K_{O_3} value observed by Pöschl et al. (2001) since soot surfaces are highly nonpolar. However, both PP and NG are polar substances, so this cannot explain the high K_{O_3} determined in our study. Kwamena et al. (2007) also suggest that a higher value of K_{O_3} may be due to a stronger binding interaction of the initial O_3 /substrate complex. In this case, the excitation of the PP/NG substrate upon irradiation may cause O_3 to bind more strongly to the surface, yielding a higher value for K_{O_3} .

The experimental data under UV irradiation are represented by a fit of Eq. (3), as shown in Fig. 8b. Using the same values for $[\text{Org}]_s$ and σ_{O_3} , the determined fit parameters are: $k_s = (8.1 \pm 2.0) \times 10^{-19} \text{ cm}^2 \text{ s}^{-1} \text{ molecule}^{-1}$ and $K_{\text{O}_3} = (1.7 \pm 0.7) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1}$. Similar to the experimental data under VIS irradiation, derived

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


k_s is similar in magnitude to k_s derived in previous studies (Pöschl et al., 2001; Shiraiwa et al., 2009). Derived K_{O_3} is similar to that obtained for the VIS uptake experiments, indicating that also for the UV case, the presence of irradiation may enhance the binding of O_3 to the organic substrate.

4 Atmospheric implications

This study shows that NG in the presence of PP acting as a photosensitiser exhibits enhanced O_3 uptake under VIS and UV irradiation with potential consequences for atmospheric chemistry. O_3 uptake by LEV with and without PP acting as a photosensitiser is not enhanced compared to the individual components under VIS and UV irradiation.

An interesting component of this study is the importance of UV radiation compared to VIS irradiation on the photosensitised heterogeneous oxidation kinetics. For example, the total UV photon flux is almost 2 orders of magnitude smaller than for the VIS regime, but the γ values are about 15% larger under UV irradiation. This indicates that photoenhanced chemical ageing will be important under UV-A irradiation, even though the UV-A photon flux from the sun is much less than that of the VIS photon flux. Furthermore, since UV-A irradiation increases with height, it can be expected that organic particles containing photosensitising compounds, such as BBA, in the upper troposphere or lower stratosphere (UT/LS) can experience significant photosensitised chemical ageing during transport.

As shown above, γ for the 1:1 PP/NG substrate can be expressed as a function of photon flux by a linear relationship. This relationship can be exploited to determine γ under VIS irradiation from the sun as $\gamma = 4.8 \times 10^{-6}$ for an irradiance of 1.0×10^{17} photons $\text{cm}^{-2} \text{s}^{-1}$ in the 400–700 region. As mentioned previously, the VIS irradiation source emits predominantly in the “red” wavelength range compared to the wavelengths emitted by the sun. As a consequence, γ determined here may be considered a lower limit since under atmospheric conditions light is received more uniformly as a function of wavelength, as shown in Fig. 2. γ under UV irradiation from the sun

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as determined from the above linear relationship is 1.97×10^{-5} for an irradiance of $5.6 \times 10^{15} \text{ photons cm}^{-2} \text{ s}^{-1}$ (300–400 nm). However, the UV irradiation source used in this study peaks between 362 and 372 nm, whereas the sun emits irradiation continually from 300–400 nm. Thus, also in this case, the derived γ may represent a lower limit.

Previous studies have shown that the number of gas molecules lost to the organic substrate will be proportional to the product of γ and the concentration of the oxidant species (Moise and Rudich, 2001; Knopf et al., 2006; Gross and Bertram, 2008; Knopf et al., 2011). Therefore, we utilise the γ values determined in this study to report oxidative power, i.e. $\gamma \cdot [\text{O}_3]$. This parameter allows one to evaluate the importance of different atmospheric oxidants in terms of gas phase loss, effect on atmospheric composition and chemistry, and oxidation potential of the condensed phase species. Shown in Fig. 9a is the oxidative power for the 1:1 PP/NG substrate as a function of O_3 concentration under VIS and UV irradiation. γ values used for calculation of the oxidative power are determined from the LH relationships given in Fig. 8a, b, and at irradiation of 5.9×10^{17} and $8.1 \times 10^{15} \text{ photons cm}^{-2} \text{ s}^{-1}$ for VIS and UV, respectively. For atmospherically relevant levels of O_3 between 1 ppb and 120 ppb, the oxidative power is greater than $1 \times 10^6 \text{ molecule cm}^{-3}$. The determined $\gamma \cdot [\text{O}_3]$ are about an order of magnitude greater than those of solid PAH surfaces reacting with 100 ppb O_3 (Gross and Bertram, 2008). At 100 ppb, $\gamma \cdot [\text{O}_3]$ is $1.8 \times 10^7 \text{ molecule cm}^{-3}$ under VIS irradiation and $2.5 \times 10^7 \text{ molecule cm}^{-3}$ under UV irradiation. $\gamma \cdot [\text{O}_3]$ can be determined for any other O_3 concentration using the fit parameters derived from the above Langmuir isotherms to calculate γ . Although the uptake of O_3 may be slow compared to the uptake of OH radical, the abundance of O_3 compared to OH yields $\gamma \cdot [\text{O}_3]$ values that are about twice that of $\gamma \cdot [\text{OH}]$ (Yokelson et al., 2009; Slade and Knopf, 2013).

$\gamma \cdot [\text{O}_3]$ is also determined as a function of photon flux, as shown in Fig. 9b. At a fixed $[\text{O}_3]$ of $\sim 20 \text{ ppb}$, $\gamma \cdot [\text{O}_3]$ is greater than $1 \times 10^6 \text{ molecule cm}^{-3}$ for all given photon fluxes.

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

$\gamma \cdot [\text{O}_3]$ for the average total VIS and UV irradiances from the sun are 1.2×10^7 and $4.9 \times 10^7 \text{ molecule cm}^{-3}$, respectively.

We can use measured γ to estimate the particle lifetime, τ , of NG, which is suggested to serve as a molecular marker for BBA particles. We assume NG is in the presence of PP acting as a photosensitiser, however we do not account for degradation of PP. Particle lifetime is calculated following Robinson et al. (2006a):

$$\tau = \frac{4}{3} r \frac{\rho N_a}{\gamma c [\text{O}_3] M}, \quad (4)$$

where r is the radius of the particle, ρ is the particle density (1.367 g cm^{-3} for NG, (Dinar et al., 2006)), and M is the molecular weight of the organic species of interest ($M = 169.13 \text{ g mol}^{-1}$ for NG). Here we assume a $0.4 \mu\text{m}$ diameter NG particle.

Shown in Fig. 10a is the lifetime of a NG particle as a function of $[\text{O}_3]$ under VIS and UV irradiation. γ values used in calculating the lifetime are determined from the LH relationships given in Fig. 8a, b, and at irradiation of 5.9×10^{17} and $8.1 \times 10^{15} \text{ photons cm}^{-2} \text{ s}^{-1}$ for VIS and UV, respectively. For atmospherically relevant levels of O_3 between 20 ppb and 120 ppb, particle lifetimes range from 39 to 88 daylight hours. The lifetime can be determined for any $[\text{O}_3]$ using the fit parameters derived from the above Langmuir-type isotherms to calculate γ . Figure 10b shows the particle lifetime as a function of photon flux, with $[\text{O}_3] \sim 20 \text{ ppb}$. The lifetimes range from 91 to 1490 daylight hours under VIS irradiation and 75 to 520 daylight hours under UV irradiation. These results show that under typical atmospheric transport times of about 1 week, the biomolecular marker NG can be significantly oxidised in the presence of a photosensitiser and VIS or UV irradiation. As discussed above, due to the spectral characteristics of the applied irradiation sources and absorbance features of NG, derived γ values may represent lower limits making these lifetime estimates upper limits. However, shielding effects due to the presence of non-reactive compounds at the surface of the particle have not been accounted for and may extend particle lifetime.

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Smoke and other absorbing aerosol have been observed in the UT/LS (Jost et al., 2004; Fromm et al., 2010). A study by Fromm et al. (2010) proposes that injection of BB emissions through pyrocumulonimbus (pyroCb) clouds could be an important source of stratospheric aerosol particles. The aerosol particles injected by pyroCb are transported throughout the UT/LS and can subsequently undergo chemical ageing. Due to the sufficiently high altitude of pyroCb smoke, the aerosol particles are exposed to much higher levels of UV-A irradiation. According to Piazena (1996), UV-A irradiation at 10 km height can be 70–150 % higher than at ground level depending on solar elevation. Coupled with the high levels of O₃ found in aged BB plumes (Delany et al., 1985; Andreae et al., 1988; Crutzen and Andreae, 1990; Ziemke et al., 2009; Konovalov et al., 2011; Dupont et al., 2012; Martins et al., 2012; Akagi et al., 2013), uptake of O₃ by BBA compounds can be significant. For example, using a solar elevation angle of ~ 70 °C which roughly corresponds to a summer afternoon in the midwestern US, UV-A irradiation at 10 km height is $9.52 \times 10^{15} \text{ photons cm}^{-2} \text{ s}^{-1}$. This corresponds to a γ value of 3.05×10^{-5} using the linear relationship of γ with photon flux. This corresponds to a NG particle lifetime of about 40 daylight hours at 100 ppb.

5 Conclusions

A novel irradiated rectangular channel flow reactor coupled to CIMS has been developed and successfully validated allowing for the investigation of the effect of irradiation on the heterogeneous oxidation kinetics of organic compounds typical of biomass burning aerosol at low pressures and applying atmospherically relevant oxidant concentrations. The effect of VIS and UV irradiation on the reactive uptake coefficients of levoglucosan and 5-nitroguaiacol with O₃ has been determined with and without the presence of Pahokee peat acting as a photosensitiser. The reactive uptake of O₃ by a 10:1:3 by mass mixture of levoglucosan, Pahokee peat, and 5-nitroguaiacol in dark, VIS, and UV irradiation conditions was also determined. The uptake of O₃ by LEV without PP yields no enhancement when exposed to VIS or UV irradiation, while the uptake

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of O_3 by NG without PP is enhanced by a factor of ~ 6 when exposed to UV irradiation, yielding $\gamma = 3.6 \times 10^{-5}$. γ of the 1:1 by mass PP/LEV substrate is not enhanced when exposed to either VIS or UV irradiation, within experimental error. However, γ of the 1:1 by mass PP/NG substrate is enhanced by about an order of magnitude under both VIS and UV irradiation, yielding values of 2.4×10^{-5} and 2.8×10^{-5} , respectively. γ of the PP/NG substrates decreases (remains constant) as the mass ratio of PP to NG decreases under VIS (UV) irradiation. γ of the 10:1:3 LEV/PP/NG substrate is also enhanced under both VIS and UV irradiation, yielding values of 2.8×10^{-5} and 1.4×10^{-5} , respectively.

We observed that the reactive uptake coefficient for constant O_3 concentration by a 1:1 PP/NG substrate increases linearly as a function of VIS and UV irradiation, allowing for extrapolation of γ to atmospherically relevant photon fluxes. Determined γ of the 1:1 PP/NG substrate under constant VIS and UV irradiation decreases as O_3 concentration increases. This behaviour is interpreted as a Langmuir–Hinshelwood type uptake mechanism and the data is represented by a Langmuir-type isotherm. From the experimental data under VIS irradiation, the second order rate constant, $k_s = (5.5 \pm 2.7) \times 10^{-19} \text{ cm}^2 \text{ s}^{-1} \text{ molecule}^{-1}$ and the O_3 adsorption/desorption equilibrium constant, $K_{O_3} = (2.3 \pm 2.0) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1}$. O_3 uptake by the 1:1 PP/NG substrate under UV irradiation yields the analogous parameters: $k_s = (8.1 \pm 2.0) \times 10^{-19} \text{ cm}^2 \text{ s}^{-1} \text{ molecule}^{-1}$ and $K_{O_3} = (1.7 \pm 0.7) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1}$.

The experimental data allows for derivation of the oxidative power as a function of the sun's photon flux and O_3 concentration, indicating that for atmospherically relevant conditions, these BBA surrogate species can be efficiently oxidised with subsequent implications for source apportionment studies. Estimated 5-nitroguaiacol particle lifetimes indicate that these BBA biomarkers can undergo significant degradation during typical atmospheric transport times of up to one week. Since the sun emits radiation with smaller wavelengths than applied irradiation sources, derived γ and lifetimes may represent lower and upper limits, respectively. In addition, since the VIS and UV flux

can increase significantly with altitude, photosensitive chemical ageing may be even more efficient in the upper troposphere/lower stratosphere.

Acknowledgements. The authors are grateful to J. Slechta for technical support. We would also like to thank A. Sedlacek for use of the fiber optic spectrometer, and J. Marecek for performing the absorbance measurements. This work was supported by the National Science Foundation, Grant AGS-0846255.

References

- Akagi, S. K., Yokelson, R. J., Burling, I. R., Meinardi, S., Simpson, I., Blake, D. R., McMeeking, G. R., Sullivan, A., Lee, T., Kreidenweis, S., Urbanski, S., Reardon, J., Griffith, D. W. T., Johnson, T. J., and Weise, D. R.: Measurements of reactive trace gases and variable O₃ formation rates in some South Carolina biomass burning plumes, *Atmos. Chem. Phys.*, 13, 1141–1165, doi:10.5194/acp-13-1141-2013, 2013. 7556, 7574
- Alvarez, E. G., Wortham, H., Strekowski, R., Zetzsch, C., and Gligorovski, S.: Atmospheric photosensitized heterogeneous and multiphase reactions: from outdoors to indoors, *Environ. Sci. Technol.*, 46, 1955–1963, doi:10.1021/es2019675, 2012. 7557, 7567
- Ammann, M., Pöschl, U., and Rudich, Y.: Effects of reversible adsorption and Langmuir–Hinshelwood surface reactions on gas uptake by atmospheric particles, *Phys. Chem. Chem. Phys.*, 5, 351–356, doi:10.1039/B208708A, 2003. 7569
- Anastasio, C., Faust, B. C., and Rao, C. J.: Aromatic carbonyl compounds as aqueous-phase photochemical sources of hydrogen peroxide in acidic sulfate aerosols, fogs, and clouds, 1. Non-phenolic methoxybenzaldehydes and methoxyacetophenones with reductants (phenols), *Environ. Sci. Technol.*, 31, 218–232, doi:10.1021/es960359g, 1997. 7558
- Andreae, M. O.: A new look at aging aerosols, *Science*, 326, 1493–1494, doi:10.1126/science.1183158, 2009. 7556
- Andreae, M. O., Browell, E. V., Garstang, M., Gregory, G. L., Harriss, R. C., Hill, G. F., Jacob, D. J., Pereira, M. C., Sachse, G. W., Setzer, A. W., Dias, P. L. S., Talbot, R. W., Torres, A. L., and Wofsy, S. C.: Biomass-burning emissions and associated haze layers over amazonia, *J. Geophys. Res.*, 93, 1509–1527, doi:10.1029/JD093iD02p01509, 1988. 7556, 7574

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Atkinson, R. and Carter, W. P. L.: Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions, *Chem. Rev.*, 84, 437–470, doi:10.1021/cr00063a002, 1984. 7566
- Baduel, C., Monge, M. E., Voisin, D., Jaffrezo, J.-L., George, C., Haddad, I. E., Marchand, N., and D'Anna, B.: Oxidation of atmospheric humic like substances by ozone: a kinetic and structural analysis approach, *Environ. Sci. Technol.*, 45, 5238–5244, doi:10.1021/es200587z, 2011. 7558
- Brigante, M., Cazoir, D., D'Anna, B., George, C., and Donaldson, D. J.: Photoenhanced uptake of NO₂ by pyrene solid films, *J. Phys. Chem. A*, 112, 9503–9508, doi:10.1021/jp802324g, 2008. 7558
- Canonica, S., Jans, U., Stemmler, K., and Hoigné, J.: Transformation kinetics of phenols in water: photosensitization by dissolved natural organic material and aromatic ketones, *Environ. Sci. Technol.*, 29, 1822–1831, doi:10.1021/es00007a020, 1995. 7558
- Canonica, S., Hellrung, B., and Wirz, J.: Oxidation of phenols by triplet aromatic ketones in aqueous solution, *J. Phys. Chem. A*, 104, 1226–1232, doi:10.1021/jp9930550, 2000. 7558
- Crutzen, P. J. and Andreae, M. O.: Biomass burning in the tropics – impact on atmospheric chemistry and biogeochemical cycles, *Science*, 250, 1669–1678, doi:10.1126/science.250.4988.1669, 1990. 7555, 7556, 7574
- Crutzen, P. J., Heidt, L. E., Krasnec, J. P., Pollock, W. H., and Seiler, W.: Biomass burning as a source of atmospheric gases CO, H₂, N₂O, NO, CH₃Cl and COS, *Nature*, 282, 253–256, doi:10.1038/282253a0, 1979. 7555
- D'Anna, B., Jammoul, A., George, C., Stemmler, K., Fahrni, S., Amman, M., and Wisthaler, A.: Light-induced ozone depletion by humic acid films and submicron aerosol particles, *J. Geophys. Res.*, 114, D12301, doi:10.1029/2008JD011237, 2009. 7558, 7563, 7566, 7567, 7568, 7569, 7587
- de Gouw, J. A. and Lovejoy, E. R.: Reactive uptake of ozone by liquid organic compounds, *Geophys. Res. Lett.*, 25, 931–934, doi:10.1029/98GL00515, 1998. 7587
- Delany, A. C., Haagensen, P., Walters, S., Wartburg, A. F., and Crutzen, P. J.: Photochemically produced ozone in the emission from large-scale tropical vegetation fires, *J. Geophys. Res.*, 90, 2425–2429, doi:10.1029/JD090iD01p02425, 1985. 7556, 7574
- Dinar, E., Mentel, T. F., and Rudich, Y.: The density of humic acids and humic like substances (HULIS) from fresh and aged wood burning and pollution aerosol particles, *Atmos. Chem. Phys.*, 6, 5213–5224, doi:10.5194/acp-6-5213-2006, 2006. 7573

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Dupont, R., Pierce, B., Worden, J., Hair, J., Fenn, M., Hamer, P., Natarajan, M., Schaack, T., Lenzen, A., Apel, E., Dibb, J., Diskin, G., Huey, G., Weinheimer, A., Kondo, Y., and Knapp, D.: Attribution and evolution of ozone from Asian wild fires using satellite and aircraft measurements during the ARCTAS campaign, *Atmos. Chem. Phys.*, 12, 169–188, doi:10.5194/acp-12-169-2012, 2012. 7556, 7574
- Dusek, U., Frank, G. P., Massling, A., Zeromskiene, K., Iinuma, Y., Schmid, O., Helas, G., Hennig, T., Wiedensohler, A., and Andreae, M. O.: Water uptake by biomass burning aerosol at sub- and supersaturated conditions: closure studies and implications for the role of organics, *Atmos. Chem. Phys.*, 11, 9519–9532, doi:10.5194/acp-11-9519-2011, 2011. 7555
- Engelhart, G. J., Hennigan, C. J., Miracolo, M. A., Robinson, A. L., and Pandis, S. N.: Cloud condensation nuclei activity of fresh primary and aged biomass burning aerosol, *Atmos. Chem. Phys.*, 12, 7285–7293, doi:10.5194/acp-12-7285-2012, 2012. 7555
- Facchini, M. C., Decesari, S., Mircea, M., Fuzzi, S., and Loglio, G.: Surface tension of atmospheric wet aerosol and cloud/fog droplets in relation to their organic carbon content and chemical composition, *Atmos. Environ.*, 34, 4853–4857, doi:10.1016/S1352-2310(00)00237-5, 2000. 7557
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from fireplace combustion of woods grown in the northeastern United States, *Environ. Sci. Technol.*, 35, 2665–2675, doi:10.1021/es001466k, 2001. 7558
- Finlayson-Pitts, B. J. and Pitts, J. N.: *Chemistry of the Upper and Lower Atmosphere*, Academic Press, San Diego, California, 2000. 7556
- Fors, E. O., Rissler, J., Massling, A., Svenningsson, B., Andreae, M. O., Dusek, U., Frank, G. P., Hoffer, A., Bilde, M., Kiss, G., Janitsek, S., Henning, S., Facchini, M. C., Decesari, S., and Swietlicki, E.: Hygroscopic properties of Amazonian biomass burning and European background HULIS and investigation of their effects on surface tension with two models linking H-TDMA to CCNC data, *Atmos. Chem. Phys.*, 10, 5625–5639, doi:10.5194/acp-10-5625-2010, 2010. 7557
- Fromm, M., Lindsey, D. T., Servranckx, R., Yue, G., Trickl, T., Sica, R., Doucet, P., and Godin-Beekmann, S.: The untold story of pyrocumulonimbus, *B. Am. Meteorol. Soc.*, 91, 1193–1209, doi:10.1175/2010BAMS3004.1, 2010. 7574
- Fu, P. Q., Kawamura, K., Chen, J., Li, J., Sun, Y. L., Liu, Y., Tachibana, E., Aggarwal, S. G., Okuzawa, K., Tanimoto, H., Kanaya, Y., and Wang, Z. F.: Diurnal variations of organic molecular tracers and stable carbon isotopic composition in atmospheric aerosols over Mt. Tai in

- the North China Plain: an influence of biomass burning, *Atmos. Chem. Phys.*, 12, 8359–8375, doi:10.5194/acp-12-8359-2012, 2012. 7556
- Gelencsér, A., Hoffer, A., Kiss, G., Tombacz, E., Kurdi, R., and Bencze, L.: In-situ formation of light-absorbing organic matter in cloud water, *J. Atmos. Chem.*, 45, 25–33, doi:10.1023/A:1024060428172, 2003. 7557
- George, C., Strekowski, R. S., Kleffmann, J., Stemmmler, K., and Ammann, M.: Photoenhanced uptake of gaseous NO₂ on solid-organic compounds: a photochemical source of HONO?, *Faraday Discuss.*, 130, 195–210, doi:10.1039/b417888m, 2005. 7557, 7558
- Gross, S. and Bertram, A. K.: Reactive uptake of NO₃, N₂O₅, NO₂, HNO₃, and O₃ on three types of polycyclic aromatic hydrocarbon surfaces, *J. Phys. Chem. A*, 112, 3104–3113, doi:10.1021/jp7107544, 2008. 7572
- Gueymard, C., Myers, D., and Emery, K.: Proposed reference irradiance spectra for solar energy systems testing, *Sol. Energy*, 73, 443–467, doi:10.1016/S0038-092X(03)00005-7, 2002. 7559, 7563
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009. 7555
- Hanson, D. R.: Reaction of ClONO₂ with H₂O and HCl in sulfuric acid and HNO₃/H₂SO₄/H₂O mixtures, *J. Phys. Chem. A*, 102, 4794–4807, doi:10.1021/jp972767s, 1998. 7565
- Hawthorne, S. B., Miller, D. J., Barkley, R. M., and Krieger, M. S.: Identification of methoxylated phenols as candidate tracers for atmospheric wood smoke pollution, *Environ. Sci. Technol.*, 22, 1191–1196, doi:10.1021/es00175a011, 1988. 7556
- Hawthorne, S. B., Krieger, M. S., Miller, D. J., and Mathiason, M. B.: Collection and quantitation of methoxylated phenol tracers for atmospheric pollution from residential wood stoves, *Environ. Sci. Technol.*, 23, 470–475, doi:10.1021/es00181a013, 1989. 7556
- Hennigan, C. J., Sullivan, A. P., Collett Jr., J. L., and Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, L09805, doi:10.1029/2010GL043088, 2010. 7556

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hennigan, C. J., Miracolo, M. A., Engelhart, G. J., May, A. A., Presto, A. A., Lee, T., Sullivan, A. P., McMeeking, G. R., Coe, H., Wold, C. E., Hao, W.-M., Gilman, J. B., Kuster, W. C., de Gouw, J., Schichtel, B. A., J. L. Collett Jr., Kreidenweis, S. M., and Robinson, A. L.: Chemical and physical transformations of organic aerosol from the photo-oxidation of open biomass burning emissions in an environmental chamber, *Atmos. Chem. Phys.*, 11, 7669–7686, doi:10.5194/acp-11-7669-2011, 2011. 7556
- Hobbs, P. V. and Radke, L. F.: Cloud condensation nuclei from a simulated forest fire, *Science*, 163, 279–280, doi:10.1126/science.163.3864.279, 1969. 7555
- Hoffmann, N.: Photochemical reactions as key steps in organic synthesis, *Chem. Rev.*, 108, 1052–1103, doi:10.1021/cr0680336, 2008. 7557
- Jacobson, M. Z.: Isolating nitrated and aromatic aerosols and nitrated aromatic gases as sources of ultraviolet light absorption, *J. Geophys. Res.*, 104, 3527–3542, doi:10.1029/1998JD100054, 1999. 7568
- Jammoul, A., Gligorovski, S., George, C., and D'Anna, B.: Photosensitized heterogeneous chemistry of ozone on organic films, *J. Phys. Chem. A*, 112, 1268–1276, doi:10.1021/jp074348t, 2008. 7557, 7558, 7563
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009. 7556
- Jost, H.-J., Drdla, K., Stohl, A., Pfister, L., Loewenstein, M., Lopez, J. P., Hudson, P. K., Murphy, D. M., Cziczo, D. J., Fromm, M., Bui, T. P., Dean-Day, J., Gerbig, C., Mahoney, M. J., Richard, E. C., Spichtinger, N., Pittman, J. V., Weinstock, E. M., Wilson, J. C., and Xueref, I.: In-situ observations of mid-latitude forest fire plumes deep in the stratosphere, *Geophys. Res. Lett.*, 31, L11101, doi:10.1029/2003GL019253, 2004. 7574

- Kaiser, J. C., Riemer, N., and Knopf, D. A.: Detailed heterogeneous oxidation of soot surfaces in a particle-resolved aerosol model, *Atmos. Chem. Phys.*, 11, 4505–4520, doi:10.5194/acp-11-4505-2011, 2011. 7570
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005. 7555
- Kessler, S. H., Smith, J. D., Che, D. L., Worsnop, D. R., Wilson, K. R., and Kroll, J. H.: Chemical sinks of organic aerosol: kinetics and products of the heterogeneous oxidation of erythritol and levoglucosan, *Environ. Sci. Technol.*, 44, 7005–7010, doi:10.1021/es101465m, 2010. 7556
- Knopf, D. A., Anthony, L. M., and Bertram, A. K.: Reactive uptake of O_3 by multicomponent and multiphase mixtures containing oleic acid, *J. Phys. Chem. A*, 109, 5579–5589, doi:10.1021/jp0512513, 2005. 7561
- Knopf, D. A., Mak, J., Gross, S., and Bertram, A. K.: Does atmospheric processing of saturated hydrocarbon surfaces by NO_3 lead to volatilization?, *Geophys. Res. Lett.*, 33, L17816, doi:10.1029/2006GL026884, 2006. 7556, 7562, 7572
- Knopf, D. A., Cosman, L. M., Mousavi, P., Mokamati, S., and Bertram, A. K.: A novel flow reactor for studying reactions on liquid surfaces coated by organic monolayers: methods, validation, and initial results, *J. Phys. Chem. A*, 111, 11021–11032, doi:10.1021/jp075724c, 2007. 7559, 7564, 7565
- Knopf, D. A., Forrester, S. M., and Slade, J. H.: Heterogeneous oxidation kinetics of organic biomass burning aerosol surrogates by O_3 , NO_2 , N_2O_5 , and NO_3 , *Phys. Chem. Chem. Phys.*, 13, 21050–21062, doi:10.1039/C1CP22478F, 2011. 7556, 7557, 7559, 7561, 7562, 7565, 7566, 7572, 7587
- Konovalov, I. B., Beekmann, M., Kuznetsova, I. N., Yurova, A., and Zvyagintsev, A. M.: Atmospheric impacts of the 2010 Russian wildfires: integrating modelling and measurements of an extreme air pollution episode in the Moscow region, *Atmos. Chem. Phys.*, 11, 10031–10056, doi:10.5194/acp-11-10031-2011, 2011. 7556, 7574
- Konovalov, I. B., Beekmann, M., D'Anna, B., and George, C.: Significant light induced ozone loss on biomass burning aerosol: evidence from chemistry-transport modeling based on

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

new laboratory studies, *Geophys. Res. Lett.*, 39, L17807, doi:10.1029/2012GL052432, 2012. 7557, 7558

Kwamena, N.-O. A., Staikova, M. G., Donaldson, D. J., George, I. J., and Abbatt, J. P. D.: Role of the aerosol substrate in the heterogeneous ozonation reactions of surface-bound PAHs, *J. Phys. Chem. A*, 111, 11050–11058, doi:10.1021/jp075300i, 2007. 7570

Levin, E. J. T., McMeeking, G. R., Carrico, C. M., Mack, L. E., Kreidenweis, S. M., Wold, C. E., Moosmüller, H., Arnott, W. P., Hao, W. M., Collett, J. L., J., and Malm, W. C.: Biomass burning smoke aerosol properties measured during Fire Laboratory at Missoula Experiments (FLAME), *J. Geophys. Res.*, 115, D18210, doi:10.1029/2009jd013601, 2010. 7555

Lin, L., Lee, M. L., and Eatough, D. J.: Review of recent advances in detection of organic markers in fine particulate matter and their use for source apportionment, *J. Air Waste Manage. Assoc.*, 60, 3–25, 2010. 7556

Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: a global perspective, *J. Geophys. Res.*, 86, 7210–7254, doi:10.1029/JC086iC08p07210, 1981. 7555

Lukács, H., Gelencsér, A., Hammer, S., Puxbaum, H., Pio, C., Legrand, M., Kasper-Giebl, A., Handler, M., Limbeck, A., Simpson, D., and Preunkert, S.: Seasonal trends and possible sources of brown carbon based on 2-yr aerosol measurements at six sites in Europe, *Geophys. Res. Lett.*, 112, D23S18, doi:10.1029/2006jd008151, 2007. 7557

Maria, S. F., Russell, L. M., Gilles, M. K., and Myneni, S. C. B.: Organic aerosol growth mechanisms and their climate-forcing implications, *Science*, 306, 1921–1924, doi:10.1126/science.1103491, 2004. 7556

Martins, V., Miranda, A., Carvalho, A., Schaap, M., Borrego, C., and Sá, E.: Impact of forest fires on particulate matter and ozone levels during the 2003, 2004 and 2005 fire seasons in Portugal, *Sci. Total Environ.*, 414, 53–62, doi:10.1016/j.scitotenv.2011.10.007, 2012. 7556, 7574

Mayol-Bracero, O. L., Guyon, P., Graham, B., Roberts, G., Andreae, M. O., Decesari, S., Facchini, M. C., Fuzzi, S., and Artaxo, P.: Water-soluble organic compounds in biomass burning aerosols over Amazonia, 2. apportionment of the chemical composition and importance of the polyacidic fraction, *J. Geophys. Res.*, 107, 8091–8106, doi:10.1029/2001JD000522, 2002. 7557

Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental ev-

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

idence for restructuring, phase transitions and kinetic limitations, *Atmos. Chem. Phys.*, 9, 9491–9522, doi:10.5194/acp-9-9491-2009, 2009. 7566

Moise, T. and Rudich, Y.: Reactive uptake of ozone by proxies for organic aerosols: surface versus bulk processes, *J. Geophys. Res.*, 105, 14667–14676, doi:10.1029/2000JD900071, 2000. 7564

Moise, T. and Rudich, Y.: Uptake of Cl and Br by organic surfaces perspective on organic aerosols processing by tropospheric oxidants, *Geophys. Res. Lett.*, 28, 4083–4086, doi:10.1029/2001GL013583, 2001. 7572

Moise, T. and Rudich, Y.: Reactive uptake of ozone by aerosol-associated unsaturated fatty acids: kinetics, mechanism, and products, *J. Phys. Chem. A*, 106, 6469–6476, doi:10.1021/jp025597e, 2002. 7565, 7587

Net, S., Nieto-Gligorovski, L., Gligorovski, S., and Wortham, H.: Heterogeneous ozonation kinetics of 4-phenoxyphenol in the presence of photosensitizer, *Atmos. Chem. Phys.*, 10, 1545–1554, doi:10.5194/acp-10-1545-2010, 2010. 7558

Nissenson, P., Knox, C. J. H., Finlayson-Pitts, B. J., Phillips, L. F., and Dabdub, D.: Enhanced photolysis in aerosols: evidence for important surface effects, *Phys. Chem. Chem. Phys.*, 8, 4700–4710, doi:10.1039/B609219E, 2006. 7558

Piazena, H.: The effect of altitude upon the solar UV-B and UV-A irradiance in the tropical Chilean Andes, *Sol. Energy*, 57, 133–140, doi:10.1016/S0038-092X(96)00049-7, 1996. 7574

Pope, C. A., Ezzati, M., and Dockery, D. W.: Fine-particulate air pollution and life expectancy in the United States, *N. Engl. J. Med.*, 360, 376–386, doi:10.1056/NEJMsa0805646, 2009. 7555

Pöschl, U., Letzel, T., Schauer, C., and Niessner, R.: Interaction of ozone and water vapor with spark discharge soot aerosol particles coated with benzo[a]pyrene: O₃ and H₂O adsorption, benzo[a]pyrene degradation, and atmospheric implications, *J. Phys. Chem. A*, 105, 4029–4041, doi:10.1021/jp004137n, 2001. 7569, 7570, 7571

Ramanathan, V., Crutzen, P. J., and Rosenfeld, D.: Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124, doi:10.1126/science.1064034, 2001. 7555

Robinson, A. L., Donahue, N. M., and Rogge, W. F.: Photochemical oxidation and changes in molecular composition of organic aerosol in the regional context, *J. Geophys. Res.*, 111, D03302, doi:10.1029/2005JD006265, 2006a. 7573

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Robinson, A. L., Subramanian, R., Donahue, N. M., Bernardo-Bricker, A., and Rogge, W. F.: Source apportionment of molecular markers and organic aerosol, 2. biomass smoke, *Environ. Sci. Technol.*, 40, 7811–7819, doi:10.1021/es060782h, 2006b. 7556
- Rogge, W. F., Hildemann, L. M., Mazurek, M. A., and Cass, G. R.: Sources of fine organic aerosol, 9. pine, oak, and synthetic log combustion in residential fireplaces, *Environ. Sci. Technol.*, 32, 13–22, doi:10.1021/es960930b, 1998. 7558
- Rudich, Y., Donahue, N. M., and Mentel, T. F.: Aging of organic aerosol: bridging the gap between laboratory and field studies, *Annu. Rev. Phys. Chem.*, 58, 321–352, doi:10.1146/annurev.physchem.58.032806.104432, 2007. 7556
- Schmidl, C., Bauer, H., Dattler, A., Hitzenberger, R., Weissenboeck, G., Marr, I. L., and Puxbaum, H.: Chemical characterisation of particle emissions from burning leaves, *Atmos. Environ.*, 42, 9070–9079, doi:10.1016/j.atmosenv.2008.09.010, 2008a. 7557
- Schmidl, C., Marr, L. L., Caseiro, A., Kotianova, P., Berner, A., Bauer, H., Kasper-Giebl, A., and Puxbaum, H.: Chemical characterisation of fine particle emissions from wood stove combustion of common woods growing in mid-European Alpine regions, *Atmos. Environ.*, 42, 126–141, doi:10.1016/j.atmosenv.2007.09.028, 2008b. 7557
- Shiraiwa, M., Garland, R. M., and Pöschl, U.: Kinetic double-layer model of aerosol surface chemistry and gas-particle interactions (K2-SURF): degradation of polycyclic aromatic hydrocarbons exposed to O₃, NO₂, H₂O, OH and NO₃, *Atmos. Chem. Phys.*, 9, 9571–9586, doi:10.5194/acp-9-9571-2009, 2009. 7570, 7571
- Shiraiwa, M., Pöschl, U., and Knopf, D. A.: Multiphase chemical kinetics of NO₃ radicals reacting with organic aerosol components from biomass burning, *Environ. Sci. Technol.*, 46, 6630–6636, doi:10.1021/es300677a, 2012. 7556, 7569
- Simoneit, B. R. T.: Biomass burning – a review of organic tracers for smoke from incomplete combustion, *Appl. Geochem.*, 17, 129–162, doi:10.1016/S0883-2927(01)00061-0, 2002. 7556
- Slade, J. H. and Knopf, D. A.: Heterogeneous OH oxidation of biomass burning organic aerosol surrogate compounds: assessment of volatilisation products and the role of OH concentration on the reactive uptake kinetics, *Phys. Chem. Chem. Phys.*, doi:10.1039/C3CP44695F, 2013. 7557, 7572
- Sosedova, Y., Rouviere, A., Bartels-Rausch, T., and Ammann, M.: UVA/Vis-induced nitrous acid formation on polyphenolic films exposed to gaseous NO₂, *Photochem. Photobiol. Sci.*, 10, 1680–1690, doi:10.1039/C1PP05113J, 2011. 7557, 7558

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Springmann, M., Knopf, D. A., and Riemer, N.: Detailed heterogeneous chemistry in an urban plume box model: reversible co-adsorption of O₃, NO₂, and H₂O on soot coated with benzo[a]pyrene, *Atmos. Chem. Phys.*, 9, 7461–7479, doi:10.5194/acp-9-7461-2009, 2009. 7570
- 5 Stemmler, K., Ammann, M., Donders, C., Kleffmann, J., and George, C.: Photosensitized reduction of nitrogen dioxide on humic acid as a source of nitrous acid, *Nature*, 440, 195–198, doi:10.1038/nature04603, 2006. 7558
- Stemmler, K., Ndour, M., Elshorbany, Y., Kleffmann, J., D'Anna, B., George, C., Bohn, B., and Ammann, M.: Light induced conversion of nitrogen dioxide into nitrous acid on submicron humic acid aerosol, *Atmos. Chem. Phys.*, 7, 4237–4248, doi:10.5194/acp-7-4237-2007, 2007. 7558
- 10 Vione, D., Maurino, V., Minero, C., Pelizzetti, E., Harrison, M. A. J., Olariu, R.-I., and Arsene, C.: Photochemical reactions in the tropospheric aqueous phase and on particulate matter, *Chem. Soc. Rev.*, 35, 441–453, doi:10.1039/B510796M, 2006. 7557
- 15 Wang, B. and Knopf, D. A.: Heterogeneous ice nucleation on particles composed of humic-like substances impacted by O₃, *J. Geophys. Res.*, 116, D03205, doi:10.1029/2010JD014964, 2011. 7556
- Yokelson, R. J., Crounse, J. D., DeCarlo, P. F., Karl, T., Urbanski, S., Atlas, E., Campos, T., Shinozuka, Y., Kapustin, V., Clarke, A. D., Weinheimer, A., Knapp, D. J., Montzka, D. D., Holloway, J., Weibring, P., Flocke, F., Zheng, W., Toohey, D., Wennberg, P. O., Wiedinmyer, C., Mauldin, L., Fried, A., Richter, D., Walega, J., Jimenez, J. L., Adachi, K., Buseck, P. R., Hall, S. R., and Shetter, R.: Emissions from biomass burning in the Yucatan, *Atmos. Chem. Phys.*, 9, 5785–5812, doi:10.5194/acp-9-5785-2009, 2009. 7557, 7572
- 20 Zahardis, J. and Petrucci, G. A.: The oleic acid-ozone heterogeneous reaction system: products, kinetics, secondary chemistry, and atmospheric implications of a model system – a review, *Atmos. Chem. Phys.*, 7, 1237–1274, doi:10.5194/acp-7-1237-2007, 2007. 7565, 7587
- 25 Zelenay, V., Monge, M. E., D'Anna, B., George, C., Styler, S. A., Huthwelker, T., and Ammann, M.: Increased steady state uptake of ozone on soot due to UV/Vis radiation, *J. Geophys. Res.*, 116, D11301, doi:10.1029/2010JD015500, 2011. 7558
- 30 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimonono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demer-

jian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007GL029979, 2007. 7555

- 5 Zhao, Z., Husainy, S., Stoudemayer, C. T., and Smith, G. D.: Reactive uptake of NO_3 radicals by unsaturated fatty acid particles, *Phys. Chem. Chem. Phys.*, 13, 17809–17817, doi:10.1039/C1CP21790A, 2011. 7565, 7587

- Ziemke, J. R., Chandra, S., Duncan, B. N., Schoeberl, M. R., Torres, O., Damon, M. R., and Bhartia, P. K.: Recent biomass burning in the tropics and related changes in tropospheric ozone, *Geophys. Res. Lett.*, 36, L15819, doi:10.1029/2009gl039303, 2009. 7556, 7574
- 10

ACPD

13, 7553–7598, 2013

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Comparison of reactive uptake coefficients, γ , determined in this study with literature values for various liquid and solid organic substrates and gas phase oxidants. Uncertainty is $\pm 1\sigma$.

Substrate	Oxidant	T/K	γ^a	γ
Oleic Acid ^h	O ₃	293	$(6.5 \pm 1.2) \times 10^{-4}$	$(7.5-73) \times 10^{-4b}$
Oleic Acid ^h	O ₃	273	$(5.8 \pm 2.5) \times 10^{-5}$	$(5.2 \pm 0.1) \times 10^{-5c}$
Canola Oil ^h	O ₃	298	$(6.2 \pm 0.6) \times 10^{-4}$	$(6.5-9.3) \times 10^{-4d}$
Linoleic Acid ^h	NO ₃	298	(0.28 ± 0.25)	$(0.53 \pm 0.12)^e$
Pahokee Peat (dark) ⁱ	O ₃	293	$(3.5 \pm 2.1) \times 10^{-6}$	$(3.6 \pm 1.9) \times 10^{-6f}$
Pahokee Peat (UV) ⁱ	O ₃	296	$(1.7 \pm 0.5) \times 10^{-5}$	$(5.5 \pm 1.4) \times 10^{-5f}$
Levoglucosan (dark) ^j	O ₃	293	$(1.7 \pm 1.5) \times 10^{-5}$	$(1.3 \pm 1.2) \times 10^{-5g}$
5-Nitroguaiacol (dark) ^j	O ₃	293	$(1.0 \pm 0.4) \times 10^{-5}$	$(8.3 \pm 6.7) \times 10^{-5g}$

^a This study.

^b Zahardis and Petrucci (2007).

^c Moise and Rudich (2002).

^d de Gouw and Lovejoy (1998).

^e Zhao et al. (2011).

^f D'Anna et al. (2009).

^g Knopf et al. (2011).

^h γ values were determined using the stepwise method.

ⁱ γ values were determined from the steady state O₃ signal.

^j γ values were determined using initial drop in O₃ signal (see text for more details).

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf

Table 2. Steady state reactive uptake coefficients, γ , of organic substrates representative of BBA compounds with O_3 under dry conditions and with atmospherically relevant O_2 levels. γ was determined in the dark and under VIS and UV irradiation. O_3 concentrations ranged between 3.5×10^{11} and 6.0×10^{11} molecule cm^{-3} and substrate temperatures ranged between 20–23 °C. Uncertainty is $\pm 1\sigma$.

Substrate	$\gamma_{\text{dark}} \times 10^{-6}$	$\gamma_{\text{VIS}} \times 10^{-6}$	$\gamma_{\text{UV}} \times 10^{-5}$
Pahokee Peat	(3.5 ± 2.1)	(8.6 ± 0.8)	(1.7 ± 0.5)
Levoglucosan	(3.6 ± 2.7)	(4.3 ± 2.1)	(0.6 ± 0.4)
Nitroguaiacol	(5.7 ± 2.5)	(6.7 ± 4.7)	(3.6 ± 1.6)
1:1 Pahokee Peat/Levoglucosan	(2.0 ± 0.9)	(6.8 ± 4.2)	(1.1 ± 0.2)
1:1 Pahokee Peat/Nitroguaiacol	(2.7 ± 1.2)	(23.8 ± 13.0)	(2.8 ± 0.8)
1:10 Pahokee Peat/Nitroguaiacol	(4.3 ± 1.8)	(17.2 ± 11.5)	(2.3 ± 0.6)
1:100 Pahokee Peat/Nitroguaiacol	(2.5 ± 0.8)	(3.8 ± 0.9)	(3.8 ± 1.2)
10:1:3 Levoglucosan/ Pahokee Peat/Nitroguaiacol	(8.5 ± 4.5)	(27.7 ± 8.7)	(1.4 ± 0.5)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 1. Schematic representation (not to scale) of the irradiated rectangular channel flow reactor (I-RCFR) coupled to the chemical ionisation mass spectrometer. Irradiation source distance above the I-RCFR is adjustable.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Photosensitised
kinetics of biomass
burning aerosol**

S. M. Forrester and
D. A. Knopf


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 2. Spectral irradiance of the VIS and UV irradiation sources used in this study are presented as red and blue lines, respectively. The black line indicates the spectral irradiance of the sun.

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 3. Total irradiances for the VIS and UV irradiation sources as a function of lamp distance from the organic substrate are shown as closed and open circles, respectively. The open circles refer to the right y-axis. Error bars denote 1σ of the mean irradiance and the mean distance.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 4. Natural logarithm of the normalised ozone signal as a function of reaction time. Squares, circles, and triangles denote exemplary reactive uptake data of O₃ by a blank glass block, the 1 : 1 by mass PP/NG substrate, and canola oil, respectively. Dotted lines represent linear fits of the data from which the observed first-order rate constants are determined.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Photosensitised
kinetics of biomass
burning aerosol**

S. M. Forrester and
D. A. Knopf




Fig. 5. Absorbances as a function of wavelength. Pahokee peat is shown in black and 5-nitroguaiacol in a 0.01 M HCl solution (pH = 2) and a 0.01 M NaOH solution (pH = 12) are shown in green and purple, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 6. Steady state reactive uptake coefficients, γ , as a function of mass ratio of Pahokee peat to 5-nitroguaiacol. Black, red, and blue open circles denote dark, VIS, and UV irradiated uptake experiments, respectively. Errors bars denote 1σ of the mean γ .

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 7. Steady state reactive uptake coefficients, γ , of the 1 : 1 by mass PP/NG substrate with O_3 as a function of photon flux shown as closed and open circles for exposure to VIS and UV irradiation, respectively. The open circles refer to the top x-axis. Errors bars denote 1σ of the mean γ . Solid and dashed lines denote linear fits of the VIS and UV irradiated uptake data, respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 8. Steady state reactive uptake coefficients, γ , of 1 : 1 by mass PP/NG substrate with O_3 as a function of O_3 concentration under VIS (a) and UV (b) irradiation. Errors bars denote 1σ of the mean γ . The solid lines represent a Langmuir-Hinshelwood reactive uptake mechanism fit to the data and dashed lines indicate corresponding 95 % confidence bands.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 9. Oxidative power, $\gamma \cdot [\text{O}_3]$, as a function of $[\text{O}_3]$ **(a)** and photon flux **(b)** for the 1 : 1 by mass PP/NG substrate. Solid and dashed lines represent oxidative power under VIS and UV irradiation, respectively. Dashed line in **(b)** refers to top x-axis.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Photosensitised kinetics of biomass burning aerosol

S. M. Forrester and
D. A. Knopf




Fig. 10. Atmospheric lifetime, τ , of a $0.4 \mu\text{m}$ 5-nitroguaiacol particle as a function of concentration **(a)** and photon flux **(b)** determined using γ of the 1 : 1 by mass PP/NG substrate. Solid and dashed lines represent derived lifetime under VIS and UV irradiation, respectively. Dashed line in **(b)** refers to top x-axis.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)