

Interactive comment on “Monitoring high-ozone events in the US Intermountain West using TEMPO geostationary satellite observations” by P. Zoogman et al.

P. Sellitto

psellitto@lmd.ens.fr

Received and published: 9 January 2014

In the Introduction, the Authors state that "This UV/Vis multispectral combination for ozone observation has not been used from space before." This is not true. There exists a published work on a multispectral inversion scheme to retrieve ozone information from SCIAMACHY (real) observations, by using UV+VIS observations:

P. Sellitto, F. Del Frate, D. Solimini, S. Casadio, Tropospheric Ozone Column Retrieval From ESA-Envisat SCIAMACHY Nadir UV/VIS Radiance Measurements by Means of a Neural Network Algorithm, IEEE Transactions on Geosciences and Remote Sensing,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

Volume 50, Issue 3, Pages 998-1011, 2012.

The improvements brought, in particular in the troposphere, by this multispectral approach, and specifically by using the VIS radiation in the Chappuis bands, are discussed in the following publication:

P. Sellitto, A. Di Noia, F. Del Frate, A. Burini, S. Casadio, D. Solimini, On the role of visible radiation in ozone profile retrieval from nadir UV/VIS satellite measurements: An experiment with neural network algorithms inverting SCIAMACHY data, *Journal of Quantitative Spectroscopy and Radiative Transfer*, Volume 113, Issue 12, Pages 1429-1436, 2012.

I imagine that these papers should be considered in this context. I suggest the Authors to take a look at those publications, add them as references to their paper and correct the wrong statement above.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 13, 33463, 2013.

ACPD

13, C10766–C10767,
2014

[Interactive
Comment](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)

C10767

