

Interactive comment on “Nighttime observation and chemistry of HO_x in the Pearl River Delta and Beijing in summer 2006” by K. D. Lu et al.

Anonymous Referee #1

Received and published: 24 January 2014

This manuscript describes nighttime observations of OH and HO₂^{*} radicals during PRIDE-PRD 2006 and CAREBEIJING 2006 field campaigns and their comparisons to theoretical model simulation results. While HO₂^{*} was in fair agreement, observed OH levels were significantly larger than simulation. After detailed discussion on the possibility of artifact, the authors suggested possibility that the combination of additional RO_x production and the recycle of OH from RO_x through HO₂ could bring the model simulation in fairly good agreement in OH. For processes, the authors implied that the additional RO_x production would be explained by ozonolysis of unmeasured reactive terpenes at ample amount, or more possibly by the downward mixing of PANs as reservoir of radicals. The topic is well suitable to the scope of the journal. Although large uncertainty needs to be taken into account, the analysis is made on the best

C11426

knowledge basis, with clear logical flow. Two major comments are that (1) the modeled (and observed, if available) PAN concentrations should be mentioned together with the estimated flux, to convince readers, and that (2) discussion on the resemblance of the night-to-night variations in the observed and modeled radical concentrations should be involved. Overall, I recommend publication after revisions with respect to the points raised above and the following minor comments.

1. Page 31316, line 8. It is better to mention that two separate cells are used for detection of each OH and HO₂^{*}. A single pump draws air from the two cells. Can the back diffusion of NO, added only to the HO₂^{*} cell, to OH cell be a problem, to measure OH and HO₂^{*} simultaneously in such a system, especially when the HO₂^{*}/OH ratio is large (>1000)?
2. Page 31318, line 11. How did the authors estimate concentrations of C₂ species at BG site, as listed in Table 1?
3. Page 31319, Section 2.2. It is unclear if the model simulation was made for individual nights and then averaged to obtain the shown average time series (Figure 3 etc) or if only one model night with average concentrations of ancillary species was simulated. In the latter case, the titration relationship between O₃ and NO will not be adequately represented in the model.
4. Page 31319, lines 16-17. List date of the seven and nine days specifically.
5. Page 31320, lines 16-17. Did the big differences in the CO and isoprene concentrations from night to night have correlation with radical concentrations, and indicate potentially important parent RO₂ that have been recycled to OH?
6. Page 31321, line 16. What was the estimated average concentration level of HCHO? Was the 24-h lifetime regarding deposition important?
7. Page 31322, line 21. It is unclear what are meant by "large oxidation rates of the sum of reactive trace gases."

C11427

8. Page 31323, line 3. Was the contribution of the O₃+NO reaction included in the nighttime O₃ turnover rate?
9. Section 4.1. Only average levels of the observed and modeled radical concentrations are compared. It is recommended to include correlation analysis between them, representing features of night-to-night variations.
10. Page 31326, line 9. Can the artefact from Criegee intermediate for example be common for LIF and CIMS instruments and thus be overlooked during the intercomparison studies?
11. Page 31328, line 18. comparison (not intercomparison)
12. Page 31330, lines 5-8. Do we expect large differences in the simulated NO₃ concentration levels in the two model runs, as suspected from the large difference in OLNN and OLND? If yes, why was it?
13. Page 31332, line 25. 10**6 (not 10**-6)
14. Section 4.4.2. Can the authors evaluate simulated PAN concentrations, through comparison to observations if available, or to the levels typically present in the nighttime boundary layer? Can the PAN entrainment from the top boundary of the upper layer be also important, to increase the flux to the lower layer?
15. Section 4.4.2. Does K_z depend on z? The value for the 50-m altitude level was used?
16. Page 31334, line 8. How was the nocturnal temperature lapse rate used?
17. Page 31334. Can the dry deposition of the radicals on the ground be effective?
18. In Lu et al. (2012) and (2013) for daytime analysis for PRIDE-PRD and CARE-BEIJING, the authors discussed that the importance of the RO₂-to-HO₂ conversion was reduced, after considering RO₂ artefact in the HO₂* measurement. On the other hand, the manuscript indicated that the RO₂-to-HO₂ and HO₂-to-OH conversion was

C11428

required in the nighttime similarly to the daytime. Are they consistent to each other?

Interactive comment on Atmos. Chem. Phys. Discuss., 13, 31311, 2013.

C11429