Atmos. Chem. Phys. Discuss., 13, C1316–C1317, 2013 www.atmos-chem-phys-discuss.net/13/C1316/2013/ © Author(s) 2013. This work is distributed under the Creative Commons Attribute 3.0 License.



ACPD 13, C1316–C1317, 2013


> Interactive Comment

Interactive comment on "Newly observed peroxides and the water effect on the formation and removal of hydroxyalkyl hydroperoxides in the ozonolysis of isoprene" by D. Huang et al.

Anonymous Referee #1

Received and published: 16 April 2013

The goal of this manuscript is to investigate the role of relative humidity (RH) on the mechanism of ozonolysis of isoprene, specifically on the formation of various peroxidic products. Ozonolysis accounts for a relatively small fraction of atmospheric oxidation of isoprene but it may become comparable in importance to oxidation by OH and NO3 under certain conditions. Because of the known reactivity of Criegee intermediate, a reactive intermediate of the ozonolysis, towards water vapor it is reasonable to expect that the product distribution should be sensitive to RH. This is indeed what the authors observe in their measurements. Furthermore, they observe several peroxide products that were not detected in previous studies of isoprene ozonolysis. The manuscript thus contains new data, and it definitively falls with the scope of this journal.





I had a number of comments in my "quick review" of this manuscript, which the authors addressed prior to the ACPD publication. I found no significant additional issues with this manuscript, other than minor editorial corrections listed below. I was especially impressed by the analytical capabilities of the method that successfully detected (and quantified) rather complex peroxides. The list of peroxides that the authors synthesized as standards for their measurements is also quite impressive. It would of course be great the unknown peroxides observed by the authors were identified also but I appreciate the difficulty of a positive assignment of all products in such a system.

Technical corrections:

Abstract: the sentence "Recent laboratory investigations of this reaction identified hydrogen peroxide (H2O2) and hydroxymethyl hydroperoxide (HMHP)." should end with "in ozonolysis of isoprene" to avoid ambiguities.

Abstract: the sentence "These standards are needed to accurately specify HAHPs, although their synthesis is a challenge." is redundant and can be deleted.

Page 5281, line 8: I would replace "cause the death of" with "harm"

Page 5281, line 13: what does 25-60% mean – fraction of molecules that have at least one peroxide group in them?

Page 5282, line 14: "paths" should b replaced with "pathways"

Page 5287, line 18: replace "linearly" with "steadily"

Page 5292, line 29: replace "regularity" with "pattern"

Interactive comment on Atmos. Chem. Phys. Discuss., 13, 5279, 2013.

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

