

Sources and geographical origins of fine aerosols in Paris (France)

M. Bressi^{1,2}, J. Sciare¹, V. Ghersi³, N. Mihalopoulos⁴, J.-E. Petit^{1,5}, J. B. Nicolas^{1,2}, S. Moukhtar³, A. Rosso³, A. Féron¹, N. Bonnaire¹, E. Poulakis⁴ and C. Theodosi⁴.

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE, UMR8212, CNRS-CEA-UVSQ, Gif-sur-Yvette, 91191, France.

²French Environment and Energy Management Agency, ADEME, 20 avenue du Grésillé, BP90406 49004, Angers Cedex 01, France.

³AIRPARIF, Surveillance de la Qualité de l'Air en Ile-de-France, Paris, 75004, France.

⁴Environmental Chemical Processes Laboratory, ECPL, Heraklion, Voutes, Greece.

⁵INERIS, DRC/CARA/CIME, Parc Technologique Alata, BP2, Verneuil-en-Halatte, 60550, France.

Correspondence to: michael.bressi@ensiacet.fr

Supplementary material for Referee 1




Fig. A.1.3.a. Comparison between the temporal variation of the PMF Road traffic factor and NOx (top) and BCff (bottom).