

Interactive
Comment

Interactive comment on “Halogen species record Antarctic sea ice extent over glacial-interglacial periods” by A. Spolaor et al.

A. Spolaor et al.

andrea.spolaor@unive.it

Received and published: 10 June 2013

REFEREE 1: ANSWER

%Brdep is anti-correlated to the temperature proxy, during the warmer periods the boundary layer dynamics and mesoscale transport may have been different. This in turn may have altered wind speeds and thus transport time scales from the Antarctic coast to TD. Have the authors considered such possibility and how would that affect the interpretation of the measurements?

We thank the referee for recognizing this critical assumption in our work. The transport patterns as well as the air mass travel time from the coast to Talos Dome have been thoroughly investigated (Scarchilli et al., 2011; Sala et al., 2008). The main at-

C3346

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



mospheric circulation seems to be quite similar between glacials and interglacials as well the dominant wind dynamics. Many papers (Petit and Delmonte, 2009; Fischer et al., 2007) suggest that no appreciable difference in wind speed and direction and in consequence transport pattern was likely similar between glacial and interglacial time periods. Dramatic changes in the wind direction and speed could have modified the travel time and our consequent data interpretation. The present literature data suggest a similar wind pattern between glacial and interglacial periods.

REFERENCES

Scarchilli, C., Frezzotti, M., and Ruti, P.: Snow precipitation at four ice core sites in East Antarctica: provenance, seasonality and blocking factors, *Clim. Dyn.*, 37, 2107-2125, 10.1007/s00382-010-0946-4, 2011.

Sala, M., Delmonte, B., Frezzotti, M., Proposito, M., Scarchilli, C., Maggi, V., Artioli, G., Dapiaggi, M., Marino, F., Ricci, P. C., and De Giudici, G.: Evidence of calcium carbonates in coastal (Talos Dome and Ross Sea area) East Antarctica snow and firn: Environmental and climatic implications, *Earth and Planetary Science Letters*, 271, 43-52, <http://dx.doi.org/10.1016/j.epsl.2008.03.045>, 2008.

Fischer, H., Siggaard-Andersen, M.-L., Ruth, U., Rothlisberger, R., and Wolff, E.: Glacial/interglacial changes in mineral dust and sea-salt records in polar ice cores: Sources, transport, and deposition, *Rev. Geophys.*, 45, RG1002, 10.1029/2005rg000192, 2007.

Petit, J. R., and Delmonte, B.: A model for large glacial–interglacial climate-induced changes in dust and sea salt concentrations in deep ice cores (central Antarctica): palaeoclimatic implications and prospects for refining ice core chronologies, *Tellus B*, 61, 768-790, 2009.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 13, 3881, 2013.

ACPD

13, C3346–C3347, 2013

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

