

Interactive comment on “Present and future nitrogen deposition to national parks in the United States: critical load exceedances” by R. A. Ellis et al.

Anonymous Referee #2

Received and published: 10 June 2013

This is an interesting and well written manuscript dealing with an important U.S. natural resource planning topic. It is appropriate for publication in ACP. I have only a few comments regarding the present period analysis. My most critical observation is a lack of caveats for GEOS-Chem and the future emission scenarios (see detailed comments below).

1. Pg 9154, lines 9-17: Emission from agricultural soils (driven in most cases by N-fertilizer) is also a major anthropogenic source of NO. Also please clarify that NO_x from “fuel combustion” includes mobile as well as non-mobile sources. This becomes important in future scenarios with projected increases in urban areas and population. The

C3348

largest fraction of NH₃ (~80% of agricultural emissions with ag emissions contributing 80%+ of total NH₃ emissions) is emitted from animal husbandry activities so I would make this more prominent (not just another source). Soil-emitted NH₃ may be quickly removed but, like NO_x, a portion is transformed into particles which can then undergo significant downstream transport. This is mentioned briefly later in the paper but should be mentioned here as well.

2. Pg 9156, lines 7-8: As noted by reviewer 1, additional information is needed to explain the source of the present day emission differences for Zhang et al.

3. Figure 2: Agreement between simulated and observed results is quite remarkable, particularly in the western U.S. where you are comparing relatively coarse model results to point measurements in complex terrain. Your results appear to be better than Zhang et al.. Some improvement is expected given your longer averaging period (annual vs monthly), but please discuss. Also, I assume the model background and observations shown are for 2006 only. If so, please correct the caption. If not, define what you are plotting for background values (e.g., average?). What observations are shown in the scatter plots, 2006, 2006-2008, avg 2006-2008? Which Parks lack NADP monitors? Would you care to speculate what the scatterplots would look like if these monitors were available (e.g., same, more scatter, less scatter, etc.).

4. Table 2: Please include “GEOS-Chem simulated deposition values for 2006” in the table caption. This is important information and should not be in a footnote.

5. Pg 9158, lines 25-26: “There is little 2006-2008 interannual variability, either in the model or in observations (Zhang et al., 2012).” On page 4547 of Zhang et al., they state that there is little interannual variability in the model results for 2006-2008, but I didn’t see any mention of interannual variability in the observations. One might expect interannual variability to be relatively low for regions dominated by dry deposition, but it can be significant in wet-dominated areas. Also, it is expected that a numerical simulation will underestimate observed variability. Unless I missed the statement regarding

C3349

observations in Zhang et al., I think your statement in this manuscript is a bit too broad without additional support.

6. You mentioned in passing that there are few or no dry deposition observations with which to evaluate your model. While correct, this is a major source of uncertainty. Since the western Parks are dominated by dry deposition, what are the implications of this uncertainty for your results? This could also be combined with reviewer 1's observation regarding the underestimation of current deposition totals in the west.

7. Insufficient caveats are provided for GEOS-Chem, but in particular caveats are missing for the the RCPs. Caveats are needed to allow the reader to place the projected results in a meaningful application context. Below are a few caveat-related questions that come to mind:

a) The future emission scenarios include land use change in response to socio-economic and energy policy changes. Did you change the land use driving the future GEOS-Chem simulations to match the RCP changes? This may not be a serious issue in the eastern U.S., where total N deposition it is usually dominated by wet removal, but underlying vegetation is a critical driver for dry deposition processes, which dominate the West. If you did not make this adjustment, what are the implications for your results?

b) Figure 1 and text discussion. After reading the RCP documentation in van Vuuren et al. (2011a & b) and Riahi et al., 2011, the series of NO_x emission maps seem reasonable. I am more curious about the NH₃ maps – assuming that the majority of NH₃ is associated with agricultural production. . . . -The RCP scenarios project land use change using transition rates determined from historical data (1500-2000) (Hurtt, et al., 2011). These rates are then projected forward in time to 2100. Is the use of these historical transition rates, particularly in the western U.S. and the southern and western edges of the Midwest that have only recently been opened to extensive irrigation (post 1950), reasonable for projections to 2050? -Land use transition rates are a function of

C3350

agricultural productivity, proximity to existing agricultural areas, proximity to current water bodies and cities and a random factor. Your graphs of both RCP 8.5 and 2.6 suggest agricultural expansion/ intensification in Arizona, New Mexico, Wyoming and Montana in 2050. Is this reasonable considering water supplies are currently limited and may become more limited under future climate conditions? -What is the source of the ammonia increase in RCP 8.5? Riahi states "aggregate arable land use in developed countries slightly decreases," and that bioenergy expands in the developing world and is focused primarily on forests biomass. He goes on to say that agricultural residues are used for bioenergy where cost effective, i.e., no dedicated bioenergy crops to drive agricultural production. Further, RCP 8.5 energy system moves toward coal-intensive technology and unconventional natural gas and oil extraction and urban population expansion in the west. The summaries in Hurtt and van Vuuren suggest that while biofuel expansion would explain NH₃ increases in RPC 2.6, there is no such driver in 8.5. It is a little difficult to tease out U.S. trends from these broad global discussions (Riahi, et al. does not explicitly address NH₃ at all). Perhaps a Table that summarizes regional changes in agricultural lands for the US would be helpful. Regardless, you need to help me to understand the source of the increased NH₃ emissions in RCP 8.5.

c) Future NO_x emissions: van Vuuren - Overview (pg 21) states that all the RCP scenarios assume NO_x emissions will continue to decline in the future in response to "rising income levels." Particularly in RCP 8.5 – this is driven by policy intervention. Is the assumption of no technical limitation on NO_x reductions reasonable? This assumption is particularly important for the western U.S., which is projected to have rather large increases in urban land use (Hurtt et al., 2011). What are the implications if these reductions are not realized?

d) This paper addresses the response of N deposition to future emissions. What do you think would happen to your results if you included future weather as well, i.e., warmer temperatures and changing precipitation patterns?

e) Finally, bidirectional ammonia flux approaches are becoming widely accepted

C3351

(Fletcher, et al., 2013, Massad et al., 2010; Bash et al., 2013; Dennis et al., 2013). Adoption of this approach would likely change both the ratio of wet to dry removal and N deposition in the western U.S. under current conditions. What are the potential implications for your study?

f) Given these considerations, how do you see this work evolving in the future?

Refereces Hurtt, G.C., Chini, L.P., Froking, S. et al., 2011. "Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands." *Climate Change*, 109: 117-161.

van Vuuren, D.P., Stehfest, E., den Elzen, M.G.J., et al., 2011. "RCP2.6: exploring the possibility to keep global mean temperature increase below 2o C." *Climatic Change*, 109: 95-116.

van Vuuren, D.P., Edmonds, J., Kainuma, M., et al., 2011. "The representative concentration pathways: an overview." *Climatic Change*, 109: 5-31.

Riahi, K., Rao, S., Krey V. et al., 2011. "RCP 8.5 – A scenario of comparatively high greenhouse gas emissions." *Climatic Change*, 109: 33-57.

Dennis, R.L., Schwede, D.B., Bash, J.O., et al., 2013. "Sensitivity of continental United States atmospheric budgets of oxidized and reduced nitrogen to dry deposition parameterizations." *Phil. Trans. R. Society B*, 368: 20130124

Bash, J.O., Cooter, E.J., Dennis, R.L., et al., 2013. "Evaluation of a regional air quality model with bidirectional NH₃ exchange coupled to an agro-ecosystem model. *Biogeosciences*, 10: 1635-1645. <http://www.biogeosciences.net/10/1635/2013>. doi:10.5194/bg-10-1635-2013/.

Flechard, C.R., Massade, R.S., Loubet, B., et al., 2013. "Advances in understanding, models and parameterisations of biosphere-atmosphere ammonia exchange" <http://www.biogeosciences-discuss.net/10/5385/2013> *Biogeosciences Discuss.*, 10, C3352

5385-5497

Massad, R.-S., E. Nemitz and M.A. Sutton, 2010. "Review and parameterization of bi-directional ammonia exchange between vegetation and the atmosphere." *Atmos. Chem. Phys.*, 10, 10359-10386, doi:10.5194/acp-10-10359-2010, 2010.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 13, 9151, 2013.