



[Interactive
Comment](#)

***Interactive comment on* “Simulation of the dispersion of the Eyjafjallajökull plume over Europe with COSMO-ART in the operational mode” by H. Vogel et al.**

Anonymous Referee #1

Received and published: 18 July 2013

Review of the article "Simulation of the dispersion of the Eyjafjallajökull plume over Europe with COSMO-ART in the operational mode" by H. Vogel et al.

The article describes model simulations of the ash dispersion from the eruption of the volcano Eyjafjallajökull on Iceland in April 2010 using the COSMO-Art model. The authors describe that their model is able to reproduce the ash distribution with good quality by comparing the simulation results with some selected observations. They show that best agreement can be reached if the source function for the emitted ash is known with good accuracy. They propose to perform a number of time-lagged model runs in order to give probability distributions for the ash concentration at a certain time

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



and height.

Overall there is not much that the reader can learn from this paper. A number of model studies already exist (as cited in the paper) that describe the difficulties in reconstructing or even forecasting the ash dispersion from a volcanic eruption. The main finding is always that the source strength of the volcanic ash is the main difficulty for an accurate forecast of the ash concentrations. The authors of this paper confirm this finding in their comparison to observations of the particle number density at Hohenpeissenberg, where the best agreement was found with a reconstructed source function (i.e. one that was already based on observations). They also confirm the finding that early observations close to the source are necessary to feed the model systems with information about the emitted mass and the size distribution of the volcanic ash. The authors finally propose that a time lagged ensemble forecast could give probability distribution functions as forecast products instead of one map derived from a deterministic forecast. This is certainly a good idea, however their ensemble includes only different starting times of the forecast. This means that only effects of the meteorological fields are considered in the probability distribution while the main uncertainty is the emission strength.

I recommend major revisions before the paper can be published. The authors should include other sources of uncertainty into their ensemble forecast and they should extend their study beyond the first eleven days after the eruption. Many aircraft and lidar observations have been done in May 2010, when the volcano was still active. These observations could serve as a test bed for the quality of the ensemble forecast that is proposed here. The authors should additionally consider a number of specific questions and comments listed in the following.

Specific comments

Page 13440, I5-7: If the model is calibrated and the emissions are modified accordingly it should show good agreement with the observations. The term “nearly perfect” is

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



inappropriate, in particular because no numbers are given for this comparison.

Page 13440, I22: Besides the temporal and the spatial distribution, information about the mass concentrations is needed most urgently.

Page 13441, I3-4: As far as I followed the publications about the Eyjafjallajökull eruption, the VAAC forecasts were not quantitative either when the eruption started. Other model systems could only give quantitative information after being compared to observations. This kind of “calibration” was the same as it is done in this paper.

Page 13442, I21-24: How are you going to determine the reliability of the forecast if you take only uncertainties in the meteorology into account and not those in the source strength?

Page 13443, I 14: What was the width of the size bins? Why is coagulation neglected? Please justify this.

Page 13444, I1: Why did you scale the source strength linearly with the source height? Why didn't you use the source function from Mastin (2009) right from the beginning?

Page 13444, I18: What do you mean by “own” analysis?

Page 13445, I7: I cannot derive from Fig.2 that the horizontal distribution of the ash plume was captured “quite well”. As far as I can follow the Meteosat picture, the plume was located in Northern Germany along the coast to the Baltic Sea. In this area, the model results reveal no ash.

Page 13446, I6: You show that the model produces thin ash layers when 80 vertical levels are used. What is the spacing of the levels in the altitude of the ash layer? How thick is the layer in the observations?

Page 13446, I11: “The simulated ash layer above the boundary layer starts to decrease”: this is unclear. What is decreasing? The concentration, the vertical thickness, the altitude?

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Page 13446, I19-23: What can I learn from this study of the modified emission profile? Couldn't you also find a modified emission profile that shows better agreement with the observations?

Page 13446, I27: What does the fact that the parameterization of deep convection does not change the results tell us? Is it not important at all or just not important in this case (and for this specific location). Shouldn't you better analyze the model results at other locations, too?

Page 13449, I22 - page 13450, I14: In this section I am missing comparisons of the simulated mass densities with observations, e.g. from the Falcon flights. Why do you restrict yourself to the number densities in the previous section while the mass is more relevant for the flight restrictions?

Page 13449, I28: It is certainly a good idea to give a probability forecast but you should include all uncertainties in your ensemble, in particular the source strength. Page 13450, I24: I assume there is a "high" missing between "resolution is" and "compared" in this sentence.

Page 13450, I15 - page 13451, I20: This is a summary of what has been done. I am missing real conclusions except the recommendations for the observations necessary to help the models to calculate the correct concentrations. These observations should not include only aircraft and lidars but also passive instruments like radiometers, either ground based or on board of satellites.

Figures:

Fig. 2: The Meteosat picture is difficult to interpret.

Fig. 3: The inlay in Fig. c) is hard to understand. What is shown here?

Fig. 5: What was changed in the emission profile (orange curve)? Does this correspond to Fig 3 c)?

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Fig. 6: How can you justify to use the size distribution from 2 May for your simulations in an earlier period?

Fig. 8: Add the date for this simulation result.

Interactive comment on Atmos. Chem. Phys. Discuss., 13, 13439, 2013.

ACPD

13, C4937–C4941, 2013

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



C4941