Atmos. Chem. Phys. Discuss., 13, C590–C591, 2013 www.atmos-chem-phys-discuss.net/13/C590/2013/ © Author(s) 2013. This work is distributed under the Creative Commons Attribute 3.0 License.



ACPD

13, C590–C591, 2013

Interactive Comment

Interactive comment on "Atmospheric nitric oxide and ozone at the WAIS Divide deep coring site: a discussion of local sources and transport in West Antarctica" by S. Masclin et al.


J. Savarino

jsavarino@ujf-grenoble.fr

Received and published: 15 March 2013

Another interesting piece of the NOx/O3 puzzle above icy terrains.

Now, considering that WAIS possesses the same snow accumulation and latitude coordinate than Summit in Greenland, it is very puzzling to see the sharp decrease of nitrate concentration in WAIS snow pits while abscent in Summit snow pits. I think this sharp difference deserves a specific discussion, considering that in the manuscript photolysis of nitrate is proposed to be the main NOx source above this site. Apparently, as previously thought, the accumulation rate is not the only parameter controlling the photodissociation of nitrate in snow (for a given solar flux).



Printer-friendly Version

Interactive Discussion

Discussion Paper



Joel Savarino

Interactive comment on Atmos. Chem. Phys. Discuss., 13, 6807, 2013.

ACPD

13, C590–C591, 2013

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

