Atmos. Chem. Phys. Discuss., 13, C7322–C7323, 2013 www.atmos-chem-phys-discuss.net/13/C7322/2013/

© Author(s) 2013. This work is distributed under the Creative Commons Attribute 3.0 License.



ACPD

13, C7322-C7323, 2013

Interactive Comment

Interactive comment on "Liquid-liquid phase separation in particles containing organics mixed with ammonium sulfate, ammonium bisulfate, ammonium nitrate or sodium chloride" by Y. You et al.

Anonymous Referee #1

Received and published: 25 September 2013

So far, liquid-liquid phase separation (LLPS) in mixed organic-inorganic aerosol particles has mostly been studied for systems with ammonium sulfate as the inorganic component. However, ambient aerosols also contain other salts. This paper investigates now mixtures with ammonium bisulfate, ammonium nitrate and sodium chloride as the inorganic component and thus covers the most common inorganic salt compositions found in tropospheric aerosols. It relates the occurrence of LLPS to O:C, which has already been established as a good predictor of LLPS for systems with ammonium sulfate as the inorganic component. This is a relevant study to better constrain the

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



particle compositions for which LLPS is present. The paper is well written and the results are clearly presented. There is one point that should be added to the discussion of LLPS of the investigated systems, namely, which of the investigated organic substances show LLPS when mixed with water even in the absence of an inorganic salt, i.e. for an OIR of infinity. This would help to better judge the salting-out effects of the investigated salts.

As a minor technical point: the authors should try to improve the contrast of the optical images shown in Fig. 1.

Interactive comment on Atmos. Chem. Phys. Discuss., 13, 20081, 2013.

ACPD

13, C7322-C7323, 2013

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

