

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Air quality in Delhi during the Commonwealth Games

P. Marrapu^{1,2}, Y. Cheng^{2,3}, G. Beig⁴, S. Sahu⁵, R. Srinivas⁴, and G. R. Carmichael^{1,2}

¹Department of Chemical and Biochemical Engineering, University of Iowa, Iowa City, 52242, USA

²Center for Global and Regional Environmental Research, University of Iowa, Iowa City, 52242, USA

³Multiphase Chemistry Department, Max Planck Institute Chemistry, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

⁴Indian Institute of Tropical Meteorology (Ministry of Earth Sciences, Govt. of India), Dr. Homi Bhabha Road, Pashan, 411008 Pune, India

⁵Forschungszentrum Julich GmbH, IEK-8:Troposphere, 52425 Julich, Germany

Received: 3 August 2013 – Accepted: 28 March 2014 – Published: 17 April 2014

Correspondence to: P. Marrapu (pallavi-marrapu@uiowa.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Air quality during The Commonwealth Games (CWG, held in Delhi in October 2010) is analyzed using a new air quality forecasting system established for the Games. The CWG stimulated enhanced efforts to monitor and model air quality in the region. The air quality of Delhi during the CWG had high levels of particles with mean values of $PM_{2.5}$ and PM_{10} at the venues of 111 and $238 \mu g m^{-3}$, respectively. Black carbon (BC) accounted for $\sim 10\%$ of the $PM_{2.5}$ mass. It is shown that BC, $PM_{2.5}$ and PM_{10} concentrations are well predicted, but with positive biases of $\sim 25\%$. The diurnal variations are also well captured, with both the observations and the modeled values showing nighttime maxima and daytime minima. A new emissions inventory, developed as part of this air quality forecasting initiative, is evaluated by comparing the observed and predicted species-species correlations (i.e., BC : CO; BC : $PM_{2.5}$; $PM_{2.5}$: PM_{10}). Assuming that the observations at these sites are representative and that *all* the model errors are associated with the emissions, then the modeled concentrations and slopes can be made consistent by scaling the emissions by: 0.6 for NO_x , 2 for CO, and 0.7 for BC, $PM_{2.5}$ and PM_{10} . The emission estimates for particles are remarkably good considering the uncertainty in the estimates due to the diverse spread of activities and technologies that take place in Delhi and the rapid rates of change.

The contribution of various emission sectors including transportation, power, domestic and industry to surface concentrations are also estimated. Transport, domestic and industrial sectors all make significant contributions to PM levels in Delhi, and the sectoral contributions vary spatially within the city. Ozone levels in Delhi are elevated, with hourly values sometimes exceeding 100 ppb. The continued growth of the transport sector is expected to make ozone pollution a more pressing air pollution problem in Delhi. The sector analysis provides useful inputs into the design of strategies to reduce air pollution levels in Delhi. The contribution for sources outside of Delhi on Delhi air quality range from $\sim 25\%$ for BC and PM to $\sim 60\%$ for day time ozone. The significant

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with more than 18 million inhabitants. Delhi hosted the Commonwealth Games (CWG), a multi sport event involving 73 countries, in October 2010. This high profile event provided an opportunity to accelerate efforts to improve air quality. To support the CWG air pollution monitoring was enhanced, a new emissions inventory was developed (Sahu et al., 2011) and new air quality forecasting efforts were initiated. In this paper we use the WRF-Chem model (Grell et al., 2005) to simulate the air quality during the GWG and evaluate its performance by comparing the predicted meteorology and concentrations of BC and other criteria pollutants with observations. We present spatial patterns of BC, PM_{2.5} and PM₁₀, CO, NO_x, and O₃ over Delhi and estimate the contributions from specific source sectors (Transportation, Power, Industrial, and Domestic). Such sector information is needed to help guide the development of effective pollution reduction measures. We also evaluate the emission estimates by analyzing observed and predicted species ratios.

2 Approach

2.1 WRF-Chem configuration and domain

The CMG air quality forecast system is based on the WRF-Chem model. Four nested domains were used in the analysis and they are shown in Fig. 1. The outer domain covered South Asia from 50 to 120° E and 0 to 55° N at a horizontal resolution of 45 km, with 131 × 131 grid cells. The next domain focused on the northern regions of India covering the Indo Gangetic plain at a resolution of 15 km with 127 × 127 grid cells. The two inner domains covered the Delhi region at resolutions of 5 km with 55 × 55 grid cells and 1.67 km with 75 × 75 grid cells. For all domains 27 vertical levels were used with a maximum height of ~ 20 km. The model configuration is summarized in Table 1. This configuration enables direct, indirect and semi direct aerosol radiative feedbacks to be included in the analysis. The Carbon Bond Mechanism version Z (CBMZ) with the MOSAIC aerosol module using 4 size bins was employed.

2.2 Emissions

A new detailed emissions inventory was prepared by IITM Pune, Maharashtra for the Delhi area at a resolution of 1.67 km^2 . The emission inventory covered the National Capital Region Delhi (NCR) ($\sim 70 \text{ km} \times 65 \text{ km}$) and the surrounding areas (in total covering $115.23 \text{ km} \times 138.6 \text{ km}$). The emission data was developed for NO_x , CO, BC, $\text{PM}_{2.5}$, PM_{10} , OC, VOC and SO_2 for four sectors (i.e. power, industrial, transport, residential). This NCR area was extensively studied through a field campaign made during March–May 2010, which collected relevant primary and secondary data with high resolution. This data included vehicular along with the type of fuel used, point source locations for manufacturing industries located within Delhi, and biofuel usage from slums, hotels and street vendors (Sahu et al., 2011). For the rest of the area beyond the NCR region emissions were based on proxies as new primary data were not generated in this region.

The emission totals by species and sector for Delhi are summarized in Table 2. The transport sector accounts for the more than 40 % of the emissions for all species with the exception of SO_2 and OC. Delhi is a densely populated city and has approximately 4 million on road vehicles and the number increases at a rate of ~ 10 percent per year (Mohan et al., 2006). The $\text{PM}_{2.5}$ and PM_{10} transport sector emissions include windblown road and construction dust sources, which comprise ~ 50 % of the total transportation sector emissions. For SO_2 power generation and industrial sources are the most important sectors. The domestic sector is large for CO, OC, BC and NMOC (non-methane organic compounds).

The spatial distribution of emissions was based on roadways, and locations of power plants, slums and major industries using various Delhi data sets (Sahu et al., 2011). The emissions were transferred into the WRF-Chem modeling system using the Emission Preprocessor Model (EPRES) designed by M. Lin (Center for Sustainability and the Global Environment at University of Wisconsin-Madison) and modified by Y. F. Cheng (CGRER at the University of Iowa). EPRES does two things, first it

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



speciates the total VOC emissions based on the chemical mechanism (i.e., CBM-Z), and second it horizontally interpolates the emission inventory onto the model grids. When interpolating the emissions, EPRES reads in the initial grid of the inventory from the WPS/WRF output and then utilizes the I/O API mass conservative interpolation subroutine to interpolate the emissions onto the WRF model grids. All the interpolated species were converted into WRF-Chem required units. After wards, the mapped emissions were distributed into vertical layers, i.e. 70 % was distributed at the surface and 30 % was uniformly distributed to the grids up to ~ 1000 m (grids 2–6 in this application). Diurnal profiles were also applied. The diurnal variability of BC emissions is shown in Fig. 7, and exhibits two peaks, one in early morning and the second in early evening associated with traffic and cooking activities.

Figure 2 shows the emission distributions of NO_x and $\text{PM}_{2.5}$ over Delhi. The emissions of all the primary species for the inner grid are shown in the Supplement Fig. S1. The spatial distribution reflects the various activities within and around the city. The NO_x and $\text{PM}_{2.5}$ emissions show intense emissions within the city and along roadways, with spatial differences reflecting different contributions by sector. Geographically Delhi is surrounded by industrially developed cities such as Gurgaon and Noida. Construction is active within and around the city and as a result many cement industries and brick kilns are situated in and around Delhi, adding to the increasing particulate emissions associated with road and construction dust. Extensive usage of wood and charcoal stoves for cooking is prevalent in the slums and in the out skirts of Delhi and high BC emissions (~ 30 % of total) occur in association with these activities (Sahu et al., 2008). The sectoral distributions of the BC emissions are shown in Fig. 3. These show clearly the locations of the power plants, industrial clusters, the distribution of slums and the major transportation networks. The venues for the CWG were clustered in the eastern half of Delhi city denoted by the completed outline in the center of the domain (see Fig. 4). These are regions most heavily impacted by domestic and industrial sources.

The INTEX-B emissions (Zhang et al., 2009) were used over the outer three domains. Biogenic emissions were calculated hourly on-line using MEGAN (Emmons

et al., 2010). The dust emissions were turned off during the run time of the model, since the post monsoon period is a period with insignificant wind-blown soil dust impacts on Delhi. No agricultural fire emissions were included, reflecting that in this post monsoon period no harvesting takes place.

2.3 Boundary conditions and simulation procedure

To consider the influence of sources outside of the largest domain, boundary conditions from the ECMWF MACC (Monitoring Atmospheric Composition and Climate) project that operates a data-assimilation and modeling system for a range of atmospheric constituents that are important for climate, air quality and surface solar radiation (<http://www.gmes-atmosphere.eu/>) were used. The meteorology was initialized using NOAA FNL data using the WPS sub program and simulations were done in five day periods each with two days of spin up time.

2.4 Observations

Ten new automatic air pollution and meteorology monitoring stations were installed for the Commonwealth Games and were placed at different venues across the city (i.e., Commonwealth Games Village (CWG), IITM Delhi (IITM), Yamuna Sports Complex (YSC), Indira Gandhi Sports Complex (ISC), M.Dhyan Chand National Stadium (MDS), Jawaharlal Nehru Sports Complex (JSC), Thyagaraj Sports Complex (TSC), University of Delhi (UD), IGI-Airport (IGI) and finally Talkotaroo Garden (TG)) as shown in Fig. 4. BC, CO, NO, NO₂, O₃, and PM_{2.5} and PM₁₀ were measured at hourly intervals over the span of two weeks from 26 September 2010 to 15 October 2010. Similar to the air quality species, meteorology parameters (i.e., wind speed (m s^{-1}), wind direction ($^{\circ}$), temperature ($^{\circ}\text{C}$), and relative humidity (%)) were also measured. Ozone was measured with a photometric UV analyzer (Thermo-49i) and NO and NO₂ with chemiluminescence analyzer (Thermo-41i). BC was measured with the

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Magee Scientific aethalometer (Model AE31) and, PM_{2.5} and PM₁₀ using BETA attenuation analyzers (Beta Met One BAM 120).

3 Results and discussions

The WRF-Chem model was run for the period 26 September–16 October 2010. The analysis in this paper focuses exclusively on the Delhi region (the inner-most grid).

3.1 Meteorology

The Commonwealth Games were held during the post monsoon season, a transition between the summer monsoon and winter. The general situation in India is that in mid September the monsoon shifts from the Southwest Monsoon to the Northeast monsoon. This brings in air masses from the Northern parts of the country (Ghude et al., 2009). During the CWG the surface winds were most frequently from the NNW and low ($< 3 \text{ m s}^{-1}$). Winds from the EES were less frequent but of higher speeds.

The predicted meteorology was compared with surface observations at two monitoring stations (results are typical of other sites). The statistics of the comparison are shown in Table 3. The model has a dry bias (under predicts RH), slightly under predicts peak day time temperatures by $1\text{--}2^\circ$ ($\sim 5\%$), and is biased high in wind speed. The model captures the low wind speeds, but the observations often are less than 1 m s^{-1} , which the model estimates at $1\text{--}3 \text{ m s}^{-1}$.

3.2 Air pollutants in Delhi

3.2.1 Aerosols

The predicted period mean surface concentrations of BC are shown in Fig. 5 (PM_{2.5}, PM₁₀ and CO show a very similar spatial distribution). Black carbon concentrations exceed $25 \mu\text{g m}^{-3}$ in the center of the domain where emissions from the residential and

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



industrial sectors are both high (see Fig. 3). $PM_{2.5}$ and PM_{10} levels are very high in this area with period mean values of 220–350 $\mu\text{g m}^{-3}$ and 350–550 $\mu\text{g m}^{-3}$, respectively, values which greatly exceed the National Ambient Air Quality standards of 60 and 100 $\mu\text{g m}^{-3}$, respectively.

The hourly surface concentrations from all the observation stations over the entire time period were combined and their distributions are presented in Fig. 6. The variability between the stations is shown by the bars that reflect the values at the stations that had the highest and lowest values. The mean value for BC from all the sites is 9.5 $\mu\text{g m}^{-3}$, with minimum and maximum site values of 4.4 and 13.1 $\mu\text{g m}^{-3}$. The overall mean values for $PM_{2.5}$ and PM_{10} (along with maximum and minimum site means) are: 111 $\mu\text{g m}^{-3}$ (77–144 $\mu\text{g m}^{-3}$) and 238 $\mu\text{g m}^{-3}$ (200–306 $\mu\text{g m}^{-3}$), respectively. The distributions of the calculated values are also shown in Fig. 6. In general the predicted values show a similar distribution, but with a positive bias ($\sim 30\%$).

Further insights into the pollutant distributions in Delhi are found by analyzing the diurnal variations. The 20 day average diurnal cycle of BC observed at Dhyanchand Stadium is shown in Fig. 7, along with the diurnal emissions used in the model and the predicted PBL height. The BC observations show a strong diurnal cycle, with minimum in the mid-afternoon when the PBL is the highest and maximum in late evening when the PBL height is at a minimum and emissions are at their highest. The evening and early morning features reflect the cooking and traffic patterns and the trapping of these surface emissions by the shallow night time mixing layer. These diurnal features are well captured by the model, which suggests that the diurnal pattern used in the emissions accurately reflect these activities. The daytime values are accurately captured, while the night time values are biased high. This suggests that the daytime PBL is reasonably well predicted, but the nighttime values may be too low. The diurnal profiles of $PM_{2.5}$ and PM_{10} exhibit similar diurnal variations as BC, with minima during mid-afternoon and peak values at night, but with less asymmetry between mid-night and 6 a.m. IST values, reflecting their stronger dependency on traffic patterns.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The predicted mean composition of $PM_{2.5}$ at Dhyanchand stadium is shown in Fig. 8. (There is only very slight variability in the percent contribution at the different monitoring stations.) BC accounts for about 8% of the fine mode mass and is greater than that for sulfate. OC accounts for $\sim 30\%$ and other primary PM (from unpaved roads, construction etc.) accounts for the largest fraction. Nitrate contribution exceeds that of sulfate.

3.2.2 Gaseous pollutants

The hourly surface observations of CO, NO, NO₂, NO_x and O₃ are also shown in Fig. 6. The mean observed CO is 1.7 ppm, with station minimum and maximum of 1.1 and 2.8 ppm. The mean NO_x value is 68 ppb, with minimum and maximum station means of 28 and 139, respectively, while the station minima and maxima NO and NO₂ are 14 and 93, and 14 and 42, respectively. The mean daytime (9.30 a.m. to 6.30 p.m. IST) ozone value is 54.5 ppb, with station minimum and maximum values of 45 and 61 ppb.

The predicted distributions of these gases are also plotted in Fig. 6. Comparison with the observations shows that CO is under predicted and NO_x is over predicted. Ozone predictions are biased high, and with a much different distribution. The smaller variability among the observation sites is also captured in the predicted values of ozone and can be explained by looking at the spatial distribution of predicted daytime average surface O₃ in Delhi (Fig. 5b). The monitoring sites are all located in the eastern half of Delhi city (see Fig. 4). The ozone surface concentrations show a broader distribution over this domain than PM with weaker gradients. Due to the rapid increase in traffic, ozone is becoming a more important problem with hourly observed and predicted values exceeding 100 ppb (the India NAAQS for 8 h average is 50 ppb ($100 \mu g m^{-3}$) (CPCB Report, 2012)).

A sensitivity simulation was performed where NO_x emissions were reduced to 1/3rd of the original values in order to see the effects on the ozone distributions. These results for ozone are also plotted in Fig. 6. Under the reduced NO_x emission case the ozone distribution is much closer to that observed, as are the distributions of NO_x and NO

and NO₂ (not shown). However the NO_x reduction is too large, as the ozone maximum values are overpredicted and the relative amounts of NO and NO₂ are now too low and high, respectively. The emissions are discussed in more detail in Sect. 3.4.

3.3 Sector contributions

We estimated the emission sector contributions to surface pollution levels in Delhi. Additional model runs were performed to isolate the contribution from specific sectors. Four different simulations were performed. An initial run was carried out to calculate the total pollution from all sources; i.e. anthropogenic, biogenic and biomass burning. Then individual runs were carried out to estimate the concentrations from each anthropogenic sector. The equations below explain the process for how the transportation sector results were determined.

$$\text{Anthropogenic (A)} = \text{Transportation (T)} + \text{Power (P)} + \text{Industry (I)} + \text{Domestic (D)} \quad (1)$$

$$\text{Initial run} = \text{I}(\text{A} + \text{Biogenic} + \text{Biomass Burning} + \text{Boundary}) \quad (2)$$

Individual run w/o Transport Sector

$$S = (\text{A} - \text{T} + \text{Biogenic} + \text{Biomass Burning} + \text{Boundary}) \quad (3)$$

$$\text{Sector contribution} = \text{I} - S, \quad (4)$$

which represents the concentrations from transport emissions only.

This process was repeated for the other three sectors and the fourth sector was estimated by difference between the four simulations (base plus three sector removed runs).

Figure 9 shows the spatial distribution of the % contribution to surface concentrations from the Transportation, Power, Industry and Residential sectors for BC. In general the largest contributions are from the transport, domestic and industry sectors. The area affected by transport is the largest, but the peak contributions are larger for the domestic and industry sectors. The contributions from industry are highest in the center

of the domain. The impact of the power sector on average surface concentrations of BC is small, but is large for SO₂, sulfate and NO_x concentrations (see Supplement Fig. S2). The sector contributions are similar for CO, PM_{2.5} and PM₁₀ as shown in the Supplement. The sector contributions for NO_x, SO₂ and ozone are also presented as SI.

Further insights into the sector contributions are found by looking at their diurnal variations. Figure 12 shows the 20 day average diurnal variation of the sectoral contributions to various criteria pollutants at the monitoring station at the Dhyanchand stadium. The contributions are shown in terms of percentages. In the case of BC residential and transportation sectors are most important at all times, but with maximum values in the late evening and early mornings. The industrial sector is also important at this site for BC. During the daytime as the mixing layer grows, air masses from higher altitudes, which are impacted by sources outside of Delhi, are entrained into the boundary layer, and the contribution from this process is shown by the black shading. These other contributions are from emissions that are outside of the inner most grid, and which are transported into this region through the boundaries. PM_{2.5} and PM₁₀ sector plots are similar to BC, with PM_{2.5} showing a larger contribution from the residential sector than PM₁₀, and PM₁₀ showing a larger contribution from transport, which includes the contribution from coarse particles from paved and unpaved roads. CO shows a larger contribution from the domestic sector and much larger daytime contribution from distant sources, which is expected due to the much longer lifetime of CO in the atmosphere than BC. In the case of NO_x, the transport sector dominates; except for the daytime periods, where other sources become important. During the daytime the power sector also contributes to surface concentrations as the growing mixed layer entrains the NO_x emissions from nearby power plants. During the nighttime the power plant plumes are above the mixed layer and are decoupled from the surface. Ozone has the largest contributions from sources outside of Delhi. This is shown more clearly in spatial plots of the contributions of outside the domain sources to the mean surface concentrations (Fig. 11). Within Delhi the outside sources are important and contribute from 20 to

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



50% depending on species. This has important implications for control strategies and indicates the need for regional perspectives.

3.4 Evaluation of emissions

The comparisons of the observations and predictions for each species were discussed earlier. Comparison of the correlations between different species can provide further information to help evaluate the emissions inventory. Illustrative results are shown in Fig. 12, where plots of hourly surface observations from all the sites combined are presented. The observations show good correlations between BC and CO, NO_x and CO, PM₁₀ and BC, and PM_{2.5} and PM₁₀. Also shown are the same correlations based on the modeled surface concentrations. If the model is perfect and the emission estimates are accurate, then the correlations between the modeled concentrations should be the same as those for the observed concentrations. The model values also show strong correlation, with higher species-species correlation coefficients than the observations. The PM_{2.5} to PM₁₀ and the PM_{2.5} to BC slopes are similar for the observations and the model concentrations. However absolute concentrations are over predicted. This suggests that the emissions are overestimated. In contrast the NO_x to CO modeled slope is too high, while the concentrations of NO_x and CO are over and under predicted, respectively. This suggest that the NO_x emissions are too high and the CO emissions too low. Assuming that the observations at these sites are representative and that all the model errors are associated with the emissions, then to make the modeled concentrations and slopes consistent with the observed values the emissions need to be scaled by the following: 0.6 for NO_x, 2 for CO, and 0.7 for BC, PM_{2.5} and PM₁₀. The emission estimates for particles are remarkably good considering the wide spread of activities and technologies taking place in Delhi and the rapid rates of change.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4 Conclusions

Stimulated by the CWG an air quality monitoring and forecasting system was established for Delhi. The monitoring program included eleven sites instrumented to obtain continuous measurements of BC, PM_{2.5}, PM₁₀, O₃ NO and NO₂. The air quality of Delhi during the CWG had high levels of particles with mean values of PM_{2.5} and PM₁₀ at the venues of 111 and 238 $\mu\text{g m}^{-3}$, respectively. BC accounted for $\sim 10\%$ of the PM_{2.5} mass. The model predictions were evaluated using the surface observations of meteorological parameters and the air pollution concentrations. The model showed a dry bias (under prediction in RH by $\sim 10\%$), slightly under predicted peak day time temperatures by $1\text{--}2^\circ$ ($\sim 5\%$), and was biased high in wind speed. The model captured the low wind speeds, but the observations often were less than 1 ms^{-1} , which the model estimates at $1\text{--}3\text{ ms}^{-1}$. BC, PM_{2.5} and PM₁₀ were over-predicted by $\sim 25\%$. The diurnal variations were well captured, with both the observations and the modeled values showing nighttime maxima and daytime minima. The daytime values compared well with the observations, but the nighttime values were over predicted, suggesting that the nighttime mixing height and/or nighttime dispersion were under predicted. Further work is needed to evaluate the mixing heights in Delhi for day and nighttime periods.

A new emissions inventory was developed to support this new air quality forecasting initiative. The emission inventory was evaluated by comparing the correlations in the observations (e.g., BC : CO; BC : PM_{2.5}; PM_{2.5} : PM₁₀). The predicted PM_{2.5} to PM₁₀ and the PM_{2.5} to BC slopes were similar to the observation-based slopes. However absolute concentrations were over predicted, suggesting that the emissions are overestimated. In contrast the NO_x to CO modeled slope was too high, while the concentrations of NO_x and CO were over and under predicted, respectively. This suggest that the NO_x emissions are too high and the CO emissions too low. Assuming that the observations at these sites are representative and that *all* the model errors are associated with the emissions, then the modeled concentrations and slopes can be made consistent by scaling the emissions by: 0.6 for NO_x, 2 for CO, and 0.7 for BC, PM_{2.5} and PM₁₀. The

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



emission estimates for particles are remarkably good considering the uncertainty in the estimates due to the diverse spread of activities and technologies that take place in Delhi and the rapid rates of change. A more complete evaluation of the modeling system and the emissions would benefit from additional observations including speciated NMOC, which would allow a more thorough evaluation of the photochemical oxidant cycle and ozone production.

The contribution of various emission sectors including transportation, power, domestic and industry to surface concentrations were also estimated. Transport, domestic and industrial sectors all make significant contributions to PM levels in Delhi, and the sectoral contributions vary spatially within the city. The transport sector is the main sector for ozone. Ozone levels in Delhi are already elevated, with hourly values sometimes exceeding 100 ppb. The continued growth of the transport sector is expected to make ozone pollution a more pressing air pollution problem in Delhi. The contribution for sources outside of Delhi on Delhi air quality was also estimated and was shown to vary spatially throughout the domain. The smallest contributions were found in the center of the domain, where on average the contributions ranged from ~ 25 % for BC and PM to ~ 60 % for day time ozone. The sector analysis provides useful inputs into the design of strategies to reduce air pollution levels in Delhi. The significant contributions from non-Delhi sources indicates that in Delhi (as has been show elsewhere) that these strategies will also need a more regional perspective.

The air quality activities established to support the CWG have been made operational, and the monitoring sites have been relocated to provide better spatial representation. Analysis of the 2 year time series of these observations will be the subject of a future paper.

Supplementary material related to this article is available online at <http://www.atmos-chem-phys-discuss.net/14/10025/2014/acpd-14-10025-2014-supplement.pdf>.

Acknowledgements. This research was supported in part through the NSF EaSM program (#1049140) and the USEPA STAR Grant program (#RD-83503701-0). The Indian Institute of Tropical meteorology, Pune, is supported by the Ministry of Earth Science, Government of India.

M. Decker and M. G. Schultz at The European Centre for Medium Weather Forecasts (ECMWF), Monitoring Atmospheric Composition and Climate (MACC) provided the chemical boundary conditions.

References

Arteta, J., Cautenet, S., Taghavi, M., and Audiffren, N.: Impact of two chemistry mechanisms fully coupled with mesoscale model on the atmospheric pollutants distribution, *Atmos. Environ.*, 40, 7983–8001, 2006.

Badarinath, K. V. S., Kharol, S. K., Sharma, A. R., and Prasad, V. K.: Analysis of aerosol and carbon monoxide characteristics over Arabian Sea during crop residue burning period in the Indo-Gangetic Plains using multi-satellite remote sensing datasets, *J. Atmos. Sol.-Terr. Phy.*, 71, 1267–1276, 2009.

Buka, I., Koranteng, S., and Osornio-Vargas, A. R.: The effects of air pollution of the health of children, *Paediatr. Child Health.*, 11, 513–516, 2006.

Carmichael, G. R., Adhikary, B., Kulkarni, S., D’Allura, A., Tang, Y., Street, D., Zhang, Q., Bond, T. C., Ramanathan, V., Jamroensan, A., and Marrapu, P.: Asia aerosol: current and year 2030 distributions and implications to human health and regional climate change, *Environ. Sci. Technol.*, 43, 5811–5817, 2009.

Chapman, E. G., Gustafson Jr., W. I., Easter, R. C., Barnard, J. C., Ghan, S. J., Pekour, M. S., and Fast, J. D.: Coupling aerosol-cloud-radiative processes in the WRF-Chem model: Investigating the radiative impact of elevated point sources, *Atmos. Chem. Phys.*, 9, 945–964, doi:10.5194/acp-9-945-2009, 2009.

Chung, C. E., Ramanathan, V., Carmichael, G., Kulkarni, S., Tang, Y., Adhikary, B., Leung, L. R., and Qian, Y.: Anthropogenic aerosol radiative forcing in Asia derived from regional models with atmospheric and aerosol data assimilation, *Atmos. Chem. Phys.*, 10, 6007–6024, doi:10.5194/acp-10-6007-2010, 2010.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- CPCB Report: National Ambient Air Quality Status & Trends in India 2010, PR Division, Central Pollution Control Board, 2012.
- Darby, L. S., McKeen, S. A., Senff, C. J., White, A. B., Banta, R. M., Post, M. J., Brewer, A. W., Marchbanks, R., Alvarez II, R. J., Peckham, S. E., Mao, H., and Talbot, R.: Ozone differences between near-coastal and offshore sites in New England: role of Meteorology, *J. Geophys. Res.*, 112, D16S91, doi:10.1029/2007JD008446, 2007.
- Davies, D. K., Ilavajhala, S., Wong, M. M., and Justice, C. O.: Fire information for Resource Management System: archiving and distributing MODIS active fire data, *IEEE T. Geosci. Remote*, 47, 72–79, 2009.
- Drimal, M., Lewis, C., and Fabianova, E.: Health risk assessment to environmental exposure to malodorous sulfur compounds in Central Slovakia, *Carpath. J. Earth Env.*, 15, 119–126, 2010.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010.
- Erismann, J. W., van Pul, A., and Wyers, P.: Parameterization of surface resistance for the quantification of atmospheric deposition of acidifying pollutants and ozone, *Atmos. Environ.*, 28, 2595–2607, 1994.
- Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology–chemistry–aerosol model, *J. Geophys. Res.*, 111, D21305, doi:10.1029/2005JD006721, 2006.
- Geng, F., Zhao, C., Tang, X., Lu, G., and Tie, X.: Analysis of ozone and VOCs measured in Shanghai: a case study, *Atmos. Environ.*, 41, 989–1001, 2007.
- Ghude, S. D., Jain, S. L., Arya, B. C., Beig, G., Ahammaed, Y. N., Kumar, A., and Tyagi, B.: Ozone in ambient air at a tropical megacity Delhi: characteristics trends and cumulative ozone exposure indices, *J. Atmos. Chem.*, 60, 237–252, 2008.
- Goyal, P., Rama Krishna, T. V. B. P. S., and Anand, S.: Assimilative capacity and dispersion of pollutants in Delhi, *India National Science Academy*, 69, 775–784, 2003.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Grell, G. A., Knoche, R., Peckham, S. E., and McKeen, S. A.: Online versus of-
 fline air quality modeling on cloud-resolving scales, *Geophys. Res. Lett.*, 31, L16117,
 doi:10.1029/2004GL020175, 2004.
- 5 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and
 Eder, B.: Fully coupled online chemistry within the WRF model, *Atmos. Environ.*, 39, 6957–
 6975, 2005.
- Guttikunda, S. K., Tang, Y., Carmichael, G. R., Kurata, G., Pan, L., Streets, D. G., Woo, J.-
 H., Thongboonchoo, N., and Fried, A.: Impacts of Asian megacity emissions on regional
 10 air quality during spring 2001, *J. Geophys. Res.*, 110, D20301, doi:10.1029/2004JD004921,
 2005.
- IPCC: Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to
 the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited
 by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and
 15 Miller, H. L., Cambridge University Press, Cambridge, UK, and New York, NY, USA, 996 pp.,
 2007.
- Kandlikar, M.: Air pollution at a hotspot location in Delhi: detecting trends, seasonal cycles and
 oscillations, *Atmos. Environ.*, 41, 5934–5947, 2007.
- Kim, D. and Stockwell, W. R.: An online coupled meteorological and air quality modeling study
 of the effect of complex terrain on the regional transport and transformation of air pollutants
 20 over the Western United States, *Atmos. Environ.*, 41, 2319–2334, 2007.
- Lau, M. and Kim, M. K.: Observational relationships between aerosol and Asian monsoon rain-
 fall, and circulation, *Geophys. Res. Lett.*, 33, L21810, doi:10.1029/2006GL027546, 2006.
- Madhav, B. G.: Transport and urban air pollution in India, *Environ. Manage.*, 36, 195–204, 2005.
- Madronich, S.: Photodissociation in the atmosphere, 1, actinic flux and the effects of ground
 25 reflections and clouds, *J. Geophys. Res.*, 92, 9740–9752, 1987.
- Mohan, M. and Anurag, K.: An analysis of the annual and seasonal trends of Air Quality Index
 of Delhi, *Environ. Monit. Assess.*, 131, 267–277, 2007.
- Mohan, M., Lalit, D., and Gurjar, B. R.: Preparation and validation of gridded emission inventory
 of criteria air pollutants and identification of emission hotspots for megacity Delhi, *Environ.*
 30 *Monit. Assess.*, 130, 323–339, 2006.
- Naseema Beegum, S., Krishna Moorthy, K., Suresh Babu, S., Satheesh, S. K., Vinoj, V., Badar-
 inath, K. V. S., Safai, P. D., Devara, P. C. S., Sacchidanand Singh, V., Dumka, U. C., and

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Pant, P.: Spatial distribution of aerosol black carbon over India during pre-monsoon season, *Atmos. Environ.*, 43, 1071–1078, 2009.
- Novaes, P., Hilario do Nascimento Saldiva, P., Matsuda, M., Macchione, M., Rangel, M. P., Kara-Jose, N., and Berra, A.: The effect of chronic exposure to traffic derived air pollution on the ocular surface, *Environ. Res.*, 110, 372–374, 2010.
- Rachna, A., Jayaram, G., Anand S., and Marimuthu, P.: Assessing respiratory morbidity through pollution status and meteorological conditions for Delhi, *Environ. Monit. Assess.*, 114, 489–504, 2006.
- Ravindra, K., Mor, S., Ameena, Kamyotra, J. S., and Kaushik, C. P.: Variation in spatial pattern of criteria air pollutants before and during Initial rain of monsoon, *Environ. Monit. Assess.*, 87, 145–153, 2003.
- Sahu, S. K., Beig, G., and Sharma, C.: Decadal growth of black carbon emissions in India, *Geophys. Res. Lett.*, 35, L02807, doi:10.1029/2007GL032333, 2008.
- Sarath, G. K., Carmichael, G. R., Calori, G., Eck, C., and Woo, J.-H.: The contribution of megacities to regional sulfur pollution in Asia, *Atmos. Environ.*, 37, 11–22, 2003.
- Sjaak, S. and Wayne, D.: Aerosols, in: *Encyclopedia of Earth*, edited by: Cutler, J., Environmental Information Coalition, National Council for Science and the Environment, Cleveland Washington DC, 2010.
- Suresh Babu, S., Satheesh, S. K., Krishna Moorthy, K., Dutt, C. B. S., Nair, V. S., Alapattu, D. P., and Kunhikrishnan, P. K.: Aircraft measurements of aerosol black carbon from a coastal location in the north-east part of peninsular India during ICARB, *J. Earth Syst. Sci.*, 117, 263–271, 2008.
- Tie, X., Madronich, S., Li, G., Ying, Z., Zhang, R., Garcia, A. R., Taylor, J. L., and Liu, Y.: Characterization of chemical oxidants in Mexico City: a regional chemical dynamical model (WRF-CHEM) study, *Atmos. Environ.*, 41, 1989–2008, 2007.
- Tulet, P., Crassier, V., Cousin, F., Suhre, K., and Rosset, R.: ORILAM, a three-moment log-normal aerosol scheme for mesoscale atmospheric model: online coupling into the Meso-NH-C model and validation on the Escompte campaign, *J. Geophys. Res.*, 110, D18201, doi:10.1029/2004JD005716, 2005.
- Xu, Y. and Carmichael, G. R.: Modelling the dry deposition velocity of sulfur dioxide and sulfate in Asia, *J. Appl. Meteorol.*, 37, 1084–1099, 1998.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Viney, A. P., Agarwal, A., Roelle, P. A., Phillips, S. B., Tong, Q., Watkins, N., and Yablonsky R.: Measurements and analysis of criteria air pollutants in New Delhi, India, *Environmental International*, 27, 35–42, 2001.

Wesley, M. L.: Parameterization of surface resistance to gaseous dry deposition in regional numerical models, *Atmos. Environ.*, 16, 1293–1304, 1989.

Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res.*, 113, D13204, doi:10.1029/2007JD008782, 2008.

Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, doi:10.5194/acp-9-5131-2009, 2009.

Zhang, Y.: Online-coupled meteorology and chemistry models: history, current status, and outlook, *Atmos. Chem. Phys.*, 8, 2895–2932, doi:10.5194/acp-8-2895-2008, 2008.

Air quality in Delhi during the CommonWealth Games

P. Marrapu et al.

Table 1. Important input settings of the WRF-Chem model used in this study.

Feature	Option	Description
Chemical Mechanism	CBMZ	Carbon Bond Mechanism v.Z with MOSAIC 4 aerosol bins (Zaveri et al., 2008)
Microphysics	Lin et al. scheme	Sophisticated scheme with ice, snow, and graupel processes.
Longwave Radiation	Rapid Radiative Transfer Model (RRTM)	Accounts for multiple bands, trace gases, and microphysics
Shortwave Radiation	Goddard shortwave	Two-stream multi-band scheme with ozone from climatology and cloud effects.
Surface Layer	MM5 similarity	Based on Monin–Obukhov with Carlson–Boland viscous sub-layer and standard similarity functions from look-up tables.
Anthropogenic Emissions	Delhi Inventory & INTEX-B	Delhi inventory at 1.67 km resolution Intercontinental Chemical Transport Experiment B (INTEX-B) data at $0.5^\circ \times 0.5^\circ$ resolution (Zhang et al., 2009).
Biogenic Emissions	MEGAN	MEGAN Model of Emissions of Gases and Aerosols from Nature, biogenic emissions on-line based upon the weather, land use data.
Boundary Conditions	MACC	Monitoring Atmospheric Composition & Climate, a global 3-D chemical transport model driven by offline meteorological fields.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Anthropogenic emissions and % sectoral contribution for Delhi.

Species	tday ⁻¹	Contribution from each anthropogenic sector			
		Transportation	Power	Industry	Domestic
SO ₂	81	12.9 %	48.7 %	25.2 %	13.2 %
NO _x	598.5	69.4 %	13.2 %	4.5 %	12.9 %
CO	1320.3	43.7 %	0.2 %	4.0 %	52.0 %
PM ₁₀	344.8	86.8 %	7.9 %	4.6 %	0.8 %
PM _{2.5}	128.6	52.6 %	9.9 %	15.3 %	22.2 %
BC	36.9	58.9 %	3.0 %	6.6 %	31.5 %
OC	35.1	30.5 %	5.6 %	10.6 %	53.3 %
NMOC	852.4	58.4 %	1.2 %	5.2 %	35.3 %

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Table 3. Comparison of predicted meteorological parameters with observations at the surface monitoring sites.

	Temp (°C)	Wind speed (ms ⁻¹)	RH (%)
Mean – Obs	29.0	1.3	53.5
Mean – Model	28.7	2.2	45.8
Bias Error	7.7	1.3	7.0
RMSE	13.6	2.9	33.9
<i>R</i>	0.9	0.5	0.7

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Nested model domains used in the Commonwealth Games forecasts and analysis. Insert shows the locations of the observations sites.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.





Fig. 2. Emission distributions of NO_x (top) and, PM_{2.5} in tyr⁻¹ over Delhi.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.




Fig. 3. Sector contributions to black carbon emissions (%) for Delhi.

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
◀ ▶
◀ ▶
[Back](#) [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

ACPD

14, 10025–10059, 2014

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Locations of the monitoring stations for the Commonwealth Games.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 5. Spatial distributions of calculated 20 Day mean concentrations of surface BC ($\mu\text{g m}^{-3}$) and daytime (9.30 a.m.–6.30 p.m. IST) ozone (ppb) over Delhi. For ozone daytime means are shown.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.




Fig. 6. Comparison of the distributions of observed and modeled hourly concentrations using data from all sites. Base corresponds to full emissions whereas control shows results where the NO_x emissions were reduced by a factor of 3.

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.





Fig. 7. Mean diurnal variation in boundary layer height, emissions and BC concentrations from model and observations at the Dhayanchand Stadium site.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

**Fig. 8.** Percent contribution of each species to total $PM_{2.5}$ at Dhyanchand Stadium.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 9. Spatial distributions of sector contributions to period mean surface BC concentrations.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.




Fig. 10. Diurnal variation in the sector contribution (in %) to period mean surface concentrations of different pollutants at the Dhayanchand monitoring station.

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)






Fig. 11. 20 day mean contributions of pollutants from outside source.

Air quality in Delhi during the Commonwealth Games

P. Marrapu et al.




Fig. 12. Comparison of correlations between different species for all the sites combined based on observations and modeled concentrations.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

⏴ ⏵

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

