

1 **Supplementary Table 1.** Percent area contributions from the major proton regions in the <sup>1</sup>H NMR spectra for the PSOM aerosol extracts.

2

<b>Functional Group Region</b>	<b>Chemical Shift (ppm)</b>	<b>% <sup>1</sup>H NMR spectral area</b>		
		<b>16-17 August 2011</b>	<b>24-25 June 2013</b>	<b>25-26 June 2013</b>
H-C-O	0.7 – 1.95	2.5	2.5	6.4
H-C-C=	1.95 – 3.2	18.3	21.0	25.5
H-C	3.2 – 4.4	78.8	76.0	67.0
<i>Calculated H/C</i>		<i>1.98</i>	<i>1.98</i>	<i>1.94</i>

3

4

5 **Supplementary Table 2.** Percent OC extraction for PSOM extracts using <sup>1</sup>H NMR and WSOM using TOC analysis.

6


<b>Aerosol Sample</b>	<b>Initial OC mass (mg)</b>	<b>Spectral Area (intensity units)</b>	<b>**Calc DOC (mg)</b>	<b>%PSOC</b>	<b>%WSOC (by TOC)</b>
16-17 August 2011	*0.045	2.4 x 10 <sup>10</sup>	0.041	90.3	37.0 ± 2.2
24-25 June 2013	0.039	1.1 x 10 <sup>10</sup>	0.021	53.8	54.3 ± 3.8
25-26 June 2013	0.049	9.1 x 10 <sup>9</sup>	0.018	36.5	60.3 ± 3.6
Glucose	0.114	2.5 x 10 <sup>9</sup>	-	-	-

7 \*There is a known error in the initial mass measurement of this aerosol, and was omitted from the results

8 \*\*Calculated by multiplying the spectral area by the glucose response factor (2.6x10<sup>11</sup> intensity units/mg H), and then converting mg H to mg C  
9 using the H/C ratio calculated in Supplementary Table 1.

10

11



12  
13  
14  
15  
16

**Supplemental Figure 1.** Full  $^1\text{H}$  NMR spectra for a) WSOM and b) PSOM for the aerosol particulate sample collected 25-26 June 2013. The strong signal (off scale) in the PSOM spectra in the aromatic region is from protons that have been exchanged in the pyridine- $\text{D}_5$  solvent, which overwhelms any possible aromatic signal from the sample.