1 Relating hygroscopicity and optical properties to chemical

2 composition and structure of secondary organic aerosol

³ particles generated from the ozonolysis of α-pinene

4 C. Denjean^{1,2}, P. Formenti¹, B. Picquet-Varrault¹, E. Pangui¹, P. Zapf¹, Y. Katrib¹, C. Giorio^{3,*},

5 A. Tapparo³, A. Monod⁴, B. Temime-Roussel⁴, P. Decorse⁵, C. Mangeney⁵ and J.F. Doussin¹

6 [1] Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), UMR-CNRS 7583,

7 Université Paris-Est-Créteil (UPEC) et Université Paris Diderot (UPD), Institut Pierre Simon

- 8 Laplace (IPSL), Créteil, France
- 9 [2] Leibniz Institute for Tropospheric Research, Permoserstr. 15, 04318, Leipzig, Germany
- 10 [3] Dipartimento di Scienze Chimiche, Università degli Studi di Padova, via Marzolo 1, 35131
- 11 Padova, Italy
- 12 [4] Aix Marseille Université, CNRS, Laboratoire Chimie de l'Environnement (LCE), FRE 3416,
- 13 13331 Marseille, France
- 14 [5] ITODYS, Université Paris Diderot, Sorbonne Paris Cité, CNRS UMR 7086, 15 rue J-A de
- 15 Baïf, 75013 Paris, France
- 16 *present address: Department of Chemistry, University of Cambridge, Lensfield road, CB2
- 17 1EW Cambridge, UK
- 18
- 19 Correspondence to: C. Denjean (denjean@tropos.de)
- 20

21 Abstract

Secondary Organic Aerosols (SOA) were generated from the ozonolysis of α -pinene in the CESAM simulation chamber. The SOA formation and aging were studied by following their optical, hygroscopic and chemical properties. The optical properties were investigated by determining the particle Complex Refractive Index (CRI). The hygroscopicity was quantified by measuring the effect of RH on the particle size (size growth factor, GF) and on the scattering coefficient (scattering growth factor, *f(RH)*). The oxygen to carbon atomic ratios (O:C) of the particle surface and bulk were used as a sensitive parameter to correlate the changes in 1 hygroscopic and optical properties of the SOA composition during their formation and aging in

2 CESAM.

- 3 The real CRI at 525 nm wavelength decreased from $1.43-1.60 (\pm 0.02)$ to $1.32-1.38 (\pm 0.02)$ during
- 4 the SOA formation. The decrease in the real CRI correlated to the O:C decrease from $0.68 (\pm 0.20)$
- 5 to 0.55 (\pm 0.16). In contrast, the GF remained roughly constant over the reaction time, with values
- 6 of 1.02-1.07 (±0.02) at 90% (±4.2%) RH. Simultaneous measurements of O:C of the particle
- 7 surface revealed that the SOA was not composed of a homogeneous mixture, but contained less
- 8 oxidised species at the surface which may limit water absorption. In addition, an apparent change
- 9 in both mobility diameter and scattering coefficient with increasing RH from 0 to 30% was
- 10 observed for SOA after 14 hours of reaction. We postulate that this change could be due to a
- change in the viscosity of the SOA from a predominantly glassy state to a predominantly liquidstate.
- 13

14 **1** Introduction

15 Organic compounds are known to account for a large fraction of atmospheric aerosol, ranging 16 between 20 and 90% of the total particle mass (Kanakidou et al., 2005). In particular, secondary organic aerosol (SOA), formed by the condensation of gas-phase oxidation products of volatile 17 18 organic compounds (VOCs), is a major constituent of the total organic aerosol (Turpin and 19 Huntzicker, 1995; Zhang et al., 2007). SOA size ranges between ten to hundreds of nanometers. 20 Particles in this size range have long atmospheric lifetimes and scatter solar radiation. SOA can 21 also change clouds properties by acting as Cloud Condensation Nuclei (CCN) (Saxena et al., 22 1995; Lohmann and Feichter, 2005; Novakov and Penner, 1993; RiveraCarpio et al., 1996; 23 Matsumoto et al., 1997).

24

There are large uncertainties in estimating the impact of SOA on climate due to their complexity and the limited range of measurements available (Kanakidou et al., 2005). SOA precursors produce a large number of oxidation products (Goldstein and Galbally, 2009), resulting in many possible chemical reaction pathways (de Gouw et al., 2005; Hallquist et al., 2009). In addition, during their lifetime in the atmosphere, SOA may undergo several physical and chemical aging processes altering their chemical composition (Kalberer et al., 2004; Baltensperger, 2005; Yasmeen et al., 2012) and size distribution (Andreae, 2009). As a result, atmospheric SOA 1 contain many organic compounds with a large variety of structures, chain lengths, functionalities

- 2 and degrees of oxidation (Kroll and Seinfeld, 2008; Jimenez et al., 2009). Therefore SOA possess
- 3 a wide range of hygroscopic and optical properties (Lambe et al., 2013; Suda et al., 2012).
- 4

5 In global climate models, the direct radiative effect of SOA is currently described by adopting a constant Complex Refractive Index (CRI) and a single size Growth Factor (GF). Depending on 6 7 the model, the adopted real part of the CRI at visible wavelengths ranges from 1.45 to 1.60 (Kinne 8 et al., 2003; Pere et al., 2011; Zaveri et al., 2010). Some models assume that SOA absorb weakly 9 solar radiation, and set the imaginary part of the CRI near 0.006, while others ignore the absorption by SOA. Concerning hygroscopic properties, a single size GF of SOA derived from 10 11 limited available data is used (O'Donnell et al., 2011; Hoyle et al., 2009). However, field 12 measurements shows that the hygroscopic and optical properties of SOA are not static and 13 depend on their origin and transport in the atmosphere (Duplissy et al., 2010; Jimenez et al., 14 2009; Chang et al., 2010; Dinar et al., 2008). As a consequence, laboratory investigations have 15 started to explore the change of SOA properties during their lifetime. The ozonolysis of α -pinene ozonolysis is one of the most well studied SOA systems (α -pinene-O₃ SOA), as α -pinene is a 16 17 significant biogenic VOC in many regions, and its ozonolysis plays an important role in SOA formation (Hallquist et al., 2009; Griffin et al., 1999; Yu et al., 1999; Kavouras et al., 1998, 18 19 1999). α -pinene-O₃ SOA is also generally considered as a model for many biogenic VOCs containing an endocyclic double bond (Guenther et al., 1995). Saathoff et al. (2003) conducted 20 21 experiments of α -pinene-O₃ SOA in a simulation chamber and observed an increase of the GF at 22 90% RH from 1.080 to 1.106. Cocker et al. (2001) reported also an increase of the GF at 85% 23 RH from 1.065 to 1.11 within 6 hours. In contrast, Warren et al. (2009) and Oi et al. (2010) 24 reported a constant GF at 90% RH for α -pinene-O₃ SOA over 6 hours of reaction. Finally, an 25 increase of the real CRI from 1.39 to 1.52 at $\lambda = 532$ nm during the formation of the α -pinene-O₃ SOA has been reported by Kim et al. (2013). To date, none of the previous studies have 26 27 simultaneously determined the hygroscopic and optical properties, and their evolution with the particle chemical composition during SOA formation and ageing. 28

29

30 To accurately quantify the SOA impacts on climate, it is critical to determine the hygroscopic 31 behaviour of both the size distribution and optical properties as well as the dependence of these 32 properties on the chemical composition. Atmospheric simulation chambers are powerful tools to 1 study the physical, chemical, optical and hygroscopic properties of SOA and follow their changes 2 along their lifecycle by simulating their formation and aging due to oxidation and photolysis in the atmosphere (Meyer et al., 2008; Henry and Donahue, 2012; Tritscher et al., 2011). In this 3 4 work, we take advantage of the long aerosol lifetime in the CESAM chamber (French acronym 5 for Experimental Multiphasic Atmospheric Simulation Chamber, Wang et al., 2011) to set-up formation and aging experiments of α -pinene-O₃ SOA in order to characterise the evolution of 6 7 both the SOA optical properties and hygroscopicity. The objective is to examine the evolution of 8 these physical properties and to relate them to the aerosol chemical composition.

9

10 **2 Methods**

Measurements have been conducted in the humidity-controlled simulation chamber CESAM (Wang et al., 2011), which permits the study of the formation and aging of SOA over long periods of time, and under various relevant atmospheric conditions (temperature, relative humidity, pressure, gas phase concentration, etc.).

15 Experiments were conducted to simultaneously measure different parameters:

- CRI at λ=525nm, a specific wavelength in the mid-visible; the Complex Refractive Index
 (CRI, m = n-ik) is an important parameter to link the physical and chemical properties of
 the SOA and its ability to interact with radiation, allowing a description of the scattering
 and absorbing characteristics of SOA.
- GF(RH), size growth factor, the ratio of the particle diameter at a given RH to the particle diameter at low RH, for one selected size of particles; this parameter is used to characterise the hygroscopic properties of the SOA and thus, it is an indicator of its ability to act as CCN.
- f(RH), scattering growth factor, the ratio of the scattering coefficient (σ_{scat}) at high RH to 25 the σ_{scat} at low RH, for the entire size distribution; this parameter allows the study of the 26 effect of water absorption on the scattering properties of SOA.
- O:C, the oxygen-to-carbon ratio, for both the bulk and the surface composition of the SOA; it is well known that particle composition, in particular that of the surface, strongly influences the water uptake ability of the particle and its CCN potential (McFiggans et al., 2006; Dusek et al., 2006; Hatch et al., 2008; Moussa et al., 2009; McIntire et al., 2010; Semeniuk et al., 2007; Lamb et al., 2011; Wong et al., 2011; Mei et al., 2013; Rickards et al., 2013). To encompass the difficulty of representing the full molecular composition, the O:C of SOA has been included in global climate models to provide a description of

the aerosol aging (Tost and Pringle, 2012). Furthermore, a number of recent studies have
 reported a positive correlation between hygroscopicity and bulk O:C for both laboratory
 and ambient SOA (Massoli et al., 2010; Jimenez et al., 2009; Chang et al., 2010; Duplissy
 et al., 2011).

5

6 **2.1** Simulation chamber and associated instruments

7 CESAM is a 4.2 m³ cylindrical stainless steel chamber which has been designed to investigate 8 both atmospheric gas-phase and aerosol-phase chemistry. As described by Wang et al. (2011), 9 the wall properties and ventilation system guarantee a lifetime for long sub-micron particles and 10 enables the study of aerosol aging for more than 20h. Water vapour can be directly injected in 11 CESAM and thus the RH of the reaction mixture can be varied *in situ* from 0 to 100%.

12

The basic experimental setup and a schematic view of the CESAM chamber are shown in Figure 14 1. Temperature and relative humidity in the chamber are monitored using a Vaisala transmitter 15 equipped with a capacitive thin-film Humicap sensor. The sensor was calibrated prior each 16 experiments. The RH accuracy was $\pm 1.9\%$ up to 90% RH and the temperature accuracy was 17 $\pm 0.1^{\circ}$ C at 20°C.

18

a-pinene, ozone and their reaction products were continuously monitored by a Fourier-Transform
InfraRed spectrometer (FTIR) from Bruker GmbH (Ettlingen, Germany) coupled to a multi
reflection cell allowing an optical path of 192 m. Additionally, ozone was monitored by a
commercial Horiba APOA 370 instrument (Kyoto, Japan) with a detection limit of 0.5 ppb and
a precision of 0.1 ppb.

24

25 The particle number size distribution was measured using a Scanning Mobility Particle Sizer 26 (SMPS including a DMA 3080 and CPC 3010, TSI). The SMPS was operated at flow rates 3/0.3 27 Lpm (sheath flow/aerosol sample flow). The aerosol flow was diluted with filtered air before entering the CPC, in order to maintain the nominal flowrate at 1 Lpm in the CPC. The dilution 28 29 air flow was sucked from the simulation chamber to avoid any pressure gradient in the SMPS. 30 The SMPS scanning time was 2 minute 15 sec in total. The resulting measured size distribution 31 ranged from 14 to 505 nm. The size calibration of the SMPS was performed using monodisperse 32 PolyStyrene Latex spheres (PSL, Duke Scientific). PSL, with diameters ranging from 100 to 500

nm, were nebulised with a constant output atomiser (TSI model 3076). The measured diameters were found to be larger than the PSL certified diameters (by about 10% for 100 nm PSL spheres), so a correction factor was applied. Size distributions were corrected by the SMPS software for both the loss by diffusion of particles in the SMPS tubing, the contribution of multicharged particles and the dilution of the aerosol flow before entering the CPC.

6

7 2.2 Optical properties measurements

8 The aerosol scattering coefficient (σ_{scat}) was measured using an integrating nephelometer (model 9 M9003, Ecotech). The nephelometer operates at 525 nm wavelength and measures light scattered 10 from particles at angles between 10° and 170°. It also measures temperature at both the sample 11 inlet and within its cell with an accuracy of ±0.6°C, and the RH within its cell with an accuracy 12 of ±3%. Prior to each series of experiments, the nephelometer was calibrated using particle-free 13 air and CO₂.

14

15 The particle light absorption coefficient (σ_{abs}) was determined by means of an aethalometer 16 (Model AE31, Magee Scientific) operated with several light sources at seven wavelengths, covering the near ultra-violet to the near infrared wavelength range (λ =370, 470, 520, 590, 660, 17 18 880 and 950 nm). The aethalometer measures the attenuation of the transmitted light through a 19 quartz fibre filter with increasing particle loading. This measurement can suffer from artefacts 20 associated with reactions with oxidants occurring on particles deposited on the surface of the 21 filter and desorption of gaseous compounds from the filter (Weintgartner et al., 2003). Thus, a 22 charcoal denuder was installed upstream of the aethalometer to remove ozone and VOCs. It has 23 also been observed that the aethalometer can suffer from biases at high RH as a result of the filter taking up water and scattering more light compared to the reference measurement (Cappa et al., 24 25 2008; Arnott et al., 2005). Therefore, the aethalometer was not used when the chamber RH was 26 higher than 1%.

27

The spectral attenuation coefficient (σ_{attn}) resulting from the attenuation of light through the sampled aerosol on the filter was obtained using equation (1):

$$\sigma_{\text{attn}} \left(\lambda, \mathbf{m} \right) = \frac{A}{Q} \frac{\Delta \text{attn}}{\Delta t} \tag{1}$$

1 where A is the spot area, Q is the volumetric flowrate and $\Delta attn is the change in light attenuation$ $2 during the time interval <math>\Delta t$. It is well known that σ_{attn} obtained with aethalometers are higher than 3 true σ_{abs} (Arnott et al., 2005; Bond et al., 1999; Bond and Bergstrom, 2006; Cappa et al., 2008; 4 Collaud Coen et al., 2010; Weingartner et al., 2003). Various systematic errors need to be 5 corrected. The detailed description of the method used to correct σ_{attn} is shown in the 6 Supplementary Material.

7

8 The details of the calculations of CRI of SOA under dry conditions (RH<1%) were given in 9 Denjean et *al.* (2014). The CRI was determined by optical closure experiments involving 10 scattering and absorption coefficients (respectively σ_{scat} and σ_{abs}) measured at 525 nm and the 11 number size distribution. Briefly, σ_{scat} and σ_{abs} were calculated according to equation (2):

$$\sigma_{\text{scat,abs}}(\lambda, m) = \int Q_{\text{scat,abs}}(D_{\text{p}}, \lambda, m) \frac{\pi}{4} D_{\text{p}}^{2} \left(\frac{dN}{d\log D_{\text{p}}}\right) d\log D_{\text{p}}$$
(2)

12 where D_p is the geometrical particle diameter, $\frac{dN}{dlogD_p}$ is the number size distribution and 13 $Q_{\text{scat,abs}}(D_p,\lambda,m)$ represent both the scattering and absorption efficiencies. $Q_{\text{scat,abs}}(D_p,\lambda,m)$ were 14 calculated using Mie scattering calculations described by Bohren and Huffman (1983).

15

16 The values of σ_{scat} and σ_{abs} of SOA were simultaneously measured and compared with those 17 calculated based on the Mie theory. To allow this comparison, all measured σ_{scat} were corrected 18 from the sample temperature and pressure and from the angular truncation error examined for 19 the nephelometer. The best-guess CRI was determined by minimizing the difference between 20 measured σ_{scat} and σ_{abs} and those obtained using Mie calculations.

21

22 2.3 Hygroscopic properties measurements

The hygroscopic properties of SOA were investigated using two complementary approaches as described in details in Denjean et al. (2014). Briefly, (1) a Hygroscopic Tandem Differential Mobility Analyzer (H-TDMA) was used to measure the RH dependency of the particles diameter for single size particles and (2) *in situ* experiments within the CESAM chamber allowed for the measurement of σ_{scat} changes after water uptake for polydisperse aerosol size distribution.

1 The hygroscopic behaviour of monodisperse SOA was measured with a H-TDMA. The 2 instrument is composed of a DMA (TSI, 3080) that selects an initial mobility diameter under dry 3 conditions D_{p.m}(RH_{dry}), an aerosol humidifier with a controlled higher RH and a DMA (TSI, 4 3080) coupled to a CPC (TSI, 3010) to measure the wet size distribution over mobility diameter 5 D_{p.m}(RH). The total sampling line used for HTDMA measurements ranged between 10 and 20 cm. Both DMAs of the H-TDMA were calibrated using monodisperse PSL particles (Duke 6 7 Scientific) ranging from 100 to 500 nm. A size shift was observed for both DMAs. The particles 8 residence time for humidification is 15 seconds and corresponds to the residence time in the 9 aerosol humidifier plus the transit time in the second DMA. In this study, the H-TDMA was typically operated at a constant high RH (\pm 1%) of 90%. The hygroscopic size growth factor 10 11 (GF) describes the relative increase in the geometric diameter of particles due to water uptake at 12 a specific RH according to equation (3):

$$GF(RH) = \frac{D_{p,m}(RH)}{D_{p,m}(40\% RH)}$$
(3)

Due to possible changes in SOA viscosities (i.e. section 4.2.), RH=40% was considered dry in this study. GF at 40% RH was measured at the beginning and the end of each series of H-TDMA measurements. Dry and humidified mobility diameters were obtained by assuming that the size distributions exhibited a lognormal profile. H-TDMA measurements were validated by measuring the humidogram of laboratory generated ammonium sulfate particles. The GF was found to agree with values calculated using the Köhler model (Denjean et *al.*, 2014).

19

20 In addition to H-TDMA measurements, the hygroscopic behaviour of polydisperse SOA was 21 measured by exposing the particles *in situ* to increasing humidity in the CESAM chamber. Water 22 vapour produced from a small glass vessel filled with ultrapure water (18.2M Ω , ELGA Maxima) 23 was injected into the chamber and mixed therein by the chamber's stainless steel fan. The RH in 24 the chamber increased linearly from 0 to 100% within approximately 1 hour. The change in σ_{scat} due to hygroscopic growth of the overall particle size distribution was monitored by the 25 26 nephelometer. The scattering growth factor (f(RH)) is the ratio between the scattering coefficient 27 at a specific RH ($\sigma_{scat}(RH)$) and the dry scattering coefficient ($\sigma_{scat}(dry)$) (equation 4):

$$f(RH) = \frac{\sigma_{scat}(RH)}{\sigma_{scat}(40\% RH)}$$
(4)

During humidification, RH was monitored within the chamber and at the inlet of the nephelometer. Using this approach, the residence time for particle humidification was a few minutes thus significantly longer than during H-TDMA measurements. It has been previously shown (Denjean et al., 2014) that the two approaches of hygroscopicity measurements could lead to different results, giving information on water transfer dynamics, possible particles reorganisation or phase transfer equilibrium.

7

8 2.4 Aerosol sampling and chemical composition analysis

9 A High Resolution time-of-flight aerosol mass spectrometer (AMS, Aerodyne) (DeCarlo et al., 2006) was used to determine the bulk composition of organic aerosols (Aiken et al., 2007; Aiken 10 11 et al., 2008). The instrument was used under standard conditions (vaporiser at 600°C and electron 12 ionisation at 70 eV). Standard AMS calibration procedures using NH₄NO₃ particles were 13 performed throughout the campaign and included Ionisation Efficiency (IE) calibration using 14 Brute Force Single Particle (BFSP) method at 350 nm particle mobility diameter as well as size 15 calibration using DMA-selected dried particle sizes over the range of interest, i.e. 100-350 nm. Single ion calibration as well as baseline and threshold were analysed prior each experiment. 16

17

18 The AMS was operated during three experiments. The instrument was switched between two 19 modalities: a single-reflectron configuration (V-mode) which offers a higher sensitivity and 20 lower resolving power (up to ~ 2100 at m/z 200) and a double-reflectron configuration (W-mode) which provides a higher resolving power (up to \sim 4300 at m/z 200) but a lower sensitivity (De 21 22 Carlo et al., 2006). Data were taken with a time resolution of 8 minutes. In V-mode, data were 23 collected in the mass spectrum (MS) mode for 5 minutes, for aerosol quantification, and in the 24 particle time of flight (PToF) mode for 1 minute for size distribution measurement (De Carlo et 25 al., 2006, Canagaratna et al. 2007). Only MS data were recorded in the W-mode (2 minutes).

26

The AMS data were analysed using Squirrel (ToF-AMS Analysis 1.51B) and Pika (ToF-AMS HR Analysis 1.10B) packages for the software Igor Pro 6.21 (Wavemetrics, Inc., Portland, OR, USA). Default Collection Efficiencies (CE) and Relative Ionisation Efficiencies (RIE) were used for quantification of SOA. Air interferences were removed by adjusting the fragmentation table (Aiken *et al.*, 2008; Allan *et al.*, 2004). High-resolution analyses (De Carlo *et al.*, 2006) were performed on V-mode data by integrating each $C_xH_yO_z$ ion in the mass range 12–180 m/z while W-mode data were used only to check for possible interferences. Elemental ratios (O:C and H:C)
 were calculated according to the procedure described by Aiken et al. (2007; 2008). Measurement
 uncertainties were estimated to be ±30%, as determined by Aiken et al. (2008).

4

5 Further chemical analyses were performed by collecting SOA on PTFE filters (Zefluor, 47mm diameter, 2 µm pore size, Pall Life Sciences), which were cut to the size of the collector using 6 7 ceramic scissors. A stainless-steel support was used for sampling and concentrating the particles 8 in a small filter area (0.9 mm²). An active charcoal denuder was installed upstream of the filter 9 to remove ozone and VOCs. Before sampling, filters were extracted three times for 10 minutes with dichloromethane (99.8%, HPLC grade) in an ultrasonic bath and baked for 5h at 250°C. 10 11 Filter samplings were performed in the chamber at different reaction times from 30 minutes to 12 17 hours at a nominal flow rate of 2 Lpm for a sampling time varying between 30 min to 2 hours, 13 depending on the total SOA volume concentration.

14

15 X-ray photoelectron spectrometry (XPS) was used to quantify the O:C of the SOA surface to a depth less than 10 nm. Measurements were performed on a VG ESCALAB 250 instrument using 16 17 monochromatic Al K_{α} radiation (1486.6 eV). The X-Ray spot size was 500 μ m. The analysis chamber of the instrument was maintained at pressures ranging between 10⁻⁸ and 10⁻¹¹ mbar. 18 19 Survey spectra of SOA were measured over an 1100 eV range at a resolution of 1 eV per step 20 and 100 eV pass energy (Figure S1a in the Supplementary Material). All peaks were referred to 21 the C_{1s} binding energy at 285.0 eV. The high-resolution C_{1s} spectrum showed the presence of C, 22 O and F. The deconvolution of the C_{1s} signal was performed in peaks corresponding to the $-CO_2$ 23 (indicative of carboxylic acids, peroxides), C-O (alcohols, aldehydes, carboxylic acids, 24 peroxides, ethers), C-C/C-H (aliphatic functional groups) and the C-F bonds of the PTFE filter 25 in order to optimise the fit (Figure S1b). Quantification of the O:C was performed using the 26 integrated peak areas of O_{1s} spectra, the peaks area of the $-CO_2$, C-O and C-C/C-H from the C_{1s} 27 spectra and the manufacturer's sensitivity factors.

28

To exclude biases in the comparison of the surface and bulk O:C ratios by different experimental methods, a control experiment was performed on pinic acid which is pure homogeneous compound containing multiple carbon-oxygen bonds and which is representative of α -pinene-O₃ SOA (theoretical value O:C value of 0.44). 0.003 M pinic acid solution (Santai Labs, purity by 'HNMR of >95%) was atomized using a constant output atomizer (TSI, model 3076) and then passed through a Nafion tube, which reduced the RH below 35%. Particles were analyzed with the AMS and simultaneously collected on PTFE filters for XPS analysis. The O:C values obtained experimentally from this single measurement by AMS and XPS were 0.34 (\pm 0.10) and 0.37 (\pm 0.08), respectively, as expected for particles homogeneous composition. In addition, both methods reproduced the expected molecular O:C of pinic acid within the uncertainty range. This control experiment sets the reference case for the joint use of XPS and AMS to resolve compositional differences in non-homogenous SOA.

8

9 **2.5 Experimental protocol**

10 Experiments were carried out without any seed particles or OH radical scavenger. Prior to each experiment, the CESAM chamber was evacuated overnight to typically 4×10^{-4} mbar. The 11 chamber was then filled to atmospheric pressure with a mixture of 200 mbar of Oxygen (Air 12 Liquide, Alphagaz class 1, purity 99.9%) and 800 mbar of Nitrogen produced from the 13 14 evaporation of a pressurised liquid nitrogen tank (Messer, purity > 99.995%, H₂O < 5ppm). Table 15 1 shows the experimental conditions of this study. Ozone was generated by a Corona discharge 16 in pure O₂ using a commercial dielectric ozone generator (MBT 802N, Messtechnik GmbH, Stahnsdorf, Germany). After ozone concentration stabilisation around 250 ppb in the chamber, 17 18 approximately 200 ppb of α -pinene was introduced. This injection was performed by flushing 19 into the chamber (in an O₂ flow) the entire volume of a glass bulb containing a known pressure 20 in a glass bulb of a known volume of pure gaseous α -pinene (Aldrich, 98%). Particles were 21 formed immediately after α -pinene injection. During the experiments, RH was lower than 1% 22 and the temperature ranged between 14°C and 27°C (for different experiments).

23

The total pressure in the chamber was maintained constant by adding gaseous nitrogen to compensate for the continuous sampling by the instruments. As a consequence, the N₂:O₂ ratio increased with the reaction time. After 20 hours of reaction, the hygroscopicity of polydisperse particles was investigated by injecting water vapour in the chamber. The RH in the chamber increased linearly from 0 to 100% within approximately 1 hour. The nephelometer was used to follow *in situ* the variation in σ_{scat} as a function of RH.

30

31 **3. Results**

3.1. Reaction profile and aerosol yield

Typical evolution of α -pinene, ozone, particle number size distribution and mass concentrations are shown in Figure 2. α -pinene was essentially consumed after 4 hours of reaction. Since no OH scavenger was used, α -pinene was oxidised by O₃ and also by OH radicals produced by the reaction. A rapid formation of SOA was observed as the α -pinene was oxidised. The aerosol mass reached its maximum of 130 ± 39 µg m⁻³ (when most α -pinene was consumed) and remained constant thereafter.

8

9 Table 1 summarises the results for the experiments conducted. The aerosol yield (*Y*) was given10 by equation 5:

$$Y = \frac{\Delta[SOA]}{\Delta[\alpha pinene]}$$
(5)

where Δ [SOA] is the SOA mass concentration produced for a given amount of α -pinene reacted, 11 12 Δ [apinene], both corrected for dilution. Since α -pinene was introduced after ozone into the chamber, and due to the very fast consumption of α -pinene, its initial concentration could not be 13 14 accurately measured. For the yield calculations, we used the SOA mass concentration and α pinene concentration after 5 minutes of reaction as initial concentrations. SOA mass 15 16 concentration was calculated from the SMPS number size distribution corrected for dilution and 17 the measured particle density, assuming homogeneous spherical particles. The SOA density of 1.2 g m⁻³ was estimated from the mobility mode and aerodynamic mode obtained from the SMPS 18 19 and AMS measurements respectively, as described by DeCarlo et al. (2004) and Katrib et al. 20 (2005). No significant change in the SOA density was found during the experiment. Furthermore, 21 the retrieved value is in agreement with previous laboratory studies (Shilling et al., 2008; Saathoff et al., 2009). Table 1 shows that the SOA mass concentration of SOA varied from 44 22 to 139 μ g m⁻³, resulting in calculated yields values ranging between 0.07 and 0.21. The 23 24 comparison between the steady-state aerosol yields determined in this study and the values 25 reported in the literature is available in the supplementary material (Figure S2).

26

27 **3.2.** Complex refractive index

The real and imaginary parts of the CRI were retrieved to describe the scattering and absorbingcharacteristics of SOA.

1 The imaginary part of the CRI was estimated from the attenuation measurements obtained with 2 the aethalometer. Since the attenuation coefficient depends on the quantity of particles accumulated on the filter, the CRI could not be calculated during the SOA formation, while the 3 4 mass concentration increased rapidly until the α -pinene was completely consumed. Figure S3 5 (Supplementary Material) shows that the absorption coefficient in the 370-950 nm wavelength range estimated after 14 hours of reaction is close to or below zero, indicating that the SOA do 6 7 not absorb radiation in the visible to near-UV region. Therefore, the imaginary part of the CRI 8 was set to zero. This result is consistent with previous studies on the absorbing properties of α -9 pinene-O₃ SOA. Cappa et al. (2008) observed no significant absorption of α -pinene-O₃ SOA at 10 532 nm with a photoacoustic spectrometer. Schnaiter et al. (2003) and Nakayama et al. (2010) 11 measured no significant difference between scattering and extinction coefficient in the visible spectrum. Nakayama et al. (2012) showed that the imaginary part of the CRI values was 12 13 negligible (< 0.003) in the 405-781 nm wavelenghts. Recently, Liu et al. (2013) reported also an imaginary part of the CRI below 10⁻⁴ in the 355-781 nm wavelenghts range. 14

15

Figure 3a shows the real part of the retrieved CRI at 525 nm as a function of reaction time. Real 16 CRI were calculated when SOA exhibited a diameter high enough to allow for the detection of 17 the full size distribution by the SMPS. Based on the measurement uncertainties of $\pm 10\%$ for σ_{scatt} , 18 19 we estimate an absolute error associated with the real CRI of ± 0.02 . The real CRI decreased from 20 1.43-1.60 at the beginning of the reaction to 1.32-1.38 after 14 hours of reaction. In contrast, no 21 significant changes in the CRI were observed during the first hour of reaction for experiments 22 E050210 and E080210. As shown in Figure 3b, the aerosol mass concentrations in these experiments were stabilised after 20 minutes of reaction, which was significantly faster than in 23 24 the other experiments. This may result from differences in the [VOC]/[O₃] concentrations (see 25 discussion in section 3.1 on the initial concentrations of α -pinene), leading to different reaction 26 kinetics (Chan et al., 2007). The real CRI appears to change significantly in most of the 27 experiments between 10 minutes and 1 hour of reaction and it is possible that these variations 28 were not detected in experiments E050210 and E080210. Values for the real part of the CRI 29 retrieved in this study at 525 nm are compared with the literature values obtained for SOA 30 produced during α -pinene + O₃ SOA formation (Table 2). Because our experiments were performed under various time scales and under various conditions and because our CRI values 31

1 evolved with SOA aging, we can globally say that our values (1.32-1.60) were in the same range

- 2 as the previous ones.
- 3

4

3.3. Hygroscopic properties

5 The first insight in the hygroscopic behaviour of SOA was brought by measuring humidograms 6 of SOA with the H-TDMA. Particles of 115 nm diameter were selected for the analysis after 2 7 hours aging and 190 nm after 14 hours aging. The GF uncertainty was estimated from the 8 uncertainty in retrieving the geometric diameters and the RH uncertainty was based on the 9 weighted average of the RH sensors uncertainties at the entrance of the H-TDMA. The GF values at 90% RH of SOA are reported in Figure 4 as a function of time. GF ranged from 1.02 (±0.02) 10 to 1.07 (±0.02) at 90±4.2% RH during and after SOA formation, in agreement with values 11 12 previously reported in the literature (Table 3).

13

14 The comparison of the mobility diameter as a function of RH after 2 and 14 hours of SOA aging 15 is shown in figure 5a. Unusual behaviour of the mobility diameter and f(RH) at the end of the 16 experiment (at t=16-20h) is evident. Mobility diameters dropped to a minimum between 20-50% 17 RH for SOA after 14 hours reaction that was not observed for "fresh" SOA. Figure 5b shows the 18 RH dependence of f(RH) of the polydisperse size distribution as measured directly in the 19 chamber during water injection after 1 hour and 16 hours of the reaction. Two RH scales are 20 used: one for the measurements performed within the chamber and the other for the 21 measurements performed using the nephelometer. In fact, a small drying of 10% has been observed between the chamber and the nephelometer. No experiment with nephelometer 22 23 measurements at the beginning and the end of the same date was available. However, the 24 experiments were performed under very similar initial conditions as shown by the reproducible 25 values of CRI and GF that are also comparable with each other. There is a first increase of f(RH) 26 as RH increases from 0 to 30%.. This can be due to a change in the physical state of SOA. In 27 fact, particles with irregular shapes or porous should be observed at higher mobility diameters in 28 the DMA than spherical and compact particles of the same mass (De Carlo et al., 2004). After 29 water uptake, the particles may become more spherical or compact, leading to a shift of the 30 mobile size distribution measured with the DMA (Milkhailov et al., 2009). Particles are expected 31 to exhibit also a higher mass concentration, and the σ_{scatt} should increase after humidification 32 (Adachi et al., 2011). This hypothesis is discussed in details in section 4.2. A second slope is

observed between 80 and 90% RH, linked to the sharp increase of GF due to particle water
absorption. All the experiments exhibited the same trend, but different values are observed at
90% RH. This can be explained by the different size distribution from one experiment to another.
In particular, the proportion of particles larger than 100 nm is different resulting in a different
capacity in absorbing water (Biskos *et al.*, 2006) and hence varying the observed f(RH).

6

An important concern in measuring the hygroscopic properties of the particles is to allow 7 8 sufficient time for particle-water vapor equilibrium. Various studies have discussed the 9 possibility that insufficient time for humidification could result in an underestimation of the particles' water content (Chan and Chan, 2005; Saxena et al., 1995; Duplissy et al., 2009; 10 11 Denjean et al., 2014). In this study, the residence time for SOA particles in the wet air stream 12 was significantly longer when particles were humidified in-situ in the chamber. It took ~1 hour in the chamber, instead of 15 seconds in the H-TDMA. These two approaches can thus be 13 14 complementary to carry information on water transfer dynamics of α -pinene-O₃ SOA. We used 15 Mie scattering calculations for homogeneous spheres to determine GF from f(RH). σ_{scat} was 16 calculated for different GF at specific RH. The optimal GF as a function of RH was determined so that the difference between measured σ_{scat} and those obtained using Mie calculations were 17 18 minimized. Particles were assumed to be homogeneous spheres of uniform CRI. The CRI 19 calculations were based on volume weighted refractive indices of SOA and water. Uncertainties 20 on the theoretical GF were estimated from the standard deviation of the measured f(RH), the 21 uncertainties on the f(RH) measurements and the uncertainties on the RH measurements. Figure 22 6 shows the comparison between measured and predicted GF values for SOA at two different reaction times: for "fresh" SOA (after 1 hour of reaction), and for "aged" SOA (after 16 hours of 23 24 reaction). For both reaction times, the model approach agrees well with the measurements above 25 30% RH. It indicates no kinetic limitations of α -pinene-O₃ SOA for water uptake. For "aged" 26 SOA (after 16 hours of reaction), the underestimation of the model below 30 % RH (Figure 6) might be due to a change in the physical state of SOA, as discussed in detail in section 4.2. 27

28

29 **3.4.** Chemical composition

The O:C of the bulk aerosol measured online by the AMS and the O:C at the surface obtained by XPS analysis on filter samples are shown in Figure 7. The bulk O:C ranged between 0.68 (\pm 0.20) to 0.55 (\pm 0.16). The uncertainties in bulk O:C given by Aiken et al. (2007) may be overestimated 1 compared to the experimental variability and even experimental reproducibility observed in this 2 study. In fact, we estimated the experimental uncertainties to be ± 0.01 from the standard 3 deviation of the experimental values.

4

5 Two potential processes affecting the bulk O:C during the SOA formation are the gas/particle phase partitioning and chemical reactions in the condensed phase. The evolution of the AMS 6 7 mass spectra during the reaction is shown in Figure S4 (Supplementary Material). The AMS 8 mass spectra were dominated by m/z 44 for 'fresh' SOA, while the strongest signal was observed 9 at m/z 29 for 'aged' SOA. Figure S5 also shows that f_{44} decreased while f_{43} increased with time. These observations indicate an increase of less oxidized semi-volatile compounds in the bulk 10 11 particle phase with aging. This trend can be explained by an increasing partitioning of less 12 oxidised semi-volatile compounds as the aerosol grows according to the structure-activity 13 correlation between the oxygenated functional groups and the species vapour pressure (Pankow 14 and Asher, 2008). Additionally, some oligomerisation/dehydration, such as aldol condensation 15 or ester formation, can alter the SOA O:C and f_{44} (Camredon et al., 2010; Ziemann and Atkinson, 2012). Additional reactions in the particle phase can take place after the SOA formation, and can 16 for example lead to the formation of oligomers (Kalberer et al., 2004; Gao et al., 2004b; Tolocka 17 et al., 2004; Gao et al., 2004a; Yasmeen et al., 2012; Reinhardt et al., 2007; Rudich et al., 2007), 18 19 but do not appear to be associated with a significant change in the SOA functionality.

20

21 Bulk O:C obtained in the literature from different techniques are shown in Table 4. The bulk O:C 22 retrieved in our study are in agreement with Shilling et al. (2008), Chhabra et al. (2010), 23 Reinhardt et al. (2007) and Tolocka et al. (2006). Values at the beginning of the reaction are, 24 however, higher than those obtained by Aiken et al. (2007) and Oi et al. (2012) who used AMS 25 to retrieve the O:C. Limited information is given by Aiken et al. (2007) about their experimental 26 conditions. In particular, temperature and reaction time scales are critical parameters which can 27 significantly affect the chemical composition of SOA (Warren et al., 2009; Shilling et al., 2008; 28 Chhabra et al., 2010). Qi et al. (2012) did not provide any uncertainty in their AMS 29 measurements, but taking into account the uncertainties of $\pm 30\%$ given by Aiken et al. (2007), 30 their O:Cs are in agreement with our values.

1 The O:C at the surface of the SOA is significantly lower than in the bulk (Figure 7) and ranged 2 between 0.33 (± 0.07) and 0.46 (± 0.08). Some volatile compounds could evaporate during the 3 XPS analysis due to the low pressure in the instrument. In our companion paper (Denjean et al., 4 2015), an increase of the O:C has been observed after heating by the evaporation of semi-volatile 5 components of SOA. Therefore, the O:C at the surface of the SOA was certainly overestimated in the present study. This suggests that the SOA composition is not homogeneous, but composed 6 7 of less oxidised species at its surface. To our knowledge, our study is the first one to investigate 8 the surface O:C of the SOA and thus no comparison with literature is possible. Nevertheless, this 9 observation is in very good agreement with one of the hypotheses by McIntire et al. (2010). The 10 authors proposed a hydrophobic shell model for hydrocarbon + O₃ SOA due to the burying of R-11 COOH compounds and other polar groups inside the particle. Here, we report direct experimental 12 measurements which support this hypothesis.

13

We examined the sensitivity of the SOA chemical composition on the surface O:C by calculating the O:C of the core. For simplicity, we assumed that the O:C of the core SOA is constant and independent on the distance from the centre of the particle. O:C of the core can be expressed as a function of the O:C of the bulk from AMS measurements, the O:C at the surface from XPS and the mass fraction of surface and core in the particle:

$$(0:C)_{core} = \frac{(0:C)_{bulk} - \frac{m_{surf}}{m_{bulk}} (0:C)_{surf}}{\frac{m_{core}}{m_{bulk}}}$$
(6)

19 XPS quantifies the O:C of the SOA surface to a depth less than 10 nm. However, this technique 20 cannot provide the precise thickness of the surface analyzed and it is possible that the O:C 21 corresponds to a thinner layer. Therefore, O:C of the core SOA was calculated for different 22 possible thicknesses of the analyzed surface O:C. Figure 8 shows the O:C of the core for fresh 23 and aged SOA. From these calculations, we observed that bulk O:C ranges from 0.68 (± 0.20) to 1.01 (± 0.40) for fresh SOA and from 0.55 (± 0.17) to 0.58 (± 0.23) for aged SOA. Based on the 24 25 elemental composition of identified compounds of α -pinene-O₃ SOA, highly oxygenated 26 compounds were found in the literature. For example, peroxypinic acid is a product of α -pinene 27 ozonolysis proposed by Docherty et al. (2005) with O:C=0.56. This simple calculation highlights 28 the importance of monitoring the surface chemical composition separately from the bulk.

2 4. Discussion

3 **4.1.** Dependence of the CRI and GF on the chemical composition

4 Our results suggest that the real part of the CRI is closely related to SOA chemical composition. Figure 9 shows that the real part of the CRI decreases substantially as the O:C of the bulk SOA 5 6 decreases. This trend is in agreement with the positive correlation between the extinction 7 coefficient at 532 nm wavelength and O:C observed by Cappa et al. (2011) for the heterogeneous 8 OH oxidation of squalane and azelaic acid particles. Nakayama et al. (2013) reported increasing 9 real part of the CRI at 532 nm with increasing O:C for SOA generated from the photooxidation 10 of toluene. Recently, Flores et al. (2014) showed that the real part of the CRI at 405 nm is positively correlated to the O:C for a mixture of biogenic VOC (α -pinene and limonene) and 11 biogenic VOC mixture with subsequent addition of an anthropogenic VOC (p-xylene-d₁₀). On 12 13 the other hand, an opposite trend was observed by Nakayama et al. (2012) with a decrease of the 14 real part of the CRI at 532 nm wavelength with increasing O:C for SOA produced from the 15 ozonolysis and photooxidation of α -pinene. Lambe et al. (2013) studied SOA formed by OHoxidation of various gas-phase precursors (naphthalene, tricyclo[5.2.1.0^{2,6}]decane, guaiacol and 16 17 α -pinene) and retrieved a decreasing real part of the CRI with increasing oxidation. These 18 different trends suggest that the change of the real CRI as a function of O:C depends strongly on 19 the organic aerosol source and aging in the atmosphere.

20

21 The decrease of bulk O:C is expected to induce a decrease in SOA hygroscopic properties 22 (Massoli et al., 2010; Jimenez et al., 2009; Chang et al., 2010; Duplissy et al., 2011). However, 23 the GF remained surprisingly constant over 20 hours of reaction (Figure 4). This trend contradicts 24 with Saathoff et al. (2003) and Cocker et al. (2001) who reported an increase of the GF from 25 1.080 to 1.106 and 1.065 to 1.1 respectively within time scale of 6 hours. However, our constant 26 GF is consistent with the constant hygroscopicity observed by Warren et al. (2009) and Qi (2010) 27 for α -pinene-O₃ SOA. The constant GF observed in our study may result from: 1) the sensitivity 28 of the instruments. The GF was studied after 2 hours of reaction when the decrease of the 29 oxidation degree was slow. The O:C of bulk SOA decreased from 0.65 to 0.60 between 2 and 10 30 hours aging (Figure 7), that should result in a GF decrease of 0.03 according to the linear GF-to-31 O:C relationship reported by Massoli et al. (2010). This expected GF variation (due to the

1 chemical composition change) is smaller than the sensitivity of the H-TDMA. 2) Other factors 2 than the bulk O:C can control the water uptake of SOA: recently, Alfarra et al. (2014) reported a 3 positive correlation between hygroscopicity of particles and their degree of oxidation for SOA 4 produced from the photooxidation of α -pinene, β -caryophyllene, linalool and myrcene, but not 5 for limonene SOA. They suggested that other factors such as solubility, surface tension, molecular weight, density and particle phase are likely to be playing important roles in 6 7 controlling GF values. We observed that the O:C of the particle surface was lower than the bulk 8 O:C and remained constant over time (see section 3.4.1.). This structure would affect the 9 heterogeneous chemistry of the particle and avoid the water absorption on particles. Broekhuizen 10 et al. (2004) observed that the oxidation of oleic acid by low concentration of ozone, shown to 11 produce oxygenated products (Moise and Rudich, 2002), did not increase the particle CCN 12 activity. Katrib et al. (2004) demonstrated that hydrophobic products were formed at the surface 13 of oleic acid particles after ozone exposure that prevented from water adsorption on particles. 14 McIntire et al. (2010) exposed particles formed from ozonolysis of surface-bound alkenes to 15 ozone and observed formation of polar groups buried inside a hydrophobic shell. This is also 16 consistent with Moussa et al. (2009) who observed that the uptake of water does not increase 17 after the oxidation of surface-bound alkenes. A core-shell structure could explain the low O:C 18 of the surface α -pinene-O₃ SOA obtained in the present study and could affect significantly the 19 heterogeneous chemistry of the particle.

20

21 **4.2.** Effect of RH and aging on the viscosity of SOA

22 An unexpected effect of aging on the dynamic of the size distribution evolution during 23 humidification has been observed (section 3.3 and Figure 5). For the 16-hour old SOA, humidification led to a decay of the mobility diameter and an increase of σ_{scatt} while such 24 behaviour was not observed for the 1-hour old reaction. Although evaporation of semi-volatiles 25 26 in the chamber could lead to a decrease of the observed mobility diameter, it cannot explain the increase of f(RH) below 40% RH. Furthermore, the observed decrease of the mobility diameter 27 28 cannot be attributed to losses of particles to the walls of the chamber, since it would lead to a decrease of f(RH). 29

30

As it is difficult to believe that a RH increase would induce a sudden loss of matter for the particulate phase, it is hypothesised that the particles experienced a rapid change of shape when

1 humidified. Being observed for aged SOA only, these behaviours support a re-shaping of 2 coagulated particles and an increase in sphericity for RH > 20-30%. This implies that, below these RH values, coagulated particles were not spherical and hence they were not liquid enough for 3 4 coalescence. Electron microscopy analysis was used to investigate the shape of SOA particles. 5 SOA particles after 1 hour and 14 hours of reaction were analysed by transmission electron microscopy (TEM) (Figure S6 in the Supplementary Material). Experimental details are given in 6 7 the Supplementary Material. In total 50 particles were analysed by TEM. Figure S6 shows an 8 example of SOA particles after 1 and 14 hours of reaction. Only spherical particles have been 9 observed in the samples. This indicates that the coagulation taking place at the beginning and 10 after 14 hours of reaction resulted in spherical coalesced particles.

11

12 A few recent studies have suggested that α -pinene-O₃ SOA is likely to be in an amorphous semi-13 solid or in an amorphous solid (glassy) state under dry conditions (RH < 30%). Virtanen et al. 14 (2011) reported that α -pinene-O₃ SOA bounces off impactor plates as if particles were glassy. 15 Other studies reported that the evaporation kinetics of α -pinene-O₃ SOA was lower than expected by models for liquid droplets, indicating a nonliquid-like state (Cappa and Wilson, 2011, Vaden 16 17 et al., 2011). These findings are in agreement with Perraud et al. (2012) who has shown that the 18 uptake of organic nitrates by α -pinene-O₃ SOA did not follow absorptive equilibrium partitioning 19 theory, as indicative of non-liquid-like behaviour. Moreover, some recent studies reported 20 viscosity changes of α -pinene-O₃ SOA with RH. Saukko et al. (2012) observed lower bounce 21 behaviour after the SOA was exposed to RH > 50%, indicating a liquid-like, less viscous phase. 22 Renbaum-Wolff et al. (2013) reported that the viscosities of α -pinene-O₃ SOA correspond to a 23 semisolid or solid for RH \leq 30%, a semisolid for 40% \leq RH < 80% and a liquid for RH \geq 80%. 24 This is consistent with our observation of the apparent step in mobility diameter and σ_{scatt} changes 25 observed between 0 and 30% RH for SOA after 16 hours of reaction (Figure 5), which may be 26 due to phase transition from a predominantly glassy state to a predominantly liquid state.

27

In addition, we find no apparent transition step between 0 and 30% RH for SOA after 1 hour of reaction. This observation mirrors the effect of aging on the physical state of SOA. We discussed previously the possible formation of oligomers in the particle phase during the formation of SOA. Roth et al. (2005) suggested that ambient organic particles could be in an extremely high viscosity, glassy state due to the presence of oligomeric constituents. There is evidence from 1 different organic materials that oligomerisation may lead to an increase in viscosity and 2 potentially to the formation of glassy states (Koop et al., 2011). Abramson et al. (2013) reported 3 decreasing evaporation kinetics during the aging of α -pinene-O₃ SOA, indicating that hardening 4 occurs with time. Our observations are consistent with an increase in the SOA oligomer content 5 with aging, leading to the transformation of SOA into glassy states under dry conditions.

6

7 These findings are of major importance as it suggests that particles may undergo reorganisation 8 after the condensation of oxidised species. It is also supported by the core-shell structure of SOA 9 observed with less oxidised species at its surface (section 3.4.1. and Figure 7). Due to possible 10 kinetic limitations, the particles viscosity can strongly affect a number of key physical and 11 chemical properties, such as water uptake, equilibrium partitioning between the gas and the 12 particle phase or heterogeneous chemistry (Koop et al., 2011; Zobrist et al., 2008, Milkhailov et 13 al., 2009, Murray 2008; Shiraiwa et al., 2013) and thus has considerable implications on the 14 understanding of the impact of SOA on climate.

15

16 **5.** Implication for direct radiative effect and conclusions

17 The imaginary part of the CRI values for α -pinene-O₃ SOA was found to be negligible in the 18 visible spectrum, indicating that these particles have a pure scattering effect. The single scattering 19 albedo ω_0 , a key parameter to estimate the influence of aerosols on the radiative balance, is 20 estimated to be equal to 1, indicating that these particles have a negative radiative effect.

21

22 In order to estimate the evolution of the direct radiative effect of SOA, the mass extinction 23 efficiency kext (dry) at 525 nm of dry SOA is shown in Figure 10. The kext values were calculated 24 as the ratio of the measured SOA σ_{scat} to their mass concentration. The k_{ext} values depend on the 25 competitive effect of aerosol size and chemical composition. The decrease of the real CRI 26 resulting from the changing particle chemical composition is expected to decrease kext while the size increase of the SOA from the nucleation mode to the accumulation mode should increase 27 k_{ext} . At the beginning of the SOA formation, we obtained k_{ext} (dry) = 0.64 ±0.27 m² g⁻¹, that is 28 significantly lower than obtained after 14 hours of reaction where $k_{ext}(dry) = 1.68 \pm 0.50 \text{ m}^2 \text{ g}^{-1}$. 29 30 The mass extinction efficiency k_{ext} is also strongly affected by the ambient relative humidity 31 (Figure 10). α-pinene-O₃ SOA did not show deliquescence below 90% RH, but a continuous 1 water uptake with increasing RH. A constant GF (90% RH) was obtained throughout the 2 experiment. From the measured f(90% RH) and GF(90% RH) obtained in this study, we can 3 estimate σ_{scat} and SOA mass concentration at 90% RH and then k_{ext} (90% RH) = 2.36 ±0.70 m² 4 g⁻¹. Water uptake by SOA at 90% RH produces an increase of k_{ext} by 40%. This simple calculation 5 highlights the importance of taking into account the actual ambient conditions that the α -pinene-6 O₃ SOA experiences during its lifetime when estimating its induced direct radiative effect in the 7 atmosphere.

8

9 For comparison, k_{ext} of organic aerosols calculated using the CRI and GF as prescribed in global 10 models are shown in Figure 10. Mie scattering calculations for homogeneous spheres were performed to calculate k_{ext}. Number size distribution representing biogenic aerosol particles in 11 12 the Amazon Basin (Martin et al., 2010) was used as an input in the Mie scattering calculations. 13 For wet conditions, the CRI calculations were based on volume-weighted CRI of values of α-14 pinene-O₃ SOA and water. k_{ext} of SOA generated from a mixture of biogenics (mostly terpenes) 15 released directly from plants (Lang-Yona et al, 2010) is also shown in Figure 10. Despite the 16 wide range of variability of the CRI values assumed in global models for organic aerosols (1.45-17 0i to 1.6-0.003i), the values for k_{ext} were very similar for all models. The k_{ext} values retrieved in the present study for SOA after 14 hours of reaction agree with the values used for global models, 18 19 while k_{ext} obtained for SOA at the beginning of the reaction are 2-7 times smaller. This work is 20 a first attempt to assess the evolution of k_{ext} of SOA during its formation and aging. Our results 21 suggest that the k_{ext} of atmospheric SOA is not static and a single size distribution and CRI does 22 not appear sufficient to accurately model its direct radiative effect.

23

24 Acknowledgements

This research work has been supported by the European Community within the seventh Framework Programme: Eurochamp-2 (EU-FP7 grant agreement n°228335). We also acknowledge the National Research Agency ANR (project Cumulus, n° ANR-10-BLAN-0617). We thank Frank Siekmann and Sylvain Ravier (Aix-Marseille Université, France) for their help in calibrating the AMS during the measurement campaign. We thank Michel Maille (LISA, France) for the analysis with electron microscopy. We gratefully acknowledge the MASSALYA instrumental platform (Aix Marseille Université, Ice.univ-amu.fr) for the analysis and

- 1 measurements used in this publication. We thank Brendan Mahon and Steven J. Campbell
- 2 (University of Cambridge, UK) for helpful revision of the manuscript.

1 References

- Abramson, E., Imre, D., Beranek, J., Wilson, J., and Zelenyuk, A.: Experimental determination
 of chemical diffusion within secondary organic aerosol particles, Physical Chemistry Chemical
- 4 Physics, 15, 2983-2991, 10.1039/C2CP44013J, 2013.
- 5 Adachi, K., Freney, E. J., and Buseck, P. R.: Shapes of internally mixed hygroscopic aerosol 6 particles after deliquescence, and their effect on light scattering, Geophys. Res. Lett., 38, L13804,
- 7 2011
- 8 Aiken, A. C., DeCarlo, P. F., and Jimenez, J. L.: Elemental Analysis of Organic Species with
- 9 Electron Ionization High-Resolution Mass Spectrometry, Analytical Chemistry, 79, 8350-8358,
 10 2007.
- 11 Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S.,
- 12 Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway,
- 13 M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen,
- 14 J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC Ratios of
- 15 Primary, Secondary, and Ambient Organic Aerosols with High-Resolution Time-of-Flight
- Aerosol Mass Spectrometry, Environmental science & technology, 42, 4478-4485,
 10.1021/es703009g, 2008.
- 18 Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A.
- 19 M., Drewnick, F., Onasch, T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A 20 generalised method for the extraction of chemically resolved mass spectra from Aerodyne
- 21 aerosol mass spectrometer data, Journal of Aerosol Science, 35, 909-922, 22 10.1016/j.jacuarej.2004.02.007.2004
- 22 10.1016/j.jaerosci.2004.02.007, 2004.
- Andreae, M. O.: A New Look at Aging Aerosols, Science, 326, 1493-1494,
 10.1126/science.1183158, 2009.
- Arnott, W. P., Hamasha, K., Moosmuller, H., Sheridan, P. J., and Ogren, J. A.: Towards aerosol
 light-absorption measurements with a 7-wavelength Aethalometer: Evaluation with a
 photoacoustic instrument and 3-wavelength nephelometer, Aerosol science and technology, 39,
 17-29, 10.1080/027868290901972, 2005.
- Baltensperger, U.: Secondary organic aerosols from anthropogenic and biogenic precursors, The
 Royal Society of Chemistry, 265-278, 2005.
- Biskos, G., Russell, L. M., Buseck, P. R., and Martin, S. T.: Nanosize effect on the hygroscopic
 growth factor of aerosol particles, Geophys. Res. Lett., 33, L07801, 2006.
- Bohren, C. F., and Huffman, D. R.: Absorption and scattering of light by small particles, Wiley,
 New York, 1983.
- 35 Bond, T. C., Anderson, T. L., and Campbell, D.: Calibration and intercomparison of filter-based
- measurements of visible light absorption by aerosols, Aerosol science and technology, 30, 582 600, 10.1080/027868299304435, 1999.
- Bond, T. C., and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative
 review, Aerosol science and technology, 40, 27-67, 10.1080/02786820500421521, 2006.
- 40 Broekhuizen, K. E., Thornberry, T., Kumar, P. P., and Abbatt, J. P. D.: Formation of cloud
- 41 condensation nuclei by oxidative processing: Unsaturated fatty acids, J. Geophys. Res.-Atmos.,
- 42 109, 10.1029/2004jd005298, 2004.

- 1 Camredon, M., Hamilton, J. F., Alam, M. S., Wyche, K. P., Carr, T., White, I. R., Monks, P. S.,
- 2 Rickard, A. R., and Bloss, W. J.: Distribution of gaseous and particulate organic composition
- 3 during dark alpha-pinene ozonolysis, Atmos. Chem. Phys., 10, 2893-2917, 10.5194/acp-10-
- 4 2893-2010, 2010.
- 5 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch,
- 6 T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M.,
- 7 Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and
- 8 microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer,
- 9 Mass Spectrometry Reviews, 26, 185-222, 10.1002/mas.20115, 2007.
- 10 Cappa, C. D., Lack, D. A., Burkholder, J. B., and Ravishankara, A. R.: Bias in Filter-Based
- 11 Aerosol Light Absorption Measurements Due to Organic Aerosol Loading: Evidence from
- 12 Laboratory Measurements, Aerosol Science and Technology, 42, 1022 1032, 2008.
- 13 Cappa, C. D., and Wilson, K. R.: Evolution of organic aerosol mass spectra upon heating:
- 14 implications for OA phase and partitioning behavior, Atmos. Chem. Phys., 11, 1895-1911,
- 15 10.5194/acp-11-1895-2011, 2011.
- 16 Cappa, C. D., Che, D. L., Kessler, S. H., Kroll, J. H., and Wilson, K. R.: Variations in organic
- aerosol optical and hygroscopic properties upon heterogeneous OH oxidation, J. Geophys. Res.,
 116, D15204, 2011.
- 19 Carrico, C. M., Kreidenweis, S. M., Malm, W. C., Day, D. E., Lee, T., Carrillo, J., McMeeking,
- 20 G. R., and Collett Jr., J. L.: Hygroscopic growth behavior of a carbon-dominated aerosol in
- 21 Yosemite National Park, Atmos. Environ., 39, 1393-1404, 2005.
- Chan, M. N., and Chan, C. K.: Mass transfer effects in hygroscopic measurements of aerosol
 particles, Atmos. Chem. Phys., 5, 2703-2712, 10.5194/acp-5-2703-2005, 2005.
- 24 Chan, A. W. H., Kroll, J. H., Ng, N. L., and Seinfeld, J. H.: Kinetic modeling of secondary 25 organic aerosol formation: effects of particle- and gas-phase reactions of semivolatile products,
- 26 Atmos. Chem. Phys., 7, 4135-4147, 2007.
- 27 Chang, R. Y. W., Slowik, J. G., Shantz, N. C., Vlasenko, A., Liggio, J., Sjostedt, S. J., Leaitch,
- 28 W. R., and Abbatt, J. P. D.: The hygroscopicity parameter (κ) of ambient organic aerosol at a
- 29 field site subject to biogenic and anthropogenic influences: relationship to degree of aerosol
- 30 oxidation, Atmos. Chem. Phys., 10, 5047-5064, 10.5194/acp-10-5047-2010, 2010.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Hansen, J. E., and
 Hofmann, D. J.: Climate forcing by anthropogenic aerosols, Science, 255, 423-430,
 10.1126/science.255.5043.423, 1992.
- 34 Chen, Q., Liu, Y. J., Donahue, N. M., Shilling, J. E., and Martin, S. T.: Particle-Phase Chemistry
- 35 of Secondary Organic Material: Modeled Compared to Measured O:C and H:C Elemental Ratios
- 36 Provide Constraints, Environmental science & technology, 45, 4763-4770, 10.1021/es104398s,
- 37 2011.
 - Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.: Elemental analysis of chamber organic aerosol
 using an aerodyne high-resolution aerosol mass spectrometer, Atmos. Chem. Phys., 10, 4111-
 - 40 4131, 10.5194/acp-10-4111-2010, 2010.
 - Chung, S. H., and Seinfeld, J. H.: Global distribution and climate forcing of carbonaceous
 aerosols, J. Geophys. Res., 107, 4407, 2002.
 - 43 Cocker, D. R., Mader, B. T., Kalberer, M., Flagan, R. C., and Seinfeld, J. H.: The effect of water 44 on gas-particle partitioning of secondary organic aerosol: II. m-xylene and 1,3,5-

- 1 trimethylbenzene photooxidation systems, Atmos. Environ., 35, 6073-6085, 10.1016/s1352-2 2310(01)00405-8, 2001.
- 3 Collaud Coen, M., Weingartner, E., Apituley, A., Ceburnis, D., Fierz-Schmidhauser, R., Flentje,
- H., Henzing, J. S., Jennings, S. G., Moerman, M., Petzold, A., Schmid, O., and Baltensperger,
 U.: Minimizing light absorption measurement artifacts of the Aethalometer: evaluation of five
- 6 correction algorithms, Atmos. Meas. Tech., 3, 457-474, 2010.
- 7 de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M.,
- 8 Fehsenfeld, F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C.,
- 9 Marchewka, M., Bertman, S. B., and Bates, T. S.: Budget of organic carbon in a polluted
- 10 atmosphere: Results from the New England Air Quality Study in 2002, J. Geophys. Res.-Atmos.,
- 11 110, 10.1029/2004jd005623, 2005.
- 12 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin,
- 13 M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-
- deployable, high-resolution, time-of-flight aerosol mass spectrometer, Analytical Chemistry, 78,
 8281-8289, 10.1021/ac061249n, 2006.
- 16 Denjean, C., Formenti, P., Picquet-Varrault, B., Katrib, Y., Pangui, E., Zapf, P., and Doussin, J.
- 17 F.: A new experimental approach to study the hygroscopic and optical properties of aerosols:
- application to ammonium sulfate particles, Atmos. Meas. Tech., 7, 183-197, 10.5194/amt-7-183-
- 19 2014, 2014.
- 20 Denjean, C., Formenti, P., Picquet-Varrault, B., Camredon, M., Pangui, E., Zapf, P., Katrib, Y.,
- 21 Giorio, C., Tapparo, A., Temime-Roussel, B., Monod, A., Aumont, B., and Doussin, J. F.: Aging
- 22 of secondary organic aerosol generated from the ozonolysis of α -pinene: effects of ozone, light
- 23 and temperature, Atmos. Chem. Phys., 15, 883-897, 10.5194/acp-15-883-2015, 2015.
- 24 Dinar, E., Riziq, A. A., Spindler, C., Erlick, C., Kiss, G., and Rudich, Y.: The complex refractive
- 25 index of atmospheric and model humic-like substances (HULIS) retrieved by a cavity ring down
- aerosol spectrometer (CRD-AS), Faraday Discussions, 137, 279-295, 10.1039/b703111d, 2008.
- Docherty, K. S., Wu, W., Lim, Y. B., and Ziemann, P. J.: Contributions of organic peroxides to
 secondary aerosol formed from reactions of monoterpenes with O-3, Environmental science &
 technology, 39, 4049-4059, 10.1021/es050228s, 2005.
- Donahue, N. M., Henry, K. M., Mentel, T. F., Kiendler-Scharr, A., Spindler, C., Bohn, B.,
 Brauers, T., Dorn, H. P., Fuchs, H., Tillmann, R., Wahner, A., Saathoff, H., Naumann, K. H.,
 Mohler, O., Leisner, T., Muller, L., Reinnig, M. C., Hoffmann, T., Salo, K., Hallquist, M.,
- 33 Frosch, M., Bilde, M., Tritscher, T., Barmet, P., Praplan, A. P., DeCarlo, P. F., Dommen, J.,
- 34 Prevot, A. S. H., and Baltensperger, U.: Aging of biogenic secondary organic aerosol via gas-
- 35 phase OH radical reactions, Proceedings of the National Academy of Sciences of the United
- 36 States of America, 109, 13503-13508, 10.1073/pnas.1115186109, 2012.
- 37 Duplissy, J., DeCarlo, P. F., Dommen, J., Alfarra, M. R., Metzger, A., Barmpadimos, I., Prevot,
- A. S. H., Weingartner, E., Tritscher, T., Gysel, M., Aiken, A. C., Jimenez, J. L., Canagaratna, M.
- 39 R., Worsnop, D. R., Collins, D. R., Tomlinson, J., and Baltensperger, U.: Relating hygroscopicity
- 40 and composition of organic aerosol particulate matter, Atmos. Chem. Phys., 11, 1155-1165,
- 41 10.5194/acp-11-1155-2011, 2011.
- Dusek, U., Reischl, G. P., and Hitzenberger, R.: CCN Activation of Pure and Coated Carbon
 Black Particles, Environmental Science & Technology, 40, 1223-1230, 2006.

- Flores, J. M., Zhao, D. F., Segev, L., Schlag, P., Kiendler-Scharr, A., Fuchs, H., Watne, Å. K., 1
- Bluvshtein, N., Mentel, T. F., Hallquist, M., and Rudich, Y.: Evolution of the complex refractive 2
- index in the near UV spectral region in aging secondary organic aerosol, Atmos. Chem. Phys. 3
- 4 Discuss., 14, 4149-4187, 10.5194/acpd-14-4149-2014, 2014.
- 5 Gao, S., Keywood, M., Ng, N. L., Surratt, J., Varutbangkul, V., Bahreini, R., Flagan, R. C., and
- 6 Seinfeld, J. H.: Low-molecular-weight and oligomeric components in secondary organic aerosol
- 7 from the ozonolysis of cycloalkenes and alpha-pinene, Journal of Physical Chemistry A, 108,
- 8 10147-10164, 10.1021/jp047466e, 2004a.
- 9 Gao, S., Ng, N. L., Keywood, M., Varutbangkul, V., Bahreini, R., Nenes, A., He, J., Yoo, K. Y.,
- 10 Beauchamp, J. L., Hodyss, R. P., Flagan, R. C., and Seinfeld, J. H.: Particle Phase Acidity and
- 11 Oligomer Formation in Secondary Organic Aerosol, Environmental Science & Technology, 38,
- 6582-6589, 2004b. 12
- 13 George, I. J., and Abbatt, J. P. D.: Chemical evolution of secondary organic aerosol from OHinitiated heterogeneous oxidation, Atmos. Chem. Phys., 10, 5551-5563, 10.5194/acp-10-5551-14
- 15 2010, 2010.
- 16 Goldstein, A. H., and Galbally, I. E.: Known and unexplored organic constituents in the Earth's 17 atmosphere, Geochimica Et Cosmochimica Acta, 73, A449-A449, 2009.
- 18 Griffin, R. J., Cocker, D. R., III, Seinfeld, J. H., and Dabdub, D.: Estimate of Global Atmospheric
- 19 Organic Aerosol from Oxidation of Biogenic Hydrocarbons, Geophys. Res. Lett., 26, 2721-2724,
- 20 1999.
- 21 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger,
- 22 L., Lerdau, M., Mckav, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Tavlor,
- J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, J. 23
- 24 Geophys. Res., 100, 8873-8892, 1995.
- 25 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen,
- 26 J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T.,
- 27 Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W.,
- McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., 28
- 29 Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic 30 aerosol: current and emerging issues, Atmos. Chem. Phys., 9, 5155-5236, 10.5194/acp-9-5155-
- 31 2009, 2009.
 - Hatch, C. D., Gierlus, K. M., Schuttlefield, J. D., and Grassian, V. H.: Water adsorption and 32
 - 33 cloud condensation nuclei activity of calcite and calcite coated with model humic and fulvic
 - 34 acids, Atmos. Environ., 42, 5672-5684, 10.1016/j.atmosenv.2008.03.005, 2008.
 - 35 Haywood, J. M., and Ramaswamy, V.: Global sensitivity studies of the direct radiative forcing due to anthropogenic sulfate and black carbon aerosols, J. Geophys. Res., 103, 1998. 36
 - Hegg, D., Larson, T., and Yuen, P. F.: A theoretical-study of the effect of relative-humidity on 37 light-scattering by tropospheric aerosols, J. Geophys. Res. Atmos., 98, 18435-18439, 38 39 10.1029/93jd01928, 1993.
 - 40 Henry, K. M., and Donahue, N. M.: Photochemical Aging of alpha-Pinene Secondary Organic
 - Aerosol: Effects of OH Radical Sources and Photolysis, Journal of Physical Chemistry A, 116, 41 5932-5940, 10.1021/jp210288s, 2012. 42
 - 43 Hoyle, C. R., Myhre, G., Berntsen, T. K., and Isaksen, I. S. A.: Anthropogenic influence on SOA 44
 - and the resulting radiative forcing, Atmos. Chem. Phys., 9, 2715-2728, 2009.

- 1 Jang, M., and Kamens, R. M.: A thermodynamic approach for modeling partitioning of
- 2 semivolatile organic compounds on atmospheric particulate matter: Humidity effects,
- 3 Environmental Science & Technology, 32, 1237-1243, 10.1021/es970773w, 1998.
- 4 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H.,
- 5 DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M.,
- 6 Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin,
- 7 C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M.,
- Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E., Dunlea, J., Huffman, J. A.,
 Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick,
- F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A.,
- 11 Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J.
- 12 R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C.,
- 13 Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic
- 14 Aerosols in the Atmosphere, Science, 326, 1525-1529, doi: 10.1126/science.1180353, 2009.
- 15 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van
- 16 Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y.,
- 17 Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati,
- 18 E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review,
- 19 Atmos. Chem. Phys., 5, 1053-1123, 10.5194/acp-5-1053-2005, 2005.
- 20 Kannosto, J., Yli-Pirila, P., Hao, L. Q., Leskinen, J., Jokiniemi, J., Makela, J. M., Joutsensaari,
- 21 J., Laaksonen, A., Worsnop, D. R., Keskinen, J., and Virtanen, A.: Bounce characteristics of
- 22 alpha-pinene-derived SOA particles with implications to physical phase, Boreal Environment
- 23 Research, 18, 329-340, 2013.
- 24 Katrib, Y., Martin, S. T., Rudich, Y., Davidovits, P., Jayne, J. T., and Worsnop, D. R.: Density
- changes of aerosol particles as a result of chemical reaction, Atmos. Chem. Phys., 5, 275-291,
 10.5194/acp-5-275-2005, 2005.
- Kavouras, I. G., Mihalopoulos, N., and Stephanou, E. G.: Formation of atmospheric particles
 from organic acids produced by forests, Nature, 395, 683-686, 1998.
- 29 Kavouras, I. G., Mihalopoulos, N., and Stephanou, E. G.: Formation and gas/particle partitioning
- 30 of monoterpenes photo-oxidation products over forests, Geophys. Res. Lett., 26, 55-58,
- 31 10.1029/1998gl900251, 1999.
- 32 Kim, H., Barkey, B., and Paulson, S. E.: Real refractive indices of α and β -pinene and toluene
- secondary organic aerosols generated from ozonolysis and photo-oxidation J. Geophys. Res.,
 115, D24212, 2010.
- Kim, H., and Paulson, S. E.: Real refractive indices and volatility of secondary organic aerosol
 generated from photooxidation and ozonolysis of limonene, α-pinene and toluene, Atmos. Chem.
- 37 Phys. Discuss., 13, 1949-1977, 10.5194/acpd-13-1949-2013, 2013.
- 38 Kinne, S., Lohmann, U., Feichter, J., Schulz, M., Timmreck, C., Ghan, S., Easter, R., Chin, M.,
- Ginoux, P., Takemura, T., Tegen, I., Koch, D., Herzog, M., Penner, J., Pitari, G., Holben, B.,
 Eck, T., Smirnov, A., Dubovik, O., Slutsker, I., Tanre, D., Torres, O., Mishchenko, M.,
 Geogdzhayev, I., Chu, D. A., and Kaufman, Y.: Monthly averages of aerosol properties: A global
- 41 Geogdznayev, I., Chu, D. A., and Kaulman, Y.: Monthly averages of aerosol properties. A global 42 comparison among models, satellite data, and AERONET ground data, J. Geophys. Res. Atmos.,
- 43 108, 10.1029/2001jd001253, 2003.
- Koop, T., Bookhold, J., Shiraiwa, M., and Poschl, U.: Glass transition and phase state of organic
 compounds: dependency on molecular properties and implications for secondary organic

- aerosols in the atmosphere, Physical Chemistry Chemical Physics, 13, 19238-19255,
 10.1039/C1CP22617G, 2011.
- 3 Kroll, J. H., and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution
- 4 of low-volatility organics in the atmosphere, Atmos. Environ., 42, 3593-3624,
- 5 10.1016/j.atmosenv.2008.01.003, 2008.
- 6 Lambe, A. T., Onasch, T. B., Massoli, P., Croasdale, D. R., Wright, J. P., Ahern, A. T., Williams,
- L. R., Worsnop, D. R., Brune, W. H., and Davidovits, P.: Laboratory studies of the chemical
 composition and cloud condensation nuclei (CCN) activity of secondary organic aerosol (SOA)
- composition and cloud condensation nuclei (CCN) activity of secondary organic aerosol (SOA)
 and oxidized primary organic aerosol (OPOA), Atmos. Chem. Phys., 11, 8913-8928,
- 10 10.5194/acp-11-8913-2011, 2011.
- 10 10.5194/acp-11-8913-2011, 2011.
- 11 Lambe, A. T., Cappa, C. D., Massoli, P., Onasch, T. B., Forestieri, S. D., Martin, A. T.,
- Cummings, M. J., Croasdale, D. R., Brune, W. H., Worsnop, D. R., and Davidovits, P.:
 Relationship between Oxidation Level and Optical Properties of Secondary Organic Aerosol,
- 14 Environmental science & technology, 47, 6349-6357, 10.1021/es401043j, 2013.
- 15 Lang-Yona, N., Rudich, Y., Mentel, T. F., Bohne, A., Buchholz, A., Kiendler-Scharr, A., Kleist,
- 16 E., Spindler, C., Tillmann, R., and Wildt, J.: The chemical and microphysical properties of
- 17 secondary organic aerosols from Holm Oak emissions, Atmos. Chem. Phys., 10, 7253-7265,
- 18 10.5194/acp-10-7253-2010, 2010.
- 19 Liu, Y. G., and Daum, P. H.: Relationship of refractive index to mass density and self-consistency
- 20 of mixing rules for multicomponent mixtures like ambient aerosols, Journal of Aerosol Science,
- 21 39, 974-986, 10.1016/j.jaerosci.2008.06.006, 2008.
- 22 Liu, P., Zhang, Y., and Martin, S. T.: Complex Refractive Indices of Thin Films of Secondary
- Organic Materials by Spectroscopic Ellipsometry from 220 to 1200 nm, Environmental Science
 & Technology, 47, 13594-13601, 10.1021/es403411e, 2013.
- Lohmann, U., and Feichter, J.: Global indirect aerosol effects: a review, Atmos. Chem. Phys., 5, 715-737, 2005.
- Martin, S. T.: Phase transitions of aqueous atmospheric particles, Chemical Reviews, 100, 34033453, 10.1021/cr990034t, 2000.
- 29 Martin, S. T., Andreae, M. O., Althausen, D., Artaxo, P., Baars, H., Borrmann, S., Chen, Q.,
- 30 Farmer, D. K., Guenther, A., Gunthe, S. S., Jimenez, J. L., Karl, T., Longo, K., Manzi, A., Müller,
- 31 T., Pauliquevis, T., Petters, M. D., Prenni, A. J., Pöschl, U., Rizzo, L. V., Schneider, J., Smith, J.
- 32 N., Swietlicki, E., Tota, J., Wang, J., Wiedensohler, A., and Zorn, S. R.: An overview of the
- Amazonian Aerosol Characterization Experiment 2008 (AMAZE-08), Atmos. Chem. Phys., 10,
 11415-11438, 10.5194/acp-10-11415-2010, 2010.
- 54 11415-11458, 10.5194/acp-10-11415-2010, 2010.
- 35 Massoli, P., Lambe, A. T., Ahern, A. T., Williams, L. R., Ehn, M., Mikkil*, J., Canagaratna, M.
- 36 R., Brune, W. H., Onasch, T. B., Jayne, J. T., Pet*j*, T., Kulmala, M., Laaksonen, A., Kolb, C.
- E., Davidovits, P., and Worsnop, D. R.: Relationship between aerosol oxidation level and hygroscopic properties of laboratory generated secondary organic aerosol (SOA) particles,
- 39 Geophys. Res. Lett., 37, L24801, 2010.
 - 40 Matsumoto, K., Tanaka, H., Nagao, I., and Ishizaka, Y.: Contribution of particulate sulfate and
 - 41 organic carbon to cloud condensation nuclei in the marine atmosphere, Geophys. Res. Lett., 24,
 - 42 655-658, 10.1029/97gl00541, 1997.
 - 43 McFiggans, G., Artaxo, P., Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S.,
 - 44 Gysel, M., Laaksonen, A., Lohmann, U., Mentel, T. F., Murphy, D. M., O'Dowd, C. D., Snider,

- J. R., and Weingartner, E.: The effect of physical and chemical aerosol properties on warm cloud
- 2 droplet activation, Atmos. Chem. Phys., 6, 2593-2649, 2006.
- 3 McIntire, T. M., Ryder, O. S., Gassman, P. L., Zhu, Z., Ghosal, S., and Finlayson-Pitts, B. J.:
- 4 Why ozonolysis may not increase the hydrophilicity of particles, Atmos. Environ., 44, 939-944,
- 5 10.1016/j.atmosenv.2009.11.009, 2010.
- Mei, F., Setyan, A., Zhang, Q., and Wang, J.: CCN activity of organic aerosols observed
 downwind of urban emissions during CARES, Atmos. Chem. Phys., 13, 12155-12169,
 10.5194/acp-13-12155-2013, 2013.
- 9 Meyer, N. K., Duplissy, J., Gysel, M., Metzger, A., Dommen, J., Weingartner, E., Alfarra, M.
- 10 R., Prevot, A. S. H., Fletcher, C., Good, N., McFiggans, G., Jonsson, A. M., Hallquist, M.,
- Baltensperger, U., and Ristovski, Z. D.: Analysis of the hygroscopic and volatile properties of
- 12 ammonium sulphate seeded and unseeded SOA particles, Atmos. Chem. Phys., 9, 721-732, 2009.
- 13 Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Poschl, U.: Amorphous and crystalline
- aerosol particles interacting with water vapor: conceptual framework and experimental evidence
 for restructuring, phase transitions and kinetic limitations, Atmos. Chem. Phys., 9, 9491-9522,
- for restructuring, phase transitions and kinetic limitations, Atmos. Chem. Phys., 9, 9491-9522,
 2009
- 17 Moise, T., and Rudich, Y.: Reactive uptake of ozone by aerosol-associated unsaturated fatty
- acids: Kinetics, mechanism, and products, Journal of Physical Chemistry A, 106, 6469-6476,
 10.1021/jp025597e, 2002.
- Moussa, S. G., McIntire, T. M., Szőri, M., Roeselová, M., Tobias, D. J., Grimm, R. L.,
 Hemminger, J. C., and Finlayson-Pitts, B. J.: Experimental and Theoretical Characterization of
 Adsorbed Water on Self-Assembled Monolayers: Understanding the Interaction of Water with
 Atmospherically Relevant Surfaces[†], The Journal of Physical Chemistry A, 113, 2060-2069,
 10.1021/jp808710n, 2009.
- Müller, T., Laborde, M., Kassell, G., and Wiedensohler, A.: Design and performance of a threewavelength LED-based total scatter and backscatter integrating nephelometer, Atmos. Meas.
 Tech., 4, 1291-1303, 10.5194/amt-4-1291-2011, 2011.
- 28 Müller, L., Reinnig, M. C., Naumann, K. H., Saathoff, H., Mentel, T. F., Donahue, N. M., and
- 29 Hoffmann, T.: Formation of 3-methyl-1,2,3-butanetricarboxylic acid via gas phase oxidation of
- 30 pinonic acid a mass spectrometric study of SOA aging, Atmos. Chem. Phys., 12, 1483-1496,
- 31 10.5194/acp-12-1483-2012, 2012
- Murray, B. J.: Inhibition of ice crystallisation in highly viscous aqueous organic acid droplets,
 Atmos. Chem. Phys., 8, 5423-5433, 10.5194/acp-8-5423-2008, 2008.
- Nakayama, T., Matsumi, Y., Sato, K., Imamura, T., Yamazaki, A., and Uchiyama, A.: Laboratory
 studies on optical properties of secondary organic aerosols generated during the photooxidation
 of toluene and the ozonolysis of alpha-pinene, J. Geophys. Res. Atmos., 115,
 10.1029/2010jd014387, 2010.
- Nakayama, T., Sato, K., Matsumi, Y., Imamura, T., Yamazaki, A., and Uchiyama, A.:
 Wavelength Dependence of Refractive Index of Secondary Organic Aerosols Generated during
- 40 the Ozonolysis and Photooxidation of alpha-Pinene, Sola, 8, 10.2151/sola.2012-030, 2012.
- 41 Nakayama, T., Sato, K., Matsumi, Y., Imamura, T., Yamazaki, A., and Uchiyama, A.:
- 42 Wavelength and NOx dependent complex refractive index of SOAs generated from the
- 43 photooxidation of toluene, Atmos. Chem. Phys., 13, 531-545, 10.5194/acp-13-531-2013, 2013.

- 1 Nemesure, S., Wagener, R., and Schwartz, S. E.: Direct shortwave forcing of climate by the
- 2 anthropogenic sulfate aerosol: Sensitivity to particle size, composition, and relative humidity, J.
- 3 Geophys. Res. Atmos., 100, 26105-26116, 10.1029/95jd02897, 1995.
- 4 Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H.,
- 5 Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L.,
- 6 Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prevot, A. S. H., Dinar, E., Rudich, Y., and
- 7 Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from
- 8 Aerosol Mass Spectrometry, Atmos. Chem. Phys., 10, 4625-4641, 10.5194/acp-10-4625-2010,
- 9 2010.s
- 10 Novakov, T., and Penner, J. E.: Large contribution of organic aerosols to cloud-condensation-
- 11 nuclei, Nature, 365, 823-826, 10.1038/365823a0, 1993.
- 12 O'Donnell, D., Tsigaridis, K., and Feichter, J.: Estimating the direct and indirect effects of
- secondary organic aerosols using ECHAM5-HAM, Atmos. Chem. Phys., 11, 8635-8659,
 10.5194/acp-11-8635-2011, 2011.
- 15 Orr Jr, C., Hurd, F. K., and Corbett, W. J.: Aerosol size and relative humidity, Journal of Colloid 16 Science, 13, 472-482, http://dx.doi.org/10.1016/0095-8522(58)90055-2, 1958.
- 17 Pajunoja, A., Malila, J., Hao, L. Q., Joutsensaari, J., Lehtinen, K. E. J., and Virtanen, A.:
- 18 Estimating the Viscosity Range of SOA Particles Based on Their Coalescence Time, Aerosol
- 19 Science and Technology, 48, I-IV, 10.1080/02786826.2013.870325, 2014.
- Pankow, J. F., Storey, J. M. E., and Yamasaki, H.: Effects of relative-humidity on gas-particle
 partitioning of semivolatile organic-compounds to urban particulate matter, Environmental
 Science & Technology, 27, 2220-2226, 10.1021/es00047a032, 1993.
- 23 Pankow, J. F., and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting
- vapor pressures and enthalpies of vaporization of multifunctional organic compounds, Atmos.
 Chem. Phys., 8, 2773-2796, 2008.
- Paulson, S. E., Chung, M., Sen, A. D., and Orzechowska, G.: Measurement of OH radical
 formation from the reaction of ozone with several biogenic alkenes, J. Geophys. Res. Atmos.,
 103, 25533-25539, 10.1029/98jd01951, 1998.
- Pere, J. C., Mallet, M., Pont, V., and Bessagnet, B.: Impact of aerosol direct radiative forcing on
 the radiative budget, surface heat fluxes, and atmospheric dynamics during the heat wave of
 summer 2003 over western Europe: A modeling study, J. Geophys. Res.-Atmos., 116,
 10.1029/2011jd016240, 2011.
- 33 Perraud, V., Bruns, E. A., Ezell, M. J., Johnson, S. N., Yu, Y., Alexander, M. L., Zelenyuk, A.,
- 34 Imre, D., Chang, W. L., Dabdub, D., Pankow, J. F., and Finlayson-Pitts, B. J.: Nonequilibrium
- 35 atmospheric secondary organic aerosol formation and growth, Proceedings of the National
- 36 Academy of Sciences, 10.1073/pnas.1119909109, 2012
- Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth
 and cloud condensation nucleus activity, Atmos. Chem. Phys., 7, 1961-1971, 10.5194/acp-71961-2007, 2007.
- Prenni, A. J., DeMott, P. J., and Kreidenweis, S. M.: Water uptake of internally mixed particles
 containing ammonium sulfate and dicarboxylic acids, Atmos. Environ., 37, 4243-4251, 2003.
- 42 Qi, L.: Can secondary organic aerosol formed in an atmospheric simulation chamber 43 continuously age?, Atmos. Environ., 2990-2996, 2010.

- 1 Qi, L., Nakao, S., and Cocker, D. R.: Aging of secondary organic aerosol from α-pinene
- 2 ozonolysis: Roles of hydroxyl and nitrate radicals, Journal of the Air & Waste Management
- 3 Association, 62, 1359-1369, 10.1080/10962247.2012.712082, 2012.
- Redmond, H., and Thompson, J. E.: Evaluation of a quantitative structure-property relationship
 (QSPR) for predicting mid-visible refractive index of secondary organic aerosol (SOA), Physical
- 6 Chemistry Chemical Physics, 13, 6872-6882, 10.1039/c0cp02270e, 2011.
- 7 Reinhardt, A., Emmenegger, C., Gerrits, B., Panse, C., Dommen, J., Baltensperger, U., Zenobi,
- 8 R., and Kalberer, M.: Ultrahigh mass resolution and accurate mass measurements as a tool to
- 9 characterize oligomers in secondary organic aerosols, Analytical Chemistry, 79, 4074-4082,
- 10 10.1021/ac062425v, 2007.
- 11 Renbaum-Wolff, L., Grayson, J. W., Bateman, A. P., Kuwata, M., Sellier, M., Murray, B. J.,
- 12 Shilling, J. E., Martin, S. T., and Bertram, A. K.: Viscosity of α -pinene secondary organic 13 material and implications for particle growth and reactivity, Proceedings of the National
- Academy of Sciences, 110, 8014-8019, 10.1073/pnas.1219548110, 2013.
- 15 Rickards, A. M. J., Miles, R. E. H., Davies, J. F., Marshall, F. H., and Reid, J. P.: Measurements
- 16 of the Sensitivity of Aerosol Hygroscopicity and the κ Parameter to the O/C Ratio, The Journal
- 17 of Physical Chemistry A, 117, 14120-14131, 10.1021/jp407991n, 2013.
- 18 RiveraCarpio, C. A., Corrigan, C. E., Novakov, T., Penner, J. E., Rogers, C. F., and Chow, J. C.:
- 19 Derivation of contributions of sulfate and carbonaceous aerosols to cloud condensation nuclei
- from mass size distributions, J. Geophys. Res. Atmos., 101, 19483-19493, 10.1029/95jd01077,
 1996.
- Roth, C. M., Goss, K. U., and Schwarzenbach, R. P.: Sorption of a diverse set of organic vapors
 to urban aerosols, Environmental Science & Technology, 39, 6638-6643, 10.1021/es0503837,
 2005.
- Rudich, Y., Donahue, N. M., and Mentel, T. F.: Aging of organic aerosol: Bridging the gap
 between laboratory and field studies, in: Annu. Rev. Phys. Chem., Annual Review of Physical
 Chemistry, Annual Reviews, Palo Alto, 321-352, 2007.
- 28 Saathoff, H., Naumann, K.-H., Schnaiter, M., Schöck, W., Möhler, O., Schurath, U.,
- 29 Weingartner, E., Gysel, M., and Baltensperger, U.: Coating of soot and (NH4)2SO4 particles by
- 30 ozonolysis products of [alpha]-pinene, Journal of Aerosol Science, 34, 1297-1321, 2003.
- 31 Saathoff, H., Naumann, K. H., Möhler, O., Jonsson, Å. M., Hallquist, M., Kiendler-Scharr, A.,
- 32 Mentel, T. F., Tillmann, R., and Schurath, U.: Temperature dependence of yields of secondary
- 33 organic aerosols from the ozonolysis of α -pinene and limonene, Atmos. Chem. Phys., 9, 1551-
- 34 1577, 10.5194/acp-9-1551-2009, 2009.
- 35 Sato, K., Takami, A., Kato, Y., Seta, T., Fujitani, Y., Hikida, T., Shimono, A., and Imamura, T.:
- AMS and LC/MS analyses of SOA from the photooxidation of benzene and 1,3,5trimethylbenzene in the presence of NOx: effects of chemical structure on SOA aging, Atmos.
- 38 Chem. Phys., 12, 4667-4682, 10.5194/acp-12-4667-2012, 2012.
- 39 Saukko, E., Lambe, A. T., Massoli, P., Koop, T., Wright, J. P., Croasdale, D. R., Pedernera, D.
- 40 A., Onasch, T. B., Laaksonen, A., Davidovits, P., Worsnop, D. R., and Virtanen, A.: Humidity-
- 41 dependent phase state of SOA particles from biogenic and anthropogenic precursors, Atmos.
- 42 Chem. Phys., 12, 7517-7529, 10.5194/acp-12-7517-2012, 2012.

- 1 Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development
- 2 of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-
- 3 aromatic volatile organic compounds, Atmos. Chem. Phys., 3, 161-180, 2003.
- Saxena, P., Hildemann, L. M., McMurry, P. H., and Seinfeld, J. H.: Organics alter hygroscopic
 behavior of atmospheric particles, J. Geophys. Res., 100, 18755-18770, 1995.
- 6 Schnaiter, M., Horvath, H., Möhler, O., Naumann, K.-H., Saathoff, H., and Schöck, O. W.: UV-
- VIS-NIR spectral optical properties of soot and soot-containing aerosols, Journal of Aerosol
 Science, 34, 1421-1444, 2003.
- 8 Science, 34, 1421-1444, 2003.
- 9 Semeniuk, T. A., Wise, M. E., Martin, S. T., Russell, L. M., and Buseck, P. R.: Water uptake
- characteristics of individual atmospheric particles having coatings, Atmos. Environ., 41, 6225-6235, 2007.
- 12 Shilling, J. E., Chen, Q., King, S. M., Rosenoern, T., Kroll, J. H., Worsnop, D. R., McKinney,
- 13 K. A., and Martin, S. T.: Particle mass yield in secondary organic aerosol formed by the dark
- 14 ozonolysis of α-pinene, Atmos. Chem. Phys., 8, 2073-2088, 10.5194/acp-8-2073-2008, 2008.
- 15 Shiraiwa, M., Yee, L. D., Schilling, K. A., Loza, C. L., Craven, J. S., Zuend, A., Ziemann, P. J.,
- 16 and Seinfeld, J. H.: Size distribution dynamics reveal particle-phase chemistry in organic aerosol
- 17 formation, Proceedings of the National Academy of Sciences, 10.1073/pnas.1307501110, 2013.
- 18 Suda, S. R., Petters, M. D., Matsunaga, A., Sullivan, R. C., Ziemann, P. J., and Kreidenweis, S.
- 19 M.: Hygroscopicity frequency distributions of secondary organic aerosols, Journal of
- 20 Geophysical Research: Atmospheres, 117, D04207, 10.1029/2011jd016823, 2012.
- 21 Tolocka, M. P., Jang, M., Ginter, J. M., Cox, F. J., Kamens, R. M., and Johnston, M. V.:
- Formation of oligomers in secondary organic aerosol, Environmental science & technology, 38,
 1428-1434, 10.1021/es035030r, 2004.
- Tolocka, M. P., Heaton, K. J., Dreyfus, M. A., Wang, S. Y., Zordan, C. A., Saul, T. D., and Johnston, M. V.: Chemistry of particle inception and growth during alpha-pinene ozonolysis,
- 26 Environmental science & technology, 40, 1843-1848, 10.1021/es051926f, 2006.
- Tost, H., and Pringle, K. J.: Improvements of organic aerosol representations and their effects in
 large-scale atmospheric models, Atmos. Chem. Phys., 12, 8687-8709, 10.5194/acp-12-86872012, 2012.
- 30 Tritscher, T., Dommen, J., DeCarlo, P. F., Gysel, M., Barmet, P. B., Praplan, A. P., Weingartner,
- 31 E., Prevot, A. S. H., Riipinen, I., Donahue, N. M., and Baltensperger, U.: Volatility and
- 32 hygroscopicity of aging secondary organic aerosol in a smog chamber, Atmos. Chem. Phys., 11,
- 33 11477-11496, 10.5194/acp-11-11477-2011, 2011.
- Turpin, B. J., and Huntzicker, J. J.: Identification of secondary organic aerosol episodes and
 quantitation of primary and secondary organic aerosol concentrations during scaqs, Atmos.
 Environ., 29, 3527-3544, 10.1016/1352-2310(94)00276-q, 1995.
- Vaden, T. D., Song, C., Zaveri, R. A., Imre, D., and Zelenyuk, A.: Morphology of mixed primary
 and secondary organic particles and the adsorption of spectator organic gases during aerosol
 formation, Proceedings of the National Academy of Sciences, 10.1073/pnas.0911206107, 2010.
- 40 Vaden, T. D., Imre, D., Beránek, J., Shrivastava, M., and Zelenyuk, A.: Evaporation kinetics and
- 41 phase of laboratory and ambient secondary organic aerosol, Proceedings of the National
- 42 Academy of Sciences, 10.1073/pnas.1013391108, 2011.

- 1 Virtanen, A., Kannosto, J., Kuuluvainen, H., Arffman, A., Joutsensaari, J., Saukko, E., Hao, L.,
- 2 Yli-Pirila, P., Tiitta, P., Holopainen, J. K., Keskinen, J., Worsnop, D. R., Smith, J. N., and
- Laaksonen, A.: Bounce behavior of freshly nucleated biogenic secondary organic aerosol particles, Atmos. Chem. Phys., 11, 8759-8766, 10.5194/acp-11-8759-2011, 2011.
- 4
 particles, Atmos. Chem. 1 hys., 11, 6757-6766, 10.5174/acp-11-6757-2011, 2011.
- Wang, J., Doussin, J. F., Perrier, S., Perraudin, E., Katrib, Y., Pangui, E., and Picquet-Varrault,
 B.: Design of a new multi-phase experimental simulation chamber for atmospheric photosmog,
- 7 aerosol and cloud chemistry research, Atmos. Meas. Tech. J1 AMT, 4, 2465-2494, 2011.
- Warren, B., Austin, R. L., and Cocker III, D. R.: Temperature dependence of secondary organic
 aerosol, Atmos. Environ., 43, 3548-3555, 2009.
- 10 Weingartner, E., Saathoff, H., Schnaiter, M., Streit, N., Bitnar, B., and Baltensperger, U.:
- 11 Absorption of light by soot particles: determination of the absorption coefficient by means of
- 12 aethalometers, Journal of Aerosol Science, 34, 1445-1463, 2003.
- 13 Wex, H., Petters, M. D., Carrico, C. M., Hallbauer, E., Massling, A., McMeeking, G. R., Poulain,
- 14 L., Wu, Z., Kreidenweis, S. M., and Stratmann, F.: Towards closing the gap between hygroscopic
- 15 growth and activation for secondary organic aerosol: Part 1 Evidence from measurements,
- 16 Atmos. Chem. Phys. Discuss., 9, 955-989, 10.5194/acpd-9-955-2009, 2009.
- 17 Wong, J. P. S., Lee, A. K. Y., Slowik, J. G., Cziczo, D. J., Leaitch, W. R., Macdonald, A., and
- 18 Abbatt, J. P. D.: Oxidation of ambient biogenic secondary organic aerosol by hydroxyl radicals:
- 19 Effects on cloud condensation nuclei activity, Geophys. Res. Lett., 38, L22805,
- 20 10.1029/2011GL049351, 2011.
- Yasmeen, F., Vermeylen, R., Maurin, N., Perraudin, E., Doussin, J. F., and Claeys, M.:
 Characterisation of tracers for aging of alpha-pinene secondary organic aerosol using liquid
 chromatography/negative ion electrospray ionisation mass spectrometry, Environmental
- 24 Chemistry, 9, 236-246, 10.1071/en11148, 2012.
- Yu, J. Z., Griffin, R. J., Cocker, D. R., Flagan, R. C., Seinfeld, J. H., and Blanchard, P.:
 Observation of gaseous and particulate products of monoterpene oxidation in forest atmospheres,
 Geophys. Res. Lett., 26, 1145-1148, 10.1029/1999gl900169, 1999.
- Zaveri, R. A., Barnard, J. C., Easter, R. C., Riemer, N., and West, M.: Particle-resolved
 simulation of aerosol size, composition, mixing state, and the associated optical and cloud
 condensation nuclei activation properties in an evolving urban plume, J. Geophys. Res. Atmos.,
- 31 115, 10.1029/2009jd013616, 2010.
- 32 Zelenyuk, A., Yang, J., Song, C., Zaveri, R. A., and Imre, D.: A New Real-Time Method for
- 33 Determining Particles' Sphericity and Density: Application to Secondary Organic Aerosol
- 34 Formed by Ozonolysis of alpha-Pinene, Environmental science & technology, 42, 8033-8038,
- 35 10.1021/es8013562, 2008.
- 36 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R.,
- Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P.
 F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa,
- N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams,
- 40 P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M.,
- 40 and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in
- 42 anthropogenically-influenced Northern Hemisphere midlatitudes, Geophys. Res. Lett., 34,
- 43 10.1029/2007gl029979, 2007.

- Ziemann, P. J., and Atkinson, R.: Kinetics, products, and mechanisms of secondary organic aerosol formation, Chemical Society Reviews, 41, 6582-6605, 10.1039/C2CS35122F, 2012.
- Zobrist, B., Soonsin, V., Luo, B. P., Krieger, U. K., Marcolli, C., Peter, T., and Koop, T.: Ultra-
- slow water diffusion in aqueous sucrose glasses, Physical Chemistry Chemical Physics, 13, 3514-3526, 10.1039/c0cp01273d, 2011.

1 Table 1. Initial Conditions, Temperatures, Relative Humidities, and Results of the Experiments.

2 All experiments started with ~ 200 ppb of α -pinene.

3

Experiments	O ₃ ^a	Temp	RH ^b	Mass concentration ^c	Yield ^d
	(ppb)	(°C)	(%)	(µg m ⁻³)	Tiela
E140910	251	23-25	<1	130	0.19
E050210	202	19-23	<1	117	0.18
E080210	232	19-20	<1	116	0.18
E090210	254	16-17	<1	118	0.15
E230910	246	22-23	<1	100	0.17
E160411	232	19-23	<1	70	0.16
E260411	247	20-25	<1	-	-
E300911	198	20-27	<1	78	0.11
E301111	197	18-20	<1	52	0.07
E091211	196	17-20	<1	81	0.11
E180411	201	21-26	<1	86	0.13
E280411	269	17-20	<1	-	-
E280911	293	22-27	<1	91	0.13
E021211	196	17-20	<1	70	0.14
E071211	162	16-19	<1	55	0.14
E200411	220	21-26	<1	97	0.15
E260911	292	23-26	<1	87	0.15
E281111	215	14-19	<1	44	0.13
E051211	185	16-19	<1	66	0.13
E120312	229	18-23	<1	139	0.21
E030512	216	20-23	<1	109	0.15
E060512	246	19-21	<1	102	0.16

4

- ⁵ ^aOzone concentrations determined from FTIR measurement
- 6 ^bRH before the injection of water at the end of the experiment

^cAerosol mass concentration estimated from the aerosol volume concentration corrected from
 dilution and by assuming a density of 1.2 g cm⁻³

9 ^dYield calculated from measured aerosol mass and the corresponding concentration of reacted α -10 pinene, both corrected from dilution. We used the SOA mass concentration and α -pinene 11 concentration after 5 minutes of reaction as initial concentrations since the initial concentration 12 of α -pinene could not be directly measured (see text).

- Table 2. Comparison of real part of the complex refractive index (CRI) of α -pinene-O₃ SOA with previous studies.

Reference	Real CRI	Residence time	λ (nm)	[α-pinene] _{initial} (ppm)	[O ₃] _{initial} (ppm)
This study	1.60 (± 0.02)	10 mn	525	0.20	0.25
	1.33 (± 0.02)	19 h			
Kim et <i>al.</i> (2010)	$1.45 (\pm 0.05)$	2.5 h	670	0.50-5.00	0.10-1.00
Kim and Paulson (2013)	1.39 (± 0.02)	< 30 mn	532	0.13-0.17	0.50
	$1.52 (\pm 0.02)$	4 h			
Liu et al. (2013)	1.498 (± 0.002)	38 s	550	4.00	52.2
Nakayama <i>et al.</i> (2010)	1.41 (± 0.02)	2-3 h	532	0.10	2.00
Nakayama <i>et al.</i> (2012)	$1.47 - 1.48 (\pm 0.02)$	2-3 h	532	0.10	1.09-2.57
Redmond and	$1.49 (\pm 0.04)$	-	532	-	0.5-1.00
Thompson (2011)					
Schnaiter et al. (2003)	1.44	1.23 h	>350	0.06	0.47
Wex et al. (2009)	1.45	2 mn	visible	-	excess

- Table 3. Comparison of size growth factor GF at 90% RH of α -pinene-O₃ SOA with previous studies.

Reference	GF(90%RH)	Residence time	[α-pinene] _{initial}	[O ₃] _{initial}
This study	1.02 - 1.07 (±0.02)	2 h	0.20	0.25
5	1.02 - 1.07 (±0.02)	20 h		
Prenni et al. (2007)	1.01 - 1.07 (±0.02)	2 h	-	excess
Qi et al. (2010)	1.09	30 mn	0.05-0.10	0.30-0.34
	1.09	7 h		
Saathoff et al. (2003)	1.08 (±0.01)	1 h	0.06	0.5
	1.11 (±0.01)	6 h		
Warren et al. (2009)	1.02 - 1.16 (±0.02)	30 mn	0.05	0.3-0.5
	1.02 - 1.16 (±0.02)	6 h		

1 Table 4. Comparison of bulk O:C of α -pinene-O ₃ SOA with previous s	studies in the liter	ature.
--	----------------------	--------

Reference	Bulk O:C of SOA	Instrumentation
This study	0.55 (±0.16) - 0.68 (±0.20)	HR-ToF-AMS
Aiken et al. (2007)	0.27 (±0.08)	HR-ToF-AMS
Shilling et al. (2008)	0.29 (±0.09) - 0.45 (±0.14)	HR-ToF-AMS
Chhabra <i>et al.</i> (2010)	0.43 (±0.13)	HR-ToF-AMS
Qi et al. (2012)	0.33	HR-ToF-AMS
Reinhardt et al. (2007)	0.4 - 0.6	FTICR-MS
Tolocka et al. (2006)	$0.37 (\pm 0.05) - 0.40 (\pm 0.12)$	NanoAMS

- 1 Figure 1: Experimental set-up of the CESAM chamber used to measure aerosol chemical,
- 2 hygroscopic and optical properties.



- 1 Figure 2. Temporal profiles of the initial phases of a typical α -pinene ozonolysis experiment
- 2 (E160410): (a) α -pinene (red), ozone (blue) and mass concentration of the SOA (green) and (b)
- 3 measured number size distribution.



- 1 Figure 3: Calculated real part of the complex refractive index at λ =525 nm of SOA as a
- 2 function of the reaction time (upper panel). The lower panel shows the corresponding mass
- 3 concentration for the initial formation phase after injection (estimated particle density set to 1.2
- $4 g m^{-3}$)



- 1 Figure 4: Size growth factor GF at 90% RH obtained from the H-TDMA measurement as a
- 2 function of the reaction time. For the calculations of GF, we used $D_{p,m}(RH_{dry})$ at 40 % RH.



- 1 Figure 5: Humidograms of (a) size growth factor measured by the H-TDMA and (b) scattering
- 2 growth factor measured by the nephelometer as a function of RH within the nephelometer
- 3 (bottom axis) and the RH within the chamber (upper axis), for "fresh" SOA (after 1 hour of
- 4 reaction) in red and for "aged" SOA (after 14 hours of reaction) in blue. For the calculations of 5 f(RH) and GF, we used $\sigma_{scat}(RH_{dry})$ and $D_{p,m}(RH_{dry})$ at 40 % RH.
- 6





1 Figure 6: Humidograms showing measured (black symbols) and predicted GF (grey line) as a

2 function of RH of SOA (a) after 1 hour of reaction and (b) after 14 hours of reaction. The grey

3 area represent the uncertainties in the calculation of GF from f(RH). . For the calculations of GF,

4 we used $D_{p,m}(RH_{dry})$ at 40 % RH.

5



- 1 Figure 7. O:C of the bulk SOA obtained from AMS measurements (blue and grey symbols) and
- 2 uncertainties in the bulk O:C values (grey area) as a function of time. These bulk O:C are
- 3 compared with those of the SOA surface determined from XPS analysis (red symbols).



- 1 Figure 8. Effect of the thickness of the SOA surface on the core O:C ratio. Values are calculated
- 2 for SOA after 1 hour of reaction (grey) and after 14 hours of reaction and aged SOA (blue).





- 1 Figure 9: Real part of the refractive indices at λ =525 nm of SOA as a function of the bulk O:C.
- 2 In our study, representative error bars represent $\pm 1\sigma$ in replicate measurement.



4 * measured at λ =405 nm

- 1 Figure 10. Mass extinction efficiency (k_{ext}) at λ =525 nm of SOA under dry (red dots) and wet
- 2 conditions (blue dots) as a function of time and bulk O:C. These kext values are compared to those
- 3 used for organic aerosols in several global models (lines) and those of SOA generated from a
- 4 mixture of biogenics released directly from plants (dashed area). Zaveri et al. (2010) assumed a
- 5 CRI of 1.45-0i and GF of 1.23, Pere et al. (2011) a CRI of 1.45-0.001i, Kinne et al. (2003) a CRI
- 6 of 1.60-0.003i and GF of 1.09 (for ULAQ model) and Hoyle et al. (2009) a CRI of 1.53-0i and
- 7 GF of 1.03. The hygroscopic growth of SOA was not taken into account by Pere et al. (2011).



Time from α -pinene injection (hour)