

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Rare temperature histories and cirrus ice number density in a parcel and one-dimensional model

**D. M. Murphy**

NOAA ESRL Chemical Sciences Division Boulder, CO USA

Received: 10 February 2014 – Accepted: 11 April 2014 – Published: 29 April 2014

Correspondence to: D. M. Murphy (daniel.m.murphy@noaa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**ACPD**  
14, 10701–10723, 2014

Discussion Paper | Discussion Paper

**Cirrus ice crystal number density**

**D. M. Murphy**

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|](#)

[|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## 1 Introduction

25 Cirrus clouds cover large areas of the Earth (Wang et al., 1994). They reflect a significant amount of sunlight, trap infrared heat, and affect the local heating rates and circulation of the upper troposphere (Liou, 1986). These radiative effects are sensitive to the number density of ice crystals (Fu and Liou, 1993; McFarquhar et al., 2000).

There are significant uncertainties in ice nucleation in cirrus clouds, that is, the mechanisms by which ice crystals form on pre-existing aerosol particles.

The most basic distinction is between homogeneous and heterogeneous freezing. Homogeneous freezing originates from a water solution droplet. Koop et al. (2000) showed that to at least a first approximation homogeneous freezing depends only on water activity and not on the identity of the solute. At equilibrium the water activity of the droplet is equal to the ambient relative humidity, so homogeneous freezing should depend only on relative humidity, an important simplification. However, as will be shown below, the number of ice crystals is significantly changed by even small departures from the Koop et al. approximation that all solutes behave the same significantly change the number of ice crystals.

Heterogeneous freezing is initiated at a solid surface or other interface. The solid may interact with the gas phase for deposition freezing, be inside a droplet for immersion freezing, or be in other configurations. Analysis of the residuals left from evaporated ice crystals shows that heterogeneous freezing is often the dominant mechanism in high altitude cirrus (Cziczo et al., 2013).

Heterogeneous freezing requires particles with specific compositions, so a key aspect is the availability of such particles. A few heterogeneous nuclei that form ice at low supersaturation can reduce the number of ice crystals compared to homogeneous freezing, whereas many heterogeneous nuclei can enhance the number (Jensen and Toon, 1997; Kärcher and Lohmann, 2003; Barahona and Nenes, 2009; Spichtinger and Gierens, 2009c). Extremely few heterogeneous nuclei form ice that sediments without a significant effect on the further evolution of the cloud.

Small-scale temperature fluctuations also strongly affect the number of ice crystals. Ice formation depends on the cooling rate as well as absolute humidity because water uptake by the ice crystals quenches the supersaturation that allows nucleation. Even though their amplitude is small, the speed of small-scale temperature fluctuations creates high cooling rates that can greatly enhance the number of ice crystals (Murphy and Gary, 1995; Hoyle et al., 2005; Jensen et al., 1998, 2013a). The shape and crystal

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



structure of the ice crystals also affects their number and especially the size distribution (Murphy, 2003; Sheridan et al., 2009).

## 2 Model description

A parcel model and a one-dimensional model are used for the calculations presented here. The parcel model is an extension of that in Murphy (2003). It tracks nucleation and growth of ice crystals from an initial aerosol size distribution. Ice is tracked in 20 size bins logarithmically spaced between 90 nm and 80  $\mu\text{m}$ . Ice is initially formed as cubic ice and anneals to hexagonal ice. Most model parameters are the same as in Murphy (2003). The annealing rate from cubic to hexagonal ice was reduced by a factor of 10 because subsequent data suggest a lower, albeit more complicated, rate (Murray and Bertram, 2006). The mass accommodation coefficient was set to 0.15 for hexagonal ice and 0.3 for cubic ice on the grounds that as a metastable phase it must be kinetically easier to form than hexagonal ice. Sensitivity tests were conducted for many of the parameters.

The one-dimensional model tracks ice crystal formation events rather than size bins. The main reason to track events is that it turns out that small amounts of sedimentation are important for the evolution of the model cloud. By tracking events the ice crystals can sediment by fractions of the vertical model spacing and there is therefore no numerical vertical diffusion of ice crystals (Jensen et al., 2010; Söhlch and Kärcher, 2010). Model physics such as the nucleation rate and ice growth equations were the same in the parcel and one-dimensional models.

Water vapor, temperature and other state parameters were tracked at a fixed vertical spacing. At each time step an event was generated if the integrated probability of nucleation exceeded 0.02 per liter. Thereafter the ice crystal diameter and sedimentation of that particle event were tracked as continuous variables whereas the water vapor for growth (evaporation) was taken from (given to) the nearest bins according to the position. For example, if a growing ice crystal event was exactly in the middle of a bin

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



all the water was taken from that bin whereas if it was at a bin boundary half the water was taken from each bin.

A detail is that the nucleation probability must be integrated over time and vertical distance in order to make nucleation events independent of the size of a time step or the water vapor grid spacing. When the integral of the nucleation probability reaches a threshold, an event is generated at the most probable location and the integrated nucleation probability reduced by one at that location. As a simple example, consider only three vertical bins in which, because of imposed cooling and water vapor, the probability of nucleation has slowly accumulated to 0.3 per 50 liters in the top and bottom bins and 0.5 in the middle bin. Because the total of 1.1 exceeds one, a nucleation event is generated in the most probable (middle) bin. Following this, the nucleation probability is 0.3 in the top and bottom bins and negative 0.5 in the middle bin. A few time steps later the integrated probability might be 0.6 in the top bin, 0.0 in the middle bin, and 0.5 in the bottom bin and the next event will take place in the top bin. During rapid nucleation, the total probability can increase in a single time step by more than one ice crystal per 50 liters. In that case ice events were initiated in more than one vertical bin and the surface area and mass of the subsequent ice were weighted according to the probability. A typical model run tracked perhaps 20,000 formation events. Ice crystals were formed as cubic ice and annealed stochastically to hexagonal ice depending on integrated probabilities analogous to those for ice nucleation. For computational reasons, every few time steps ice events with essentially identical sizes and (sedimented) vertical positions were combined.

### 3 Parcel model results

Parcel model simulations were run for frost point temperatures ranging from 185 to 230 K. At each temperature 80 simulations were run with various seeds for the random number generator that initializes the fractal small-scale temperature fluctuations. As in Murphy (2003), fractal temperature fluctuations with periods shorter than 2 min-

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



utes were removed. A 3.2K cooling with a 12-hour, half-sine pattern was added to the temperature fluctuations; otherwise many random fluctuations would never generate a model cloud. The same set of temperature fluctuations were repeated at each initial temperature. All simulations were recalculated for a variety of model parameters and 5 assumptions about homogeneous and heterogeneous ice nucleation processes.

One striking result is that various temperature histories generated a very wide range of ice crystal number densities for the same model assumptions. Figure 1 shows a histogram of the maximum ice number density with homogeneous freezing for temperature histories with ice saturation of 1.25 at 196 K (frost point of 197.4 K) along 10 with a subset of the model temperature trajectories. Jensen et al. (2010) also found wide distributions of ice number density depending on the relative phase of waves creating temperature fluctuations. Low ice number densities can be generated by temperature histories that spend a very short amount of time above the ice nucleation point (Spichtinger and Krämer, 2013). Wide distributions of number density are also 15 evident in various portions of one- and two-dimensional model fields (Lin et al., 2005; Spichtinger and Gierens, 2009b).

The absolute number of ice crystals in the model is sensitive to several of the assumed parameters, especially the accommodation coefficient. However, the results here emphasize trends with temperature and probability distributions due to temperature fluctuations. Such trends and distributions are much less sensitive to model 20 parameters.

Figure 2 shows results from the parcel model for a range of temperatures. The homogeneous nucleation calculations show many more ice crystals at low temperatures. This is because there is less water vapor at lower temperatures causing slower growth 25 of the ice crystals. With slower growth, more time elapses before the ice surface area grows enough to reduce the supersaturation and suppress further nucleation. With a longer nucleation event, more ice crystals form. This is a very fundamental temperature dependence and occurs when a variety of model parameters are changed (blue lines). The slope of the temperature dependence is similar to Kärcher (2002, Fig. 3)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

or Spichtinger and Gierens (2009a), but represents the median of a wide distribution rather than a single vertical velocity.

The inset in Fig. 2a explores which are the most sensitive parameters for ice crystal number density. The sensitivity to the upper side of the default parameters is almost a mirror image of those shown. Ice number is sensitive to the accommodation coefficient of water and extremely insensitive to the absolute nucleation rate. Ice number is moderately sensitive to displacing the homogeneous nucleation point by a small amount of supersaturation. “Faster  $T$  fluctuations” means that the fractal temperature fluctuations were smoothed at one one rather than 2 min

Of interest is the sensitivity to the slope of the nucleation rate. The Koop et al. (2000) nucleation rate is a very steep function of relative humidity at the nucleation threshold but is not quite a step function. Artificially increasing or lowering the slope with supersaturation has a much larger impact on the number of ice crystals than the nucleation rate itself. This is potentially important on the side of reducing the slope. If the Koop et al. result that water activity controls freezing is only a first approximation, then particles with different composition should freeze at slightly different supersaturations. This would be equivalent to a reduced slope, and the parcel model shows that the ice number is quite sensitive to this. Because the Koop et al. (2000) nucleation rate is extremely steep, the 0.67 slope case shown in Fig. 2a means that the relative humidity at freezing is the same within about 1 % for particles of various compositions. Yet that reduction in slope reduced the number of ice crystals by over a factor of two.

Figure 2b shows results from several heterogeneous cases. As expected, the ice number density is sensitive to the number of heterogeneous nuclei. Also as expected, if there are too few nuclei at very low temperature then the heterogeneous nuclei do not deplete the water vapor enough to prevent homogeneous freezing (Jensen and Toon, 1997; Kärcher and Lohmann, 2003). Figure 2b also compares cases where the heterogeneous nuclei are essentially identical or freeze over a range of relative humidities (“diverse nuclei” in the figure). Over these and other assumptions about heterogeneous freezing, the number of ice crystals always increased at lower temperatures.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 2c shows some published results on observed ice number density. There are only a few studies since the recognition that ice shattering on aircraft probes invalidates much of the older data (Jensen et al., 2009). There are several points of comparison. First, the results here support the contention in Jensen et al. (2013b) that the high ice number densities observed in thin layers near 190 K can be explained by homogeneous freezing. Second, it is extremely hard in a parcel model to reproduce the low end of the observed ice number densities below about 190 K, regardless of the number of heterogeneous ice nuclei. It might be possible if the accommodation coefficient were unity and other parameters were adjusted, but then the model could not match observations at warmer temperatures. Third, the upper limit of ice number densities observed in Krämer et al. (2009) above about 210 K cannot be reproduced by the parcel model. A possible explanation is that the upper limit may be due to outflow cirrus that are not described by the parcel model.

Fourth and most important, neither homogeneous nor heterogeneous freezing in a parcel model can reproduce the sign of the observed slope in either Krämer et al. (2009) or Jensen et al. (2012). No simple tuning of the parcel model parameters can change the sign of the slope with temperature. The slope is a fundamental consequence of less water vapor at lower temperatures. The only way to reverse the slope would be if some parameters such as the number of ice nuclei or the accommodation coefficient were themselves functions of temperature. Even so, the temperature dependence of the parameters would have to be very strong to overcome the basic increase in ice number at lower temperature.

#### 4 One-dimensional model

The failure of a parcel model to reproduce the observed temperature dependence of ice crystal number suggests that there may be important ice cloud microphysics not captured by a parcel model. The most obvious candidate is sedimentation of ice crystals. At least a one-dimensional model is necessary to investigate sedimentation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 3 shows the configuration of the one-dimensional model runs. An initial saturation ratio of 1.1 is imposed except for a subsaturated region near the bottom of the model domain so that falling ice crystals can evaporate. Then one of the same temperature trajectories used in the parcel model is imposed over a 100 m thick layer with a maximum amplitude at the center of the layer and a smooth transition to zero above and below that. A slightly supersaturated layer remains below the imposed temperature fluctuation. The right-hand profiles in Fig. 3 show the saturation profile just as ice starts to nucleate and grow near the center of the cooled layer, causing a dip in saturation ratio at the 200 m vertical tick.

Figure 4 shows sample results from the one-dimensional model for one of the imposed temperature profiles. There are differences but more importantly similarities when different assumptions about ice nucleation are made.

One difference is that the cloud in homogeneous cooling forms later in time because it requires a higher ice supersaturation. Second, the number of ice crystals formed (contours) is much lower when heterogeneous ice nuclei are present. Third, the homogeneous freezing case is less striated than the heterogeneous cases.

There are important similarities. Although the peak ice crystal number densities are a factor of roughly 25 different between the top two panels, the peak ice mass densities are within 40 %. For all cases, layers and fall streaks are generated that are far narrower than the vertical scale of the imposed cooling (Fig. 3). On this plot of altitude vs. time, the slopes of the fall streaks are also similar, meaning that the fall speeds are similar.

A final similarity is that in this model the fall streaks are often generated in layers that are slightly displaced from the layers that produce large numbers of ice crystals. This is most visible in the diverse nuclei (bottom) panel but there is also a nearly complete but less visible displacement between high number and fall streak layers in the homogeneous freezing (top) panel. Only with a few good ice nuclei do the most productive nucleation events generate fall streaks, and even then the fall rate of the number density (contours) is perhaps half the fall rate of the mass (color).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In hindsight there must be often a difference between layers and times that produce high number density and those that produce fall streaks. The high number density ice crystals are small and usually don't fall fast enough to reach supersaturated regions where they can continue to grow. It is worth thinking about how this relates to the 5 probability distributions of the parcel model in Fig. 1. Because the imposed cooling has different amplitudes at different altitudes each layer will reach a given nucleation threshold at a different point in the time history. Some will reach it during a rapid jump in temperature, some at a slow point. It is similar to using different temperature histories. Figure 1 shows that there are always a few temperature histories that produce low ice 10 numbers, even in homogeneous freezing that usually creates high number densities.

These rare freezing events that produce fall streaks are disproportionately important because the falling crystals influence a much larger volume of air than those that don't fall. Exactly how much more important will depend on the initial vertical temperature and humidity profiles. The model does give a key result: these rare low number density 15 events can still generate fall streaks even if they are between or falling through high number density layers.

Figure 5 shows ice number densities from the one-dimensional model analogous to the parcel model in Fig. 2. The ice number densities within the upper, cooled layer follow the same patterns as the parcel model. This is true for both the temperature dependence of each case and the relative magnitudes of the various cases. The absolute 20 values are lower in the one-dimensional model because the vertical averaging includes both high number density layers and layers with few ice crystals.

The averaging layer below the cooled region is looking at fall streaks. Here the number densities are much lower. The heterogeneous cases have a slightly smaller temperature dependence: a factor of about 5 over the temperature range instead of a factor 25 of 10. The homogeneous case is completely different below the imposed cooling than within it. It is not clear if the maximum near 200 K is robust or a feature of the initial profile in the model. But the absence of very high number densities below 200 K should be robust. With very little water available at cold temperatures, only regions with few

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

particles can grow large enough crystals to produce fall streaks. The sensitivity to the accommodation coefficient is also much smaller in the fall streaks than in the upper layer.

The one-dimensional model produced remarkably fine vertical structure from a much smoother temperature profile. A vertical resolution for water vapor concentration of about 2 m was found to be necessary even though the falling ice crystals were tracked to fractions of the grid spacing. The generation of fine vertical structure in cirrus clouds has been seen previously. The sedimenting NAT crystals in Fueglistaler et al. (2002) required fine vertical resolution at the base of the cloud. They used 1 cm (!) vertical resolution near cloud base although no tests were done to see if such fine resolution was necessary (S. Fueglistaler, personal communication, 2013). Lin et al. (2005) found that 1 m vertical spacing was necessary in a cirrus cloud model with broad uplift. Spichtinger and Gierens (2009a) and Jensen et al. (2012) found that 10 m was necessary, again without the boundary conditions changing that rapidly in the vertical.

There are two reasons for the fine vertical structure. First, ice formation is very sensitive to the cooling rate as a nucleation threshold is crossed, so small variations in temperature history can produce large variations in ice number density. Second, sedimentation acts to amplify small vertical differences in ice crystal size into much larger fall streaks.

The one-dimensional model is still missing some important physics. Simply looking at cirrus clouds shows that fall streaks are horizontally patchy and curved, neither of which can be captured in a one-dimensional model. Wind shear is important to cirrus clouds (Spichtinger and Gierens, 2009b). Radiation-induced turbulence affects the generation of large particles and fall streaks (Gu and Liou, 2000). Neither process is easy to capture in one-dimension.

The vertical profiles of saturation and cooling in the one-dimensional model are obviously arbitrary. Trying to find a more realistic profile is limited both by the inherent limitations of working in one dimension and a lack of observations. There are few observations of upper tropospheric temperatures on time scales of a minute and hori-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

zontal scales well below a kilometer. Even when an aircraft can measure a slice of temperature profiles, translating the observed temperatures into those experienced by an air parcel depends on unmeasured phase relationships between waves of different frequencies (Bacmeister et al., 1999). Similar issues can arise for ground-based remote sensing of a single point.

## 5 Discussion and conclusions

The one-dimensional model results presented here are in many respects similar to a model of nitric acid trihydrate (NAT) crystals in the polar stratosphere (Fueglistaler et al., 2002). In an environment with much slower crystal growth than even the coldest

10 cirrus cloud, Fueglistaler et al. found that sedimentation out of a “mother cloud” into a supersaturated layer could produce a few large NAT crystals. The number of sedimenting NAT crystals was insensitive to assumptions about the nucleation mechanism. The reason was the supersaturated region below the cloud. Any crystal that managed to make its way into this layer grew and accelerated down from the cloud. The physics 15 of those falling crystals was determined by mixing at the bottom of the cloud and the growth conditions below the cloud, not the nucleation conditions in the cloud. This work suggests that a similar process often occurs for cirrus clouds.

There is a fundamental difference in sedimentation from cirrus clouds compared to 20 warm liquid clouds. The air around a warm cloud is subsaturated whereas the high supersaturations required for ice nucleation mean that the air around a newly formed cirrus cloud is supersaturated (Diao et al., 2013). Mixing and entrainment at the edges of a warm cloud causes evaporation of water droplets. In contrast, mixing and entrainment at the base or edges of a cirrus cloud causes growth of ice crystals near the edge 25 while simultaneously reducing their number density and hence competition for vapor. This can induce sedimentation from the base and periphery of the cloud, very unlike a warm cloud.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Sedimentation has several roles in cold cirrus clouds (Spichtinger and Gierens, 2009b). Depending on the environment, sedimentation can limit (Kärcher, 2002) or extend (Luo et al., 2003) the lifetime of very thin cirrus clouds. For some ranges of temperature fluctuations a balance can be achieved between nucleation and sedimentation (Barahona and Nenes, 2011). Here, sedimentation is a mechanism for amplifying the importance of conditions that produce only a few ice crystals.

The very low ice number densities sometimes observed below 200 K can be produced in fall streaks, as also found by Jensen et al. (2012). The picture developed here of a cold cirrus cloud (below perhaps 210 K) is that the reason aircraft observations usually measure low ice number density is that the falling crystals sweep out a much larger volume than the ones that stay put. Jensen et al. (2013a) suggested that a few heterogeneous nuclei could initiate fall streaks even if there aren't enough such nuclei to suppress homogeneous nucleation. The model here generalizes that: fall streaks are produced by any time or place that makes a few large particles. These regions of few crystals might be from heterogeneous nucleation, temperature blips that happened to produce only a few ice crystals from homogeneous nucleation, or mixing at the bottom or edge of a cloud. The number and mass of ice crystals in such fall streaks is fairly insensitive to how they were produced.

There may be other observational tests of the importance of sedimentation besides comparing to observed ice number densities. Using statistics on the size of regions containing supersaturation and/or ice crystals, Diao et al. (2013) estimated that mature or evaporating cirrus are about seven times more prevalent than nucleation regions. Although their mature and evaporation classifications are not exactly comparable to fall streaks, it does support the notion that throughout most of a cirrus cloud the number of ice crystals is not determined by nucleation at that spot. A lidar with high spatial resolution could probably observe whether the fall streaks are originating from locations slightly displaced from the high number density layers. Such an observation might constrain the nucleation mechanism: in the model, only nucleation on a few good ice nuclei created fall streaks coincident with the highest number densities.

## Cirrus ice crystal number density

D. M. Murphy

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



At least for homogeneous freezing, the one-dimensional model predicts a wide range in ice number density between various regions of the cloud. In that case the visible reflectance could still be determined by the zones of numerous, small ice crystals that scatter light more efficiently even though other properties such as median ice number density were determined mostly by sedimentation.

Finally, the results here contain both bad and good news for large-scale modeling of cirrus clouds. The bad news is that the requirement for extremely fine vertical resolution implies that even mesoscale models have far too coarse of resolution to directly model cirrus clouds. The importance of sedimentation from rare temperature trajectories implies that parameterizations based on parcel models (e.g. Kärcher and Lohmann, 2003; Barahona and Nenes, 2009) probably miss essential physics.

The good news for large-scale modeling is that in the one-dimensional model the ice number density below cloud is much less sensitive to details of the microphysics than in a parcel model. There are new parameters required to describe sedimentation, such as the vertical extent of any supersaturated region below the cloud. But such parameters should be amenable to calculation by a three-dimensional model, although the required vertical resolution (perhaps tens of meters) is still more suitable for a local model than a global model.

*Acknowledgements.* This work was supported by NOAA base and climate funding.

## 20 References

Bacmeister, J. T., Eckermann, S. D., Tsias, A., Carslaw, K. S., and Peter, T.: Mesoscale temperature fluctuations induced by a spectrum of gravity waves: a comparison of parameterizations and their impact on stratospheric microphysics, *J. Atmos. Sci.*, 56, 1913–1924, 1999.

Barahona, D. and Nenes, A.: Parameterizing the competition between homogeneous and heterogeneous freezing in ice cloud formation – polydisperse ice nuclei, *Atmos. Chem. Phys.*, 9, 5933–5948, doi:10.5194/acp-9-5933-2009, 2009.

## Cirrus ice crystal number density

D. M. Murphy

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Barahona, D., and A. Nenes, Dynamical states of low temperature cirrus, *Atmos. Chem. Phys.*, 11, 3757–3771, 2011,  
<http://www.atmos-chem-phys.net/11/3757/2011/>.

5 Cziczo, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M., Zondlo, M. A., Smith, J. B., Twohy, C. H., and Murphy, D. M.: Clarifying the dominant sources and mechanisms of cirrus cloud formation, *Science*, 340, 1320–1324, 2013.

10 Diao, M., Zondlo, M. A., Heymsfield, A. J., Beaton, S. P., and Rogers, D. C.: Evolution of ice crystal regions on the microscale based on in situ observations, *Geophys. Res. Lett.*, 40, 3473–3478, 2013.

15 Fu, Q., and Liou, K. N., Parameterization of the radiative properties of cirrus clouds, *J. Atmos. Sci.*, 50, 2008–2025, 1993.

Fueglistaler, S., Luo, B. P., Voigt, C., Carslaw, K. S., and Peter, T.: NAT-rock formation by mother clouds: a microphysical model study, *Atmos. Chem. Phys.*, 2, 93–98, doi:10.5194/acp-2-93-2002, 2002.

20 15 Hoyle, C. R., Luo, B. P., and Peter, T., The origin of high ice crystal number densities in cirrus clouds, *J. Atmos. Sci.*, 62, 2568–2579, 2005.

Gensch, I. V., Bunz, H., Baumgardner, D. G., Christensen, L. E., Fahey, D. W., Herman, R. L., Popp, P. J., Smith, J. B., Troy, R. F., Webster, C. R., Weinstock, E. M., Wilson, J. C., Peter, T., and Krämer, M., Supersaturations, microphysics, and nitric acid partitioning in a cold cirrus cloud observed during CR-AVE 2006: an observation-modeling intercomparison study, *Environ. Res. Lett.*, 3, 1–9, 2008.

25 Gu, Y. and Liou, K. N.: Interactions of radiation, microphysics, and turbulence in the evolution of cirrus clouds, *J. Atmos. Sci.*, 57, 2463–2479, 2000.

Jensen, E. J., and Toon, O. B., The potential impact of soot particles from aircraft exhaust on cirrus clouds, *Geophys. Res. Lett.*, 24, 249–252, 1997.

30 Jensen, E. J., Toon, O. B., Tabazadeh, A., Sachse, G. W., Anderson, B. E., Chan, K. R., Twohy, C. W., Gandrud, B., Aulenbach, S. M., Heymsfield, A. J., Hallett, J., and Gary, B.: Ice nucleation processes in upper tropospheric wave-clouds observed during SUCCESS, *Geophys. Res. Lett.*, 25, 1363–1366, 1998.

Jensen, E. J., Lawson, P., Baker, B., Pilson, B., Mo, Q., Heymsfield, A. J., Bansemer, A., Bui, T. P., McGill, M., Hlavka, D., Heymsfield, G., Platnick, S., Arnold, G. T., and Tanelli, S.: On the importance of small ice crystals in tropical anvil cirrus, *Atmos. Chem. Phys.*, 9, 5519–5537, doi:10.5194/acp-9-5519-2009, 2009.

## Cirrus ice crystal number density

D. M. Murphy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

|◀

▶|

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cirrus ice crystal  
number density

D. M. Murphy

Jensen, E. J., Pfister, L., Bui, T.-P., Lawson, P., and Baumgardner, D.: Ice nucleation and cloud microphysical properties in tropical tropopause layer cirrus, *Atmos. Chem. Phys.*, 10, 1369–1384, doi:10.5194/acp-10-1369-2010, 2010.

Jensen, E. J., Pfister, L., and Bui, T. P.: Physical processes controlling ice concentrations in cold cirrus near the tropical tropopause, *J. Geophys. Res.*, 117, D11205, doi:10.1029/2011JD017319, 2012.

Jensen, E. J., Lawson, R. P., Bergman, J. W., Pfister, L., Bui, T. P., and Schmitt, C. G.: Physical processes controlling ice concentrations in synoptically forced, midlatitude cirrus, *J. Geophys. Res.-Atmos.*, 118, 5348–5360, doi:10.1002/jgrd.50421, 2013a.

Jensen, E. J., Diskin, G., Lawson, R. P., Lance, S., Bui, T. P., Hlavka, D., McGill, M., Pfister, L., Toon, O. B., and Gao, R.: Ice nucleation and dehydration in the Tropical Tropopause Layer, *P. Natl. Acad. Sci. USA*, 110, 2041–2046, 2013b.

Kärcher, B.: Properties of subvisible cirrus clouds formed by homogeneous freezing, *Atmos. Chem. Phys.*, 2, 161–170, doi:10.5194/acp-2-161-2002, 2002.

Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: heterogeneous freezing, *J. Geophys. Res.*, 108, 4402, doi:10.1029/2002JD003220, 2003.

Koop, T., Luo, B. B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, 2000.

Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, *Atmos. Chem. Phys.*, 9, 3505–3522, doi:10.5194/acp-9-3505-2009, 2009.

Lin, R.-F., Starr, D. O., Reichardt, J., and DeMott, P. J.: Nucleation in synoptically forced cirrostratus, *J. Geophys. Res.*, 110, D08208, doi:10.1029/2004JD005362, 2005.

Liou, K.-N.: Influence of cirrus clouds on weather and climate processes: a global perspective, *Mon. Weather Rev.*, 114, 1167–1199, 1986.

Luo, B. P., Peter, Th., Wernli, H., Fueglistaler, S., Wirth, M., Kiemle, C., Flentje, H., Yushkov, V. A., Khattatov, V., Rudakov, V., Thomas, A., Borrmann, S., Toci, G., Mazzinghi, P., Beuermann, J., Schiller, C., Cairo, F., Di Don-Francesco, G., Adriani, A., Volk, C. M., Strom, J., Noone, K., Mitev, V., MacKenzie, R. A., Carslaw, K. S., Trautmann, T., Santacesaria, V., and Stefanutti, L.: Ultrathin Tropical Tropopause Clouds (UTTCs): II. Stabilization mechanisms, *Atmos. Chem. Phys.*, 3, 1093–1100, doi:10.5194/acp-3-1093-2003, 2003.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	

<a href="#">Printer-friendly Version</a>
<a href="#">Interactive Discussion</a>



McFarquhar, G. M., Heymsfield, A. J., Spinhirne, J., and Hart, B., Thin and subvisual tropopause tropical cirrus: Observations and radiative impacts, *J. Atmos. Sci.*, 57, 1841–1853, 2000.

Murphy, D. M.: Dehydration in cold clouds is enhanced by a transition from cubic to hexagonal ice, *Geophys. Res. Lett.*, 30, 2230, doi:10.1029/2003GL018566, 2003.

5 Murphy, D. M. and Gary, B. L.: Mesoscale temperature fluctuations and polar stratospheric clouds, *J. Atmos. Sci.*, 52, 1753–1760, 1995.

Murray, B. J. and Bertram, A. K.: Formation and stability of cubic ice in water droplets, *Phys. Chem. Chem. Phys.*, 8, 186–192, 2006.

10 Sheridan, L., Harrington, J. Y., Lamb, D., and Sulia, K.: Influence of ice crystal aspect ratio on the evolution of ice size spectra during vapor depositional growth, *J. Atmos. Sci.*, 66, 3732–3743, 2009.

Söhlch, I., and Kärcher, B., A large-eddy model for cirrus clouds with explicit aerosol and ice microphysics and Lagrangian ice particle tracking, *Q. J. R. Meteorol. Soc.*, 136, 2074–2093, 2010.

15 Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1a: Model description and validation, *Atmos. Chem. Phys.*, 9, 685–706, doi:10.5194/acp-9-685-2009, 2009a.

Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1b: Structuring cirrus clouds by dynamics, *Atmos. Chem. Phys.*, 9, 707–719, doi:10.5194/acp-9-707-2009, 2009b.

20 Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 2: Competition of different nucleation mechanisms, *Atmos. Chem. Phys.*, 9, 2319–2334, doi:10.5194/acp-9-2319-2009, 2009c.

Spichtinger, P., and Krämer, M., Tropical tropopause ice clouds: a dynamic approach to the mystery of low crystal numbers, *Atmos. Chem. Phys.*, 13, 9801–9818, 2013, <http://www.atmos-chem-phys.net/13/9801/2013/>.

25 Wang, P.-H., McCormick, M. P., Poole, L. R., Chu, W. P., Yue, G. K., Kent, G. S., and Skeens, K. M., Tropical high cloud characteristics derived from SAGE II extinction measurements, *Atmos. Res.*, 34, 53–83, 1994.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



## Discussion Paper | Cirrus ice crystal number density

D. M. Murphy




**Fig. 1.** Histogram of ice number density generated by homogeneous freezing in a parcel model. Also shown are the first 8 of the 80 temperature trajectories, all with same amplitude and fractal slope of temperature fluctuations. Doubling the accommodation coefficient of water on the growing ice crystals systematically reduces the number density. However, the phase and slope of small-scale fluctuations near the homogeneous freezing threshold creates a histogram that is much wider than the effect of changing this or other model parameters.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Discussion Paper | Cirrus ice crystal number density

D. M. Murphy

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Title Page

## Abstract

## Introduction

## Conclusions

## References

## Tables

## Figures

ANSWER

Page 10

1

1

10

1

Back


Close

Full Screen / Esc

[Printer-friendly Version](#)

## Interactive Discussion






**Fig. 3.** Vertical profiles showing how the one-dimensional model was initialized and cooling imposed. The imposed cooling used a smooth vertical profile over 100 m but fluctuated in time. One can picture the half-sine temperature perturbation wiggling with an amplitude that followed one of the temperature trajectories in Fig. 1.

## Cirrus ice crystal number density


D. M. Murphy

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fig. 4.** One-dimensional model results for three assumptions about ice nucleation and the same imposed cooling. Contours show ice crystal concentration in  $\text{L}^{-1}$ . Color shows ice mass concentration with the same scale for all three panels.

## Discussion Paper | Cirrus ice crystal number density

D. M. Murphy



**Fig. 5.** Ice number densities in the one-dimensional model. The values are the median over 20 temperature histories of the maximum in time averaged over vertical range either in or below the layer with imposed cooling. These ranges are indicated in Fig. 4. The axis ranges are the same as in Fig. 2.

<a href="#">Title Page</a>	<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>	
<a href="#">Tables</a>	<a href="#">Figures</a>	
<a href="#">◀</a>	<a href="#">▶</a>	
<a href="#">◀</a>	<a href="#">▶</a>	
<a href="#">Back</a>	<a href="#">Close</a>	
<a href="#">Full Screen / Esc</a>		

<a href="#">Printer-friendly Version</a>
<a href="#">Interactive Discussion</a>