

# Observation and analysis of speciated atmospheric mercury in Shangri-la, Tibetan Plateau, China

H. Zhang<sup>1,2</sup>, X. W. Fu<sup>1</sup>, C.-J. Lin<sup>1,3,4</sup>, X. Wang<sup>1,2</sup>, X. B. Feng<sup>1\*</sup>

<sup>1</sup> State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang 550002, PR China (E-mail: [fengxinbin@vip.skleg.cn](mailto:fengxinbin@vip.skleg.cn))

<sup>2</sup> University of Chinese Academy of Sciences, Beijing 100049, PR China

<sup>3</sup> Department of Civil Engineering, Lamar University, Beaumont, Texas 77710, United States

<sup>4</sup> College of Energy and Environment, South China University of Technology, Guangzhou 510006, China

\*: Corresponding author

## Abstract

This study reports the concentrations and potential sources of speciated atmospheric mercury at the Shangri-La Atmosphere Watch Regional Station (SAWRS), a pristine high-altitude site (3580 m above sea level) in Tibetan Plateau, China. Total gaseous mercury (TGM, defined as the sum of Gaseous Elemental Mercury, GEM, and Gaseous Oxidized Mercury, GOM), GOM and particulate-bound mercury (PBM) were monitored from November 2009 to November 2010 to investigate the characteristics and potential influence of the Indian summer monsoon (ISM) and westerlies on the atmospheric transport of mercury. The mean  $\pm$  standard deviation concentrations of TGM, PBM and GOM were  $2.55 \pm 0.73$  ng m<sup>-3</sup>,  $37.78 \pm 31.35$  pg m<sup>-3</sup> and  $7.90 \pm 7.89$  pg m<sup>-3</sup>, respectively. A notable seasonal pattern of TGM concentrations was observed with higher concentrations in the beginning and end of the ISM. High TGM concentrations were associated with the transport of dry air that carried regional anthropogenic emissions from both Chinese domestic and foreign sources. Low PBM and GOM levels during the ISM period were attributed to the enhanced wet scavenging. Backward trajectories analysis indicates that high atmospheric mercury (> 2.5 ng m<sup>-3</sup>) concentrations at the SAWRS were influenced by the ISM and westerlies, which carried Hg plumes from Burma, Bengal bay, north India and western China to the SAWRS.

Keywords: Atmospheric mercury, Shangri-La, Long-range transport, Indian summer monsoon.

## 1 Introduction

Mercury (Hg) is a persistent toxic pollutant released to the atmosphere from both anthropogenic sources

1 and natural sources (Pirrone et al., 1996;Pirrone et al., 2001;Carpi and Lindberg, 1998). Due to its volatility,  
2  $\text{Hg}^0$  can be transported over a long distance. Hg interconverts among GEM, GOM and PBM and enters  
3 terrestrial and aquatic ecosystems in remote area via dry and wet deposition (Johansson et al., 2001).  
4 Deposited Hg can be converted into methyl-mercury, a neurotoxin that can bioaccumulates and biomagnify  
5 in the food chain and causes severe public health concerns (Lindqvist, 1991;Sunderland et al., 2009). In the  
6 atmosphere, GEM as the predominant specie accounts for >90% of total Hg (Iverfeldt and Lindqvist,  
7 1986;Schroeder and Munthe, 1998). The global background of GEM in the Northern Hemisphere is 1.5-1.7  
8  $\text{ng m}^{-3}$  and 1.1-1.3  $\text{ng m}^{-3}$  in the Southern Hemisphere and typical concentrations of GEM, GOM and PBM  
9 measured at remote sites are in the ranges of 1–4  $\text{ng m}^{-3}$ , 3–70  $\text{pg m}^{-3}$  and 7–100  $\text{pg m}^{-3}$ , respectively. In  
10 the Arctic, the GEM, PBM and GOM median concentrations are 1.6  $\text{ng m}^{-3}$ , 11.3 and 3.2  $\text{pg m}^{-3}$ ,  
11 respectively. In Antarctica, the mean annual GEM concentration of  $0.93 \pm 0.19 \text{ ng m}^{-3}$  is in good agreement  
12 with recent southern-hemispheric measurements (Iverfeldt and Lindqvist, 1986;Lin and Pehkonen,  
13 1999;Valente et al., 2007;Steffen et al., 2014;Pfaffhuber et al., 2012).

14  
15 Observations of atmospheric Hg are important to understand its global biogeochemical cycling. The  
16 long-term and continuous measurements at remote sites are particularly valuable because such datasets  
17 yield valuable evidence of seasonal trend and transport pattern of a given region (Li and Lee,  
18 2014;Pirrone et al., 2010b;Driscoll et al., 2013). In addition, the data provides evidence for constraining  
19 atmospheric models of Hg. To address the data need, there has been ongoing efforts on establishing  
20 monitoring networks, including the atmospheric mercury network (AMNet) in North America and the  
21 global mercury observation system (GMOS) (Sprovieri et al., 2013;Gay et al., 2013). However, the data at  
22 remote sites are still insufficient to describe the temporal and spatial variability of Hg on a global scale.  
23 Monitoring of speciated atmospheric Hg at remote sites is challenging due to issues related to  
24 instrumental maintenance, accessibility to the sites and other logistic limitations. To date, the availability  
25 of speciated atmospheric Hg observations is inadequate for a thorough understanding of Hg chemical  
26 transport in different regions.

27  
28 Shangri-La is located remotely in the southeastern corner of Tibetan Plateau that extends from  $25^\circ$  to  $45^\circ\text{N}$   
29 and from  $70^\circ$  to  $105^\circ\text{E}$ . This Plateau covers approximately one-quarter of the land area in Mainland China  
30 with a mean elevation of more than 4,000 m above sea level. Given the locations between South and East  
31 Asia, the levels of air pollutants at Shangri-La are potentially influenced by the two largest source regions  
32 in the world. Recently, there is an increasing number of studies highlighting large and increasing Hg  
33 emissions in fast developing countries (Pirrone et al., 2010). China is the largest Hg emitter in the world,  
34 despite its continued effort in Hg emission reduction (Streets et al., 2005;Pacyna et al., 2006). India, with

the world's fourth biggest coal reserve, is another rapidly growing economy whose Hg use and emissions have been increasing over the last few decades (Sharma, 2003). Therefore, it is very significant to study the long-range transport of air pollutants between China and India. Actually, recent assessments suggested that air pollutants emissions in India are of global importance, especially to the Tibetan plateau where the long-range transport of black soot accelerates the glacier melt (Xu et al., 2009). Also, it has been suggested that Hg from South Asia can be transported to Tibetan plateau and deposit on glaciers through Indian summer monsoon and has led to the higher Hg concentrations in the snow packs (Loewen et al., 2007; Zhang et al., 2012). In addition, Hg emissions from the biomass burning in Southeast Asia represent a significant yet uncertain regional source (Pirrone et al., 2010b). In light of these perspectives, Shangri-La appears to be a strategic observational site for studying the long-range transport of Hg emitted from these sources.

In this study, we report the characteristics of the speciated atmospheric Hg at the SAWRS and analyze the potential source regions that contributed to the observed concentrations. In particular, we assess the role of the ISM and the westerlies on the long-range transport of mercury. This study is part of the Global Mercury Observation System (GMOS, <http://www.gmos.eu/>), which is aimed to establish a global mercury monitoring network for ambient concentrations and deposition of mercury through ground-based observational platforms as well as oceanographic aircraft campaigns. The data within the GMOS network will provide valuable global monitoring data for modeling assessment in understanding the global biogeochemical cycling of Hg, and for establishing the global source-receptor relationship of Hg emissions. The program involves 24 partners in the world (Gay et al., 2013).

## **2 Materials and methods**

### **2.1 Measurement site description**

Fig. 1 shows the locations of SAWRS and large Asian cities. The SAWRS (28°01'N, 99°44'E, 3580 m above sea level) is a remote highland site located in Hengduan Mountains area southeast of the Tibetan Plateau. Kawa Karpo and Gongga Mountain both are more than 6000 m a.s.l. and almost 150 Km away from SAWRS. Kawa Karpo is located in the northwest of SAWRS and Gongga Mountain is northeast of SAWRS. SAWRS is near a mountaintop and surrounded by alpine forest and 30 km away from north of Shangri-La city. The closest large point sources are in Kunming city, capital of Yunnan province, which is ~500 km to the southeast. Other large cities (Chengdu, Guiyang and Chongqing) are to east of Shangri-La. India and Bengal are located to the west of SAWRS and Southeast Asia (in this study, it defined as the region 92°E-140°E, 10°S-23°26'N). It has a cold and dry winter that lasts 5 months from late October to April before onset of ISM. Its annual average temperature is 5.1 °C. The average temperature is -1.4 °C

from December to February and 12.4 °C from June to August (Table 1). The population density in Shangri-La is approximately 10 persons per square kilometer. As SAWRS is surrounded by the forest, there are no large-scale industrial activities and fossil fuel consumption in the area. Domestic biofuel burning for cooking and house heating is the main emission source, which release <10 g km<sup>-2</sup>y<sup>-1</sup> of Hg annually (AMAP/UNEP, 2013).

## **2.2 Sampling methods and analysis**

### **2.2.1 Measurements of atmospheric TGM, PBM and GOM**

TGM was monitored between November 2009 and November 2010. The inlet of the heated Teflon sampling tubing for TGM monitoring was set up at 10 m above ground. Measurement of the TGM concentrations was performed with an automated Hg vapor analyzer Tekran Model 2537A (Schroeder et al., 1995). The instrument allows continuous measurement of TGM (GEM+GOM) in ambient air through a cycle of gold trap amalgamation, thermal desorption and detection by cold vapor atomic fluorescence spectrometry ( $\lambda=253.7$  nm). The precision, accuracy, stability of the instrument have been documented extensively (Gay et al., 2013b). Tekran 2537A automatically calibrates for Hg<sup>0</sup> every 25 h using an internal permeation source, which provides approximately 1 pg s<sup>-1</sup> of Hg<sup>0</sup> at 50 °C into a zero air stream. External calibration using Tekran 2505 with manual injections of known concentrations of Hg<sup>0</sup> were performed every 4 months, and the accuracy of the analyzer based on manual injection calibrations is 96.84%. The precision is <2% with a detection limit <0.1 ng m<sup>-3</sup>. PBM in ambient air was removed using a 47mm diameter Teflon filter (pore size 0.2  $\mu$ m). For sampling at the reduced ambient pressure at the site, a lower sampling rate of 0.75 l min<sup>-1</sup> (at standard temperature and pressure, as compared to the typical 1 l min<sup>-1</sup>) at sampling interval of 5 minutes was used during the entire campaign (Fu et al., 2012a).

A denuder-based system was used for sampling PBM and GOM and it was not separated from the TGM measurements using the Hg vapor analyzer. The annular denuder tubes are made of quartz with the inner wall coated with KCl for GOM sampling. An integrated elutriator/acceleration jet and a glass impactor plate capable of removing coarse particles (>2.5  $\mu$ m) were installed before the inlet of the denuder. The PBM and GOM samples were collected continuously and analyzed immediately after the sampling cycle. A fresh glass impactor was replaced when finished one sampling cycle. PBM (defined as Hg associated with particles <2.5  $\mu$ m) was collected on 47-mm diameter quartz filters housed in a Teflon filter holder downstream of the denuder. Each quartz filter was used for one sampling and analysis cycle to prevent from possible deformation and contamination, An unused KCl-coated denuder was also installed in a separate sampling line with an impactor. Once a sampling period was completed, the prepared PBM and GOM sampling lines were installed swiftly for the next sampling cycle. The typical replacement time of

each denuder and filter was very short (almost 10 minutes) (Xiao et al., 1997; Landis et al., 2002; Feng et al., 2000). Both the PBM filter and denuder were positioned vertically. The inlets were located at 1.5 m above ground. An electronic temperature-controlled heating tape was utilized to maintain the temperature of the denuder sampling train at 40–50 °C above the ambient temperature to prevent water condensation over the KCl coating and the wall loss of GOM in the sampling tube (Feng et al., 2000). The sampling flow rate (10 L min<sup>-1</sup>) for both GOM and PBM was controlled by mass-flow controllers (MKS, USA) calibrated to standard atmospheric pressure (0 °C and 1013 mbar). Both PBM and GOM samples were collected every 2h. Using this sampling protocol, twelve GOM and twelve PBM samples can be collected a day.

Analytical procedures for PBM and GOM described by Landis et al. were followed. The GOM and PBM were sampled simultaneously and each analysis cycle was two hours. During each cycle, we had approximately one hour for PBM and GOM detection after thermal pyrolysis and about one hour for preparing the next sampling installation for swift sample switch. The denuder and filter after the air sampling were analyzed immediately. After sampling, Hg contents are determined via pyrolysis followed by Tekran 2537A detection. First, the Tekran 2537A pulled Hg-free air (0.5 L min<sup>-1</sup>) through the trap and denuders for two sample cycles (10 min) to confirm the zero background for ensuring no leaks in the system. Then, the trap and denuders were rapidly heated to 900 and 500 °C using a pyrolyzer for three heating cycles (15 min) to convert PBM and GOM into Hg<sup>0</sup>, which is analyzed by the Tekran 2537A. Typically, a signal of <2 pg was observed in the third heating cycle, which is indistinguishable from the system blank. The PBM traps and denuders were then cooled by a cooling fan in the third step.

Four sampling campaigns were carried out for PBM and GOM measurements: January 17- February 1, May 8-18, July 10-25, and October 28 to November 11, 2010. The selected periods represented four seasonal observations. Before sampling, the denuders were cleaned by pyrolysis to obtain the blank (1.67±0.66 pg, N=12). The quartz filter was cleaned by heating at 850 °C for 30 min. A somewhat higher blank (6.62±2.69 pg, N=20) was observed and used for correcting the PBM concentrations by subtracting the mean blank from the detected concentration.

## **2.2 Meteorological data and backward trajectory calculation**

Meteorological parameters including hourly wind direction (WD), wind speed (WS), air temperature (AT), relative humidity (RH) and rainfall (RF) were continuously monitored by the SAWRS. The Indian monsoon indices (IMI), defined as the difference of 850 hPa zonal winds between a southern region of 5 °N–15 °N, 40 °E–80 °E and a northern region of 20 °N–30 °N, 70 °E–90 °E (Wang and Fan, 1999; Wang et

al., 2001), were obtained online at the Asia-Pacific Air Data-Research Center (<http://apdrc.soest.hawaii.edu/projects/monsoon/realtime-monidx.html>). IMI reflects both the intensity of the tropical westerly monsoon and the lower-tropospheric vorticity anomalies associated with the ISM trough. When IMI of  $> 0$  was observed continuously, the study region was expected to be controlled by ISM. Moreover, the levels of IMI also indicate the intensity of air flow and moisture originated from Indian Ocean to inland. Zero IMI indicates weak air movement. Negative IMI indicates northerly wind that push the air back the Indian Ocean. Five-day backward trajectories were calculated using HYSPLIT (Wang et al., 2009) to assess the long-range transport pathways. The Global Data Assimilation System (GDAS) meteorology archived by the Air Resource Laboratory of National Oceanic and Atmospheric Administration (NOAA) were used as the input (<ftp://arlftp.arlhq.noaa.gov/pub/archives/gdas1/>). The spatial and temporal resolution of the met data is  $1^\circ \times 1^\circ$  and 6-hour, respectively. All back trajectories ended at the SAWRS ( $28^\circ 01' \text{ N}$ ,  $99^\circ 44' \text{ E}$ ) at an arrival height of 500 m above ground. The back trajectories endpoints were recorded at 6-h intervals (02:00, 08:00, 14:00, 20:00 LT).

Cluster analysis was performed to obtain the synoptic transport pattern to the site via HYSPLIT4 (<http://www.arl.noaa.gov/ready/hysplit4.html>). Clustering started by assigning each trajectory to its own cluster, so that there was the same number of clusters as the number of trajectories (i.e., one trajectory in each cluster). For each iteration step, the cluster number was reduced by one as two clusters were merged together. For every combination of trajectory pairs, the cluster spatial variance (SV), defined as the sum of the squared distances between the endpoints of the cluster's component trajectories and the mean of the trajectories in that cluster, was computed. Then the total spatial variance (TSV), defined as the sum of all cluster SV and increased for each additional inclusion of a trajectory, was calculated. The clustering process continued until the TSV suddenly increased by a large value, indicating that the included trajectory in the given iteration was not similar to the included trajectories included in a cluster. Such processes continued until the iterative step just before the large increase in the change of TSV gave the final number of clusters. Once the number of clusters was determined, the endpoints of trajectories in the same cluster were averaged and shown in the map (Wang et al., 2009; Dorling et al., 1992). This cluster analysis has been used proverbially in previous studies (Rozwadowska et al., 2010; Landis et al., 2002; Fu et al., 2012a).

Potential Source Contribution Function (PSCF) was performed to analyze and estimate the influence of the potential sources. The PSCF value for the  $ij_{th}$  cell is then defined as:

$$PSCF_{ij} = \frac{M_{ij}}{N_{ij}} \times W_{ij} \quad (1)$$

Where  $M_{ij}$  represents the number of trajectory endpoints associated with 85 or higher percentiles of the observed TGM,  $N_{ij}$  is the number of endpoints associated with 50 or higher percentiles of the observed TGM. To reduce the effect of small values of  $N_{ij}$ , the PSCF values were multiplied by a weight function  $W_{ij}$  to reflect the uncertainty in the values for these cells (Polissar et al., 2001). The endpoints reaching the ground were excluded in the PSCF calculation. Overall, more than 27,000 endpoints were included for the calculation. The weighting function was defined as:

$$W_{ij} = \begin{cases} 1.0 & n_{ij} > 3Avg \\ 0.7 & 1.5Avg < n_{ij} \leq 3Avg \\ 0.4 & Avg < n_{ij} \leq 1.5Avg \\ 0.2 & n_{ij} \leq Avg \end{cases} \quad (2)$$

where  $Avg$  is the average endpoints in each grid cells.

### 3 Results and discussion

#### 3.1. Distribution and influences of potential regional sources.

The 5-min time series plot of the TGM concentrations is shown in Fig. 2. It ranged from 1.03 to 13.59 ng m<sup>-3</sup> with a mean of 2.55±0.73 ng m<sup>-3</sup> (Table 1). Spikes of high TGM concentrations were observed from late April through early May, from late June through early July and in late September. The three high TGM events occurred during the months when the ISM was prevailing, suggesting the ISM played an important role in the high TGM observed at the SAWRS (more in Section 3.3). The levels of TGM at the SAWRS were much lower than those observed in southwestern Chinese cities (9.7±10.2 ng m<sup>-3</sup> in Guiyang(Fu et al., 2011); 6.74±0.37 ng m<sup>-3</sup> in Chongqing, (Yang et al., 2009)). It was also lower than the background concentrations observed at Mt. Leigong in Guizhou province (2.80 ± 1.51 ng m<sup>-3</sup>, (Fu et al., 2010) and Mt. Gongga in Sichuan province (3.98±1.62 ng m<sup>-3</sup>, (Fu et al., 2008)), but higher than those observed at Mt. Changbai (1.60 ± 0.51 ng m<sup>-3</sup>) in Northeast China and at Mt. Waliguan (WLG) Baseline Observatory (1.98 ± 0.98 ng m<sup>-3</sup>) in Tibetan Plateau (Fu et al., 2012a; Fu et. al, 2012b). The TGM mean concentration at the SAWRS was higher than the global background (1.5-1.7 ng m<sup>-3</sup>) in the Northern Hemisphere and 1.1-1.3 ng m<sup>-3</sup> in the Southern Hemisphere (Lindberg et al., 2007). As a remote background site, the TGM level was mainly influenced by the long-range transport and possibly weak local sources. SAWRS is located between East Asia and South Asia, which are regions with large Hg emissions (Fig. 1). Like some previous studies, the TGM concentrations measured in Korea and Japanese background sites, due to the nearby emission sources and the prevailing westerly wind, were also much higher (4.61 ng m<sup>-3</sup> at Global Atmospheric Watch Station in An-Myun Island of Korea, (Nguyen et al., 2007); 2.04 ng m<sup>-3</sup> at Cape Hedo Station in Okinawa Island of Japan, (Jaffe et al., 2005)). The GEM level of Lulin Atmospheric Background Station in Taiwan (1.73 ng m<sup>-3</sup>) was relatively lower compared to Shangri-La. The mean TGM concentration at SAWRS was higher than the background sites in North

America and Europe (mean  $1.5 \text{ ng m}^{-3}$  from the North America Atmospheric Mercury Network,  $1.4\text{--}1.8 \text{ ng m}^{-3}$  at Mace Head and Zingst of Europe, (Kock et al., 2005; Gay et al., 2013b)). The elevated background level of TGM at the SAWRS is likely caused by strong regional sources in Asia.

The measured TGM concentration exhibited a strong dependence on wind direction. Higher TGM levels ( $2.5\text{--}2.7 \text{ ng m}^{-3}$ ) were frequently associated with northerly winds that carried domestic emissions from West China to the site (Fig. 3). The elevated TGM concentrations ( $2.5\text{--}2.6 \text{ ng m}^{-3}$ ) associated with the southerly winds could be caused by a combination of the regional emission sources in South Asia and Shangri-La County. The wind flows from the east and west gave low TGM concentrations ( $2.3\text{--}2.5 \text{ ng m}^{-3}$ ). The mountain ranges in Shangri-La area run in the north-south direction. The air masses in the east-west direction needed to cross high mountains to reach the SAWRS. When the air flow climb up to Shangri-La, the cumulus process could cause dilution of the air masses and therefore gave lower TGM concentrations (Discussion in section 3.3). Fig. 4 displays the average diurnal trend of WD and WS along with those of TGM, GOM and PBM. The WD was predominantly southerly and changed from southeast (daytime) to southwest (nighttime). WS and TGM were slightly higher in daytime. Since the SAWRS is located at a small mountaintop in Hengduan Mountain area, the terrain could affect local air circulation. The diurnal mountain valley breezes played a significant role in the vertical mixing. During daytime, air near the valley heats up faster than the air at the low altitude. The air mass rapidly rose and brought the Hg trapped in the boundary layer to the monitoring site. At nighttime, the cool air from the high mountains sank and carried the clean air from higher altitude to the site (Fu et al., 2010).

The diurnal pattern of GOM at the SAWRS showed higher concentrations in the afternoon. The relative humidity in the afternoon (14:00~20:00) was 81.47% which was much higher than the night and morning (21:00-13:00) of 58.24%. Meanwhile, the mean GOM concentrations in the afternoon was  $9.22 \text{ pg m}^{-3}$  and the mean GOM of the other time period was  $7.34 \text{ pg m}^{-3}$ . A possible reason is the lower relative humidity in air during the afternoon hours, which decreases the scavenging of GOM into water vapor. Another explanation is the oxidation of  $\text{Hg}^0$  caused by the stronger solar radiation (Lindberg et al., 2002; Goodsite et al., 2004; Fain et al., 2009). The measured PBM did not show a distinct diurnal trend. The observed diurnal pattern of TGM and GOM reflected the influence from local meteorology and from the photochemical productions.

### 3.2 Seasonal patterns of TGM, GOM and PBM

The seasonal statistics of the measured TGM, GOM, PBM, WS, AT, RH and RF are shown in Table 1. The TGM concentration in winter was lower ( $2.28 \text{ ng m}^{-3}$ ) compared to the level in the other three



seasons. At 95% confidence, except spring and winter (Sig.=0.19, by two- tail test) among other season, TGM concentrations are statistically different (Sig. =0.00 for all). In addition, temperature, RH and RF were lower in winter after the retreat of the ISM. During the period, dry air masses, typically associated with high WS, from Tibetan plateau caused the lower observed TGM. The TGM level was highest in spring ( $2.76 \text{ ng m}^{-3}$ ) and peaked in late April (Fig. 2) when RF, AT and RH increased with the onset of the ISM. The high TGM events were caused by the regional transport from the heavily industrial regions in Northwest China (more details in Section 3.3), with a peak observed concentration of  $13.59 \text{ ng m}^{-3}$ . Elevated TGM concentrations (mean= $2.71 \text{ ng m}^{-3}$ ) were higher than the mean value (mean= $2.55 \text{ ng m}^{-3}$ ) and also observed in autumn (Fig. 2), which was associated with lower AT, RH and RF. During the ISM period, the mean TGM level was  $2.51 \text{ ng m}^{-3}$ , which was accompanied by the higher RH and RF.

The variation of the monthly means of TGM, WS, AT and RH is shown in Fig.5. During the ISM season (May to September), a negative correlation between mean TGM concentration and the RH was most evident ( $r=-0.83$ ,  $p=0$ ). The likely reason is that during ISM period, clean Indian Ocean air mass and water vapor can flow into inland and lead to high RH and low TGM. In addition, during rainfall formation, cumulus process also could cause dilution of Hg in atmosphere. The RH itself may not have a significant effect on GEM. However, we would like to point out that high RH can decrease the Hg emission from surface and enhance wet deposition of Hg, which could contribute a relative lower TGM level (Seo et al., 2012; Poissant and Casimir, 1998). The TGM concentrations in the late autumn and winter months were also relatively lower than other months (except August) due to the more stagnant winds that limited the regional transport. The variability of WS was not significant during the campaign and not likely to cause the observed seasonal variation.

Both GOM (Fig. 6a) and PBM (Fig. 6b) exhibited substantial seasonal variability. Except spring and summer (Sig.=0.53, by two- tail test), among other season, GOM concentrations differences are statistically significant (Sig. =0.00 for all). The GOM concentrations were higher in autumn (mean:  $7.46 \text{ pg m}^{-3}$ ) and peaked in winter ( $14.48 \text{ pg m}^{-3}$ ), about five times of the mean level in spring ( $2.82 \text{ pg m}^{-3}$ ). A similar pattern was observed at Storm Peak Laboratory (3220 m a.s.l.) in the Rocky Mountains of USA (Fain et al., 2009). The elevated GOM has been attributed to the much lower RH, which was also the case at the SAWRS. However, importantly, the SAWRS was built in the heavily forested valleys, and the air flow in Shangri-La was dominated by westerlies which perform the highest wind speed of  $2.34 \text{ m/s}$  in the winter, especially the south tributary of westerlies, the air flow from south and southeast Asia could transport to SAWRS along with the valleys, therefore the GOM level at SAWRS could be effected primarily by local photochemical transformation or meteorological conditions, compared to the

1 subsidence of free troposphere.

2 The PBM concentrations were high in autumn, with a peak concentration of  $59.14 \text{ pg m}^{-3}$ . Moreover, only  
3 concentration in autumn is statistically different from other seasons (Sig.=0 for all, at 0.05 level). In the  
4 absence of volcanic sources, PBM concentrations are usually associated with Hg emission from  
5 anthropogenic activities (Friedli et al., 2003; Finley et al., 2009). Therefore the local anthropogenic  
6 sources could contribute to the high PBM level at SAWRS.

7  
8 Additionally, Back-trajectory analysis suggested that the high PBM during the campaign was associated  
9 with the air mass from Burma, Vietnam, Laos, Kampuchea and Thailand. Due to increasing consumptions  
10 of resource, these areas are expected to release remarkable Hg to the atmosphere. This can be supported  
11 by a previous study which showed highly elevated TGM concentrations in the Southeast Asia (Sheu et al.,  
12 2013). Given the frequent air flows from the Southeast Asia to southwest china, elevated atmospheric Hg  
13 concentrations could be likely resulted from Hg emissions in Southeast Asia. The GOM and PBM levels  
14 were relatively lower from late spring to summer during the ISM period. PBM and GOM concentrations  
15 were found to be negatively correlated with RF ( $r = -0.142$  and  $-0.186$ , respectively). This may indicate  
16 the role of increasing wet scavenge processes that played in the lower GOM and PBM concentrations  
17 during the ISM period.

### 19 **3.3 Air mass back trajectories analysis**

20 Fig. 7 shows the time series of WD, RF and TGM along with the calculated Indian Monsoon Index (IMI)  
21 during ISM months (April through September). The values IMI represent seasonal rainfall anomalies, and  
22 are highly correlated with the transport associated with ISM (Wang et al., 2001). An IMI value of  $>0$   
23 suggests air transport from the south Asia. Such a transport pattern was consistent from late May through  
24 mid-September. During this period, WD was mainly from the southeast. The precipitation events occurred  
25 predominantly during the ISM period ( $\text{IMI} > 1$ ). However the high Hg peaks were not appeared frequently,  
26 this could be related to the location and topography of Shangri-La, which is in the Tibetan Plateau, when  
27 the air flow from south and southeast Asia climb up to Shangri-La, the air flow will move speed  
28 slackened and formed cloud, perhaps the cumulus process could cause dilution of the air masses, and we  
29 can see that the wind speed was low of  $1.63 \text{ m/s}$  in summer. Therefore the wet deposition of Hg mainly  
30 happened before the air flow arrive in Shangri-La. Additionally, in ISM period, the wet air masses were  
31 mainly form Indian ocean and could dilute the mercury concentration in air. This two reasons might be  
32 explain why there were not many high Hg peaks appeared in ISM period. The three high Hg events in  
33 Fig.7 were further analyzed with the back trajectories of the air masses.

Fig. 8 shows the back trajectories during April 26 to May 2 (first block in Fig.7). The air masses originated from Siberia moved over north Xinjiang, east Qinghai, Gansu and west Sichuan provinces. These are industrial areas in Northwest China that have major Hg emission sources (Fig. 1), mainly from the use of fossil fuels (Wang et al., 2000;Fu et al., 2008;Fu et al., 2012a;Fu et al., 2012c). From June 19 to July 2 (second block in Fig.7), the air mass was mainly from the Bengal and Burma because of the ISM (Fig. 9), suggesting trans-boundary transport possibly caused by anthropogenic sources in the region and the industrial cities (Fig. 1) such as Yangon (Burma's capital), Bangkok (Thailand's capital). Fig. 10 shows the back trajectories during September 17-22 (third block in Fig.7). Most trajectories in this period were very short and reached ground level (the endpoints were not shown after grounding). The elevated concentration in this event was likely to be contributed by the domestic emissions.

### 3.4 Cluster analysis of back trajectories

The back trajectories arriving at the SAWRS over the study period were grouped into four clusters (Fig. 11) to show the synoptic air transport pattern and understand the regional transport pathways. Cluster analysis was performed to obtain the transport pattern. The cluster spatial variance (CSV) was calculated as the sum of the spatial variance of all trajectories within its cluster and the total spatial variance (TSV) was calculated as the sum of the CSV over all clusters. The cluster analysis in this study showed the prevailing air flow arriving at Shangri-La and facilitated the identification of potential Hg source regions. Cluster 1 (19%) represented air masses originating from North Africa and had the highest height of trajectories (Fig. 12). This indicated that the air masses moved fast and passing through Middle Asia and then over Xinjiang, Qinghai provinces and Tibet of China. Cluster 2 (18%) and Cluster 3 (59%) were the air masses originating from South Asia and north India. Cluster 4, although relatively infrequent (4%), was associated with the highest TGM concentrations (mean =  $3.9 \text{ ng m}^{-3}$ ) due to the passing of air masses over known source regions in Sichuan province. The area has a high background TGM level ( $3.98 \text{ ng m}^{-3}$ ) caused by industrial and domestic coal combustion, smelting industries, cement production, and biomass burning (Fu et al., 2008). Air masses of Cluster 3, the most frequent transport sector, also had high TGM concentrations (mean =  $2.6 \text{ ng m}^{-3}$ ). Cluster 3 had the lowest height of trajectories and could easily transport mercury emitted from surface to SAWRS (Fig. 12). The most likely source region was South Asia's biomass burning activities. TGM concentration ( $2.4 \text{ ng m}^{-3}$ ) associated with Cluster 2, also frequent, could be linked to transport of atmospheric Hg from Pakistan and northern India including New Delhi. Anthropogenic Hg emission in India has drawn increasing attention in recent years (Qureshi et al., 2013). Air masses in Cluster 1 showed the lowest mean TGM concentration ( $2.3 \text{ ng m}^{-3}$ ) and originated from the west Asia. This cluster was also associated with high wind speed caused by the strong prevailing westerly in cold season.

The back trajectories of the upper quartile of TGM observations at the SAWRS were displayed in Fig. 13. It is clear that the frequent transport from South and Southeast Asia during summer and autumn dominated the high TGM observation at the SAWRS. The winter and spring trajectories originated from Middle Asia, North Africa and Mediterranean and then passed through Middle East and Tibetan plateau were more likely caused by the domestic sources in China. The relatively shorter trajectories from South Asia suggest that the transport from this sector may be caused by mild air movement of the ISM (summer and early autumn). For lower quartile of TGM observations, most back trajectories show the same directions as upper quartile of TGM observations (Fig.14). However, back trajectories associated with the lower quartile of TGM observations were comparatively higher and longer. These indicated that the air masses were traveling well above the planetary boundary layer where ground based emission may not be incorporated in the air masses during low TGM period.

The potential sources and transport pathways of TGM at the SAWRS is shown in Fig. 15 as a PSCF distribution plot. It suggests that Burma, Bengal bay, north India, west Sichuan Province and west Yunnan Province were likely the source regions. Trans-boundary transport of aerosols released from forest fires and agricultural burning in the Bengal bay region had been attributed to the seasonal changes of air movement caused by monsoons (Reid et al., 2013). Our PSCF analysis seems to support such trans-boundary transport events. Two high PSCF values in India were also identified, including the surrounding areas of New Delhi and Uttar Pradesh. Both are considered large urban regions with high industrial activities. Domestic source regions in Sichuan and Yunnan were also identified. The atmospheric mercury concentrations observed at the SAWRS were influenced by both domestic and foreign emission sources and highly dependent on the seasonal winds that carry the air masses from different source regions to the observational site.

#### **4 Conclusions**

Measurements of TGM, GOM and PBM at the SAWRS, a high-elevation background site in Tibetan plateau of China, were investigated from November 2009 to November 2010. The mean TGM ( $2.55 \text{ ng m}^{-3}$ ) was higher than the background values observed in the Northern Hemisphere ( $1.5\text{--}1.7 \text{ ng m}^{-3}$ ). Significant seasonal variability in TGM concentration was observed, which was attributed to meteorological factors including WD, WS, RH and RF as well as long-range transport caused by the ISM and westerlies. The ISM enhanced TGM pollution at SAWARS occasionally by carrying TGM-enriched air from South Asia, while the moist air during the ISM significantly decreased GOM and PBM levels. The westerlies carried the Hg emission from the domestic sources in China to the site. The TGM

concentration was higher during the daytime mainly due to upslope transport of polluted air from lower-altitude areas around the site. High GOM levels were highly correlated with dry air in autumn and winter, and in-situ photochemical production might have contributed to occasionally high GOM concentration. Back trajectory analysis showed that the SAWRS was strongly affected by ISM and westerlies during the study period, which carried Hg-enriched air masses to the site from South Asia and industrial regions in China in different seasons.

**Acknowledgements.** This work is supported by National “973” Program (2013CB430003), and Natural Science Foundation of China (41273145, 41003051). We also thank the staff of Meteorological Bureau of Diqing Tibetan Autonomous Prefecture for field sampling assistance.

## References

- AMAP/UNEP: Technical Background Report for the Global Mercury Assessment 2013, in, Arctic Monitoring and Assessment Programme, Oslo, Norway/UNEP Chemicals Branch Geneva, Switzerland, 2013.
- Carpi, A., and Lindberg, S. E.: Application of a Teflon (TM) dynamic flux chamber for quantifying soil mercury flux: Tests and results over background soil, *Atmospheric Environment*, 32, 873-882, 1998.
- Dorling, S. R., Davies, T. D., and Pierce, C. E.: Cluster-Analysis - a Technique for Estimating the Synoptic Meteorological Controls on Air and Precipitation Chemistry - Results from Eskdalemuir, South Scotland, *Atmospheric Environment Part a-General Topics*, 26, 2583-2602, 1992.
- Driscoll, C. T., Mason, R. P., Chan, H. M., Jacob, D. J., and Pirrone, N.: Mercury as a Global Pollutant: Sources, Pathways, and Effects, *Environmental science & technology*, 47, 4967–4983, doi:10.1021/es305071v, 2013.
- Fain, X., Obrist, D., Hallar, A., Mccubbin, I., and Rahn, T.: High levels of reactive gaseous mercury observed at a high elevation research laboratory in the Rocky Mountains, *Atmospheric Chemistry and Physics*, 9, 8049-8060, doi:10.5194/acp-9-8049-2009, 2009.
- Feng, X., Sommar, J., Gårdfeldt, K., and Lindqvist, O.: Improved determination of gaseous divalent mercury in ambient air using KCl coated denuders, *Fresenius' journal of analytical chemistry*, 366, 423-428, 2000.
- Finley, B., Swartzendruber, P., and Jaffe, D.: Particulate mercury emissions in regional wildfire plumes observed at the Mount Bachelor Observatory, *Atmospheric Environment*, 43, 6074-6083, 2009.
- Friedli, H., Radke, L., Prescott, R., Hobbs, P., and Sinha, P.: Mercury emissions from the August 2001 wildfires in Washington State and an agricultural waste fire in Oregon and atmospheric mercury budget estimates, *Cy.*, 17, 1039, doi:10.1029/2002GB001972, 2003.
- Fu, X., Feng, X., Zhu, W., Wang, S., and Lu, J.: Total gaseous mercury concentrations in ambient air in the eastern slope of Mt. Gongga, South-Eastern fringe of the Tibetan plateau, China, *Atmospheric Environment*, 42, 970-979, 2008.
- Fu, X., Feng, X., Dong, Z., Yin, R., Wang, J., Yang, Z., and Zhang, H.: Atmospheric gaseous elemental mercury(GEM) concentrations and mercury depositions at a high-altitude mountain peak in south China, *Atmospheric Chemistry and Physics*, 10, 2425-2437, doi:10.5194/acp-10-2425-2010, 2010.
- Fu, X., Feng, X., Qiu, G., Shang, L., and Zhang, H.: Speciated atmospheric mercury and its potential source in Guiyang, China, *Atmospheric Environment*, 45, 4205-4212, 2011.
- Fu, X., Feng, X., Liang, P., Zhang, H., Ji, J., and Liu, P.: Temporal trend and sources of speciated atmospheric mercury at Waliguan GAW station, Northwestern China, *Atmospheric Chemistry and Physics*, 12, 1951-1964, doi:10.5194/acp-12-1951-2012, 2012a.

1 Fu, X., Feng, X., Shang, L., Wang, S., and Zhang, H.: Two years of measurements of atmospheric total  
2 gaseous mercury (TGM) at a remote site in Mt. Changbai area, Northeastern China, *Atmospheric*  
3 *Chemistry and Physics*, 12, 4215-4226, doi:10.5194/acp-12-4215-2012, 2012b.

4 Fu, X., Feng, X., Sommar, J., and Wang, S.: A review of studies on atmospheric mercury in China,  
5 *Science of the Total Environment*, 421, 73-81, 2012c.

6 Gay, D. A., Schmeltz, D., Prestbo, E., Olson, M., Sharac, T., and Tordon, R.: The Atmospheric Mercury  
7 Network: measurement and initial examination of an ongoing atmospheric mercury record across North  
8 America, *Atmospheric Chemistry and Physics*, 13, 11339-11349, 2013.

9 Goodsite, M. E., Plane, J., and Skov, H.: A theoretical study of the oxidation of Hg<sup>0</sup> to HgBr<sub>2</sub> in the  
10 troposphere, *Environmental science & technology*, 38, 1772-1776, 2004.

11 Graydon, J. A., St. Louis, V. L., Lindberg, S. E., Hintelmann, H., and Krabbenhoft, D. P.: Investigation of  
12 mercury exchange between forest canopy vegetation and the atmosphere using a new dynamic chamber,  
13 *Environmental science & technology*, 40, 4680-4688, 2006.

14 Iverfeldt, Å., and Lindqvist, O.: Atmospheric oxidation of elemental mercury by ozone in the aqueous  
15 phase, *Atmospheric Environment* (1967), 20, 1567-1573, 1986.

16 Jaffe, D., Prestbo, E., Swartzendruber, P., Weiss-Penzias, P., Kato, S., Takami, A., Hatakeyama, S., and  
17 Kajii, Y.: Export of atmospheric mercury from Asia, *Atmospheric Environment*, 39, 3029-3038, 2005.

18 Johansson, K., Bergbäck, B., and Tyler, G.: Impact of atmospheric long range transport of lead, mercury  
19 and cadmium on the Swedish forest environment, *Water, Air, & Soil Pollution: Focus*, 1, 279-297, 2001.

20 Kock, H., Bieber, E., Ebinghaus, R., Spain, T., and Thees, B.: Comparison of long-term trends and  
21 seasonal variations of atmospheric mercury concentrations at the two European coastal monitoring  
22 stations Mace Head, Ireland, and Zingst, Germany, *Atmospheric Environment*, 39, 7549-7556, 2005.

23 Landis, M. S., Stevens, R. K., Schaedlich, F., and Prestbo, E. M.: Development and characterization of an  
24 annular denuder methodology for the measurement of divalent inorganic reactive gaseous mercury in  
25 ambient air, *Environmental Science & Technology*, 36, 3000-3009, 2002a.

26 Landis, M. S., Vette, A. F., and Keeler, G. J.: Atmospheric mercury in the Lake Michigan basin: Influence  
27 of the Chicago/Gary urban area, *Environmental Science & Technology*, 36, 4508-4517, 2002b.

28 Li, J., and Lee, S. M.: Progress of Global Atmospheric Mercury Field Observations, *Journal of Clean*  
29 *Energy Technologies*, 2, 252-257, doi:10.7763/JOCET.2014.V2.135, 2014..

30 Lin, C. J., and Pehkonen, S. O.: The chemistry of atmospheric mercury: a review, *Atmospheric*  
31 *Environment*, 33, 2067-2079, 1999.

32 Lindberg, S.: Forests and the global biogeochemical cycle of mercury: the importance of understanding  
33 air/vegetation exchange processes, in: *Global and Regional Mercury Cycles: Sources, Fluxes and Mass*  
34 *Balances*, Springer, 359-380, 1996.

35 Lindberg, S., Bullock, R., Ebinghaus, R., Engstrom, D., Feng, X., Fitzgerald, W., Pirrone, N., Prestbo, E.,  
36 and Seigneur, C.: A synthesis of progress and uncertainties in attributing the sources of mercury in  
37 deposition, *AMBIO: A Journal of the Human Environment*, 36, 19-33, 2007.

38 Lindberg, S. E., Brooks, S., Lin, C.-J., Scott, K. J., Landis, M. S., Stevens, R. K., Goodsite, M., and  
39 Richter, A.: Dynamic oxidation of gaseous mercury in the Arctic troposphere at polar sunrise,  
40 *Environmental science & technology*, 36, 1245-1256, 2002.

41 Lindqvist, O.: Mercury in the Swedish environment: recent research on causes, consequences and  
42 corrective methods, *Water, air and soil pollution*, 55, 1-261, 1991.

43 Loewen, M., Kang, S., Armstrong, D., Zhang, Q., Tomy, G., and Wang, F.: Atmospheric transport of  
44 mercury to the Tibetan Plateau, *Environmental Science & Technology*, 41, 7632-7638, 2007.

45 Nguyen, H. T., Kim, K. H., Kim, M. Y., Hong, S., Youn, Y. H., Shon, Z. H., and Lee, J. S.: Monitoring of  
46 atmospheric mercury at a global atmospheric watch (GAW) site on An-Myun Island, Korea, *Water, Air, &*  
47 *Soil Pollution*, 185, 149-164, 2007.

48 Obrist, D., Hallar, A., McCubbin, I., Stephens, B. B., and Rahn, T.: Atmospheric mercury concentrations  
49 at Storm Peak Laboratory in the Rocky Mountains: Evidence for long-range transport from Asia,  
50 boundary layer contributions, and plant mercury uptake, *Atmospheric Environment*, 42, 7579-7589, 2008.

51 Pacyna, E. G., Pacyna, J. M., Steenhuisen, F., and Wilson, S.: Global anthropogenic mercury emission

inventory for 2000, *Atmospheric Environment*, 40, 4048-4063, 2006.

Poissant, L., and Casimir, A.: Water-air and soil-air exchange rate of total gaseous mercury measured at background sites, *Atmospheric Environment*, 32, 883-893, 1998.

Pfaffhuber, K., Berg, T., Hirdman, D., and Stohl, A.: Atmospheric mercury observations from Antarctica: seasonal variation and source and sink region calculations, *Atmospheric Chemistry and Physics*, 12, 3241-3251, 2012.

Pirrone, N., Keeler, G. J., and Nriagu, J. O.: Regional differences in worldwide emissions of mercury to the atmosphere, *Atmospheric Environment*, 30, 2981-2987, 1996.

Pirrone, N., Costa, P., Pacyna, J., and Ferrara, R.: Mercury emissions to the atmosphere from natural and anthropogenic sources in the Mediterranean region, *Atmospheric Environment*, 35, 2997-3006, 2001.

Pirrone, N., Cinnirella, S., Feng, X., Finkelman, R., Friedli, H., Leaner, J., Mason, R., Mukherjee, A., Stracher, G., and Streets, D.: Global mercury emissions to the atmosphere from anthropogenic and natural sources, *Atmospheric Chemistry and Physics*, 10, 5951-5964, doi:10.5194/acp-10-5951-2010, 2010.

Polissar, A. V., Hopke, P. K., and Harris, J. M.: Source regions for atmospheric aerosol measured at Barrow, Alaska, *Environmental science & technology*, 35, 4214-4226, 2001.

Qureshi, A., Vadenbo, C., and Hellweg, S.: Anthropogenic mercury flows in India and impacts of emission controls, *Environmental science & technology*, 47, 8105-8113, doi:10.1021/es401006k, 2013..

Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J., Campbell, J. R., Christopher, S. A., Di Girolamo, L., and Giglio, L.: Observing and understanding the Southeast Asian aerosol system by remote sensing: An initial review and analysis for the Seven Southeast Asian Studies (7SEAS) program, *Atmospheric Research*, 122, 403-468, 2013.

Rozwadowska, A., Zielinski, T., Petelski, T., and Sobolewski, P.: Cluster analysis of the impact of air back-trajectories on aerosol optical properties at Hornsund, Spitsbergen, *Atmospheric Chemistry and Physics*, 10, 877-893, 2010.

Seo, Y.-S., Han, Y.-J., Choi, H.-D., Holsen, T. M., and Yi, S.-M.: Characteristics of total mercury (TM) wet deposition: scavenging of atmospheric mercury species, *Atmospheric Environment*, 49, 69-76, 2012.

Schroeder, W., Keeler, G., Kock, H., Roussel, P., Schneeberger, D., and Schaedlich, F.: International field intercomparison of atmospheric mercury measurement methods, *Water, Air, & Soil Pollution*, 80, 611-620, 1995.

Schroeder, W. H., and Munthe, J.: Atmospheric mercury--an overview, *Atmospheric Environment*, 32, 809-822, 1998.

Schwesig, D., and Krebs, O.: The role of ground vegetation in the uptake of mercury and methylmercury in a forest ecosystem, *Plant and Soil*, 253, 445-455, 2003.

Sharma, D. C.: Concern over mercury pollution in India, *The Lancet*, 362, 1050, 2003.

Sheu, G.-R., Lin, N.-H., Lee, C.-T., Wang, J.-L., Chuang, M.-T., Wang, S.-H., Chi, K. H., and Ou-Yang, C.-F.: Distribution of atmospheric mercury in northern Southeast Asia and South China Sea during Dongsha Experiment, *Atmospheric Environment*, 78, 174-183, 2013.

Sprovieri, F., Gratz, L., and Pirrone, N.: Development of a Ground-Based Atmospheric Monitoring Network for the Global Mercury Observation System (GMOS), *E3S Web of Conferences*, 17007, Rome, Italy, 22-27 September 2012, doi:10.1051/e3sconf/20130117007, 2013.

Steffen, A., Bottenheim, J., Cole, A., Ebinghaus, R., Lawson, G., and Leaitch, W.: Atmospheric mercury speciation and mercury in snow over time at Alert, Canada, *Atmospheric Chemistry and Physics*, 14, 2219-2231, doi: 10.5194/acp-14-2219-2014, 2014.

Streets, D. G., Hao, J., Wu, Y., Jiang, J., Chan, M., Tian, H., and Feng, X.: Anthropogenic mercury emissions in China, *Atmospheric Environment*, 39, 7789-7806, 2005.

Sunderland, E. M., Krabbenhoft, D. P., Moreau, J. W., Strode, S. A., and Landing, W. M.: Mercury sources, distribution, and bioavailability in the North Pacific Ocean: Insights from data and models, *Cy.*, 23, GB2010, doi:10.1029/2008GB003425, 2009.

Valente, R. J., Shea, C., Lynn Humes, K., and Tanner, R. L.: Atmospheric mercury in the Great Smoky Mountains compared to regional and global levels, *Atmospheric Environment*, 41, 1861-1873, 2007.

Wang, B., and Fan, Z.: Choice of South Asian summer monsoon indices, *Bulletin of the American*

Meteorological Society, 80, 629-638, 1999.  
 Wang, B., Wu, R., and Lau, K.: Interannual Variability of the Asian Summer Monsoon: Contrasts between the Indian and the Western North Pacific-East Asian Monsoons\*, Journal of Climate, 14, 4073-4090, 2001.  
 Wang, Q., Shen, W., and Ma, Z.: Estimation of mercury emission from coal combustion in China, Environmental science & technology, 34, 2711-2713, 2000.  
 Wang, Y. Q., Zhang, X. Y., and Draxler, R. R.: TrajStat: GIS-based software that uses various trajectory statistical analysis methods to identify potential sources from long-term air pollution measurement data, Environmental Modelling & Software, 24, 938-939, 2009.  
 Xiao, Z., Sommar, J., Wei, S., and Lindqvist, O.: Sampling and determination of gas phase divalent mercury in the air using a KCl coated denuder, Fresenius' journal of analytical chemistry, 358, 386-391, 1997.  
 Xu, B., Cao, J., Hansen, J., Yao, T., Joswila, D. R., Wang, N., Wu, G., Wang, M., Zhao, H., and Yang, W.: Black soot and the survival of Tibetan glaciers, Proceedings of the National Academy of Sciences, 106, 22114-22118, 2009.  
 Yang, Y., Chen, H., and Wang, D.: Spatial and temporal distribution of gaseous elemental mercury in Chongqing, China, Environmental monitoring and assessment, 156, 479-489, 2009.  
 Zhang, Q., Huang, J., Wang, F., Mark, L., Xu, J., Armstrong, D., Li, C., Zhang, Y., and Kang, S.: Mercury Distribution and Deposition in Glacier Snow over Western China, Environmental Science & Technology, 46, 5404-5413, 2012.


---




**Table 1:** The seasonal statistics of the measured TGM、GOM、PBM、WS、AT、RH and RF from November 2009 through November 2010 at the SAWRS site. N is the number of samples.

		TGM (ng m <sup>-3</sup> )	GOM (pg m <sup>-3</sup> )	PBM (pg m <sup>-3</sup> )	AT (°C)	RH (%)	WS (m s <sup>-1</sup> )	RF (mm)
Spring Mar–May	Mean	2.76	2.82	32.82	4.85	78.06	1.98	34.07
	±St.Dev	0.87	3.18	25.24	3.32	12.88	0.63	
	Median	2.6	1.05	26.93	4.05	79.5	1.9	
	Range	1.43-13.59	0.33-16.69	0.48-112.82	-0.4-13	37-98	0.7-3.4	
	N	18706	54	54	2160	2160	2160	90
Summer Jun–Aug	Mean	2.51	3.79	25.46	12.37	85.97	1.63	183.97
	±St.Dev	0.71	3.80	18.37	1.59	7.91	0.61	
	Median	2.35	2.8	22.14	12.4	87	1.5	
	Range	1.05-6.11	0.24-18.81	1.50-86.97	7.2-15.6	65-99	0.5-3.4	
	N	17662	90	90	2280	2280	2280	95
Autumn Sep–Nov	Mean	2.71	7.46	59.14	6.25	84.46	1.5	38.17
	±St.Dev	0.79	6.18	39.31	3.61	8.39	0.44	
	Median	2.59	5.23	45.83	6.3	84.5	1.5	
	Range	1.03-6.99	0.44-31.09	10.93-185.06	-0.3-11.7	63-99	0.4-2.66	
	N	13161	106	106	2280	2280	2280	95
Winter Dec–Feb	Mean	2.28	14.48	21.22	-1.37	54.68	2.34	0
	±St.Dev	0.42	9.18	15.86	1.64	16.98	0.72	
	Median	2.22	12.66	15.23	-1.11	58	2.25	
	Range	1.18-4.51	0.6-39.85	2.01-65.9	-6.6-1.6	19-84	0.9-4.3	
	N	20003	98	98	2256	2256	2256	94
Total	Mean	2.55	7.90	37.78	5.57	75.97	1.86	
	±St.Dev	0.73	7.89	31.35	5.57	16.81	0.68	
	Median	2.39	4.78	30.42	5.1	79	1.8	
	Range	1.03-13.59	0.24-39.85	0.48-185.59	-6.6-15.6	19-99	0.4-4.3	
	N	69532	348	348	8976	8976	8976	374


1 **Fig. 1:** Map showing the location of SAWRS, anthropogenic Hg emissions ( $\text{g km}^{-2} \text{y}^{-1}$ ) and major cities in  
 2 Asia (AMAP/UNEP, 2013).



**Fig. 2:** The 5-min time series distribution of TGM concentrations at SAWRS in the campaign, spikes of high TGM concentrations were observed from late April through early May, from late June through early July and in late September, respectively.




1 **Fig. 3:** The concentration wind rose at SAWRS, TGM concentration exhibited a strong dependence on  
 2 wind direction. Higher TGM levels were frequently associated with northerly winds.




3  
4  
5

1 **Fig. 4:** The diurnal variation of TGM, GOM and PBM with WD,WS and RH.




2

1 **Fig. 5:** The variation of the monthly means of TGM, WS, AT and RH. High TGM concentration in ISM  
2 period with high AT and RH.



3  
4


1 **Fig. 6:** Seasonal variability of GOM and PBM during four sampling campaigns.



2


3

**Fig. 7:** Time series of WD, RF and TGM and the calculated Indian Monsoon Index (IMI) from April to September 2010. An IMI value  $>0$  indicates air movement from south toward north, which is also associated with greater precipitation. WD measured at the SAWRS was mostly from the southeast. Zero IMI indicates weak air movement. Negative IMI indicates northerly wind that push the air back the Indian Ocean. Highlighted in red represent the causes of these events are not only impacted by local airflow but also long-term transportation; while blue just represent the local airflow impact.






**Fig. 8:** The back trajectories of air masses from April 26 to May 2. The high pressure mean that the height of trajectories are low, therefore the air masses can easily transport mercury emitted from surface to SAWRS.




1 **Fig. 9:** The back trajectories of air mass from June19 to July 2.




2  
3

1 **Fig. 10:** The back trajectories of air mass from September 17 to 22. The trajectory points reaching the  
2 ground level are not shown in the figure.




3  
4

**Fig. 11:** The back trajectories arriving at the SAWRS over the study period were grouped into four clusters. Most of air masses originated from Southeast Asia (Cluster 3, 59%), considerable air masses originated from North Africa and south Asia (Cluster 1, 19%; Cluster 2, 18%), only 4% of air masses originated from Middle Asia and Russia.




1    **Fig. 12:** The three-dimensional height of all the clusters from different direction at SAWRS.




2  
3  
4  
5

**Fig. 13:** The back trajectories of the upper quartile of TGM observations at the SAWRS in the four seasons. The air masses with high Hg from South and Southeast Asia in summer and autumn, while some air masses with high Hg were from northwest China. The rest of the air masses were from west Asia, Middle Asia and Tibetan plateau in spring and winter.





**Fig. 14:** The back trajectories associated with the lower quartile of TGM observations in the four seasons. The air masses with low Hg from South and Southeast Asia in autumn and summer. The rest of the air masses were from west Asia, Middle Asia and Tibetan plateau.



**Fig. 15:** The potential sources and transport pathways of TGM at the SAWRS identified by the PSCF. Burma, Bengal bay, north India, western Sichuan Province and western Yunnan Province were likely the potential source regions for Shangri-La.

