

Dear Editor:

Thank you for your feedback. We have revised and clarified our responses to make it more clearer that the reviewers' comments have been addressed and the text modifications corresponding to the points raised.

All of the responses have lead to changes in the manuscript text and we have highlighted the specific changes in the responses.

We have added the various evaluations of the model for Central Asia, North America and the Arctic region, in terms of transport/meteorology and aerosols using the same model configuration. The transport is indeed good by all the measures that we have evaluated and is fit for purpose for the general analysis that we have carried out in this paper. The aerosol predictions have appreciable skill but have limitations. We have modified the closing statement as follows.

“These studies indicate that the simulated transport patterns are able to capture the main important features of the hemispheric flows impacting Central Asia, North America and the Arctic region, at the scale discussed. The prediction of aerosol mass and composition at the hemispheric scales are able to capture important aspects of horizontal gradients and variability, but has considerably higher uncertainties associated largely with emission estimates (in particular forest fires/biomass burning and natural dust emissions) and wet removal processes (Bates et al., 2006).”

Please find the updated response to the reviewers' comments included here.

We thank the two anonymous reviewers for their careful review of our manuscript. We have addressed all the issues/concerns raised by reviewers. Below is our response that addresses the specific comments from each of the two reviewers. [Our response (in italicized font) is preceded by the reviewers' comments for clarity]. Changes in the paper text in response to reviewers' comments are indicated by “**green**” highlighted text.

**Interactive comment on “Source sector and region contributions to BC and PM_{2.5} in Central Asia” by S. Kulkarni et al.
Anonymous Referee #1**

This manuscript describes an effort to quantify concentrations and sources of PM in Central Asia. This region appears to not be specifically target by air quality model applications, so this work seems to be a good contribution in that sense. Regional contributions were obtained using the simple “zero-out” method. Future emissions scenarios were also considered.

I suggest the following items for the authors to consider:

1) How are the winter heating emissions currently quantified? This would be good to know since the authors point to this as one of the reasons for underestimations in the PM concentrations.

Authors' Response: *Seasonal variations of residential heating/cooking emissions were not included as the base inventory did not have this information for CA as little information is available for this region. This has been explicitly stated in section 2.3.1.*

2) The BC surface mean values was found to be around 0.1 ug/m³. That seems very low. Do you have confidence in your apportionment models at such low values?

Authors' Response:

We thank the reviewer for this comment. This modeling system has shown appreciable skill in capturing the large gradients and spatial patterns in regional distribution of BC over the areas surrounding CA as evident in figure 4a. In addition, this particular modeling system (i.e. same domain configuration and settings) has been used to characterize the summertime sectoral and regional contributions to BC over continental US and identified the residential and biomass burning emissions outside CONUS were among the major source categories contributing to BC levels over the region (Huang et al., 2012).

Since the surface BC and the associated source apportionment estimates are based on the chemical transport model simulations in this study, we are confident that we can track relative source contributions at these levels. We have extensively added more model evaluation to the text in Section 3.

3) It is not clear to me why the authors chose to have different source regions for anthropogenic emissions, biomass burning emissions, and dust emissions instead. Perhaps the value gained in the analysis overcomes the confusion, but it is not clear.

Authors' Response:

We have included the following text in section 2.4 to clarify this information.

“The CA region is strategically located and aerosols at these sites are shown to reflect impact of varied dust (including CA, Western China, Africa and Middle East), biomass burning (including Europe, Siberia, South Asia and North America) and anthropogenic sources (including Europe, South, East and Central Asia, and Russia) depending on the time period.

So we selected different regions for the anthropogenic, dust and biomass burning in order to capture the impact and the associated seasonality of these main regions on the regional distribution of aerosols in this region. We have more anthropogenic regions to better understand how regional changes in emissions may impact CA PM levels.”

4) Why were residential and transportation emissions portioned between model layers 1 and 2? 10% seems like a lot to inject higher up. How high was level 2?

Authors' Response: *Residential and transportation emissions are considered near surface emissions so we portioned them from surface to ~ 100m (AGL altitude of level 2). We have also included this information in section 2.3.1*

5) How were back (and forward) trajectories calculated? This does not seem to be described in any deal.

Authors' Response: *They were calculated using the three dimensional wind fields simulated by WRF model (that were also used as input to the STEM simulations). Details describing the trajectory calculations have been added in section 2.2.4 of the manuscript.*

Anonymous Referee #2

This manuscript describes modeling analysis of PM seasonal cycles, source type and region contribution, and future emission scenarios over Central Asia (CA). The WRF/STEM model is used to conduct a hemispheric scale simulation at 60km x 60km resolution. Observations at two surface sites in CA and AOD retrieved from MODIS are used to evaluate the WRF-STEM model prediction. The analysis is focused on the source attributions to PM in CA, suggesting dust and residential heating as two major contributors of PM for this region. Additional analysis is conducted on the impact of CA emissions on other regions of the Northern Hemisphere. Future emission scenarios for 2030 are adopted from the WMO/UNCEP report on short-lived climate forcers. Putting these emissions scenarios in the model, the authors find significant increase of PM and BC over CA if current legislation is kept unchanged for pollution abatement measures.

Given little attention has been given to Central Asia, this paper represents new analysis. The paper is well written and well organized. It is suitable for publication in ACP once several revisions have been made, as described below.

Major comments:

1. The model simulation and analysis are largely performed on the scale of the whole Northern Hemisphere, including other Asian regions with large emissions such as China, Europe, and North America. The emissions from CA will account for only a small portion of emissions in NH, thus the source attribution and long-range transport analysis will depend on the model's ability in capturing sources seasonality and abundance in other regions. However, the model evaluation is conducted only based on two surface sites in CA. Almost all the maps of model predictions shown in the paper are of the hemispheric scale, but such results are not evaluated nor do the authors discuss the uncertainties of them. The inconsistency between the scale of the model/results presented and the scale of the focused region is the main shortcoming of this paper.

Authors' Response:

We thank the reviewer for this comment. The primary focus of this paper is Central Asia and we focus our analysis on model performance on the two sites located in this region. We would like to point out that our analysis also includes the long range transport into and out of the region and this brings in the hemispheric scale.

We have included more information on model prediction skills (outside CA region) by discussing the analysis/results describing the comparison of simulated BC values with surface BC observations from IMPROVE network in section 3.1 and quoted here.

“We evaluated the simulated BC values with IMPROVE BC observations for the simulation time period (i.e. April 2008 – July 2009) used in this study. The IMPROVE BC analysis is limited to 27 sites on the US west coast (i.e. west of 120 W) to be consistent with the focus of this paper (i.e., long range transport of pollutants on a hemispheric scale). The mean observed and

modeled BC for the West Coast sites are ~ 0.16 and $0.19 \mu\text{g m}^{-3}$ ($n = 3759$ points) with mean bias and RMSE of $0.03 \mu\text{g m}^{-3}$ and $\sim 32\%$ respectively. The mean simulated/observed BC at surface sites is ~ 1.2 , which compares well with the corresponding ratio of 1.6 reported in the global model performance over North America region by Koch et al., 2009. The detailed analysis of the IMPROVE BC sites and associated seasonality will be addressed in a future paper. The rest of this section will focus on discussing the regional aerosol distributions in the CA region.”

The model has been evaluated on a hemispheric scale using MODIS AOD in the domain of interest for this analysis and is described in section 3.2.

The evaluation of this model framework that use the same domain configuration and settings in other regions including arctic region and continental US are summarized below and now included in the model performance evaluation described in section 3.

“We have evaluated this model framework in other regions outside CA including Arctic region and continental US previously, which are summarized below. The domain configuration and settings of the WRF/STEM modeling system used in this study have been applied to provide meteorology and air quality forecasting during the ARCTAS field campaign in 2008 (Dallura et al., 2011) as mentioned earlier in sections 2.2.3 and 2.3.1 respectively. The study by Dallura et al., used WRF v2.2 and pre-emission ARCTAS emissions data, (http://mic.greenresource.cn/arctas_premission) real time biomass burning emissions from the RAQMS modeling group (<http://raqms-ops.ssec.wisc.edu/>) and evaluated the performance of the WRF-STEM modeling framework (used in this study) through the case study of the ARCTAS mission DC-8 flight conducted on July 9 2008. The WRF model compared well with observed meteorological variables (including temperature, pressure, wind speed and wind direction) along the flight and correctly simulated the various meteorological processes (i.e. cyclonic circulation, low pressure system, anticyclones and geostrophic wind flows) that facilitate transport of pollutants into the Arctic region. This study also compared CO along the flight and found that the simulated values were able to capture the magnitude and temporal variability seen in observed values. The study also found that the fires from North Asia and anthropogenic CO from China were the major sources contributing to observed CO levels along the flight suggesting that the midlatitude pollution and summertime forest fires/biomass burning transport events were well represented in this modeling system.

Huang et al, 2012 used a subset of this domain (covering the region of continental North America) to simulate summertime sectoral and regional contributions to BC over continental US using the WRF/STEM modeling framework. Note that the Huang et al., 2012 study used the same WRF meteorology, ARCTAS emissions and RAQMS biomass burning emissions used in the Dallura et al., 2011 study, but used the full-chemistry version of the STEM model. The comparison of the simulated aerosol BC from the tracer STEM model with corresponding BC values from the full chemistry version of the STEM model showed that the simulated tracer BC was able to capture trend and the same relative variability seen in the full chemistry predictions. Huang et al., 2012 evaluated the model performance aloft and at surface using measurements from ARCTAS mission DC8 flights conducted on June 20 and 22 and 24 that sampled over California and IMPROVE surface site network. Huang et al., 2012 found that the predicted BC

values captured many of the vertical features seen in the flight observations with a tendency to over predict surface BC and > 4km. The comparison of predicted BC with observed values from the IMPROVE network showed that the model was able to capture the gradients in BC values with high concentrations in western and eastern NA regions with slight positive and negative biases over the mountain regions and the eastern US/California.

These studies indicate that the simulated transport patterns are able to capture the main important features of the hemispheric flows impacting Central Asia, North America and the Arctic region, at the scale discussed. The prediction of aerosol mass and composition at the hemispheric scales are able to capture important aspects of horizontal gradients and variability, but has considerably higher uncertainties associated largely with emission estimates (in particular forest fires/biomass burning and natural dust emissions) and wet removal processes (Bates et al., 2006)."

2. In discussion of the emissions and future emission scenarios, it will be more informative for the readers to obtain a quantitative understanding of the emissions changes and the resulting changes in PM concentrations. I suggest the authors compile the results in a table, for example, listing how much BC emissions are increased in the current legislation scenario in 2030 and the resulting changes in BC concentrations. The table should be focused on CA. Currently these results are shown in the supplementary materials and by figures which are hard to read.

Authors' response:

We appreciate the reviewer's comment. We have compiled a summary table that shows the change in emissions and concentration over the Central Asia region defined in the domain. This summary is included in section 3.8 that discusses the future emission scenarios along with the following text.

"To get a regional perspective of how the future emission changes would specifically impact the CA region (See Fig. 1 for the CA region definition used in this study), we have summarized the changes in emissions and the corresponding concentrations in Table 1. The reference 2030 emissions scenario reports an increase in emissions of BC, SO₂ and PM_{2.5} over Central Asia by ~ 22, 17 and 14 % respectively w.r.t to base 2005 levels. However, the corresponding BC, SO₄ and anthropogenic portion of PM_{2.5} decrease by ~ 5, 12 and 5 % respectively. This suggest that on average the concentration levels go down even though CA emissions increase due to long range transport of pollutants into CA region from the surrounding regions, but locations within CA still increase when dominated by local sources. The Low GWP emission scenario predicts decrease in BC and PM_{2.5} emissions by 31% and 10% respectively with the corresponding decrease in concentrations by ~ 37% and 10 % respectively, while SO₂ emissions remained same as the reference 2030 scenario. The lowest + 450 ppm scenario shows a consistent decreasing trend in emissions and concentration of all the species. This analysis suggests that the impact of the changes in major source region emissions and their subsequent transport to Central Asia will play a major role in determining the future aerosol levels in CA."

Table: 1 Summary of changes in Central Asia region (See figure 1 for Central Asia region definition) emissions and concentration under the future emission scenarios (See Sect. 3.9 for more details)

Emission Scenario	Emissions Gg yr ⁻¹			Emissions change (%) w.r.t to base 2005		
	BC	SO ₂	PM ₂₅	BC	SO ₂	PM ₂₅
Base 2005	39	1130	197			
Reference 2030	47	1326	224	22.1	17.3	13.9
Low GWP 2030	27	1325	177	-31.1	17.3	-10.1
Lowest GWP + 450	21	1058	115	-46.9	-6.4	-41.5

	Concentration (µg m ⁻³)			Concentration change (%) w.r.t to base 2005		
	BC	SO ₄	PM ₂₅	BC	SO ₄	PM _{2.5}
Base 2005	0.17	3.04	4.26			
Reference 2030	0.16	2.65	4.03	-4.2	-12.6	-5.3
Low GWP 2030	0.11	2.64	3.85	-36.6	-13	-9.5
Lowest GWP + 450	0.10	2.04	3.06	-38.5	-32.9	-28

Minor comments:

1. pg 11357, line 21: % changed to percentage

Done

2. pg 11357, line 26: “are” changed to “is”

Done

3. pg 11360, line 8: remove “is” after in turn

Done

References:

Bates, T. S., Anderson, T. L., Baynard, T., Bond, T., Boucher, O., Carmichael, G., Clarke, A., Erlick, C., Guo, H., Horowitz, L., Howell, S., Kulkarni, S., Maring, H., McComiskey, A., Middlebrook, A., Noone, K., O'Dowd, C. D., Ogren, J., Penner, J., Quinn, P. K., Ravishankara, A. R., Savoie, D. L., Schwartz, S. E., Shinozuka, Y., Tang, Y., Weber, R. J., and Wu, Y.: Aerosol direct radiative effects over the northwest Atlantic, northwest Pacific, and North Indian Oceans: estimates based on in-situ chemical and optical measurements and chemical transport modeling, *Atmos. Chem. Phys.*, 6, 1657-1732, doi:10.5194/acp-6-1657-2006, 2006.

D'Allura, A., Kulkarni, S., Carmichael, G. R., Finardi, S., Adhikary, B., Wei, C., Streets, D., Zhang, Q., Pierce, R. B., Al-Saadi, J. A., Diskin, G., and Wennberg, P.: Meteorological and air quality forecasting using the WRF-STEM model during the 2008 ARCTAS field campaign, *Atmos. Environ.*, 45, 6901–6910, doi:10.1016/j.atmosenv.2011.02.073, 2011.

Huang, M., G. Carmichael, S. Kulkarni, D. Streets, Z. Lu, Q. Zhang, B. Pierce, Y. Kondo, J. Jimenez-Palacios, M. J. Cubison, B. E. Anderson, and A. Wisthaler: Sectoral and geographical contributions to summertime continental United States (CONUS) black carbon spatial distributions, *Atmos. Environ.*, 51, 165-174, doi:10.1016/j.atmosenv.2012.01.021, 2012

Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Berntsen, T., Bond, T. C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O., Easter, R., Fahey, D. W., Feichter, J., Fillmore, D., Freitag, S., Ghan, S., Ginoux, P., Gong, S., Horowitz, L., Iversen, T., Kirkevåg, A., Klimont, Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N., Myhre, G., Penner, J. E., Perlwitz, J., Pitari, G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G., Schwarz, J. P., Seland, Ø., Stier, P., Takegawa, N., Takemura, T., Textor, C., van Aardenne, J. A., and Zhao, Y.: Evaluation of black carbon estimations in global aerosol models, *Atmos. Chem. Phys.*, 9, 9001-9026, doi:10.5194/acp-9-9001-2009, 2009