



Horizontal
distributions of
aerosol constituents

K. Hara et al.

Horizontal distributions of aerosol constituents and their mixing states in Antarctica during the JASE traverse

K. Hara¹, F. Nakazawa², S. Fujita², K. Fukui^{2,3}, H. Enomoto^{4,2}, and S. Sugiyama⁵

¹Department of Earth System Science, Faculty of Science, Fukuoka University, Fukuoka, Japan

²National Institute of Polar Research, Tokyo, Japan

³Tateyama Caldera Sabo Museum, Toyama, Japan

⁴Kitami Institute of Technology, Kitami, Japan

⁵Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan

Received: 26 March 2014 – Accepted: 22 April 2014 – Published: 7 May 2014

Correspondence to: K. Hara (harakei@fukuoka-u.ac.jp)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Measurements of aerosol number concentrations and direct aerosol sampling were conducted on continental Antarctica during the traverse of Japanese–Swedish joint Antarctic expedition (JASE) from 14 November 2007 until 24 January 2008. Aerosol concentrations in background conditions decreased gradually with latitude in inland regions during the traverse. The lowest aerosol number concentrations were 160 L^{-1} in $D_p > 0.3\ \mu\text{m}$, and 0.5 L^{-1} in $D_p > 2\ \mu\text{m}$. In contrast, aerosol concentrations reached 3278 L^{-1} in $D_p > 0.3\ \mu\text{m}$, and 215 L^{-1} in $D_p > 2\ \mu\text{m}$ under strong wind conditions. The estimated aerosol mass concentrations were $0.04\text{--}5.7\ \mu\text{g m}^{-3}$. Single particle analysis of aerosol particles collected during the JASE traverse was conducted using a scanning electron microscope equipped with an energy dispersive X-ray spectrometer. Major aerosol constituents were sulfates in fine mode, and sulfate, sea-salts, modified sea-salts, and fractionated sea-salts in coarse mode. K-rich sulfates, Mg-rich sulfate, Ca-rich sulfates, and minerals were identified as minor aerosol constituents. Sea-salt particles were modified greatly with SO_4^{2-} and NO_3^- in coarse mode, and dominantly with SO_4^{2-} in fine mode during the traverse. Most sea-salt particles in the continental region were modified with sulfate and methanesulfonate near the coast, although NO_3^- contributed markedly to sea-salt modification in inland areas during summer. Mg-rich sea-salt particles and Mg-free sea-salt particles were present in coarse and fine modes from the coast to inland areas. These sea-salt particles might be associated with sea-salt fractionation on the snow surface of continental Antarctica.

1 Introduction

Atmospheric aerosol constituents in Antarctic regions have been measured for more than three decades to elucidate the regions' aerosol chemistry, to monitor Earth background levels, and to interpret ice core records (e.g., Savoie et al., 1992, 1993; Minikin

ACPD

14, 11393–11445, 2014

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thin-film method) to demonstrate that sulfuric acid was dominant in aerosol constituents in Antarctic troposphere during summer. In addition, single particle analysis using laser microprobe mass spectrometry showed that CH_3SO_3^- was mixed internally with sulfate particles (probably H_2SO_4 particles) near the surface on the Antarctic coasts during summer (Wouters et al., 1990; Hara et al., 1995).

Sea-salts are dominant during winter–spring (e.g., Hara et al., 2004). Sea-salt particles were distributed widely in ultrafine–coarse mode throughout the year at Syowa Station (Hara et al., 2010a, 2011a) and were distributed in fine–coarse mode during the summer at Aboa Station (Kerminen et al., 2000; Teinilä et al., 2000). From elemental analysis of individual particles using EDX, Mouri et al. (1999) and Hara et al. (2005, 2013) reported that sea-salt particles near the surface were modified with SO_4^{2-} and CH_3SO_3^- during the summer, and were modified with NO_3^- in August at Syowa Station. Furthermore, single particle analysis of aerosol particles collected using tethered-balloon operations exhibited seasonal and vertical features of aerosol constituents and their mixing states in near surface–lower free troposphere (2.5 km), sea-salt modification, and sea-salt fractionation (Hara et al., 2013). In addition to sea-salt modification, sea-salt particles were fractionated through precipitation of several salts such as mirabilite ($\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$) and hydrohalite ($\text{NaCl} \cdot 2\text{H}_2\text{O}$) in sea ice formation and frost flower appearance under colder conditions (e.g., Wagenbach et al., 1998; Rankin et al., 2000, 2002; Hara et al., 2004, 2012). Sea-salt fractionation on sea-ice (including frost flowers) by depletion of Na-salts engenders Mg-enrichment in sea-salt particles during winter–spring on the Antarctic coast (Hara et al., 2010, 2012, 2013). Furthermore, Hara et al. (2013) pointed out the likelihood that sea-salt fractionation (Mg separation in sea-salt particles) occurs during summer. The following questions, however, remain; (1) “where does sea-salt fractionation occur on the Antarctic regions during summer?”, and (2) “what are the specific processes of sea-salt fractionation (Mg separation)?”.

Although useful and important knowledge related to aerosol chemical properties (e.g., concentrations, and mixing states) has been obtained gradually from previous investigations using bulk and single-particle analysis of aerosol particles along the

Horizontal distributions of aerosol constituents

K. Hara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Antarctic coasts, a great dearth of knowledge remains for aerosol chemical properties and chemical processes taking place on the Antarctic continent. To elucidate spatiotemporal features of glaciological environment and atmospheric quality in Droning Maud Land, East Antarctica, scientific traverse using snow vehicles was conducted by a Japanese Swedish Antarctic Expedition (JASE) in the austral summer of 2007/2008 (Fujita et al., 2011). In this study, measurements of the aerosol number concentrations and direct aerosol sampling were conducted during the JASE traverse to characterize and elucidate the horizontal features of aerosol constituents and their mixing states in atmosphere near the surface on the Antarctic continent during summer. Herein, we mainly discuss (1) the horizontal distributions of aerosol constituents and mixing states on the Antarctic continent during summer, (2) sea-salt modification, and (3) sea-salt fractionation.

2 Aerosol measurements and data analysis

2.1 JASE traverse

Figure 1 depicts the JASE traverse route in Droning Maud Land, East Antarctica. On this campaign, the Japanese team traveled from S16 on the Antarctic continent (near Syowa Station) to a meeting point on the Antarctic plateau via Dome F Station. The Swedish team traveled from Wasa Station to the meeting point via Kohnen Station. Aerosol measurements and direct aerosol sampling for this study were conducted during the Japanese team traverse. The Japanese traverse team left from S16 on 14 November 2007, and arrived at the meeting point on 27 December 2007 in the incoming traverse. After some joint scientific works at the meeting point for several days, then, the outgoing traverse (return to S16) began on 30 December 2007. During the outgoing traverse, the Japanese team traveled from the meeting point to Dome F on the southern side of the ice divide for glaciological measurements. The team approached S16 on 24 January 2008. Details of the traverse were described in Fujita et al. (2011).

2.2 Meteorological measurements during the JASE traverse

Continuous measurements of air pressure (F4711; Yokogawa Analytical Systems Inc.), air temperature (KDC-A01-S001 and KADEC21-U4; Kona System Inc.), wind speed, and wind direction (KADEC21-KAZE; Kona System Inc.) were performed during the JASE traverse. Meteorological sensors were fixed on the snow vehicle, located approximately at 3 m above the snow surface. Daily meteorological observations (air, pressure, air temperature, wind direction, wind speed, weather, visibility, degree of cloud cover, and cloud type) were made around 15:00 LT. Weather conditions were observed also during aerosol measurements conducted at camp sites.

2.3 Aerosol measurements

Aerosol measurements and direct aerosol sampling were conducted during the JASE traverse. For safe operation of aerosol measurements, measurements of the aerosol size distribution, and direct aerosol sampling were not made during the traverse during the daytime but were conducted only at camp sites.

2.3.1 Measurements of aerosol number concentrations

Aerosol number concentrations were measured using a portable optical particle counter (OPC, KR12A; Rion Co. Ltd.) during direct aerosol sampling at every camp site. The measurable size range was $D_p > 0.3, 0.5, 0.7, 1.0, 2.0, \text{ and } 5.0 \mu\text{m}$. The OPC packed in an insulator box was set at ca. 1 m above the snow surface by tripod, located on the windward side of the camp site to avoid local contamination (mainly exhaust from snow vehicles). Aerosol number concentrations were recorded every 23–25 s (corresponding to 1 L air sucking) during the measurements. The ambient number concentrations were converted to the concentration under standard conditions (0°C and 1013.25 hPa).

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.2 Direct aerosol sampling and analysis

Aerosol particles were collected on carbon-coated collodion thin film supported by Ni-microgrid (square-300 mesh; Veco Co.) using a two-stage aerosol impactor. The cut-off diameter of the impactor was 2.0 and 0.2 μm at a flow rate of 1 L min^{-1} . The impactor was set at ca. 1 m above the snow surface next the OPC. Because of the lower aerosol number concentrations on the Antarctic continent, direct aerosol sampling was conducted for 28–86 min (mean, 60 min) depending on the aerosol number concentration. After direct sampling, aerosol samples were kept in air tight boxes including desiccant until analysis in our laboratory in Japan to prevent humidification that can engender morphological change and chemical reactions.

Individual aerosol particles on the sample substrate were observed and analyzed in this study with a scanning electron microscope equipped with an energy dispersive X-ray spectrometer (SEM-EDX; Quanta FEG-200F, FEI; XL30; EDAX Inc.). Analytical conditions were the following: 20 kV accelerating voltage and 30 s counting time. To avoid analytical bias of localization of aerosol constituents in each particle, the rectangular or square area almost covering a particle was scanned using an electron beam in EDX analysis. Details of analytical procedures and data quality were described in Hara et al. (2005, 2013). We analyzed 2690 particles in coarse mode (mean: 58 particles sample^{-1}) and 45 044 particles in fine mode (mean: 961 particles sample^{-1}). Although we attempted to analyze as many coarse particles as possible, the lower aerosol number concentrations in coarse mode limit the number of the analyzed aerosol particles in this study.

ACPD

14, 11393–11445, 2014

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 Meteorological conditions during the JASE campaign

3.1.1 Air temperature, wind speed and weather near the surface

Figure 2 shows variations of latitude, elevation, air temperature, and wind speed during the JASE traverse of the Japanese team. The air temperature decreased gradually with latitude in 69–73° S in both incoming and outgoing traverse, although the highest temperature near the coast was ca. -7.6°C in the incoming traverse and -2.9°C in the outgoing traverse. By contrast, the air temperature was around -30°C at latitudes higher than 73° S in this study. The lowest air temperature was -43°C . In addition, air temperature showed strong diurnal variation. The diurnal temperature range was approximately 8.6°C on average (maximum, 16°C).

Wind speed often showed diurnal feature during the traverse. In addition to diurnal features of wind speed, the wind speed increased because of turbulence, for example on 18–22 November, 27–30 November, 7–8 December, and 20–23 December. Although the diurnal maximum of air temperature was observed at noon time (LT), those of wind speed were lagged at latitudes lower than 73° S. Similarly, similar diurnal variations of wind speed were observed on Antarctic continent (Allison, 1998; Van As et al., 2005) and Antarctic coasts (Sato and Hirasawa, 2007). However, the diurnal maxima were mutually synchronized well at latitudes higher than 74° S, when diurnal features occur. According to a manual weather check at every camp site, weather was mostly clear and fine. When wind speed was greater than ca. 6 ms^{-1} , drifting snow occurred during the traverse, for example on 28–29 November, 7 December, and 21–23 December.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1.2 Air mass history during the JASE traverse

To elucidate the history and origins of air masses observed in this study, the 5 day backward trajectory was computed using the NCEP reanalysis dataset and kinetic mode in this study (Draxler and Rolph, 2013). Figure 3 depicts the backward trajectories from 200 m a.g.l. over each camp site. As depicted in Fig. 3a, air masses from the camp sites located in 69–71° S in the incoming traverse were transported westward along with Antarctic coastal line during the prior 5 days. Vertical features of the backward trajectories imply that these air masses traveled in the upper boundary and lower free troposphere. In contrast to the transport pathway in 69–71° S, air masses at the camp sites in > 71° S traveled in the free troposphere over the Antarctic continent during the prior 5 days. Air masses in the traverse from Dome F to the meeting point flowed over the Antarctic plateau during the prior 5 days (Fig. 3b). The vertical motions in these air masses were classified into (1) transport from the free troposphere and (2) travel in the boundary layer (near surface). Compared to the weather at camp sites, the first type (transport from free troposphere) and second type (transport in the boundary layer) corresponded to fine days, and conditions with strong winds and drifting snow, respectively. Backward trajectories in the outgoing traverse from the meeting point to Dome F indicate transport in the boundary layer–lower free troposphere over the Antarctic continent (Fig. 3c). The transport pathway during the outgoing traverse from Dome F to S16 was clearly divisible into transport from coastal areas and transport over the continent (Fig. 3d). Although air masses at latitudes higher than 73° S came traveled over the continent, air masses in > 75° S and 73–75° S were transported from near the South Pole, and were transported along ice ridges through near Dome A and Dome C in East Antarctica, respectively. Furthermore, air masses in the outgoing traverse moved near ground level (probably in boundary layer) for past 5 days, in contrast to the large variability of vertical motions in the incoming traverse from S16 to Dome F. These features of transport pathways of air masses on the Antarctic continent during

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the JASE traverse showed good agreement with analysis of the air mass origins over the Antarctic continent by Suzuki et al. (2013).

3.2 Number concentrations, mass concentrations, and size distribution of aerosol particles during the JASE traverse

5 Figures 4 and 5 depict features of wind speed, the aerosol number concentrations, Junge slope, and aerosol mass concentrations during direct aerosol sampling at each camp site during the JASE traverse. During the incoming traverse, the aerosol number concentrations decreased gradually with latitude and elevation, except high number concentrations in strong winds. Higher aerosol number concentrations were identified
10 under conditions with wind speeds higher than 6 ms^{-1} in this study. Therefore, we use aerosol concentrations at wind speeds lower than 6 ms^{-1} as “background aerosol concentrations” in this study. For instance, the background number concentrations in $D_p > 0.3 \mu\text{m}$ and $D_p > 2 \mu\text{m}$ changed, respectively, from ca. 1330 L^{-1} near the coast to ca. 160 L^{-1} , and from 36 L^{-1} to 0.5 L^{-1} . The number concentrations on the Antarctic
15 continent during the JASE traverse were lower than those in the lower free troposphere over Syowa Station in summer (Hara et al., 2011b). The aerosol number concentrations increased considerably under conditions with strong wind and drifting snow. Considering that air masses on the Antarctic continent during JASE were not transported from coastal areas during the prior 5 days, this increase suggests that strong winds engender
20 the release of aerosol particles and small snow/ice flakes from the snow surface. Particularly the number concentrations in coarse particles having diameter larger than $2.0 \mu\text{m}$ increased remarkably by one–two orders higher in strong wind conditions, as shown in Figs. 4 and 5. The aerosol number concentrations and their horizontal features in the outgoing traverse were similar to those in the incoming traverse.

25 For comparison of the aerosol size distribution, we estimated the “Junge-slope” in this study. The aerosol size distribution in fine–coarse mode can be approximated as

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the following equation (Junge, 1963).

$$\frac{dN}{d\log D_p} = \alpha D_p^{-\beta}$$

Therein, α and β respectively represent a constant and the Junge-slope. In this study, the Junge-slope (β) was estimated as 0.3–5.0 μm in diameter. The Junge-slope was 2.22–3.03 (mean, 2.70; median, 2.75) in background conditions (wind speed lower than 6 ms^{-1}) in the incoming traverse. In contrast, the Junge-slope decreased to less than 2.5 under conditions with strong winds ($> 6 \text{ms}^{-1}$) or drifting snow. The Junge-slope in the outgoing traverse was 2.73–3.37 (mean, 3.03; median, 2.92) under background conditions. Similar to the Junge slope in the incoming traverse, the Junge slope decreased in strong winds during the outgoing traverse. Consequently, aerosol size distributions depended closely on release of aerosol particles from snow surfaces by wind blowing. In addition, the Junge-slopes in the outgoing traverse were slightly larger than those in the incoming traverse. This difference derived from the reduction of aerosol release (especially in coarse mode) from the snow surface under calm wind conditions during the outgoing traverse. Considering that weather conditions are usually stable in December–January on the Antarctic region (e.g., Sato and Hirasawa, 2007), these differences between the incoming traverse and outgoing traverse might reflect seasonal features of aerosol number concentrations and the size distribution on the Antarctic continent during early summer (or end-spring) through mid-summer. Compared to Junge-slope over Syowa Station (range, 2.2–3.2; median, 2.5, Hara et al., 2011b), the Junge-slope was slightly higher on the Antarctic continent during the JASE traverse. The difference of the Junge-slopes might result from horizontal features of the number concentrations, particularly in coarse mode, from coasts to inland areas. Indeed, the number concentrations in coarse mode in JASE traverse were several factors – one order lower than those over Syowa Station.

Here, we attempt to estimate the mass concentrations using aerosol number concentrations, assuming spherical shape and density of aerosol particles. The density

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



corresponded to the density of sulfates (ca. 1.8 g cm^{-3}) and NaCl (2.2 g cm^{-3}). Major aerosol constituents were sulfate particles during JASE traverse (details are shown in later Sects. 3.4 and 3.5). Therefore, we used the density of 1.8 g cm^{-3} to estimate the mass concentration in this study. The number concentrations of aerosol particles smaller than $0.3 \mu\text{m}$ were not included in the estimation. Therefore, the estimated mass concentrations can be underestimated slightly. In addition, aerosol number concentrations were observed using OPC above snow surface. Thus, number concentrations were measured under ambient conditions (not dry conditions), although OPC packed in an insulator box. Therefore, the estimated mass concentrations included masses of water in aerosol particles in this study. In exception of higher mass concentrations under conditions with drifting snow and strong winds, the aerosol mass concentrations gradually decreased from ca. $0.16 \mu\text{g m}^{-3}$ (near coasts) to $0.04 \mu\text{g m}^{-3}$ (near Dome F Station). The mass concentrations on the Antarctic continent during JASE traverse were similar to the mass concentrations of water soluble aerosol constituents at Kohnen Station (Weller and Wagenbach, 2007), and slightly higher than the mass concentrations water soluble aerosol constituents at Dome C (Preunkert et al., 2008; Udisti et al., 2010). The aerosol mass concentrations increased considerably to larger than several $\mu\text{g m}^{-3}$ under conditions with drifting snow and strong winds. The highest mass concentrations were approximately $5.7 \mu\text{g m}^{-3}$. Higher mass concentrations were observed in high number concentrations in coarse mode. Therefore, aerosol particles from snow surface by erosion under strong winds might make significant contribution to high aerosol mass concentrations.

Figure 3 shows that air masses on the Antarctic continent were transported over the Antarctic plateau during the prior 5 days. Considering that coarse particles can be removed efficiently from the atmosphere through dry deposition during transport, isolation from coastal regions might account for gradient features of aerosol number concentrations in coarse mode. In other words, coarse particles are supplied only slightly from coastal regions into the Antarctic plateau during summer. The aerosol number concentrations increased markedly in all size ranges ($D_p > 0.3 \mu\text{m}$) under strong wind

Because EDX can provide only elemental information, we cannot distinguish SO_4^{2-} from CH_3SO_3^- . Hereinafter, we designate these aerosol particles as “sulfate particles”.

In fact, S and K were detected from aerosol particles as shown in Fig. 7b. Aerosol particles containing S and K had no satellite structure. Similar aerosol particles were identified in aerosol samples taken in the lower free troposphere–upper free troposphere in Antarctic region (Hara et al., 2010, 2013). K-rich sulfate particles cannot be formed through gas-to-particle conversion from aerosol precursors derived from oceanic bioactivity. Therefore, K-rich sulfate particles might be non-biogenic aerosol particles. As discussed by Okada et al. (2001, 2008) and Niemi et al. (2005), K-rich sulfate particles and nss-K in aerosol particles are released from combustion processes such as biomass burning. Because of the negligible source strength of biomass burning on the Antarctic regions, the particles might be transported from outside of the Antarctica via the free troposphere.

Peaks of Mg and S were obtained from aerosol particles as portrayed in Fig. 7c. Atomic ratios of Mg and S of the aerosol particles containing only Mg and S were compatible with the ratios of MgSO_4 . Mg-rich sulfate particles in the lower troposphere over Syowa Station were identified dominantly in air masses transported from the Antarctic continent (Hara et al., 2013). Hara et al. (2013) reported that Mg-rich sulfate particles in the Antarctic atmosphere were associated with sea-salt fractionation and sea-salt modification. Details of Mg-rich sulfate particles and sea-salt fractionation are discussed in Sect. 3.7.

Aerosol particles containing Ca and S were also observed in this study (Fig. 7d). Because of the atomic ratios of Ca and S, particles of this type might be CaSO_4 particles. CaSO_4 particles were identified in the lower troposphere on the Antarctic region (Hara et al., 2010, 2013). Actually, CaSO_4 is well-known as a major mineral constituent (gypsum). Moreover, Marion and Farren (1999) and Gelifus et al. (2013) pointed out CaSO_4 (gypsum) formation by sea-salt fractionation on sea-ice in polar regions. However, the origins of CaSO_4 particles in the Antarctic atmosphere remain unclear.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Peaks of Na, Mg, S, and Cl were obtained aerosol particles as shown in Fig. 7e. These elements are major seawater constituents. When artificial NaCl particles were analyzed using SEM-EDX, the peak height of Cl was slightly higher than that of Na. Consequently, the particle in Fig. 7e might be identified as partly Cl-depleted sea-salt particles. Also, Na, Mg, and S were detected from aerosol particle in Fig. 7f. Compared to aerosol particle in Fig. 7e, the peak height of S relative to Na was higher in particle in Fig. 7f. Cl in sea-salt particles can be liberated through heterogeneous reactions with acidic sulfur species such as SO_4^{2-} and CH_3SO_3^- in Antarctic troposphere during summer (e.g., Mouri et al., 1999; Hara et al., 2005, 2013). Therefore, the particles in Fig. 7f might be sea-salt particle modified with acidic sulfur species such as SO_4^{2-} and CH_3SO_3^- . Additionally, the height of Mg relative to Na was higher than that of sea-salt particles in Fig. 7d. The particle in Fig. 7f might be identified as a wholly Cl-depleted sea-salt particle with slight Mg-enrichment. Although Na, Mg, and S were detected from aerosol particle in Fig. 7g, the peak height of S was lower than particles in Figs. 7e and f. Acidic species other than SO_4^{2-} and CH_3SO_3^- might contribute significantly to Cl depletion from sea-salt particles. Also, the particle in Fig. 7g might be identified as a wholly Cl-depleted sea-salt particle. Hereinafter, we designate wholly Cl-depleted particles as “modified sea-salt particles”. Less Cl-depleted sea-salt particles and partially Cl-depleted sea-salt particles were divided into “sea-salt particles” in this study. Modification of sea-salt particles is discussed in Sect. 3.6. Although Mg is a major sea-salt constituent, Mg was not detected from aerosol particles containing Na, K, and Cl, as portrayed in Fig. 7h. A particularly strong K peak relative to Na peak was identified clearly from the particle in Fig. 7h. Na is a dominant constituent in sea-salt particles. Therefore, particles of this type were identified as “sea-salt-like particles” in this study. The sea-salt-like particles were observed only at sampling sites on the way from Dome F to the meeting point in the incoming traverse on 21–23 December 2007 when aerosol particles were collected under conditions with drifting snow.

In fact, Si, Al, and S were detected from solid particles as shown in Fig. 7i. Al and Si were major constituents. Therefore, particles of this type might be identified as mineral

particles. Furthermore, stain and satellite structures were observed around irregular solid core in some mineral particles. Considering that S was detected in mineral particles (irregular solid particles) satellite structure, these particles were likely to be mineral particle coated acidic sulfates and H₂SO₄. In addition, internal mixtures of minerals and sea-salt particles were often identified in this study (Fig. 7j).

3.5 Horizontal distributions of aerosol constituents during the JASE traverse

To compare quantitatively horizontal distributions of each aerosol constituent and mixing states, we estimated the relative abundance (number fraction) of each type of aerosol constituent. Figures 8 and 9 show horizontal distributions of relative abundance of aerosol mixing states in coarse and fine modes during the incoming traverse (14 November–27 December 2007) and the outgoing traverse (27 December 2007–24 January 2008).

3.5.1 Sea-salts

In coarse mode, major aerosol particles were sea-salt particles, modified sea-salt particles and sulfate particles. The relative abundance of sea-salt particles in coarse mode was larger than 40% near the coast in the incoming traverse. Relative abundance of sea-salt particles in coarse mode decreased gradually in inland areas in exception of the high abundance at sampling sites in 77–76° S on the way from Dome F Station to the meeting point in the incoming traverse. With the gradual decrease of the relative abundance of sea-salt particles, the relative abundance of the modified sea-salt particles increased gradually up to 40% on the Antarctic plateau. By contrast, the modified sea-salt particles were dominant in fine mode in the incoming traverse. Because of the predominance of sulfate particles in fine mode, the relative abundance of sea-salt particles and the modified sea-salt particles were less than 6.8% even at latitudes lower than 72° S and less than 2.8% at latitudes higher than 72° S during the incoming traverse. In addition, the relative abundance of sea-salt particles increased markedly at

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sampling sites in 77–76° S on the way from Dome F to the meeting point. As described above, sea-salt-like particles (Na–K–Cl) were observed dominantly in coarse mode at these sites. According to the 5 day backward trajectory as depicted in Fig. 3, transport from the coasts was found only at 69–71° S during the incoming traverse, whereas air masses in the inland area (> 72° S) were transported over the continent during the prior 5 days. Therefore, isolation from the coastal regions might cause gradual distributions of sea-salt particles and modified sea-salt particles in the continent. The transport pathway, however, cannot account for the high relative abundance of sea-salt particles and the modified sea-salt particles, for example at 72–73° S and at 77–76° S. As depicted in Fig. 4, the aerosol number concentrations increased considerably, particularly in coarse mode, under strong wind conditions in the incoming traverse to the meeting point. Consequently, high aerosol number concentrations and their high relative abundance might result from redistribution of aerosol particles such as sea-salt particles from the snow surface. Therefore, the horizontal distribution of relative abundance of sea-salt particles and the modified sea-salt particles on the Antarctic continent during summer might be associated with the transport pathway and release from the snow surface via wind-blowing.

Relative abundance of sea-salt particles and the modified sea-salt particles in coarse mode ranged mostly in lower than 60 % during the outgoing traverse. Although the total relative abundance of sea-salt particles and the modified sea-salt particles in coarse mode was similar to that in the incoming traverse except the high relative abundance under strong wind conditions, the modified sea-salt particles were dominant in the outgoing traverse. However, the relative abundance of sea-salt particles and the modified sea-salt particles in fine mode was mostly less than 0.5 % in inland areas during the outgoing traverse. In some samples taken on the Antarctic plateau, sea-salt particles and the modified sea-salt particles were not detected in fine mode. Compared to the horizontal distributions of sea-salt particles and the modified sea-salt particles in fine mode during the incoming traverse, relative abundances of sea-salt particles and the modified sea-salt particles were remarkably lower. Mass concentrations of sea-salts

(Na⁺) in aerosol particles show a minimum at Syowa and Dome F stations during summer (Hara et al., 2004). In addition, the relative abundance of sea-salt particles and the modified sea-salt particles decreased considerably over Syowa Station (Hara et al., 2013). Therefore, the difference of the relative abundance between in the incoming traverse and in the outgoing traverse might correspond to seasonal features of sea-salt particles from the end of spring toward summer on the Antarctic continent.

3.5.2 Sulfates

The following aerosol particles containing sulfates were identified in this study: (1) sulfate particles, (2) modified sea-salt particles, (3) sulfate particles containing K, (4) MgSO₄ particles, and (5) CaSO₄ particles. Because horizontal distributions of the modified sea-salt particles were described in Sect. 3.5.1, their description is excluded from this section.

Figures 8 and 9 show that sulfate particles were dominant in both coarse and fine modes. Particularly, the mean relative abundance of sulfate particles in fine mode was 98.5 % in the incoming traverse and 99.5 % in the outgoing traverse. In some samples taken on the Antarctic plateau, the relative abundance of sulfate particles in fine mode reached 100 % in incoming and outgoing traverses. Because of the number of analyzed particles in each sample in fine mode (mean, 958 particles/one sample), the upper limit of relative abundance of aerosol particles other than sulfate particles might be less than 0.1 % in these samples (100 % abundance of sulfate particles). Higher relative abundance of sulfate particles in the outgoing traverse is likely to result from (1) increase of the number concentrations of sulfate particles through new particle formation and particle growth, and (2) decrease of the number concentration of sea-salt particles and the modified sea-salt particles during the summer. Indeed, mass concentrations of nss-SO₄²⁻ and sea-salts (Na⁺) respectively showed a maximum and minimum in the summer (January–February) in Antarctic coast and inland (e.g., Minikin et al., 1998; Hara et al., 2004; Weller and Wagenbach, 2007; Preunkert et al., 2008; Udisti et al., 2012). The relative abundance of sulfate particles in fine mode was 82.9–

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



98.2% (mean, 93.1%) in the boundary layer and 96.2–99.7% (mean, 98.1%) in the lower free troposphere over Syowa Station, Antarctica during mid-November through January (Hara et al., 2013). Consequently, the relative abundance of sulfate particles in fine mode near the surface on the Antarctic continent was similar to that in the lower free troposphere over the Antarctic coast (Syowa Station). The relative abundance of sulfate particles in coarse mode often exceeded 40% on the Antarctic continent during the JASE traverse. Such a high relative abundance of coarse sulfate particles was not observed in the boundary layer but often in the lower free troposphere over Syowa Station (Hara et al., 2013). Figure 5 shows that the aerosol number concentrations in coarse and fine modes were lower near the surface on the Antarctic continent. Consequently, sulfate particles can be grown to coarse mode through coagulation and condensation of condensable vapors (e.g., H_2SO_4 gas) under conditions with the low number concentrations of pre-existing particles on the Antarctic continent.

The relative abundance of K-rich sulfate particles was not detected (n.d.) –2.0% in coarse mode and n.d. –0.9% in fine mode in this study. Although K-rich sulfate particles were obtained in a few samples in coarse mode, these particles were observed often in fine mode in this study. Moreover, K-rich sulfate particles were often observed in the incoming traverse. In addition, the relative abundance of K-rich sulfate particles in the incoming traverse was higher than that in the outgoing traverse. This difference implies seasonal features of K-rich sulfate particles in the inland area during late-spring – summer. K-rich sulfate particles were distributed also in boundary layer – upper free troposphere over Syowa Station (Hara et al., 2010, 2013). As discussed above and by Hara et al. (2013), K-rich sulfate particles might have originated from biomass burning and fossil fuel combustion. Figure 3 shows that the backward trajectory exhibited direct transport from outside of the Antarctic Circle was not found for the prior 5 days. Moreover, source strength of combustion of fossil fuel is extremely low on the Antarctic regions. Particularly, biomass burning does not occur on the Antarctic regions. Therefore, sulfate particles containing K are likely to be transported to the Antarctic continent via free troposphere during summer for the prior 5 days. The presence of combustion-

origin aerosol particles (e.g., sulfate particles containing K) was supported by high BC concentrations at the South Pole in the summer (Bodhaine, 1995).

The relative abundance of Mg-rich sulfate particles (probably MgSO_4) examined in this study was n.d. -1.8% in coarse mode and n.d. -0.5% in fine mode. During the incoming traverse, Mg-rich sulfate particles were found frequently in fine mode at sampling sites in $69.2\text{--}72.5^\circ\text{S}$, although those were observed in some aerosol samples collected at plateau sites. However, Mg-rich sulfate particles were identified mainly on the Antarctic plateau during the outgoing traverse. The relative abundance of Mg-rich sulfate particles was lower in the outgoing traverse than that in the incoming traverse. Mg-rich sulfate particles were distributed mostly in fine mode over Syowa Station (Hara et al., 2013). Previous investigation (Hara et al., 2013) indicated that Mg-rich sulfate particles were associated with sea-salt fractionation (Mg separation) and modification (Cl loss by heterogeneous reactions). High Mg-enrichment was observed in aerosols, surface snow, and drifting snow at Dome F during summer (Hara et al., 2014). Consequently, the presence of Mg-rich sulfate particles in the atmosphere near the surface on the Antarctic continent in this study strongly suggests that sea-salt fractionation occurs on the Antarctic continent during summer. Details of sea-salt fractionation are discussed in Sect. 3.7. and by Hara et al. (2014).

Relative abundance of Ca-rich sulfate particles (probably CaSO_4) was n.d. -8.7% in coarse mode and n.d. -0.2% in fine mode in incoming traverse. However, the relative abundance of Ca-rich sulfate particles was n.d. in coarse mode and n.d. -0.1% in outgoing traverse. Although Ca-rich sulfate particles were not detected in most samples in both traverses, Ca-rich sulfate particles were identified mainly near coasts ($< 72^\circ\text{S}$). Although Ca-rich sulfates (e.g. gypsum) are mineral components, most Ca-rich sulfates were not mixed with Al and Si, which are major elements of mineral particles. Consequently, horizontal distributions and external mixing states of Ca-rich sulfates imply the important contribution of non-mineral origins such as sea-salt fractionation on sea-ice as suggested by Geilfus et al. (2013).

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5.3 Minerals

In spite of a smaller source area of mineral particles on the Antarctic continent, aerosol particles containing Al and Si were identified during the JASE traverse. Here, aerosol particles containing Si and Al were classified into “mineral particles”. The relative abundance of mineral particles was n.d. –14.6% (mean, 3.8%) in coarse mode and n.d. –0.4% (mean, 0.1%) in fine mode during the incoming traverse, although the relative abundance was n.d. –5.7% (mean, 1.8%) in coarse mode and n.d. –0.1% (mean, 0.01%) in fine mode during the outgoing traverse. Most mineral particles were internally mixed with sea-salts or sulfates in this study. Although the mineral particles were observed mainly in coarse mode, SEM observation showed that the size of irregular solid cores containing Al and Si ranged in sub-micrometer in this study. The size of irregular solid cores was coincident with the size distribution of water-insoluble particles (mainly mineral particles) in ice cores taken in Antarctica (e.g., Ram and Gayley, 1988; Delmonte et al., 2004, 2007). In contrast, external mixing states of mineral particles were often present in the boundary layer and free troposphere over Syowa Station located at the coast (Hara et al., 2006, 2013). The difference of mixing states of mineral particles suggests that mixing states of mineral particles changed gradually to internal mixtures through coagulation in cloud processes, and through heterogeneous reactions during their transport to inland areas. As described above, a change of mixing state of mineral particles can lead to particle size greater than the external mixing states of mineral particles. Because of the higher dry deposition velocity of coarse aerosol particles, internal mixing among minerals, sulfate, and sea-salts during transport might enhance the efficient deposition of minerals onto the snow surface on the Antarctic continent.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SO_4^{2-} and CH_3SO_3^- . Previous investigations (Mouri et al., 1999; Hara et al., 2005, 2013) revealed the high contribution of acidic sulfur species such as SO_4^{2-} and CH_3SO_3^- to sea-salt modification in fine mode in the atmosphere on the Antarctic coastal regions during summer.

3.6.2 Horizontal features of sea-salt modification on the Antarctic continent

Figures 11 and 12 show horizontal features of the atomic ratios of Cl/Na and S/Na of sea-salt particles and the modified sea-salt particles in coarse and fine modes during the incoming and outgoing traverses, respectively. The internal mixing particles between sea-salts and minerals were excluded from the box plots. Latitudinal gradient of Cl/Na ratios in coarse mode was observed in 69–71° S in the incoming traverse. High Cl depletion was identified in most aerosol samples obtained for the Antarctic plateau, except some samples taken at 76.5–76° S in the incoming traverse from Dome F to the meeting point. In spite of the latitudinal gradient of Cl/Na ratios in coarse mode in 69–71° S in the incoming traverse, the S/Na ratio increased slightly. Similarly, median S/Na ratios of sea-salt particles and the modified sea-salt particles in coarse mode were distributed approximately in 0.2 during the JASE traverse. When sea-salt particles are modified solely with SO_4^{2-} , S/Na ratios in sea-salt particles are 0.5. As suggested in ternary plots in Fig. 10. Therefore, NO_3^- contributed dominantly to sea-salt modification in coarse mode on the Antarctic continent. By contrast, S/Na ratios in fine sea-salt particles exceeded mostly 0.5 during the traverse. The high S/Na ratios in each sea-salt particle in fine mode imply that sulfates were formed on the fine sea-salt particles through heterogeneous reactions with gaseous sulfur species such as H_2SO_4 and SO_2 .

As described above, the acids contributing to sea-salt modification differed between those in coarse sea-salt particles on the Antarctic coasts and those on the continent during summer. The air mass history and origins of coarse sea-salt particles, NO_3^- , and its precursors on the Antarctic continent must be discussed to elucidate high con-

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ratios in the modified sea-salt particles were also obtained in boundary layer and lower free troposphere over Syowa Station during summer (Hara et al., 2013). Because of the longer residence time of fine aerosol particles, the modified sea-salt particles in fine mode with high S/Na ratio might be supplied by transport from coastal regions to the continent.

3.7 Sea-salt fractionations on the Antarctic continent during summer

3.7.1 Sea-salt fractionations in coarse and fine mode

Figure 13 shows examples of ternary plots of sea-salts (Na, Mg, and S) and Mg-rich sulfates in coarse and fine modes. Internal mixtures of sea salts and minerals were removed from the ternary plots. In the ternary plots, sea-salt particles with bulk seawater ratio are distributed around the red star (bulk seawater ratio). When the sea-salt particles are modified by sulfate and not fractionated, they are distributed around the stoichiometric line between the red star (seawater ratio) and the blue star (modified sea-salt ratio with sulfate). With sea-salt fractionation by precipitation of Na-salts such as mirabilite ($\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$) and hydrohalite ($\text{NaCl} \cdot 2\text{H}_2\text{O}$) (Hara et al., 2012), Mg in sea-salt particles can be enriched gradually. For cases in which sea-salt fractionation (replacement between Na and Mg) occurs without sea-salt modification by sulfate, sea-salt particles are distributed around the stoichiometric line between the red star (bulk seawater ratio) and the cyan star (MgCl_2). When sea-salt fractionation and sea-salt modification by sulfate occur stoichiometrically and simultaneously, sea-salt particles are distributed around the stoichiometric line between the red star (seawater ratio) and green star (MgSO_4).

Because sea-salt particles were modified dominantly during JARE (as shown in Figs. 10–12), most sea-salt particles and the modified sea-salt particles in coarse mode were distributed in 60–90% of the Na atomic ratio. The Mg ratios in coarse mode, however, were greater than the stoichiometric line of sea-salt modification (between the red star and blue star). Some of the modified sea-salt particles had Mg ratio ≈ 0 , even in coarse

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mode. The Mg ratios in fine mode were distributed at Mg ratio ≈ 0 and around the stoichiometric line between the red star (seawater) and green star (MgSO_4). Here, we designate sea-salt particles with Mg ratio ≈ 0 as “Mg-free sea-salt particles”. In addition to Mg-free sea-salts, MgSO_4 particles were often observed in this study, as shown in Figs. 7–9 and 13. Similar distributions were observed in sea-salt particles and the modified sea-salt particles collected over Syowa Station (Hara et al., 2013). Mg/Na ratios cannot be changed by sea-salt modification. Sea-salt fractionation in sea-ice regions during winter through spring (e.g., precipitation of mirabilite and hydrohalite) can promote Mg enrichment in sea-salt particles, as reported by Hara et al. (2012). Consequently, sea-salt fractionation in sea-ice regions cannot account for the presence of Mg-Free sea-salt particles during spring–summer. Because Mg-rich sulfate particles over Syowa Station were identified in the air masses from the Antarctic continent (Hara et al., 2013), discussion must address horizontal features of Mg-rich sulfates, Mg-rich sea-salt particles and Mg-free sea-salt particles on the Antarctic regions. Then it will be possible to assess the possibility of sea-salt fractionation on the Antarctic continent.

3.7.2 Horizontal features of sea-salt fractionation on the Antarctic continent

To elucidate sea-salt fractionation on the Antarctic continent during summer, we must compare horizontal features of sea-salt constituents on the Antarctic continent. Figures 14 depict horizontal features of Mg/Na ratios in coarse mode during the JASE traverse. Horizontal features of Mg/Na ratios in fine mode were added to “Supplement”. Internal mixtures of sea salts and minerals were removed from the plots. Median Mg/Na ratios in coarse mode were approximately 0.32 in the incoming traverse and 0.31 in the outgoing traverse. Because of Mg/Na ratio ≈ 0.11 in bulk seawater (e.g., Wilson, 1975), Mg might be enriched in most sea-salt particles in coarse mode. As shown in the ternary plots (Fig. 13), Mg/Na ratios varied greatly because of presence of Mg-rich sea-salt particles, Mg-free sea-salt particles, and MgSO_4 particles. In addition, Mg-free sea-salt particles (Mg/Na ratio ≈ 0) in coarse mode were identified often on the Antarctic continent, particularly on the Antarctic plateau. For example,

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mg-free sea-salt particles in coarse mode were dominant in three samples taken on 21–23 December 2007 (76.6–76.0° S) in the traverse from Dome F to the meeting point, as shown in Fig. 14. The presence of Mg-free sea-salt particles in coarse mode corresponded often to occurrence of drifting snow and high aerosol number concentrations in coarse mode in the incoming traverse. Mg-rich sea-salt particles, Mg-free sea-salt particles, and MgSO₄ particles were present also in fine mode on the Antarctic continent, although the low relative abundance of sea-salt particles and the modified sea-salt particles prevented us from elucidating the horizontal features of Mg/Na ratios on the Antarctic plateau. These horizontal features of Mg/Na ratios during the JASE traverse imply strongly that the fractionated sea-salt particles were distributed widely throughout the Antarctic continent during summer. Mg-free sea-salt particles were identified mainly in fine mode and rarely in coarse mode over Syowa Station (Hara et al., 2013). Figure 3 shows that air masses on the Antarctic plateau were isolated from coastal regions for the prior 5 days. The lower relative abundance of Mg-free sea-salt particles in coarse mode over Syowa Station suggests that Mg-free sea-salt particles and other fractionated sea-salt particles such as Mg-rich sulfates and Mg-rich sea-salts had not originated from coastal regions, considering the efficient dry deposition of aerosol particles in coarse mode. Aerosol particles on the Antarctic plateau, especially in coarse mode, were supplied to a marked degree from snow surface by wind blowing under strong conditions, as described above. Furthermore, Mg-rich sulfates and Mg-free NaCl were present in surface snow (Iizuka et al., 2012). Therefore, the fractionated sea-salt particles (Mg-rich sea-salt particles, Mg-free sea-salt particles, and MgSO₄ particles) might be originated from surface snow on the Antarctic continent. Considering that Mg-free sea-salt particles were often observed under strong wind conditions (especially on 21–23 December), strong wind blowing might be necessary for the release of Mg-free sea-salt particles.

To explain the presence of Mg-rich sea-salt particles, Mg-free sea-salt particles and MgSO₄ particles on the snow surface of the Antarctic continent, it is necessary to discuss Mg separation processes. Redistribution of chemical constituents can occur

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



through (1) seawater freezing (Marion and Farren, 1999; Hara et al., 2012), and (2) phase transformation by deliquescence and efflorescence (e.g., Ge et al., 1998; Wise et al., 2009; Woods et al., 2010). Sea-salt fractionation in seawater freezing depends on the temperature (Marion and Farren, 1999; Hara et al., 2012). The air temperature on the Antarctic plateau was often below -30°C even during summer, as presented in Fig. 2 and previous investigations (King et al., 2006; Hirasawa et al., 2013). Air temperature might constitute an important condition for some sea-salt precipitation (e.g., ca. -34°C and -36°C for KCl and $\text{MgCl}_2 \cdot 12\text{H}_2\text{O}$) as discussed by Marion and Farren (1999). Unlike sea-salts in seawater, however, sea-salts in snow on the Antarctic content were supplied solely by deposition of sea-salt particles that had been transported from coastal regions. When Mg separation is controlled only by lower temperature, Mg-free sea-salt particles, Mg-rich sea-salt particles, and Mg-rich sulfate particles can be present on the Antarctic regions. However, Mg-free sea-salt particles were not observed over Syowa Station during winter in the air masses transported from the continent and coastal regions (Hara et al., 2013). This result implies that Mg separation was controlled not only by lower air temperature but also by other factors.

Figure 2 show that the air temperature near the surface had diurnal change also in continental areas. Furthermore, strong diurnal change of air temperatures and solar radiation engender water sublimation on the snow surface of the Antarctic continent (e.g., Kameda et al., 1997; Motoyama et al., 2005). With strong diurnal change of the air temperature, the relative humidity might exhibit a diurnal change near the snow surface on the Antarctic continent during summer. Actually, relative humidity reached ca. 100% during the local night time and reached a minimum in the afternoon at Kohonen Station during summer (Van As et al., 2005). The diurnal variation of the relative humidity can engender (1) phase transformation by deliquescence of available sea-salts on surface snow, (2) condensation and re-freezing of water vapor (e.g., formation of surface hoar) during local night time, and (3) enhancement of quasi-liquid layer and super-cold liquid on the surface snow. Laboratory experiments conducted for earlier studies (e.g., Ge et al., 1998; Wise et al., 2009; Woods et al., 2010) revealed that

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



chemical constituents with lower deliquescence relative humidity (DRH) can be localized in the outer layer (surface) around a solid core through phase transformation by deliquescence. Although relative humidity was minimal in the afternoon at Kohlen Station (Van As et al., 2005), the minimum relative humidity ($\sim 89\%$) was often higher than the deliquescence relative humidity (DRH) of plausible sea-salts. Some examples are the following: NaCl, 75 % (Tang and Munkelwitz, 1993); Na₂SO₄, 84 % (Tang and Munkelwitz, 1994); NaNO₃, 75 % (Tang and Munkelwitz, 1994); MgCl₂·6H₂O, 33 % (Kelly and Wexler, 2005); MgSO₄, 42 % (Wang et al., 2008); and KCl, 84 % (Tang, 1980). Most plausible sea-salts can be wholly deliquescent even under minimum relative humidity ($\sim 89\%$) on the Antarctic continent. Therefore, phase transformation by deliquescence/efflorescence might not be key processes for Mg separation. With diurnal features of air temperature and relative humidity, re-condensation of water vapor on the snow surface during the local night time might induce enhancement of super-cold water in nanometer-to-micrometer scales, quasi-liquid layer, and re-freezing of water vapors and super-cold water such as hoar formation. Particularly, re-freezing processes on surface snow might cause sea-salt fractionation as well as seawater freezing. Therefore, we propose that the repetition of a diurnal cycle of relative humidity and water sublimation under colder conditions can engender Mg separation on the surface snow on the Antarctic continent during summer.

4 Concluding remarks

Measurements of aerosol size distribution and direct aerosol sampling were made in the Queen Maud Land, Antarctica during the JASE traverse from 14 November 2007 until 24 January 2008. OPC measurements revealed that aerosol number concentrations decreased gradually with latitude under background conditions (without drifting snow or strong winds) on the Antarctic continent during summer. The estimated aerosol mass concentrations in size range larger than $0.3\mu\text{m}$ were $0.04\text{--}5.7\mu\text{g m}^{-3}$. When strong winds and drifting snow occurred, aerosol number concentrations increased

drastically, especially in coarse mode. Air masses were isolated from the Antarctic coasts during the prior 5 days. Therefore, coarse aerosol particles (mainly sea-salt particles) might be released from the snow surface by blowing winds.

Single-particle analysis using SEM-EDX revealed that major aerosol particles were sea-salt particles, modified sea-salt particles, and sulfate particles in coarse mode, and that the sulfate particles and modified sea-salt particles were dominant in fine mode during the JASE traverse. The K-rich sulfates, Mg-rich sulfates, Ca-rich sulfates, and minerals were minor aerosol constituents in coarse and fine modes. Although SO_4^{2-} and CH_3SO_3^- contributed to sea-salt fractionation in coarse and fine modes over Syowa Station during summer (Hara et al., 2005, 2013), sea-salt particles were modified greatly with SO_4^{2-} and NO_3^- in coarse mode, and dominantly with SO_4^{2-} in fine mode in this study. Precursors of particulate NO_3^- (e.g., HNO_3 , and NO_x) might have originated from photochemical recycle of NO_3^- in surface snow and then oxidation in the atmosphere. Median atomic ratios of Mg/Na in sea-salt particles and modified sea-salt particles in coarse and fine modes were often higher than the bulk seawater ratio. In addition, Mg-free sea-salt particles were identified on the Antarctic continent. Because the Mg/Na ratio cannot be changed by sea-salt modification, the presence of Mg-rich sea-salt particles, Mg-free sea-salt particles, and Mg-rich sulfate particles might be associated with sea-salt fractionation. Mg-free sea-salt particles in coarse mode were often identified under conditions with strong winds and drifting snow. Therefore, fractionated sea-salt particles were likely to have been released from the snow surface. This study proposed and assessed the hypothesis that sea-salt fractionation (Mg separation in sea-salts) occurs in surface snow on the Antarctic continent.

Supplementary material related to this article is available online at
[http://www.atmos-chem-phys-discuss.net/14/11393/2014/
acpd-14-11393-2014-supplement.pdf](http://www.atmos-chem-phys-discuss.net/14/11393/2014/acpd-14-11393-2014-supplement.pdf).

Acknowledgements. The JASE traverse was organized by several organizations both in Sweden and Japan. The National Institute of Polar Research (NIPR), Tokyo and the Swedish Polar

Horizontal
distributions of
aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Research Secretariat (SPRS) managed the logistics in Antarctica. Science management was a collaborative effort of NIPR, Stockholm University, the Royal Institute of Technology in Stockholm and individuals from several universities and institutes in Japan. The JASE traverse is one research project undertaken by the Japanese Antarctic Research Expedition “Studies on systems for climate change and ice sheet change, by introducing new observational methods and technologies”. We thank JARE48 wintering members and JARE49 members for logistic support of the JASE traverse. Special thanks are extended to the logistics members, S. Gunnarsson, H. Kaneko, T. Karlberg, P. Ljusberg and K. Taniguchi and the medical doctors, S. Eriksson and N. Shiga, for their very generous support during the traverse. This research was supported by the Swedish Research Council (VR) and by a Grant-in-Aid for Scientific Research (A) 20241007 from the Japan Society for the Promotion of Science (JSPS).

References

- Allison, I.: Surface climate of the interior of the Lambert Glacier basin, Antarctica, from automatic weather station data, *Ann. Glaciol.*, 27, 515–520, 1998.
- Artaxo, P., Rabello, M. L. C., Maenhaut, W., and van Grieken, R.: Trace elements and individual particle analysis of atmospheric aerosols from the Antarctic Peninsula, *Tellus*, 44, 318–334, 1992.
- Bartels-Rausch, T., Jacobi, H.-W., Kahan, T. F., Thomas, J. L., Thomson, E. S., Abbatt, J. P. D., Ammann, M., Blackford, J. R., Bluhm, H., Boxe, C., Domine, F., Frey, M. M., Gladich, I., Guzmán, M. I., Heger, D., Huthwelker, Th., Klán, P., Kuhs, W. F., Kuo, M. H., Maus, S., Moussa, S. G., McNeill, V. F., Newberg, J. T., Pettersson, J. B. C., Roeselová, M., and Sodeau, J. R.: A review of air–ice chemical and physical interactions (AICI): liquids, quasi-liquids, and solids in snow, *Atmos. Chem. Phys.*, 14, 1587–1633, doi:10.5194/acp-14-1587-2014, 2014.
- Bertler, N., Mayewski, P. A., Aristarain, A., Barrett, P., Becagli, S., Bernardo, R. T., Cunde, X., Curran, M., Dahe, Q., Dixon, D., Ferron, F. A., Fischer, H., Frey, M., Frezzoi, M., Fundel, F., Genthon, C., Gragani, R., Hamilton, G., Handley, M., Hong, S., Isaksson, E., Jiawen, R., Kamiyama, K., Kanamori, S., Karkas, E., Karlöf, L., Kaspari, S., Kreutz, K., Kurbatov, A., Meyerson, E., Motoyama, H., Mulvaney, R., Mingjun, Z., Oerter, H., Osterberg, E., Proposito, M., Pyne, A., Ruth, U., Simoes, J. C., Smith, B., Sneed, S., Teinila, K., Traufetter, F.,

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Udisti, R., Virkkula, A., Watanabe, O., Williamson, B., Wolff, E., and Zhongqin, L.: Snow Chemistry Across Antarctica, *Ann. Glaciol.*, 41, 167–179, 2005.
- Bodhaine, B. A.: Aerosol absorption measurements at Barrow, Mauna Loa and the South Pole, *J. Geophys. Res.*, 100, 8967–8975, 1995.
- 5 Davis, D., Nowak, J. B., Chen, G., Buhr, M., Arimoto, R., Hogan, A., Eisele, F., Mauldin, L., Tanner, D., Shetter, R., Lefer, B., and McMurry, P.: Unexpected high levels of NO observed at South Pole, *Geophys. Res. Lett.*, 28, 3625–3628, 2001.
- Davis, D., Chen, G., Buhr, M., Crawford, J., Lenschow, D., Lefer, B., Shetter, R., Eisele, F., Mauldin, L., and Hogan, A.: South Pole NO_x chemistry: an assessment of factors controlling variability and absolute levels, *Atmos. Environ.*, 38, 5375–5388, 2004.
- 10 Delmonte, B., Petit, J. R., Andersen, K. K., Basile-Doelsch, I., Maggi, V., Ya Lipenkov, V.: Dust size evidence for opposite regional atmospheric circulation changes over east Antarctica during the last climatic transition, *Clim. Dynam.*, 23, 427–438, doi:10.1007/s00382-004-0450-9, 2004.
- 15 Delmonte, B., Petit, J. R., Basile-Doelsch, I., Jagoutz, E., Maggi, V.: Late quaternary interglacials in East Antarctica from ice-core dust records, in: *Developments in Quaternary Sciences*, edited by: Sirocko, F., Claussen, M., Fernanda, M., Goñi, S., and Litt, T., vol. 7, Elsevier, 53–73, ISSN 1571-0866, ISBN 9780444529558, 2007.
- Dibb, J., Huey, L., Slusher, D., and Tanner, D.: Soluble reactive nitrogen oxides at South Pole during isCAT 2000, *Atmos. Environ.*, 38, 5399–5409, doi:10.1016/j.atmosenv.2003.01.001, 2004.
- 20 Draxler, R. R. and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY Website, available at: <http://ready.arl.noaa.gov/HYSPLIT.php> (last access: 7 May 2014), NOAA Air Resources Laboratory, Silver Spring, MD, 2013.
- 25 Eisele, F., Davis, D. D., Helmig, D., Oltmans, S. J., Neff, W., Huey, G., Tanner, D., Chen, G., Crawford, J., Arimoto, R., Buhr, M., Mauldin, L., Hutterli, M., Dibb, J., Blake, D., Brooks, S. B., Johnson, B., Roberts, J. M., Wang, Y., Tan, D., and Flocke, F.: Antarctic Tropospheric Chemistry Investigation (ANTCI) 2003 overview, *Atmos. Environ.*, 42, 2749–2761, doi:10.1016/j.atmosenv.2007.04.013, 2008.
- 30 Frey, M. M., Savarino, J., Morin, S., Erbland, J., and Martins, J. M. F.: Photolysis imprint in the nitrate stable isotope signal in snow and atmosphere of East Antarctica and implications for

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reactive nitrogen cycling, *Atmos. Chem. Phys.*, 9, 8681–8696, doi:10.5194/acp-9-8681-2009, 2009.

Fujita, S., Holmlund, P., Andersson, I., Brown, I., Enomoto, H., Fujii, Y., Fujita, K., Fukui, K., Furukawa, T., Hansson, M., Hara, K., Hoshina, Y., Igarashi, M., Iizuka, Y., Imura, S., Ingvander, S., Karlin, T., Motoyama, H., Nakazawa, F., Oerter, H., Sjöberg, L. E., Sugiyama, S., Surdyk, S., Ström, J., Uemura, R., and Wilhelms, F.: Spatial and temporal variability of snow accumulation rate on the East Antarctic ice divide between Dome Fuji and EPICA DML, *The Cryosphere*, 5, 1057–1081, doi:10.5194/tc-5-1057-2011, 2011.

Fujita, S., Fukui, K., Nakazawa, F., Enomoto, H., and Sugiyama, S.: Report of the Japanese-Swedish joint Antarctic traverse: II. Report of field expedition in 2007/2008 season, Submitted to Antarctic record, in review, 2014 (in Japanese with English abstract).

Ge, Z., Wexler, A. S., and Johnston, M. V.: Deliquescence behavior of multicomponent aerosols, *J. Phys. Chem. A*, 102, 173–180, 1998.

Geilfus, N.-X., Galley, R. J., Cooper, M., Halden, N., Hare, A., Wang, F., Søgaard, D. H., and Rysgaard, S.: Gypsum crystals observed in experimental and natural sea ice, *Geophys. Res. Lett.*, 40, 6362–6367, doi:10.1002/2013GL058479, 2013.

Hara, K., Kikuchi, T., Furuya, K., Hayashi, M., and Fujii, Y.: Characterization of Antarctic aerosol particles using laser microprobe mass spectrometry, *Environ. Sci. Technol.*, 30, 385–391, 1995.

Hara, K., Osada, K., Kido, M., Hayashi, M., Matsunaga, K., Iwasaka, Y., Yamanouchi, T., Hashida, G., and Fukatsu, T.: Chemistry of sea-salt particles and inorganic halogen species in the Antarctic regions: compositional differences between coastal and inland stations, *J. Geophys. Res.*, 109, D20208, doi:10.1029/2004JD004713, 2004.

Hara, K., Osada, K., Kido, M., Matsunaga, K., Iwasaka, Y., and Hashida, G.: Seasonal variations of sea-salt constituents and sea-salt modification at Syowa station, Antarctica, *Tellus B*, 57, 230–246, 2005.

Hara, K., Osada, K., Yabuki, M., Hashida, G., Yamanouchi, T., Hayashi, M., Shiobara, M., Nishita-Hara, C., and Wada, M.: Haze episodes at Syowa Station, coastal Antarctica: where did they come from?, *J. Geophys. Res.*, 115, D14205, doi:10.1029/2009JD012582, 2010a.

Hara, K., Hirasawa, N., Yamanouchi, T., Wada, M., Herber, A., and ANTSYO-II members, Spatial distributions and mixing states of aerosol particles in the summer Antarctic troposphere, *Antarctic Record*, 54, 704–730, 2010b (in Japanese with English abstract).

**Horizontal
distributions of
aerosol constituents**

K. Hara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Hara, K., Osada, K., Nishita-Hara, C., Yabuki, M., Hayashi, M., Yamanouchi, T., Wada, M., and Shiobara, M.: Seasonal features of ultrafine particle volatility in the coastal Antarctic troposphere, *Atmos. Chem. Phys.*, 11, 9803–9812, doi:10.5194/acp-11-9803-2011, 2011a.
- Hara, K., Osada, K., Nishita-Hara, C., and Yamanouchi, T.: Seasonal variations and vertical features of aerosol particles in the Antarctic troposphere, *Atmos. Chem. Phys.*, 11, 5471–5484, doi:10.5194/acp-11-5471-2011, 2011b.
- Hara, K., Osada, K., Yabuki, M., and Yamanouchi, T.: Seasonal variation of fractionated sea-salt particles on the Antarctic coast, *Geophys. Res. Lett.*, 39, L18801, doi:10.1029/2012GL052761, 2012.
- Hara, K., Osada, K., and Yamanouchi, T.: Tethered balloon-borne aerosol measurements: seasonal and vertical variations of aerosol constituents over Syowa Station, Antarctica, *Atmos. Chem. Phys.*, 13, 9119–9139, doi:10.5194/acp-13-9119-2013, 2013.
- Hirasawa, N., Nakamura, H., Motoyama, H., Hayashi, M., and Yamanouchi, T.: The role of synoptic-scale features and advection in prolonged warming and generation of different forms of precipitation at Dome Fuji station, Antarctica, following a prominent blocking event, *J. Geophys. Res. Atmos.*, 118, 6916–6928, doi:10.1002/jgrd.50532, 2013.
- Huey, L. G., Tanner, D. J., Slusher, D. L., Dibb, J. E., Arimoto, R., Chen, G., Davis, D., Buhr, M. P., Nowak, J. B., Mauldin III, R. L., Eisele, F. L., and Kosciuch, E.: CIMS measurements of HNO₃ and SO₂ at the South Pole during ISCAT 2000, *Atmos. Environ.*, 38, 5411–5421, doi:10.1016/j.atmosenv.2004.04.037, 2004.
- Iizuka, Y., Tsuchimoto, A., Hoshina, Y., Sakurai, T., Hansson, M., Karlin, T., Fujita, K., Nakazawa, F., Motoyama, H., and Fujita, S.: The rates of sea salt sulfatization in the atmosphere and surface snow of inland Antarctica, *J. Geophys. Res.*, 117, D04308, doi:10.1029/2011JD016378, 2012.
- Jones, A. E., Wolff, E. W., Ames, D., Bauguitte, S. J.-B., Clemitshaw, K. C., Fleming, Z., Mills, G. P., Saiz-Lopez, A., Salmon, R. A., Sturges, W. T., and Worton, D. R.: The multi-seasonal NO_y budget in coastal Antarctica and its link with surface snow and ice core nitrate: results from the CHABLIS campaign, *Atmos. Chem. Phys.*, 11, 9271–9285, doi:10.5194/acp-11-9271-2011, 2011.
- Jourdain, B. and Legrand, M.: Seasonal variations of atmospheric dimethylsulfide, dimethylsulfoxide, sulfur dioxide, methanesulfonate, and non-sea-salt sulfate aerosols at Dumont d'Urville (coastal Antarctica) (December 1998 to July 1999), *J. Geophys. Res.*, 106, 14391–14408, 2001.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jourdain, B. and Legrand, M.: Year-round records of bulk and size-segregated aerosol composition and HCl and HNO₃ levels in the Dumont d'Urville (coastal Antarctica) atmosphere: implications for sea-salt aerosol fractionation in the winter and summer, *J. Geophys. Res.*, 107, 4645, doi:10.1029/2002JD002471, 2002.

5 Junge, C. E.: Aerosols, in: *Air Chemistry and Radioactivity*, Academic Press, 111–208, 1963.

Kameda, T., Azuma, N., Furukawa, T., Ageta, Y., and Takahashi, S.: Surface mass balance, sublimation and snow temperatures at Dome Fuji Station, Antarctica, in 1995, *Proc. NIPR Symp. Polar Meteorol. Glaciol.*, 11, 24–34, 1997.

10 Kelly, J. T. and Wexler, A. S.: Thermodynamics of carbonates and hydrates related to heterogeneous reactions involving mineral aerosol, *J. Geophys. Res.*, 110, D11201, doi:10.1029/2004JD005583, 2005.

Kerminen, V. M., Teinilä, K., and Hillamo, R.: Chemistry of sea-salt particles in the summer Antarctic atmosphere, *Atmos. Environ.*, 34, 2817–2825, 2000.

15 King, J. C., Argentini, S. A., and Anderson, P. S.: Contrasts between the summertime surface energy balance and boundary layer structure at Dome C and Halley stations, Antarctica, *J. Geophys. Res.*, 111, D02105, doi:10.1029/2005JD006130, 2006.

Legrand, M., Sciare, J., Jourdain, B., and Genthon, C.: Subdaily variations of atmospheric dimethylsulfide, dimethylsulfoxide, methanesulfonate, and non-sea-salt sulfate aerosols in the atmospheric boundary layer at Dumont d'Urville (coastal Antarctica) during summer, *J. Geophys. Res.*, 106, 14409–14422, 2001.

20 Marion, G. M. and Farren, R. E.: Mineral solubilities in the Na–K–Mg–Ca–Cl–SO₄–H₂O system: a re-evaluation of the sulfate chemistry in the Spencer–Møller–Weare model, *Geochim. Cosmochim. Ac.*, 63, 1305–1318, doi:10.1016/S0016-7037(99)00102-7, 1999.

25 Minikin, A., Legrand, M., Hall, J., Wagenbach, D., Kleefeld, C., Wolff, E., Pasteur, E., and Ducroz, F.: Sulfur-containing species (sulfate and methanesulfonate) in coastal Antarctic aerosol and precipitation, *J. Geophys. Res.*, 103, 10975–10990, 1998.

Motoyama, H., Hirasawa, N., Satow, K., and Watanabe, O.: Seasonal variations in oxygen isotope ratios of daily collected precipitation and wind drift samples and in the final snow cover at Dome Fuji Station, Antarctica, *J. Geophys. Res.*, 110, D11106, doi:10.1029/2004JD004953, 2005.

30 Mouri, H., Nagao, I., Okada, K., Koga, S., and Tanaka, H.: Individual-particle analyses of coastal Antarctic aerosols, *Tellus B*, 51, 603–611, 1999.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Neff, W., Helmig, D., Grachev, A., and Davis, D.: A study of boundary layer behavior associated with high NO concentrations at the South Pole using a minisodar, tethered balloon, and sonic anemometer, *Atmos. Environ.*, 42, 2762–2779, doi:10.1016/j.atmosenv.2007.01.033, 2008.
- Preunkert, S., Jourdain, B., Legrand, M., Udisti, R., Becagli, S., and Cerri, O.: Seasonality of sulfur species (dimethylsulfide, sulfate, and methanesulfonate) in Antarctica: inland vs. coastal regions, *J. Geophys. Res.*, 113, D15302, doi:10.1029/2008JD009937, 2008.
- Rankin, A. M., Auld, V., and Wolff, E. W.: Frost flowers as a source of fractionated sea salt aerosol in the polar regions, *Geophys. Res. Lett.*, 27, 3469–3472, doi:10.1029/2000GL011771, 2000.
- Rankin, A. M., Wolff, E. W., and Martin, S.: Frost flowers: implications for tropospheric chemistry and ice core interpretation, *J. Geophys. Res.*, 107, 4683, doi:10.1029/2002JD002492, 2002.
- Rankin, A. M. and Wolff, E. W.: A year-long record of size-segregated aerosol composition at Halley, Antarctica, *J. Geophys. Res.*, 108, 4775, doi:10.1029/2003JD003993, 2003.
- Sato, K. and Hirasawa, N.: Statistics of Antarctic surface meteorology based on hourly data in 1957–2007 at Syowa Station, *Polar Sci.*, 1, 1–15, 2007.
- Savoie, D. L., Prospero, J. M., Larsen, R. J., and Saltzman, E. S.: Nitrogen and sulfur species in aerosols at Mawson, Antarctica, and their relationship to natural radionuclides, *J. Atmos. Chem.*, 14, 181–204, 1992.
- Savoie, D. L., Prospero, J. M., Larsen, R. J., Huang, F., Izaguirre, M. A., Huang, T., Snowdon, T. H., Custals, L., and Sanderson, C. G.: Nitrogen and sulfur species in Antarctic aerosols at Mawson, Palmer Station, and Marsh (King George Island), *J. Atmos. Chem.*, 17, 95–122, 1993.
- Suzuki, K., Yamanouchi, T., Kawamura, K., and Motoyama, H.: The spatial and seasonal distributions of air-transport origins to the Antarctic based on 5 day backward trajectory analysis, *Polar Sci.*, 7, 205–213, 2013.
- Teinilä, K., Kerminen, V.-M., and Hillamo, R.: A study of size-segregated aerosol chemistry in the Antarctic atmosphere, *J. Geophys. Res.*, 105, 3893–3904, doi:10.1029/1999JD901033, 2000.
- Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, *Atmos. Environ.*, 52, 109–120, 2012.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Van As, D., Van Den Broeke, M., and Van De Wal, R.: Daily cycle of the surface layer and energy balance on the high Antarctic Plateau, *Antarct. Sci.*, 17, 121–133, 2005.
- Wagenbach, D., Ducroz, F., Mulvaney, R., Keck, L., Minikin, A., Legrand, M. J., Hall, S., and Wolff, E. W.: Sea-salt aerosol in coastal Antarctic regions, *J. Geophys. Res.*, 103, 10961–10974, 1998.
- Wang, L.-Y., Ding, F., Zhang, Y.-H., Zhao, L.-J., and Hu, Y.-A.: Anomalous hygroscopic growth of fine particles of MgSO_4 aerosols investigated by FTIR/ATR spectroscopy, *Spectrochim. Ac. Pt. A*, 71, 682–687, doi:10.1016/j.saa.2008.01.027, 2008.
- Weller, R. and Wagenbach, D.: Year-round chemical aerosol records in continental Antarctica obtained by automatic samplings, *Tellus B*, 59, 755–765, 2007.
- Weller, R., Minikin, A., Wagenbach, D., and Dreiling, V.: Characterization of the inter-annual, seasonal, and diurnal variations of condensation particle concentrations at Neumayer, Antarctica, *Atmos. Chem. Phys.*, 11, 13243–13257, doi:10.5194/acp-11-13243-2011, 2011.
- Wilson, T. R.: Salinity and the major elements of sea-water, in *Chemical Oceanography*, edited by: Riley, J. P. and Skirrow, G., Academic Press, San Diego, Calif., 365–413, 1975.
- Wise, M. E., Freney, E. J., Tyree, C. A., Allen, J. O., Martin, S. T., Russell, L. M., and Buseck, P. R.: Hygroscopic behavior and liquid-layer composition of aerosol particles generated from natural and artificial seawater, *J. Geophys. Res.*, 114, D03201, doi:10.1029/2008JD010449, 2009.
- Woods, E., Chung, D., Lanney, H. M., and Ashwell, B. A.: Surface morphology and phase transitions in mixed $\text{NaCl}/\text{MgSO}_4$ aerosol particles, *J. Phys. Chem. A*, 114, 2837–2844, 2010.
- Wouters, L., Artaxo, P., and van Grieken, R.: Laser microprobe mass analysis of individual antarctic aerosol particles, *Int. J. Environ. Anal. Chem.*, 38, 427–438, 1990.
- Yamato, M., Iwasaka, Y., Ono, A., and Yoshida, M.: On the sulfate particles in the submicron size range collected at Mizuho Station and in east Queen Maud Land, Antarctica, *Polar Meteorol. Glaciol.*, 1, 82–90, 1987a.
- Yamato, M., Iwasaka, Y., Okada, K., Ono, A., Nishio, F., and Fukabori, M.: Evidence for the presence of submicron sulfuric acid particles in summer Antarctic atmosphere: preliminary results, *Proc. NIPR Symp. Polar Meteorol. Glaciol.*, 1, 74–81, 1987b.

Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 1. Traverse routes of Japanese team during JASE campaign. Black lines represent traverse routes between S16 and Dome F in incoming and outgoing traverse. Blue line represents traverse route from Dome F to meeting point in incoming traverse. Red line indicates traverse route from meeting point to Dome F in outgoing traverse. Traverse route of Swedish team was depicted in Fujita et al. (2011).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Horizontal
distributions of
aerosol constituents

K. Hara et al.




Fig. 2. Variations of latitudes, elevation, air temperature, and relative humidity during the JASE traverse. Cyan stars represent dates and locations of aerosol measurements and direct sampling.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 3a. 5 day backward trajectories from every camp site in **(a)** incoming traverse from S16 to Dome F, and **(b)** traverse from Dome F (DF) to meeting point (MP), **(c)** traverse from meeting point to S16 in outgoing traverse. Black lines represent traverse route. Altitudes indicate height above ground.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Fig. 3b. Continued.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3c. Continued.

Horizontal distributions of aerosol constituents

K. Hara et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 4. Horizontal features of air temperature, elevation, wind speed, aerosol number concentrations and Junge slope during incoming traverse from S16 to the meeting point. Red boxes indicate conditions with drifting snow.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 5. Horizontal features of air temperature, elevation, wind speed, aerosol number concentrations and Junge slope during outgoing traverse from the meeting point to S16. Red boxes indicate conditions with drifting snow.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. SEM images of aerosol particles in (a) coarse mode and (b) fine mode.

Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 7. EDX spectra of aerosol particles collected during the JASE traverse. Asterisks show background peaks derived from the sample substrate.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 8. Horizontal features of relative abundance of each aerosol constituent in **(a–b)** coarse and **(c–d)** fine modes during the incoming traverse.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 9. Horizontal features of relative abundance of each aerosol constituent in **(a–b)** coarse and **(c–d)** fine modes during the outgoing traverse.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 10. Ternary plots (Na–S–Cl) of sea-salt particles in coarse and fine modes during the JASE traverse. Red, and blue stars respectively denote atomic ratios of (1) bulk seawater, and (2) wholly Cl depleted sea-salt particles with sulfates. Black lines represent stoichiometric lines among constituents.

Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 11. Horizontal features of atomic ratio of Cl/Na and S/Na in coarse and fine modes during the incoming traverse. In box plots, the top bar, top box line, black middle box line, bottom box line, and bottom bar respectively denote values of 90%, 75%, 50% (median), 25%, and 10%. The red line shows mean values.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 12. Horizontal features of atomic ratio of Cl/Na and S/Na in coarse and fine modes during the outgoing traverse. In box plots, the top bar, top box line, black middle box line, bottom box line, and bottom bar respectively denote values of 90 %, 75 %, 50 % (median), 25 %, and 10 %. The red line shows mean values.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 13. Ternary plots (Na–Mg–S) of sea-salt particles in coarse and fine modes during the JASE traverse. Red, blue, cyan, and green stars respectively denote atomic ratios of (1) bulk seawater, (2) wholly Cl depleted sea-salt particles with sulfates, (3) MgCl_2 , and (4) MgSO_4 . Blue, pink and red lines represent stoichiometric lines among constituents.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Horizontal distributions of aerosol constituents

K. Hara et al.




Fig. 14. Horizontal features of atomic ratio of Mg/Na in coarse mode during the incoming traverse. In box plots, the top bar, top box line, black middle box line, bottom box line, and bottom bar respectively denote values of 90 %, 75 %, 50 % (median), 25 %, and 10%. The red line shows mean values.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)