



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Airborne characterization of smoke marker ratios from prescribed burning

A. P. Sullivan¹, A. A. May¹, T. Lee¹, G. R. McMeeking¹, S. M. Kreidenweis¹, S. K. Akagi², R. J. Yokelson², S. P. Urbanski³, and J. L. Collett Jr.¹

¹Colorado State University, Department of Atmospheric Science, Fort Collins, Colorado 80523, USA

²University of Montana, Department of Chemistry, Missoula, MT 59812, USA

³USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory, Missoula, MT 59808, USA

Received: 7 April 2014 – Accepted: 14 April 2014 – Published: 9 May 2014

Correspondence to: A. P. Sullivan (sullivan@atmos.colostate.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

A Particle-into-Liquid Sampler – Total Organic Carbon and fraction collector system was flown aboard a Twin Otter aircraft sampling prescribed burning emissions in South Carolina in November 2011 to obtain smoke marker measurements. The fraction collector provided 2 min time-integrated off-line samples for carbohydrate (i.e., smoke markers levoglucosan, mannosan, galactosan) analysis by high-performance anion-exchange chromatography with pulsed amperometric detection. Each fire location appeared to have a unique Δ levoglucosan/ Δ water-soluble organic carbon (WSOC) ratio (RF01/RF02/RF03/RF05 = $0.163 \pm 0.007 \mu\text{g C} \mu\text{g C}^{-1}$, RF08 = $0.115 \pm 0.011 \mu\text{g C} \mu\text{g C}^{-1}$, RF09A = $0.072 \pm 0.028 \mu\text{g C} \mu\text{g C}^{-1}$, RF09B = $0.042 \pm 0.008 \mu\text{g C} \mu\text{g C}^{-1}$). These ratios were comparable to those obtained from controlled laboratory burns and suggested that the emissions sampled during RF01/RF02/RF03/RF05 were dominated by the burning of grasses, RF08 by leaves, RF09A by needles, and RF09B by marsh grasses. These findings were further supported by the Δ galactosan/ Δ levoglucosan ratios (RF01/RF02/RF03/RF05 = $0.067 \pm 0.004 \mu\text{g} \mu\text{g}^{-1}$, RF08 = $0.085 \pm 0.009 \mu\text{g} \mu\text{g}^{-1}$, RF09A = $0.101 \pm 0.029 \mu\text{g} \mu\text{g}^{-1}$) obtained as well as by the ground-based fuel and filter sample analyses during RF01/RF02/RF03/RF05. Differences between Δ potassium/ Δ levoglucosan ratios obtained for these prescribed fires vs. laboratory-scale measurements suggest that some laboratory burns may not accurately represent potassium emissions from prescribed burns. The Δ levoglucosan/ Δ WSOC ratio had no clear dependence on smoke age or fire dynamics suggesting that this ratio is more dependent on the type of fuel being burned. Levoglucosan was stable over a timescale of at least 1.5 h and could be useful to help estimate the air quality impacts of biomass burning.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The smoke marker approach is the most common method used to estimate the contribution of primary biomass burning to the total organic carbon aerosol concentration (e.g., Schauer et al., 1996; Schauer and Cass, 2000; Fraser et al., 2003; Rinehart et al., 2006). In this approach, a compound produced as part of the smoke (i.e., smoke marker) is monitored as a plume is transported downwind. If the smoke marker is conserved, or decays at a known rate, during transport and the ratio of the smoke marker to the total organic carbon is known at the source, then a downwind measurement of the smoke marker's concentration can be used to apportion the contribution of primary biomass burning emissions to the total organic carbon.

Generally, source sample smoke marker ratios are obtained from ground-based studies utilizing integrated filter sampling with sample collection over the duration of the entire fire (e.g., Hays et al., 2002; Fine et al., 2004; Lee et al., 2005; Mazzoleni et al., 2007). One of the main reasons for this approach is that traditional methods used to measure smoke markers, such as gas chromatography-mass spectrometry (GC-MS), require a large amount of mass for analysis. Therefore, ground-based sampling would provide the best means to collect a large amount of sample as this is generally not feasible from an aircraft platform that can quickly fly through a plume.

However, being able to measure smoke marker concentrations from an aircraft platform could be useful. Sampling a smoke plume right after emission as well as following it downwind during transport could help to better characterize source smoke marker ratios and any evolution due to plume dilution and aging. In addition, collecting multiple samples from the same fire, which is often not feasible with filter sampling due to the time required to collect sufficient mass, would be possible.

As a first attempt to make airborne smoke marker measurements, a PILS-TOC (Particle-into-Liquid Sampler – Total Organic Carbon) and fraction collector system was flown aboard a Twin Otter aircraft in November 2011 as it sampled emissions from prescribed burning activities taking place in South Carolina. This study was the last

ACPD

14, 11715–11747, 2014

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



field deployment in a series of measurements of prescribed burning emissions from the southeastern US (Burling et al., 2010, 2011; Akagi et al., 2013; Yokelson et al., 2013). The idea behind the chosen instrumentation was to provide a 3 s time-integrated measurement of water-soluble organic carbon (WSOC), whose two main sources are biomass burning and secondary organic aerosol (SOA), from the PILS-TOC (Sullivan et al., 2004, 2006), then take advantage of the high sensitivity and low limit of detection of high-performance anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD) to analyze the fraction collector samples off-line to provide a near real-time measurement of carbohydrates (i.e., smoke markers). PAD is an electrochemical technique where hydroxyl groups are electroanalytically oxidized on the surface of a gold electrode. This approach offers numerous advantages including extraction of an aerosol sample directly in water (i.e., no derivatization or organic solvents are needed) and the ability to directly analyze an aqueous sample for smoke markers. This technique has been applied to biomass burning source samples as well as studies examining ambient aerosol contributions by biomass burning (Gao et al., 2003; Gorin et al., 2006; Engling et al., 2006; Puxbaum et al., 2007; Sullivan et al., 2008, 2011a, b).

In this work, measurements of WSOC and various smoke markers, including levoglucosan, mannosan, galactosan, and water-soluble potassium, for five different prescribed burns (six research flights) are presented. Smoke marker ratios for the various prescribed burns are discussed and compared with results from biomass burning source samples collected from controlled laboratory burns and on the ground. The influence of plume aging and fire dynamics on smoke marker ratios is also investigated.

2 Methods

2.1 Particle collection

A Particle-into-Liquid Sampler (PILS) was used to collect ambient particles into purified water, providing the liquid sample for analysis (Orsini et al., 2003). Upstream of

and required no precleaning before use. The fraction collector program, which was manually started at take-off, was set to allow continuous collection of 2 min time-integrated samples. Based on the liquid flowrates used for the PILS, each test tube collected approximately 1.6 mL of sample. At the completion of each flight, the test tubes were capped, packed in coolers with ice packs, and shipped back to Colorado State University to be stored in a 2 °C cold room until analysis.

2.2 Off-line analysis

Each fraction collector test tube was brought back to room temperature before analysis. Two 600 µL aliquots were transferred to separate polypropylene vials for carbohydrate and cation measurements.

The carbohydrate analysis was performed on a Dionex DX-500 series ion chromatograph with an ED-50 electrochemical detector operating in integrating amperometric mode using waveform A and a GP-50 gradient pump. The detector contains an ED-50/ED-50A electrochemical cell. This cell includes a pH-Ag/AgCl (silver/silver chloride) reference electrode and “standard” gold working electrode. The separation was performed using a Dionex CarboPac PA-1 column (4 mm × 250 mm) employing a sodium hydroxide gradient. The complete run time was 59 min and an injection volume of 50 µL was used. More details of the method can be found in Sullivan et al. (2011a, b). Of the carbohydrates that can be detected by this method, levoglucosan, mannosan, and galactosan were found in all samples. Glucose and arabinose were only occasionally detected and will not be discussed further. The limit of detection (LOD) for the various carbohydrates was calculated to be less than approximately 0.10 ng m⁻³.

Water-soluble potassium was measured using a second Dionex DX-500 ion chromatograph. This system included an isocratic pump, self-regenerating cation SRS-ULTRA suppressor, and conductivity detector. A Dionex IonPac CS12A analytical column (3 mm × 150 mm) using 20 mM methanesulfonic acid at a flowrate of 0.5 mL min⁻¹ was used for the separation. The injection volume and analysis time were 50 µL and 17 min, respectively. Unlike for the carbohydrates, a blank correction was necessary for

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the water-soluble potassium. Concentrations were corrected by using the average of all particle-free background samples (i.e., with the actuated valve before the PILS in the closed position) collected during a specific flight. The LOD for water-soluble potassium was $0.02 \mu\text{g m}^{-3}$.

2.3 Airborne mission and other measurements

These research flights were part of a combined ground-based and airborne-based study to examine the emissions from prescribed burning in the southeastern US. The Twin Otter was operated out of Columbia, SC from 29 October 2011 through 11 November 2011. Measurements were made of several chemical and physical aerosol particle properties, and of reactive and stable trace gases. Table 1 provides a list of the flights when the PILS-TOC and fraction collector sampler operated. The first flights (RF01/RF02/RF03/RF05) focused on prescribed burning at Fort Jackson, SC. Once these burns were completed, airborne sampling only of other prescribed burns taking place in SC began (RF08 and RF09). A typical flight involved first characterizing the emissions right at the source with numerous passes through the smoke (Fig. 1). Then the smoke was sampled downwind mostly by crossing back and forth through the plume further and further downwind until it could not be distinguished from the background air. On flights RF03, RF08, and RF09, sampling of the smoke downwind was achieved.

All aircraft aerosol instruments sampled from a LTI (Low Turbulence Inlet) (Wilson et al., 2004). Following the LTI was a nonrotating MOUDI impactor with a 50 % transmission efficiency at $1 \mu\text{m}$ and 1 atmosphere ambient pressure (Marple et al., 1991). The combined flow through the inlet and MOUDI was approximately 20 LPM and was then split to the individual instruments.

In the analysis presented in this paper we focus on characterizing source smoke marker ratios from prescribed burning. Other measurements presented here include 3-D location and windspeed for estimated time since emission values collected with a wing-mounted Aircraft Integrated Meteorological Measuring System probe

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fraction collector samples that directly overlapped with a peak in the CO concentrations are considered. Given the longer response time constant of the PILS (due to being a liquid system, Sorooshian et al., 2006) vs. the other instruments, we initially focus on internal comparisons of species concentrations measured by the PILS.

The absolute WSOC, levoglucosan, mannosan, galactosan, and potassium concentrations along with altitude data for this subset of data from all flights is given in Table 2. Throughout for all flights excess ratios (denoted by Δ) will be presented and were determined as the difference in the concentration when in and outside of a smoke plume. Concentrations used for all flights for WSOC, levoglucosan, mannosan, galactosan, and potassium outside of a smoke plume were $2.00 \mu\text{g C m}^{-3}$, $0.03 \mu\text{g m}^{-3}$, $0.03 \mu\text{g m}^{-3}$, $0.03 \mu\text{g m}^{-3}$, and $0.30 \mu\text{g m}^{-3}$, respectively.

The correlation of Δ levoglucosan with Δ WSOC is shown in Fig. 4a. All the burns occurring at Fort Jackson (RF01/RF02/RF03/RF05) appeared to have a similar ratio, based on the slope of the linear correlation, of $0.163 \pm 0.007 \mu\text{g C } \mu\text{g C}^{-1}$. In addition, there appeared to be no concentration dependence or significant altitude dependence to this ratio (Fig. 4b). The smoke sampled during RF08 had lower Δ levoglucosan/ Δ WSOC ratios (approximately $0.115 \pm 0.011 \mu\text{g C } \mu\text{g C}^{-1}$) than the ratios for the Fort Jackson burns. Interestingly, the fire sampled during RF09 appeared to have two distinct groups of Δ levoglucosan/ Δ WSOC ratios (denoted RF09A and RF09B). Group B had only a few samples, so linear regression statistics were not reliable. Therefore, throughout for RF09B the average ratio \pm standard deviation was calculated. The Δ levoglucosan/ Δ WSOC ratio was $0.042 \pm 0.008 \mu\text{g C } \mu\text{g C}^{-1}$ for RF09B, which was lower than the ratio of $0.072 \pm 0.028 \mu\text{g C } \mu\text{g C}^{-1}$ for RF09A, suggesting that the vegetation may have been different at the two ends of the fire being sampled during RF09. In South Carolina it is very common for marshy bays to be mixed in with a forested area (B. Manks, personal communication, 2011).

The importance of fuel type combusted can be further illustrated by comparing the airborne smoke marker ratios to those from typical biomass burning source samples collected from controlled laboratory burns ($0.149 \pm 0.012 \mu\text{g C } \mu\text{g C}^{-1}$ for grasses,

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Water-soluble potassium has long been used as an inorganic marker for biomass burning. As can be seen in Fig. 5c, the prescribed burn observations contain quite a bit of scatter in the Δ potassium to Δ levoglucosan ratio, even for a particular burn location. For example, attempting a linear fit to the data from the burns at Fort Jackson (RF01/RF02/RF03/RF05) yields a very low R^2 value of 0.13. Although, there is somewhat of a better correlation for RF08 ($R^2 = 0.41$). Poor correlation between potassium and levoglucosan concentrations in biomass burning smoke is not surprising. The presence of a small amount of inorganic substances, such as potassium, in a fuel can cause changes in the product yields of levoglucosan during cellulose pyrolysis, with potassium suppressing the formation of levoglucosan (Radlein et al., 1991; Richards et al., 1991; Patwardhan et al., 2010; Eom et al., 2012). In addition, potassium is predominately emitted from the flaming phase of a fire, whereas levoglucosan is emitted across both smoldering and flaming fire phases (Ward et al., 1991; Echalar et al., 1995; Lee et al., 2010). Changes in the mix of flaming and smoldering combustion in a laboratory or prescribed burn, therefore, can readily yield large differences in the emitted abundances of potassium.

3.3 Role of aging and fire dynamics

The time since emission (i.e., the smoke age) was estimated for all possible downwind aircraft smoke marker samples from RF03, RF08, RF09A, and RF09B as the distance from the source divided by the average wind speed for the sampling altitude (Akagi et al., 2013). Pseudo-Lagrangian sampling was accomplished for the majority of these downwind samples, meaning the aircraft was sampling the source of the fire at their estimated time of emission (see Akagi et al., 2013 for more details). Changes in plume composition occur with plume aging, due both to plume dilution (which can influence gas-to-particle partitioning) and photochemical reactions, but very little data quantitatively examines the impact (if any) of these processes on smoke marker ratios. Figure 6a shows the Δ levoglucosan/ Δ WSOC ratio as a function of time since emission. Over the range of smoke plume ages (up to approximately 1.5 h), the observations give

no clear indication that the ratio changes in a consistent manner as the plume ages. Low ratios in RF09B, for example, remain low, while higher initial ratios in RF03 remain high. These observations suggest that the Δ levoglucosan/ Δ WSOC source smoke marker ratio is stable for at least 1–1.5 h as the plume dilutes and ages.

We can also make use of a subset of observations from the AMS to look at plume characteristics for the different flights. Of particular interest from a source apportionment perspective is the ratio of the key AMS biomass burning marker at m/z 60 to AMS total organic aerosol (OA). Figure 7 plots $\Delta m/z$ 60 vs. Δ OA concentrations for flights RF03, RF05, RF08, and RF09. (Concentrations used for all flights for OA and m/z 60 outside of a smoke plume were $4.00 \mu\text{g m}^{-3}$ and $0.003 \mu\text{g m}^{-3}$, respectively.) Despite the differences in fuel type and Δ levoglucosan/ Δ WSOC ratios across burns described above, this AMS ratio shows a very tight relationship ($R^2 = 0.99$). A Δ OA concentration of $3.35 \mu\text{g m}^{-3}$ associated with no m/z 60 suggests a background OA concentration from other sources onto which the smoke aerosol is added. There is some change apparent in this ratio as a function of plume age (Fig. 6b). A bigger spread of ratios across fresh, individual plumes appears to rapidly converge to a more consistent ratio in less than approximately 0.5 h of plume aging. The consistency of this ratio across burns and fuel types reinforces the quality of m/z 60 as a quantitative biomass burning source marker for use with AMS data sets. Although m/z 60 is often identified with levoglucosan, it can be comprised of a variety of structurally similar molecules (Aiken et al., 2009; Mohr et al., 2009; Lee et al., 2010). While differences in levoglucosan emissions across fuel type are apparent from the PILS measurements discussed above, emissions of the larger suite of structurally similar molecules that fragment to yield m/z 60 appear to be more constant across fuel types.

The $\Delta m/z$ 60/ Δ OA concentration ratio is plotted as a function of the Δ OA concentration in Fig. 8a. Below Δ OA concentrations of approximately $50 \mu\text{g m}^{-3}$, the ratio is strongly related with Δ OA concentration. At higher concentrations, the ratio levels off. If a simple two factor Positive Matrix Factorization (PMF) solution is applied to this AMS OA data it can provide background OA and biomass burning OA (BBOA)

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations (May et al., 2014). If we then plot the ratio of $\Delta m/z 60/\text{BBOA}$ as a function of total ΔOA (Fig. 8b) we observe much less concentration dependence. This suggests that changes in the $\Delta m/z 60/\Delta \text{OA}$ ratio as a function of ΔOA concentration are largely driven by changes in the concentration of background (non-biomass burning) OA. Some of the background OA concentration change is likely due to gas-to-particle partitioning changes that are themselves a function of OA concentration. In contrast to the AMS observations, there is no clear relationship between the PILS Δ levoglucosan/ Δ WSOC ratio and the Δ WSOC concentration (Fig. 8c).

The modified combustion efficiency (MCE) can be used to investigate the role of fire dynamics. MCE is calculated as the excess carbon dioxide divided by the sum of the excess carbon monoxide and carbon dioxide ($\Delta \text{CO}_2/(\Delta \text{CO} + \Delta \text{CO}_2)$) on a molar basis (Ward and Radke, 1993). Therefore, a higher MCE value indicates emissions were dominated by flaming combustion whereas a lower value indicates more extensive contributions from smoldering. As seen in Fig. 9, there appears to be no clear dependence of the Δ levoglucosan/ Δ WSOC on MCE. Overall, this finding is similar to the pattern observed for controlled laboratory burns, which covered the same dynamic range of MCE values, although the prescribed burns include a greater variety of MCE values for a particular fuel type. Therefore, it appears that source smoke marker ratios for prescribed burns are more dependent on the fuel type being burned than on differences in fire behavior.

4 Summary

Concentrations of smoke markers (e.g., levoglucosan and galactosan) are generally measured from ground-based, integrated filter samples. Here we took advantage of a new approach that permitted the first measurements of these compounds from an airborne platform. A PILS-TOC and fraction collector system was flown aboard a Twin Otter aircraft sampling prescribed burning emissions in South Carolina in November 2011. The PILS-TOC provided a 3 s time-integrated measurement of WSOC. The

fraction collector provided 2 min time-integrated samples to be analyzed off-line for carbohydrates (i.e., smoke markers) by high-performance anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD). The HPAEC-PAD had ample sensitivity to detect levoglucosan, mannosan, and galactosan in the short-duration fraction collector samples. Comparisons with other measurements aboard the Twin Otter show that the 2 min time resolution was adequate to characterize smoke markers in the smoke plumes.

The ratio of Δ levoglucosan to Δ WSOC varied across fires (RF01/RF02/RF03/RF05 = $0.163 \pm 0.007 \mu\text{g C } \mu\text{g C}^{-1}$, RF08 = $0.115 \pm 0.011 \mu\text{g C } \mu\text{g C}^{-1}$, RF09A = $0.072 \pm 0.028 \mu\text{g C } \mu\text{g C}^{-1}$, and RF09B = $0.042 \pm 0.008 \mu\text{g C } \mu\text{g C}^{-1}$). Available information about fire fuel type in the burns and a comparison of levoglucosan/WSOC ratios with laboratory burns of specific fuel types indicate the Δ levoglucosan/ Δ WSOC ratio differences are related to the mix of fuel types combusted in each fire. Comparison of prescribed vs. laboratory burn Δ galactosan/ Δ levoglucosan ratios yield a consistent finding about the type of fuel involved in each fire. This was not, however, the case for Δ mannosan/ Δ levoglucosan ratios, which could be due to a fuel element burned in the field, but not the laboratory. A poor correlation of Δ potassium to Δ levoglucosan concentrations and large differences between values of this ratio observed in controlled laboratory burns and prescribed burns suggest that the laboratory burns may not yield representative potassium emissions. This result is not surprising given the known, strong dependence of potassium emissions on fire flaming/smoldering ratios. No clear dependence of the Δ levoglucosan/ Δ WSOC on fire conditions, as represented by MCE, was observed.

AMS measurements of smoke plumes aboard the aircraft also yielded interesting findings. A strong correlation between the AMS biomass burning marker at $\Delta m/z$ 60 vs. Δ OA was observed. This ratio did not vary with fuel type, but was positively correlated with Δ OA at concentrations below $50 \mu\text{g m}^{-3}$. PMF analysis suggested that the concentration dependence of this ratio was largely driven by changes in the aerosol content of non-biomass burning OA.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Overall, this study demonstrated: (1) a new capability for airborne, in-plume measurements of levoglucosan and other smoke marker concentrations, (2) a clear relationship between the Δ levoglucosan/ Δ WSOC ratio and fuel type, and (3) the utility of AMS measurements of OA and m/z 60 as a quantitative method for apportioning biomass burning aerosol contributions to ambient aerosol, for several biomass fuel types sampled in this study.

Acknowledgements. We acknowledge funding from the Joint Fire Science Program under Project JFSP 11-1-5-12 and the Strategic Environmental Research and Development Program (SERDP) project RC-1649. We thank R. J. Weber for generously providing some of the parts used in the PILS rack. We also thank E. J. T. Levin for his help in assembling the rack. We additionally acknowledge the Twin Otter Science Team, especially our pilot B. Mank and mechanic S. Woods. Lastly, special thanks to J. Maitland and his team at Fort Jackson who conducted those burns and the Columbia Dispatch Office of the South Carolina Forestry Commission for providing information on the additional burns.

References

- Aiken, A. C., Salcedo, D., Cubison, M. J., Huffman, J. A., DeCarlo, P. F., Ulbrich, I. M., Docherty, K. S., Sueper, D., Kimmel, J. R., Worsnop, D. R., Trimborn, A., Northway, M., Stone, E. A., Schauer, J. J., Volkamer, R. M., Fortner, E., de Foy, B., Wang, J., Laskin, A., Shutthanandan, V., Zheng, J., Zhang, R., Gaffney, J., Marley, N. A., Paredes-Miranda, G., Arnott, W. P., Molina, L. T., Sosa, G., and Jimenez, J. L.: Mexico City aerosol analysis during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite (T0) – Part 1: Fine particle composition and organic source apportionment, *Atmos. Chem. Phys.*, 9, 6633–6653, doi:10.5194/acp-9-6633-2009, 2009.
- Akagi, S. K., Yokelson, R. J., Burling, I. R., Meinardi, S., Simpson, I., Blake, D. R., McMeeking, G. R., Sullivan, A., Lee, T., Kreidenweis, S., Urbanski, S., Reardon, J., Griffith, D. W. T., Johnson, T. J., and Weise, D. R.: Measurements of reactive trace gases and variable O₃ formation rates in some South Carolina biomass burning plumes, *Atmos. Chem. Phys.*, 13, 1141–1165, doi:10.5194/acp-13-1141-2013, 2013.

**Smoke marker ratios
from prescribed
burning**

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Burling, I. R., Yokelson, R. J., Griffith, D. W. T., Johnson, T. J., Veres, P., Roberts, J. M., Warneke, C., Urbanski, S. P., Reardon, J., Weise, D. R., Hao, W. M., and de Gouw, J.: Laboratory measurements of trace gas emissions from biomass burning of fuel types from the southeastern and southwestern United States, *Atmos. Chem. Phys.*, 10, 11115–11130, doi:10.5194/acp-10-11115-2010, 2010.
- 5 Burling, I. R., Yokelson, R. J., Akagi, S. K., Urbanski, S. P., Wold, C. E., Griffith, D. W. T., Johnson, T. J., Reardon, J., and Weise, D. R.: Airborne and ground-based measurements of the trace gases and particles emitted by prescribed fires in the United States, *Atmos. Chem. Phys.*, 11, 12197–12216, doi:10.5194/acp-11-12197-2011, 2011.
- 10 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281–8289, 2006.
- Echalar, F., Gaudichet, A., Cachier, H., and Artaxo, P.: Aerosol emissions by tropical forest and savanna biomass burning: characteristic trace elements and fluxes, *Geophys. Res. Lett.*, 22, 3039–3042, 1995.
- 15 Eom, I.-Y., Kim, J.-Y., Kim, T.-S., Lee, S.-M., Choi, D., Choi, I.-G., and Choi, J.-W.: Effect of essential inorganic metals on primary thermal degradation of lignocellulosic biomass, *Biore-source Technol.*, 104, 687–694, 2012.
- 20 Hays, M. D., Geron, C. D., Linna, K. J., Smith, N. D., and Schauer, J. J.: Speciation of gas-phase and fine particle emissions from burning of foliar fuels, *Environ. Sci. Technol.*, 36, 2281–2295, 2002.
- Eatough, D. J., Wadsworth, A., Eatough, D. A., Crawford, J. W., Hansen, L. D., and Lewis, E. A.: A multiple system, multi-channel diffusion denuder sampler for the determination of fine-particulate organic material in the atmosphere, *Atmos. Environ.*, 27, 1213–1219, 1993.
- 25 Engling, G., Carrico, C. M., Kreidenweis, S. M., Collett Jr., J. L., Day, D. E., Malm, W. C., Lincoln, E., Hao, W. M., Iinuma, Y., and Herrmann, H.: Determination of levoglucosan in biomass combustion aerosol by high-performance anion-exchange chromatography with pulsed amperometric detection, *Atmos. Environ.*, 40, S299–S311, 2006.
- 30 Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the fireplace combustion of wood types grown in the midwestern and western United States, *Environ. Eng. Sci.*, 21, 387–409, 2004.

**Smoke marker ratios
from prescribed
burning**

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Fraser, M. P., Yue, Z. W., and Buzcu, B.: Source appointment of fine particulate matter in Houston, TX, using organic molecular markers, *Atmos. Environ.*, 37, 2117–2123, 2003.
- Gao, S., Hegg, D. A., Hobbs, P. V., Kirchstetter, T. W., Magi, B. I., and Sadilek, M.: Water-soluble organic components in aerosols associated with savanna fires in southern Africa: identification, evolution, and distribution, *J. Geophys. Res.*, 108, 8491, doi:10.1029/2002JD002324, 2003.
- Gorin, C. A., Collett Jr., J. L., and Herckes, P.: Wood smoke contribution to winter aerosol in Fresno, CA, *JAPCA J. Air Waste Ma.*, 56, 1584–1590, 2006.
- Lee, S., Baumann, K., Schauer, J. J., Sheesley, R. J., Naeher, L. P., Meinardi, S., Blake, D. R., Edgerton, E. S., Russell, A. G., and Clements, M.: Gaseous and particulate emissions from prescribed burning in Georgia, *Environ. Sci. Technol.*, 39, 9049–9056, 2005.
- Lee, T., Sullivan, A. P., Mack, L., Jimenez, J. L., Kreidenweis, S. M., Onasch, T. B., Worsnop, D. R., Malm, W., Wold, C. E., Hao, W. M., and Collett Jr., J. L.: Chemical smoke marker emissions during flaming and smoldering phases of laboratory open burning of wildland fuels, *Aerosol Res. Lett.*, 44, i–v, 2010.
- Marple, V. A., Rubow, K. L., and Behm, S. M.: A microorifice uniform deposit impactor (MOUDI): description, calibration, and use, *Aerosol Sci. Tech.*, 14, 434–446, 1991.
- May, A. A., Lee, T., McMeeking, G. R., Akagi, S., Sullivan, A. P., Urbanski, S., Yokelson, R. J., and Kreidenweis, S. M.: Investigation of Chemical and Physical Perturbations to Organic Aerosol Present in Biomass Burning Plumes over Prescribed Fires in South Carolina, *Atmos. Chem. Phys.*, in preparation, 2014.
- Mazzoleni, L. R., Zielinska, B., and Moosmüller, H.: Emissions of levoglucosan, methoxy phenols, and organic acids from prescribed burns, laboratory combustion of wildland fuels, and residential wood combustion, *Environ. Sci. Technol.*, 41, 2115–2122, 2007.
- Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of primary organic aerosol emissions from meat cooking, trash burning, and motor vehicles with high-resolution aerosol mass spectrometry and comparison with ambient and chamber observations, *Environ. Sci. Technol.*, 43, 2443–2449, 2009.
- Orsini, D. A., Ma, Y., Sullivan, A., Sierau, B., Baumann, K., and Weber, R. J.: Refinements to the particle-into-liquid sampler (PILS) for ground and airborne measurements of water-soluble aerosol composition, *Atmos. Environ.*, 37, 1243–1259, 2003.



**Smoke marker ratios
from prescribed
burning**

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Patwardhan, P. R., Satrio, J. A., Brown, R. C., and Shanks, B. H.: Influence of inorganic salts on the primary pyrolysis products of cellulose, *Bioresource Technol.*, 101, 4646–4655, 2010.
- Puxbaum, H., Caseiro, A., Sánchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencsér, A., Legrand, M., Preunkert, S., and Pio, C.: Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background, *J. Geophys. Res.*, 112, D23S05, doi:10.1029/2006JD008114, 2007.
- 5 Radlein, D., Piskorz, J., and Scott, D. S.: Fast pyrolysis of natural polysaccharides as a potential industrial process, *J. Anal. Appl. Pyrol.*, 19, 41–63, 1991.
- Richards, G. N. and Zheng, G.: Influence of metal ions and of salts on products from pyrolysis of wood: applications to thermochemical processing of newsprint and biomass, *J. Anal. Appl. Pyrol.*, 21, 133–146, 1991.
- 10 Rinehart, L. R., Fujita, E. M., Chow, J. C., Magliano, K., and Zielinska, B.: Spatial distribution of PM_{2.5} associated organic compounds in central California, *Atmos. Environ.*, 40, 290–303, 2006.
- 15 Schauer, J. J. and Cass, G. R.: Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers, *Environ. Sci. Technol.*, 34, 1821–1832, 2000.
- Schauer, J. J., Rogge, W. F., Hildemann, L. M., Mazurek, M. A., and Cass, G. R.: Source apportionment of airborne particulate matter using organic compounds as tracers, *Atmos. Environ.*, 30, 3837–3855, 1996.
- 20 Sorooshian, A., Brechtel, F. J., Ma, Y., Weber, R. J., Corless, A., Flagan, R. C., and Seinfeld, J. H.: Modeling and characterization of a Particle-into-Liquid Sampler (PILS), *Aerosol Sci. Tech.*, 40, 396–409, 2006.
- Sullivan, A. P., Weber, R. J., Clements, A. L., Turner, J. R., Bae, M. S., and Schauer, J. J.: A method for on-line measurement of water-soluble organic carbon in ambient aerosol particles: recent results from an urban site, *Geophys. Res. Lett.*, 31, L13105, doi:10.1029/2004GL019681, 2004.
- 25 Sullivan, A. P., Peltier, R. E., Brock, C. A., de Gouw, J. A., Holloway, J. S., Warneke, C., Wollny, A. G., and Weber, R. J.: Airborne measurements of carbonaceous aerosol soluble in water over northeastern United States: method development and an investigation into water-soluble organic carbon sources, *J. Geophys. Res.*, 111, D23S46, doi:10.1029/2006JD007072, 2006.
- 30

**Smoke marker ratios
from prescribed
burning**

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Sullivan, A. P., Holden, A. S., Patterson, L. A., McMeeking, G. R., Kreidenweis, S. M., Malm, W. C., Hao, W. M., Wold, C. E., and Collett Jr., J. L.: A method for smoke marker measurements and its potential application for determining the contribution of biomass burning from wildfires and prescribed fires to ambient PM_{2.5} organic carbon, *J. Geophys. Res.*, 113, D22302, doi:10.1029/2008JD010216, 2008.
- Sullivan, A. P., Frank, N., Onstad, G., Simpson, C. D., and Collett Jr., J. L.: Application of high-performance anion-exchange chromatography – pulsed amperometric detection for measuring carbohydrates in routine daily filter samples collected by a national network: 1. Determination of the impact of biomass burning in the upper midwest, *J. Geophys. Res.*, 116, D08302, doi:10.1029/2010JD014166, 2011a.
- Sullivan, A. P., Frank, N., Kenski, D. M., and Collett Jr., J. L.: Application of high-performance anion-exchange chromatography – pulsed amperometric detection for measuring carbohydrates in routine daily filter samples collected by a national network: 2. Examination of sugar alcohols/polyols, sugars, and anhydrosugars in the upper midwest, *J. Geophys. Res.*, 116, D08303, doi:10.1029/2010JD014169, 2011b.
- Sullivan, A. P., Kreidenweis, S. M., Schichtel, B. A., and Collett Jr., J. L.: Source Smoke Marker Ratios from Controlled Laboratory Burns vs. Prescribed Burns and Wildfires, *Atmos. Chem. Phys.*, in preparation, 2014.
- Ward, D. E. and Radke, L. F.: Emission measurements from vegetation fires: a comparative evaluation of methods and results, in: *Fire in the Environment: the Ecological, Atmospheric, and Climatic Importance of Vegetation Fires*, edited by: Crutzen, P. J. and Goldammer, J. G., Wiley, Chichester, England, 53–76, 1993.
- Ward, D. E., Setzer, A. W., Kaufman, Y. J., and Rasmussen, R. A.: Characteristics of smoke emissions from biomass fires of the Amazon region – BASE – a experiment, in: *Global Biomass Burning: Atmospheric, Climatic, and Biospheric Implications*, edited by: Levine, J. S., MIT Press, Cambridge, MA, 394–402, 1991.
- Wilson, J. C., Lafleur, B. G., Hilbert, H., Seebaugh, W. R., Fox, J., Gesler, D. W., Brock, C. A., Huebert, B. J., and Mullen, J.: Function and performance of a low turbulence inlet for sampling supermicron particles from aircraft platforms, *Aerosol Sci. Tech.*, 38, 790–802, 2004.
- Yokelson, R. J., Goode, J. G., Ward, D. E., Susott, R. A., Babbitt, R. E., Wade, D. D., Bertschi, I., Griffith, D. W. T., and Hao, W. M.: Emissions of formaldehyde, acetic acid, methanol, and other trace gases from biomass fires in North Carolina measured by air-

borne Fourier transform infrared spectroscopy, J. Geophys. Res., 104, D23, 30109–30126, doi:10.1029/1999JD900817, 1999.

5 Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., de Gouw, J., Akagi, S. K., Urbanski, S. P., Veres, P., Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, T. J., Hosseini, S., Miller, J. W., Cocker III, D. R., Jung, H., and Weise, D. R.: Coupling field and laboratory measurements to estimate the emission factors of identified and unidentified trace gases for prescribed fires, Atmos. Chem. Phys., 13, 89–116, doi:10.5194/acp-13-89-2013, 2013.

**Smoke marker ratios
from prescribed
burning**

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Information for each research flight with the PILS-TOC and fraction collector system including flight number, date and sampling time, and the location of the fire being sampled.

Flight Number*	Date and Time (LT)	Fire Location	Coordinates (degrees)
RF01	30 Oct 11 12:30–14:00	Fort Jackson, SC Block 6	34°1'29", 80°52'16"
RF02	30 Oct 11 15:00–17:10	Fort Jackson, SC Block 6	34°1'29", 80°52'16"
RF03	1 Nov 11 12:00–15:00	Fort Jackson, SC Block 9b	34°0'15", 80°52'37"
RF05	2 Nov 11 13:00–17:00	Fort Jackson, SC Block 22b	34°5'4", 80°46'23"
RF08	8 Nov 11 12:00–16:00	Francis Marion National Forest, SC	33°12'55", 79°28'34"
RF09	10 Nov 11 11:00–13:00	Midway, SC Bamberg Burn	33°14'5", 80°56'41"

* For RF04 the PILS system was not operational. RF06 was a flight over Columbia to examine urban emissions and did not sample any burning. RF07 had limited access to a prescribed burn in Georgetown, SC due to it being conducted on private land, leading to few smoke impacted fraction collector samples. RF09 is denoted as RF09A and RF09B throughout to indicate the two different ends of this burn.

Smoke marker ratios
from prescribed
burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Concentrations of 2 min averaged absolute WSOC, levoglucosan, mannosan, galactosan, and potassium for each plume directly sampled by the fraction collector system. Also included is the average altitude for each plume as well as estimated time since emission and Modified Combustion Efficiency (MCE) determined from the AFTIR measurements when available. ND = not detected and NA = not applicable.

Date and Time (LT)	WSOC (μgCm^{-3})	Levoglucosan ($\mu\text{g m}^{-3}$)	Mannosan ($\mu\text{g m}^{-3}$)	Galactosan ($\mu\text{g m}^{-3}$)	Potassium ($\mu\text{g m}^{-3}$)	Altitude (m)	Time Since Emission (h)	MCE (gCgC^{-1})
30 Oct 11 12:42:18–12:44:18	6.08	2.04	0.60	0.20	0.54	590.12		
30 Oct 11 12:52:18–10:54:18	7.36	1.75	0.60	0.22	0.52	304.43		
30 Oct 11 12:58:18–13:00:18	6.32	2.66	0.77	0.27	0.66	321.80		
30 Oct 11 13:02:18–13:04:18	16.71	8.10	1.89	0.62	0.72	308.02		
30 Oct 11 13:06:18–13:08:18	28.27	13.83	3.10	0.96	0.52	249.70		
30 Oct 11 13:10:18–13:12:18	10.30	3.23	0.82	0.31	ND	280.45		0.96
30 Oct 11 13:14:18–13:16:18	24.54	11.21	2.43	0.79	0.44	218.53		
30 Oct 11 13:18:18–13:20:18	32.29	14.57	3.19	1.02	0.46	227.58		
30 Oct 11 13:22:18–13:24:18	13.21	3.80	0.96	0.36	0.34	250.43		
30 Oct 11 13:26:18–13:28:18	8.67	1.89	0.63	0.25	2.34	417.38		
30 Oct 11 13:30:18–13:32:18	13.86	5.18	1.21	0.43	0.12	569.80		
30 Oct 11 13:42:18–13:44:18	22.62	8.40	2.00	0.69	0.20	291.52		0.93
30 Oct 11 13:44:18–13:46:18	43.69	13.60	3.11	1.08	ND	273.89		
30 Oct 11 13:48:18–13:50:18	8.82	3.30	0.84	0.34	0.12	281.85		0.88
30 Oct 11 16:55:40–16:57:40	53.82	26.25	5.49	2.68	NA	240.30		
30 Oct 11 17:05:40–17:07:40	3.51	1.04	0.35	0.22	ND	409.28		
1 Nov 11 12:10:18–12:12:18	40.38	13.85	2.80	0.82	0.86	269.48	0.02	0.93
1 Nov 11 12:12:18–12:14:18	80.85	28.57	5.69	1.56	3.19	307.93	0.02	0.90
1 Nov 11 12:18:18–12:20:18	74.14	24.98	5.42	1.53	1.01	297.43		
1 Nov 11 12:30:18–12:32:18	22.43	5.45	1.08	0.34	0.67	397.02		
1 Nov 11 12:40:18–12:42:18	22.66	8.05	1.56	0.48	0.49	643.72	0.03	0.87
1 Nov 11 12:42:18–12:44:18	45.75	12.07	2.41	0.72	0.70	685.25	0.06	0.90
1 Nov 11 12:48:18–12:50:18	29.83	11.00	2.18	0.67	0.86	958.52	0.13	0.92
1 Nov 11 12:52:18–12:54:18	28.93	8.98	1.76	0.54	0.47	1053.30		
1 Nov 11 13:02:18–13:04:18	5.04	1.16	0.16	0.10	0.61	703.53		
1 Nov 11 13:14:18–13:16:18	15.02	3.64	0.64	0.20	0.38	487.93		
1 Nov 11 13:16:18–13:18:18	14.96	4.31	0.77	0.23	0.40	486.98	0.72	
1 Nov 11 13:18:18–13:20:18	10.86	3.22	0.60	0.18	0.67	478.93	1.22	
1 Nov 11 13:36:18–13:38:18	5.92	1.33	0.19	0.07	0.38	1238.35	1.54	
1 Nov 11 13:56:18–13:58:18	23.22	11.03	2.19	0.72	0.49	494.98	0.09	0.77
1 Nov 11 14:12:18–14:14:18	37.67	16.39	3.18	0.91	0.85	944.40	0.04	0.93
1 Nov 11 14:20:18–14:22:18	5.66	2.73	0.35	ND	0.44	1684.88	0.07	0.93
1 Nov 11 14:32:18–14:34:18	3.48	1.18	0.10	ND	0.32	1183.07		
1 Nov 11 14:48:18–14:50:18	4.23	1.64	0.17	ND	0.67	368.12		0.98
1 Nov 11 14:50:18–14:52:18	1.74	0.42	ND	ND	0.78	294.02		

Table 2. Continued.

Date and Time (LT)	WSOC (μgCm^{-3})	Levoglucosan (μgm^{-3})	Mannosan (μgm^{-3})	Galactosan (μgm^{-3})	Potassium (μgm^{-3})	Altitude (m)	Time Since Emission (h)	MCE (gCgC^{-1})
2 Nov 11 13:07:46–13:09:46	13.14	4.81	1.22	0.49	1.29	618.85		
2 Nov 11 13:29:46–13:31:46	15.30	4.70	1.21	0.47	2.26	349.10		0.92
2 Nov 11 13:31:46–13:33:46	30.08	9.19	2.23	0.73	2.42	264.32		
2 Nov 11 13:41:46–13:43:46	15.87	5.11	1.09	0.46	1.43	595.72		
2 Nov 11 14:09:46–14:11:46	42.08	15.16	3.35	1.12	ND	421.67		
2 Nov 11 14:15:46–14:17:46	27.86	12.77	2.72	0.99	1.43	399.45		0.88
2 Nov 11 14:39:46–14:41:46	19.06	9.40	2.06	0.76	1.55	305.38		
2 Nov 11 14:41:46–14:43:46	35.84	12.20	2.76	1.00	ND	235.32		0.94
2 Nov 11 15:17:46–15:19:46	31.61	9.84	2.03	0.81	ND	1248.65		
2 Nov 11 15:23:46–15:25:46	16.61	4.38	0.93	0.48	ND	1179.45		
2 Nov 11 15:31:46–15:33:46	36.71	9.67	2.16	0.82	1.32	1188.73		0.94
2 Nov 11 15:33:46–15:35:46	24.12	5.50	1.15	0.45	1.37	1162.23		
2 Nov 11 15:35:46–15:37:46	15.37	3.22	0.76	0.33	1.63	1169.90		0.94
2 Nov 11 15:41:46–15:43:46	34.27	14.11	3.04	1.15	1.82	1081.77		0.91
2 Nov 11 16:41:46–16:43:46	27.08	8.07	1.32	0.54	ND	304.35		0.90
8 Nov 11 12:28:52–12:30:52	5.24	1.30	0.14	0.11	1.24	406.93	0.04	0.94
8 Nov 11 12:30:52–12:32:52	6.89	1.20	0.24	0.18	0.66	133.55	0.00	0.93
8 Nov 11 12:38:52–12:40:52	30.22	7.41	1.10	0.72	2.84	130.13	0.03	0.93
8 Nov 11 12:56:52–12:58:52	25.12	5.85	1.05	0.51	1.58	121.62	0.02	0.95
8 Nov 11 13:02:52–13:04:52	14.53	3.44	0.48	0.26	2.17	96.42	0.02	0.94
8 Nov 11 13:10:52–13:12:52	26.85	8.26	1.42	0.75	3.53	139.30		
8 Nov 11 13:14:52–13:16:52	16.12	2.76	0.45	0.30	0.99	129.32	0.01	0.92
8 Nov 11 13:18:52–13:20:52	9.11	1.90	0.28	0.19	1.86	122.83		
8 Nov 11 13:22:52–13:24:52	7.23	1.60	0.25	0.23	0.88	173.30	0.01	0.94
8 Nov 11 13:24:52–13:26:52	26.71	6.13	1.14	0.59	1.19	136.88	0.03	0.94
8 Nov 11 13:28:52–13:30:52	20.29	3.71	0.56	0.32	1.38	138.68		
8 Nov 11 14:18:52–14:20:52	10.93	2.94	0.51	0.47	1.27	737.47	0.83	
8 Nov 11 14:44:52–14:46:52	4.46	1.02	0.19	0.26	1.19	423.23		
8 Nov 11 14:48:52–14:50:52	22.10	6.07	1.11	0.71	1.27	163.37	0.03	0.93
8 Nov 11 14:50:52–14:52:52	14.69	4.84	0.88	0.58	0.77	159.43	0.07	0.90
10 Nov 11 11:11:56–11:13:56	14.07	2.50	0.30	0.33	ND	252.98	0.00	0.96
10 Nov 11 11:15:56–11:17:56	13.68	2.11	0.21	0.29	0.87	256.03	0.26	0.94
10 Nov 11 11:25:56–11:27:56	9.24	1.72	0.23	0.27	0.67	314.86	0.24	0.95
10 Nov 11 11:31:56–11:33:56	10.69	1.51	0.15	0.20	1.07	307.85	0.23	0.98
10 Nov 11 11:35:56–11:37:56*	8.99	0.64	0.02	0.14	0.89	314.86	0.21	0.97
10 Nov 11 11:45:56–11:47:56*	7.26	0.50	ND	ND	0.76	303.89	0.44	0.99
10 Nov 11 11:49:56–11:51:46*	6.70	0.54	0.04	0.17	ND	326.14	0.48	0.98
10 Nov 11 12:27:56–12:29:56*	6.51	0.72	0.21	0.12	0.82	466.95	1.51	
10 Nov 11 12:31:56–12:33:56	12.14	2.21	0.40	0.28	3.09	469.09	0.28	0.80

* Denotes plumes from fire B during the RF09 flight.

Smoke marker ratios
from prescribed
burning

A. P. Sullivan et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. WSOC to OC, levoglucosan to WSOC on a carbon mass basis, galactosan to levoglucosan, mannosan to levoglucosan, and potassium to levoglucosan ratios from controlled laboratory burns. Ratios were determined as the slope of a linear regression between the two species using data from the Fire Lab at Missoula Experiments (FLAME) (Sullivan et al., 2014).

Fuel Type	WSOC/OC ($\mu\text{g C } \mu\text{g C}^{-1}$)	Levoglucosan/ WSOC ($\mu\text{g C } \mu\text{g C}^{-1}$)	Galactosan/ Levoglucosan ($\mu\text{g } \mu\text{g}^{-1}$)	Mannosan/ Levoglucosan ($\mu\text{g } \mu\text{g}^{-1}$)	Potassium/ Levoglucosan ($\mu\text{g } \mu\text{g}^{-1}$)
Grasses	0.81 ± 0.02	0.149 ± 0.012	0.060 ± 0.005	0.051 ± 0.005	0.211 ± 0.026
Leaves*	0.54 ± 0.02	0.095 ± 0.006	0.094 ± 0.009	0.027 ± 0.008	no correlation
Needles*	0.54 ± 0.02	0.064 ± 0.008	0.119 ± 0.010	0.249 ± 0.016	0.079 ± 0.009
Marsh Grasses	0.78 ± 0.07	0.017 ± 0.014	0.095 ± 0.038	0.006 ± 0.002	no correlation

* Includes the burning of live and dead material.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Example of a typical flight path. This flight path is from RF08 and is colored by CO to indicate the location of the burn and where the smoke plume was intercepted downwind.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.




Fig. 2. Time series of 1 s altitude along with absolute CO, levoglucosan, and WSOC from RF01.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Smoke marker ratios
from prescribed
burning

A. P. Sullivan et al.




Fig. 3. Time series of 2 min absolute (a) levoglucosan and WSOC and (b) galactosan, levoglucosan, and mannosan from RF01.

Smoke marker ratios
from prescribed
burning

A. P. Sullivan et al.




Fig. 4. Correlation of (a) Δ levoglucosan vs. Δ WSOC on a carbon mass basis for all flights with the data segregated by fire location and (b) Δ levoglucosan vs. Δ WSOC on a carbon mass basis for only the Fort Jackson prescribed burns colored by altitude. In plot a, the fit through the filter samples collected on the ground during the burns conducted at Fort Jackson is also provided. Uncertainties with the least square regressions are one standard deviation.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.




Fig. 5. Correlation of (a) Δ galactosan vs. Δ levoglucosan, (b) Δ mannosan vs. Δ levoglucosan, and (c) Δ potassium vs. Δ levoglucosan for all flights with the data segregated by fire location. Uncertainties with the least square regressions are one standard deviation.

Smoke marker ratios
from prescribed
burning

A. P. Sullivan et al.




Fig. 6. (a) Δ levoglucosan/ Δ WSOC ratio on a carbon mass basis and (b) $\Delta m/z\ 60/\Delta$ OA ratio as a function of time since emission for all flights with the data segregated by fire location.




Fig. 7. Correlation of $\Delta m/z$ 60 vs. Δ OA for all flights with the data segregated by fire location. Uncertainty with the least square regression is one standard deviation.

Smoke marker ratios from prescribed burning

A. P. Sullivan et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smoke marker ratios
from prescribed
burning

A. P. Sullivan et al.




Fig. 8. (a) $\Delta m/z 60/\Delta OA$ ratio as a function of ΔOA , (b) $\Delta m/z 60/BBOA$ ratio as a function of ΔOA , and (c) $\Delta \text{levoglucosan}/\Delta \text{WSOC}$ ratio on a carbon mass basis as a function of ΔWSOC for all flights with the data segregated by fire location.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 9. Δ levoglucosan/ Δ WSOC ratio on a carbon mass basis for (a) RF01/RF02/RF03/RF05 and controlled laboratory burns involving grasses, (b) RF08 and controlled laboratory burns involving leaves, (c) RF09A and controlled laboratory burns involving needles, and (d) RF09B and controlled laboratory burns involving marsh grasses as a function of modified combustion efficiency. In each plot the squares represent the prescribed burns and circles controlled laboratory burns.