



Supplement of

Kinetics of the reactions of isoprene-derived hydroxynitrates: gas phase epoxide formation and solution phase hydrolysis

M. I. Jacobs et al.

Correspondence to: M. J. Elrod (mjelrod@oberlin.edu)

Supplemental Information for

Gas Phase and Solution Phase Study of Atmospherically Relevant Isoprene-derived Hydroxynitrates

M. I. Jacobs, W. J. Burke, and M. J. Elrod

Department of Chemistry and Biochemistry, Oberlin College, Oberlin, Ohio, USA

Correspondence to: M. J. Elrod (mjelrod@oberlin.edu)

1 Synthesis of hydroxynitrates

1.1 Synthesis of 4-hydroxy-3-nitrooxy isoprene (4,3-HNI)

The synthesis of 4,3-HNI required the preparation of 3,4-epoxyisoprene (2-(prop-1-en-2-yl)oxirane), which was synthesized according to a procedure first reported by Harwood *et al.* (Harwood *et al.*, 1990). In a 1-L round bottom flask, 60 g of powdered NaOH and 50 g of trimethylsulfonium methyl sulphate (Oakwood Products, Inc.) were added to 600 mL of CH₂Cl₂. The solution was vigorously stirred while 20 mL of methacrolein (Sigma-Aldrich) were added in 300 μ L increments every 3 minutes. The reaction mixture was washed with 200 mL of DI H₂O thrice to remove the solid NaOH. Anhydrous MgSO₄ (roughly 5 g) was added to the resulting organic phase to dry the solution. After filtering off the MgSO₄, a majority of the solvent was removed at atmospheric pressure (50 °C water bath) using a fractionating (Vigreux) column yielding approximately 20 mL of residue. Atmospheric pressure distillation was performed on the residue and the liquid (6.80 g) with a boiling range of 65-80 °C was collected. A ¹H NMR spectrum confirmed the presence and purity of the desired epoxide.

4,3-HNI was synthesized from 3,4-epoxy isoprene via reaction with fuming HNO₃ (Muthuramu *et al.*, 1993). In a 1-L round bottom flask, 2.75 g of 3,4-epoxyisoprene were added to 200 mL of CH₂Cl₂ and cooled to 0 °C. Fuming HNO₃ (10 mL) was added drop-wise using a burette over the course of 20 minutes. After addition of HNO₃, the reaction mixture was refluxed in a 50 °C water bath for 15 minutes. After refluxing, NaHCO₃ was added slowly until all of the acid had been neutralized. The solid was filtered off and the vast majority of the solvent was removed under reduced pressure (150-400 mbar, 30 °C). The residue was purified with flash chromatography using a 14-in. silica gel (Baker, 40 μ m packing, 60 Å pore diameter) column, with a hexanes and ethyl acetate (3:2 ratio) eluent. Nitrate species were detected using

diphenylamine as previously described (Treves et al., 2000). ^1H and ^{13}C NMR confirmed the presence and purity of 4,3-HNI. The final remaining solvent was removed using the vacuum system at low pressure (150 mtorr, room temperature), and the final product was stored at $-80\text{ }^\circ\text{C}$.

1.2 Synthesis of 1-hydroxy-4-nitrooxy isoprene (1,4-HNI)

1,4-HNI was synthesized from 2-methyl-2-vinylloxirane (Sigma-Aldrich) via reaction with fuming HNO_3 using a modified method to minimize the presence of water (Lockwood et al., 2010). A 1.5:1 molar ratio of 2-methyl-2-vinylloxirane to fuming HNO_3 was used. In a 500-mL round bottom flask, 2-methyl-2-vinylloxirane (3.00 g, 35.7 mmol) was added to 200 mL of CHCl_3 under a N_2 flow at $0\text{ }^\circ\text{C}$. Fuming HNO_3 (1.00 mL, 24.0 mmol) was added drop-wise to the reaction flask over the course of 10 minutes. The remaining procedure matched that of 4,3-HNI as described above, except a 2:1 hexane to ethyl acetate mobile phase was used for flash chromatography.

1.3 Synthesis of 1-hydroxy-2-nitrooxybut-3-ene (1,2-HNB)

1,2-HNB was synthesized using the 4,3-HNI and 1,4-HNI synthetic procedures detailed above. 2.0 g of 3,4-epoxy-1-butene (98%, Sigma-Aldrich) was dissolved in 200 mL CH_2Cl_2 . 5.0 mL of fuming HNO_3 was added drop-wise over the course of 15 minutes. The remaining procedures were identical to 1,4-HNI synthesis.

1.4 NMR assignments

All ^1H chemical shifts for deuterated chloroform and aqueous solutions were determined relative to the CHCl_3 and HDO peaks at 7.26 and 4.79 ppm, respectively. The primary contaminant for the synthesized hydroxynitrates in each of the spectra is ethyl acetate, which was not removed entirely until gas phase experiments were performed. Peaks from ethyl acetate are

labeled as EA while peaks from the relevant hydroxynitrate are labeled according to a numbering scheme where the isoprene backbone is numbered 1 through 4 (using the isoprene IUPAC naming scheme) with the methyl group being carbon number 5. The protons on each isomer are labeled a-e with proton “a” attached to carbon “1”, proton “c” attached to carbon “3”, etc. Each isomer is labeled on the ^1H and ^{13}C spectra for reference.

1.4.1 4,3-HNI structure assignment

^1H and ^{13}C spectra of 4,3-HNI were collected in CDCl_3 and the annotated spectra are given below (Figures S1 and S2). The structural position of the nitrate group was confirmed using HMQC. The HMQC spectrum (Figure S3) shows that ^1H peak at 5.32 ppm is a nitrate proton and not an alkene proton due to the coupling with the ^{13}C peak at 86.69 ppm (which is too low for an alkene carbon). The triplet and doublet splitting patterns of the nitrate peak and the alcohol peak (at 3.78 ppm), respectively, confirms that the secondary nitrate was exclusively formed in the synthetic procedure. The final ^1H and ^{13}C assignments for 4,3-HNI are presented in Table S1.




Figure S1. ^1H NMR spectrum of 4,3-HNI collected in CDCl_3 .




Figure S2. ^{13}C NMR spectrum of 4,3-HNI collected in CDCl_3 (EA indicates ethyl acetate solvent impurity).




Figure S3. HMQC NMR spectrum of 4,3-HNI collected in CDCl_3 . The red box indicates the coupling between peaks at 5.32 ppm in the ^1H spectrum and 86.69 ppm from the ^{13}C spectrum, which confirms those carbon and hydrogen nuclei as being adjacent to the nitrate functional group.

1.4.2 E and Z 1,4-HNI structure assignment

^1H and ^{13}C spectra of 1,4-HNI were collected in CDCl_3 and the annotated spectra are given below (Figures S4 and S5). The structural position of the nitrate group was once again confirmed using HMQC. The HMQC spectrum (Figure S6) shows that the ^1H peak at 5.00 ppm is a nitrate proton and not an alkene proton due to the coupling with the ^{13}C peaks at 69.3 and 68.9 ppm (which are too low for an alkene carbon). Furthermore, a COSY spectrum (Figure S7) showed coupling between the ^1H nitrate peak at 5.00 ppm and the ^1H alkene peaks at 5.46 and 5.65 ppm, which demonstrates the nitrate group is adjacent to an alkene carbon atom. The fact that no other peaks are observed indicates that the other possible isomers (e.g. 2,1- and 1,2-HNI) were not formed. Because the two isomers were not synthesized in equal yields, the assignment of the E vs. Z isomer was made using the relative sizes of the peaks—primarily E-1,4-HNI was

formed, so the larger product peaks were attributed to E-1,4-HNI. The final assignments for (E) 1,4-HNI and (Z) 1,4-HNI are presented in Table S1.




Figure S4. ^1H NMR spectrum of (E,Z) 1,4-HNI collected in CDCl_3




Figure S5. ^{13}C NMR spectrum of (E,Z) 1,4-HNI collected in CDCl_3




Figure S6. HMQC NMR spectrum of E and Z 1,4-HNI collected in CDCl_3 . The red box indicates the coupling between peaks at 5.00 ppm in the ^1H spectrum and 68.9 and 69.3 ppm from the ^{13}C spectrum, which confirms those carbon and hydrogen nuclei as being adjacent to the nitrate functional group.




Figure S6. COSY NMR spectrum of E and Z 1,4-HNI collected in CDCl_3 . The red box indicates the coupling between the 5.00 ppm ^1H peaks and the 5.46 and 5.65 ppm ^1H peaks, which demonstrates the nitrate group is adjacent to an alkene carbon atom in both of the isomers.

H label	C label	4,3-HNI		(E) 1,4-HNI		(Z) 1,4-HNI	
		^1H	^{13}C	^1H	^{13}C	^1H	^{13}C
A	1	5.10(ds)	115.16	4.07(s)	67.11	4.22(s)	61.64
	2		138.52		145.21		145.30
C	3	5.32(t)	86.69	5.65(t)	114.03	5.46(t)	117.36
D	4	3.78(d)	61.69	5.00(d)	68.88	5.00(d)	69.28
E	5	1.88(s)	19.18	1.76(s)	13.94	1.87(s)	21.38

Table S1. ^1H and ^{13}C NMR (in CDCl_3 solvent) assignments for all three of the synthesized isoprene derived hydroxynitrates.

A previous study has reported the synthesis of 1,2-HNI (a tertiary nitrate) and 2,1-HNI using a similar procedure as that reported here (Lockwood et al., 2010). The mechanism for the four possible hydroxynitrate isomers from 2-methyl-2-vinyloxirane is provided in Figure S7. Previous work from our laboratory has demonstrated that hydrolysis of the tertiary nitrate occurs rapidly (with a lifetime of only a couple of minutes) and because of this, isolating the tertiary

nitrate using conventional chromatographic techniques is expected to be very difficult (Darer et al., 2011; Hu et al., 2011). The hydroxynitrate synthetic procedures reported herein were designed to minimize the amount of water added to the reaction to avoid hydrolysis of the products. Nonetheless, based on the NMR data collected, we did not detect the 1,2- nor 2,1-HNI isomers at any point in the synthetic process.




Figure S7. Mechanism for E and Z 1,4-HNI and 1,2-HNI synthesis from reaction of 2-methyl-2-vinyloxirane with fuming nitric acid.

1.4.3 1,2-HNB assignment

¹H and ¹³C spectra of 1,2-HNI were collected in CDCl₃ and the annotated spectra are given below (Figures S8 and S9). The peak assignments for 1,2-HNB were made based on the NMR spectra for the other two hydroxynitrates, and the specific shifts are given in Table S2.




Figure S8. ^1H NMR spectrum of 1,2-HNB collected in CDCl_3




Figure S9. ^{13}C NMR spectrum of 1,2-HNB collected in CDCl_3

H label	C label	1,2-HNB	
		¹ H	¹³ C
a	1	3.81	62.44
b	2	5.45	84.57
c	3	5.85	130.11
d	4	5.47	121.56

Table S2. ¹H and ¹³C NMR assignments (in CDCl₃ solvent) for 1,2-HNB.

References

Darer, A. I., Cole-Filipiak, N. C., O'Connor, A. E. and Elrod, M. J.: Formation and stability of atmospherically relevant isoprene-derived organosulfates and organonitrates., *Environ. Sci. Technol.*, 45, 1895–902, doi:10.1021/es103797z, 2011.

Harwood, L. M., Casy, G. and Sherlock, J.: A Simple Laboratory Procedure for Preparation of (1-methylethenyl)oxirane (3,4-epoxyisoprene), *Synth. Commun.*, 20(9), 1287–1292, 1990.

Hu, K. S., Darer, A. I. and Elrod, M. J.: Thermodynamics and kinetics of the hydrolysis of atmospherically relevant organonitrates and organosulfates, *Atmos. Chem. Phys.*, 11, 8307–8320, doi:10.5194/acp-11-8307-2011, 2011.

Lockwood, A. L., Shepson, P. B., Fiddler, M. N. and Alaghmand, M.: Isoprene nitrates: preparation, separation, identification, yields, and atmospheric chemistry, *Atmos. Chem. Phys.*, 10, 6169–6178, doi:10.5194/acp-10-6169-2010, 2010.

Muthuramu, K., Shepson, P. B., Street, K. and York, N.: Preparation, Analysis, and Atmospheric Production of Multifunctional Organic Nitrates, *Environ. Sci. Technol.*, 27, 1117–1124, 1993.

Treves, K., Shragina, L. and Rudich, Y.: Henry's Law Constants of Some β-, γ-, and δ- Hydroxy Alkyl Nitrates of Atmospheric Interest, *Environ. Sci. Technol.*, 34, 1197–1203, 2000.