

**Measured  
morphology of  
biomass burning  
CCN**

M. Giordano et al.

# Experimentally measured morphology of biomass burning aerosol and its impacts on CCN ability

**M. Giordano<sup>1,2</sup>, C. Espinoza<sup>1,2</sup>, and A. Asa-Awuku<sup>1,2</sup>**

<sup>1</sup>Department of Chemical and Environmental Engineering, University of California – Riverside, Riverside, California, USA

<sup>2</sup>College of Engineering – Center for Environmental Research and Technology (CE-CERT), Riverside, CA, USA

Received: 10 April 2014 – Accepted: 17 April 2014 – Published: 16 May 2014

Correspondence to: A. Asa-Awuku (akua@engr.ucr.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

This study examines the morphological properties of freshly emitted and atmospherically aged aerosols from biomass burning. The impacts of particle morphology assumptions on hygroscopic predictions are examined. Chamber experiments were conducted at the UC-Riverside Center for Environmental Research and Technology (CE-CERT) Atmospheric Processes Lab using two biomass fuel sources, manzanita and chamise. Morphological data was obtained through the use of an aerosol particle mass analyzer (APM), scanning mobility particle sizer (SMPS) system and transmission electron microscopy (TEM). Data from these instruments was used to calculate both a dynamic shape factor and a fractal-like dimension for the biomass burning emissions. This data was then used with  $\kappa$ -Köhler theory to adjust the calculated hygroscopicity for experimentally determined morphological characteristics of the aerosol. Laboratory measurement of biomass burning aerosol from two chaparral fuels show that particles are non-spherical with dynamic shape factors greater than 1.15 for aerosol sizes relevant to cloud condensation nuclei (CCN) activation. Accounting for particle morphology can shift the hygroscopicity parameter  $\kappa$  by 0.15 or more. To our knowledge, this work provides the first laboratory chamber measurements of morphological characteristics for biomass burning cloud condensation nuclei and provides experimental particle shape evidence to support the variation in reported hygroscopicities of the complex aerosol.

## 1 Introduction

Aerosols have important health effects, affect regional visibility, are a key factor in the earth's climate via radiative forcing mechanisms, and play a vital role in atmospheric chemistry (Seaton et al., 1995; Haywood and Boucher, 2000; Ramanathan et al., 2001; Pöschl, 2005; Forster et al., 2007). For all impacts, particle composition and size are important. Particles are ubiquitous but can be non-uniform and non-spherical.

ACPD

14, 12555–12589, 2014

### Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Measured  
morphology of  
biomass burning  
CCN**

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As such, a number of instruments exist to measure these critical aerosol properties. One property that remains difficult to characterize through conventional techniques is particle morphology. Morphology is used here as a term to describe the shape of a 3-dimensional particle and has a direct impact on particle size measurements (DeCarlo et al., 2004). The true volume of a particle is difficult to accurately measure without knowledge of the particle morphology. Reducing uncertainties in particle morphology will increase the accuracy of characterizations and parameterizations that rely on knowing the true volume of a particle, e.g. particle density and hygroscopicity.

Biomass burning emissions are a major source of non-uniform and non-spherical complex aerosol emissions (2–5 petagrams of Carbon year<sup>-1</sup>) to the atmosphere (Crutzen and Andreae, 1990; Andreae et al., 2004; Reid et al., 2005; van der Werf et al., 2006). Both the developed and developing world are subject to biomass burning events through agricultural burning, wildfires, and domestic burning applications (Forster et al., 2007). Particles formed during the combustion process are non-uniform. The aerosol formed during biomass burning events is a complex mixture of volatile, semi-volatile, and non-volatile components that are able to seed clouds, modify cloud properties, and indirectly impact radiative forcing (Forster et al., 2007; Reutter et al., 2009; Liu and Wang, 2010). The impact of a potential wildfire or prescribed burn depends on the fuel type, the burning regime (e.g. smoldering, flaming, or mixed), and the ageing time of the atmospheric emissions. This work further explores the role of biomass burning aerosol as Cloud Condensation Nuclei (CCN). We obtain information about the shape and true volume of biomass burning particles using a semi-continuous method to investigate the impact of particle size, changes in shape that occur during the evolution of photochemical ageing and changes in perceived hygroscopicity as a result of particle volume and sizing assumptions.

## 2 Theoretical considerations

### 2.1 Biomass burning cloud condensation nuclei

Several recent works on biomass burning aerosol have focused on their ability to act as cloud condensation nuclei (CCN) (e.g.: Novakov and Corrigan, 1996; Martins et al., 2009; Petters et al., 2009; Carrico et al., 2010; Rose et al., 2010, 2011; Hennigan et al., 2011; Engelhart et al., 2012; Giordano et al., 2013; Latham et al., 2013). The CCN ability of inorganic compounds can be described well by classical Köhler theory (Köhler, 1936). The theory has since been expanded to better model organic/inorganic mixed particles with varying degrees of accuracy and complexity (e.g. Shulman et al., 1996; Raymond and Pandis, 2002). One extension,  $\kappa$ -Köhler theory, has become a common method for reporting CCN ability and estimating particle hygroscopicity. Its popularity is due to its empirical grounding, ease of understanding, ability to represent organic/inorganic mixed particles and translate to both the sub- and supersaturated regimes. This has led to a large body of published  $\kappa$ -Köhler theory values. The  $\kappa$  parameterization of hygroscopicity posits that particle diameter and the supersaturation at the point of activation effectively capture CCN ability (Petters and Kreidenweis, 2007):

$$\kappa = \frac{4A^3}{27D_d^3 \ln^2 s_c}; \text{ where } A = \frac{4\sigma_{s/a} M_w}{RT \rho_w} \quad (1)$$

where  $\rho_w$  is the density of water,  $M_w$  is the molecular weight of water,  $\sigma_{s/a}$  is the surface tension of the solution/air interface,  $R$  is the universal gas constant,  $T$  is the temperature,  $D_d$  is the dry particle diameter, and  $s_c$  is the critical supersaturation required to activate a particle with size  $D_d$ .  $D_d$  is the measured mobility diameter and is calculated as the 50 % efficiency for which half of all particles activate at  $s_c$ , the instrument supersaturation. From Eq. (1),  $\kappa$  is interpolated using a linear regression of experimental  $D_d$  and calibrated  $s_c$ .

### Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Thus, the CCN activity can be represented by a single hygroscopicity parameter,  $\kappa$ . A range of  $\kappa$  biomass burning literature values has been reported. Multiple factors affect this range. Different fuels can have a wide variability in CCN activity. The particles can be very non-hygroscopic (e.g.  $\kappa < 0.1$  for various softwoods) to very hygroscopic (e.g.  $\kappa > 0.5$  for sage and sawgrass) (Petters et al., 2009; Carrico et al., 2010). Differences in burning conditions contribute to the range of  $\kappa$ . Additionally, the oxidative aging of these emissions can affect the observed  $\kappa$  values (Hennigan et al., 2011; Engelhart et al., 2012). The presence of surface active compounds can also affect the range of  $\kappa$  (Giordano et al., 2013).

$\kappa$ -Köhler theory applies a few assumptions. First, ideal droplet water activity conditions are assumed when the growing cloud droplet exists as an aqueous solution and all available solute is dissolved. Second,  $\kappa$ -Köhler theory applies the ideal surface tension of water,  $\sigma_{s/a} = 0.072 \text{ J m}^{-2}$ . Third, and most important for application to biomass burning, the diameter of the particle cubed,  $D_a^3$ , is assumed to estimate the volume of a spherical particle. A number of studies have investigated the applicability of the first two assumptions and come to varying conclusions on their importance. However, the assumption of particle sphericity for hygroscopicity has only been theoretically manipulated and not experimentally measured, especially as a function of particle aging (Dusek et al., 2011). Here we provide simultaneous morphological and hygroscopicity measurements of fresh and aged biomass aerosol. These measurements together provide a more accurate parameterization of particle hygroscopicity.

## 2.2 Volume equivalent and electrical mobility diameters

CCN activity measurements require either mass or volume information in addition to droplet activation data. Particle diameter alone can introduce 84 to 96 % variation in CCN concentrations (Dusek et al., 2006). Particle volume information is commonly obtained from mobility particle sizing instrumentation and is combined with a cloud condensation nuclei counter (CCNC). This experimental method selects the electrical mobility diameter of a particle. The mobility-selected particle then flows to the

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



supersaturated column of the CCNC where, if above a critical size, it activates to form a cloud droplet. When combined with  $\kappa$ -Köhler theory, the method assumes that a particle's electrical mobility is equivalent to its volume equivalent diameter. For a spherical particle this is true: the mobility diameter equals the volume-equivalent diameter. However, for non-spherical particles, electrical mobility diameter deviates from the volume equivalent diameter. The phenomena has also been observed in CCN experiments using mass-selected particles measured with a Kanomax APM (Kuwata and Kondo, 2009). The volume equivalent diameter ( $d_{ve}$ ) is defined as the diameter of a spherical particle of the same volume as the particle of interest (Baron and Willeke, 2001). For a non-spherical particle,  $d_{ve}$  is the diameter of the particle if it were compacted or melted to form a sphere while preserving internal void spaces (empty regions of the particle that are isolated from the surrounding gas).

The electrical mobility diameter,  $d_m$ , can be measured with a differential mobility analyzer (DMA) and is defined as the diameter of a sphere with the same migration velocity in a constant electric field as the considered particle (Knutson and Whitby, 1975). A force balance occurs in the annular region of the DMA to mobility-select a certain "size" particle. The particle is first charged and flown into an annulus with a constant electric field being applied between the inner and outer walls. As the particle moves in the electric field, the charge on the particle is attracted toward the inner annular wall. A drag force between the particle and gas is equal and opposite to the electrical force when the particle reaches a terminal migration velocity. Careful design of the annulus allows for a particle of a desired electrical mobility to exit the DMA in a monodisperse size stream.

Assuming a particle and its volume equivalent sphere have the same charge, the relationship between the two diameters is as follows:

$$\frac{d_m}{C_C(d_m)} = \frac{d_{ve}\chi}{C_C(d_{ve})} \quad (2)$$

where  $C_C$  is the Cunningham slip correction factor and  $\chi$  is the dynamic shape factor. The dynamic shape factor is a correction to account for the increased drag on a non-

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spherical particle (Fuchs, 1964). The shape factor is equal to one for a sphere and can be greater than one for non-spherical particles. The flow regime of the suspending fluid relative to the particle dictates the nature of the forces acting on the particle. Hence, both  $C_C$  and  $\chi$  are functions of the Knudsen number. The flow regime here is in the transition regime because both particle sizing and the CCNC commonly operate near standard ambient temperature and pressure (approx. 298 K, 1 atm) and the particles of interest as CCN are between 10–200 nm. This introduces an added level of complexity to volume calculations, as the dynamic shape factor behavior is more readily available in the continuum and free molecular regimes (Dahneke, 1973a, b; Kousaka et al., 1996; Baron et al., 2001). However, other work on non-spherical particles and their dynamic shape factors have assumed that the shape factor in the transition regime is approximately equal to the shape factor in the free-molecular regime (DeCarlo et al., 2004). The analysis presented here also makes this assumption. Our analysis also neglects particle orientation effects on the shape factor. Because  $\chi$  can be greater than one for non-spherical particles, the use of electrical mobility for volume measurement will likely over predict particle volume. An over prediction in particle volume in Eq. (1) will underestimate calculated  $\kappa$ -values thus misrepresenting the cloud forming ability of non-spherical aerosols.

### 2.3 Fractal and fractal-like dimensions

Combustion processes often generate aggregates of small ( $d < 10$ –30 nm) particles. The fractal dimension ( $D_f$ ) is a semi-empirical relationship often used to describe aggregates.  $D_f$  is defined as the relationship between the number of primary particles in an aggregate to the aggregate's radius of gyration. The relation can be written as a power law expression (Mandelbrot, 1982):

$$N = C \left( \frac{R_g}{a} \right)^{D_f} \quad (3)$$

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where  $N$  is the number of primary particles in the aggregate,  $R_g$  is the radius of gyration of the aggregate,  $a$  is the radius of the primary particles, and  $C$  is a constant. Park et al. (2003) use this relationship with the assumption that the primary particle diameters are approximately constant to show that  $N$  then varies proportionately to particle mass. Park et al. also assume that the radius of gyration is linearly proportional to the mobility diameter in the continuum and free-molecular regimes (Schmidt-Ott et al., 1990) to define a “fractal-like” dimension, here referred to as  $D_f'$ :

$$m = C' D_m^{D_f'} \quad (4)$$

where  $m$  is the mass of the aggregate and  $C'$  is a constant. Here, a  $D_f'$  of 3 implies a spherical particle while  $D_f' < 3$  indicates a non-spherical particle. This relationship, and its analogue in density, is directly applicable to data obtained from the Kanomax Aerosol Particle Mass Analyzer (APM). Park et al. used APM data to determine the fractal dimension of diesel exhaust particles. They obtained fractal dimensions of  $2.33 < D_f' < 2.84$  depending on engine load. Work done by Nakao et al. (2011) used this relationship to show SOA formation from diesel exhaust have similar  $D_f'$ . Zhang et al. (2008) exposed flame-generated soot to subsaturated sulfuric acid vapor and obtained a fresh (unexposed)  $D_f'$  of 2.1 and an aged (exposed)  $D_f'$  of 2.8.

In this work, we obtain information about the shape and true volume of biomass burning particles using a semi-continuous method. We do this for both freshly emitted and photochemically aged emissions. We provide evidence to support the work of others that demonstrate that, (1) the particles emitted are non-spherical, (2) the degree of non-sphericity changes with photochemical aging, and (3) using the true volume-equivalent diameter to calculate  $\kappa$  greatly shifts the predicted hygroscopicity. We also discuss how the limitations of the current methods for measuring particle volume affects CCN measurements.



### 3 Experimental methods and instrumentation

A series of biomass burning experiments were carried out at the University of California, Riverside College of Engineering – Center for Environmental Research and Technology (UCR CE-CERT) atmospheric processes lab from the spring to the summer of 2013. Two shrubs common to the California chaparral biome, chamise (*Adenostoma fasciculatum*) and manzanita (*Arctostaphylos glandulosa*), were burned in an outdoor wood-burning stove. Both fuels were collected in the San Bernardino National Forest (approximately 33.84°, -116.88° and the surrounding area). The collection site is approximately 3600–3900 feet in elevation. Chaparral species such as chamise, manzanita, scrub oak, and ceanothus surround the area. Both manzanita and chamise burned in the flaming regime. The flame integrated modified combustion efficiencies (MCE) of both fuels were greater than 0.97 (where  $MCE = \frac{\Delta CO_2}{(\Delta CO_2 + \Delta CO)}$ ) (Ward and Hao, 1991).

A sample line from the stack of the stove drew smoke from the fire and diluted it with a Venturi tube injection system. The diluted sample was then injected into a 12 m<sup>3</sup> Teflon chamber to a final particle concentration ~ 100 µg m<sup>-3</sup>. The chamber concentration is within range of ambient and plume-like concentrations also used in previous chamber studies (Hennigan et al., 2011). The sample was injected with a 10 : 1 total flow to sample flow ratio. As short of a copper transfer line as possible (approximately 7 m in length) was used for the smoke injection. The enclosure is surrounded by 170 ultraviolet (UV) lights with peak intensity of 350 nm (350 BL, Sylvania®). UV lights were turned on one hour after injection to promote photochemistry and remained on until the experiment concluded. The chamber was flushed and filled with filtered air prior to injection. The temperature of the chamber ranged approximately from 20–27 °C during the experiments. The experiments were operated until the chamber was empty (up to 6 h). This series of chamber burns produced data similar to that of ambient studies and previous biomass burning studies conducted at CE-CERT (Hennigan et al., 2011; Giordano et al., 2013).

ACPD

14, 12555–12589, 2014

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Scanning Mobility CCN Analysis (SMCA) was used to obtain fast size-resolved CCN activity of the aging system (Moore et al., 2010). The sampled aerosol was size-classified using a Scanning Mobility Particle Sizer (SMPS, TSI 3080/3081). The monodisperse stream from the classifier was split and one stream entered a condensation particle counter (TSI 3772) while the other stream was sampled with a DMT Continuous Flow Streamwise Thermal Gradient Chamber CCN counter (CCNC) (Roberts and Nenes, 2005; Rose et al., 2008). The CCNC instrument was operated at a 10 : 1 sheath-to-aerosol ratio with a sample flow of  $0.5 \text{ L min}^{-1}$ . The total concentration (CN) of sized particles measured by the CPC determined the ratio of CCN/CN. For each supersaturation,  $s_c$ , the activation diameter,  $D_d$ , was defined as the diameter for which CCN/CN = 0.5, where CN is the number of condensation nuclei measured by the CPC. The experimental set-up was calibrated with ammonium sulfate  $((\text{NH}_4)_2\text{SO}_4)$  aerosol. CCNC calibration information is provided in Supplement (Table S1).

A schematic of the experimental setup to measure particle mass and calculate particle effective density is shown in Fig. 1. Particles were selected by mass in the Kanomax APM, then sized by electrical mobility in a downstream DMA column, and the DMA exit stream was split between the condensation particle counter (CPC, TSI 3084) and an electrostatic TEM precipitator (FHNW TEM sampler). The APM-DMA setup was used for improved time resolution over the DMA-APM setup (McMurry et al., 2002; Khalizov et al., 2009; Xue et al., 2009). The system is described in more detail in Malloy et al. (2009). Here, the APM selected 3 different particle masses between the (spherical, unit density equivalent) 50–100 nm particle size range. TEM grids were used to collect three masses of particles relevant to CCN activation, corresponding to approximately 50 nm, 75 nm, and 100 nm particles (assuming unit density). Using the effective density analogue of Eq. (4), the fractal-like dimension was obtained by fitting the power function (Park et al., 2003; Xue et al., 2009; Nakao et al., 2011):

$$\rho_{\text{eff}} = C d_{\text{m}}^{D_{\text{f}}' - 3} \quad (5)$$

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where  $\rho_{\text{eff}}$  is the effective density of particles,  $C$  a constant,  $d_m$  is the mobility diameter of the particles, and  $D_f'$  the fractal-like dimension. During the beginning and the end of the experiment, the TEM precipitator was turned on. The APM was operated in a single mass selection until the charge across the TEM grid was 60 pA, indicating a sufficient capture of particles. After an experiment, the TEM grids were stored in a grid holder away from heat and light. The grids were analyzed using a Philips Tecnai 12 Transmission Electron Microscope.

The pictures of the particles obtained from the TEM were further analyzed using the freeware image analysis program ImageJ (<http://imagej.nih.gov/ij/>). The shape factor,  $\chi$ , is determined from mobility diameter data and a volume calculation of each particle. The mean volume of the agglomerates for each APM mass selection was calculated as follows (Park et al., 2004):

$$\bar{V} = \frac{\sum_i N_i \frac{\pi}{6} \bar{d}_p^3}{n} \quad (6)$$

where  $N$  is the number of primary particles in the agglomerate,  $d_p$  is the mean primary particle diameter, and  $n$  the number of agglomerates. By using two-dimensional projected images of three dimensional objects from the TEM, previous studies have demonstrated that the projected area can be used to calculate  $N$  or  $R_g$  in fractal agglomerates (e.g. Meakin et al., 1989; Koylu et al., 1995):

$$N = k_a \left( \frac{A_a}{A_p} \right)^\alpha \quad (7)$$

where  $A_a$  is the projected area of the agglomerate,  $A_p$  is the mean projected area of primary particles,  $k_a$  is a constant, and  $\alpha$  is a projected area exponent. Numerical simulations have been used to calculate the empirical factors of Eq. (7),  $k_a$  and  $\alpha$ . In this study, we use the values of Oh and Sorensen (1997) who found that  $\alpha = 1.19$  and  $k_a = 1.81$  well described agglomerates whose primary particles were allowed to overlap.

$A_a$ ,  $A_p$ , and  $d_p$  were determined through image analysis using ImageJ. Figure 2 shows how the three values were selected.

Once the mean volume of an aggregate at a certain mass was found with the above analysis, the diameter of the sphere with that volume was used as the volume-equivalent diameter for that particle. The mean peak mobility diameter measured from the SMPS during the collection of the TEM grid was used as the mobility diameter for application in Eq. (2) to calculate  $\chi$ .

## 4 Results and discussion

### 4.1 Morphology of freshly emitted biomass burning aerosol

The APM-SMPS-TEM system (Fig. 1) measured the freshly emitted properties of biomass burning aerosol. The TEM grids were then photographed and analyzed as illustrated in Fig. 2. The primary particle diameter was both calculated from the projected area of the primary particles and measured directly. If the directly measured and calculated diameters deviated greater than 50 %, then the calculated value from the projected area was used and is reported here.

To measure the area and diameter of primary particles, only particles clearly distinguishable were selected for analysis. This usually meant that particles on the ends of fractal chains were selected as opposed to those in the center of the agglomerates. Previous work on flame-generated agglomerates has shown that the primary particle diameters follow a normal distribution with a mean between 28.5 to 34.4 nm (Koylu and Faeth, 1992; Lee et al., 2001). Figure 3 shows the primary particle diameter distribution from particles at the three selected masses from the APM-SMPS. The total number of individual primary particles analyzed is 475. The data is fit with a normal Gaussian distribution whose mean lies at  $25.7 \pm 2.1$  nm. This mean projected primary particle diameter, while slightly smaller than other reported values, is within the 25 % standard deviation reported in the literature for primary particle size diameters (Koylu and Faeth,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1992). This value is smaller than China et al.'s (2013) projected primary particle diameters of 37–56 nm but a number of differences exist between these studies. Primarily, China et al. measured particles much larger than those examined here (aerodynamic diameters > 250 nm) through use of scanning electron microscopy (SEM). Their study was also an ambient study, conducted downstream of a fire plume. The China et al. fire likely had different fuel sources and different burning conditions than those presented here. The distribution here may also be shifted to the lower end from the abundance of 10–20 nm primary particles that appeared mostly in the 0.9 fg (50 nm) particles from the APM. Over 80 % of the primary particle diameters below 20 nm were from these particles. Over 80 % of the primary particles observed below 20 nm were from 0.9 fg agglomerates. However, primary particles below 20 nm only made up 40 % of counts for the total number of primary particles from the 0.9 fg agglomerates. When < 20 nm primary particles are omitted, the distribution of primary particle size shifts to a mean of 27 nm. When only the particles from the 0.9 fg agglomerates are considered, the distribution becomes a double-peaked normal distribution centered on 14 and 19 nm.

In this study, the mean primary particle projected area and diameter shown in Fig. 3 were used in the application of Eq. (7). The analysis was applied for each of the three selected agglomerate masses from the APM. The application of mean primary projected area introduces a slight overestimation for the 50 nm agglomerates and a slight underestimation for the 75 and 100 nm agglomerates. However, the percent change on the calculated volume as a whole is small. For example, the bi-modal 50 nm primary particle distribution, if half of the total agglomerate projected area is assumed to be contributed by one peak, the calculated volume is forced upward by ~ 35 %. For the 75 and 100 nm particles, the calculated volume is forced downward by ~ 7 %.

Table 1 shows the calculated application of Eqs. (6) and (7) to the TEM images obtained for the 50 nm, 75 nm, and 100 nm equivalents selected from the APM-SMPS system. Figure 4 graphically demonstrates the size dependence of the shape factor to the mobility diameter. The shape factor for these particles ranges from 1.06 to 1.42

# Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

indicating a 6 % to 42 % overestimation of the true volume of a particle when using the electrical mobility diameter.

## 4.2 Morphology of aged biomass burning aerosol

To measure the effect of atmospheric aging on biomass burning aerosol's morphology, the APM-SMPS system was used as described in Nakao et al. (2011). The TEM was not used in this section because saturation of the TEM grids required extensive time, longer than that necessary to observe changing morphology that can occur with aging.

Figure 4 shows the evolution of the fractal-like dimension ( $D_f'$ ) as a function of experimental time. In this analysis a fractal-like dimension of 3 indicates a spherical particle and  $D_f' < 3$  indicates a non-spherical particle. Calibration with ammonium sulfate, an inorganic crystalline solute and not quite spherical particle ( $D_f' = 2.7$ ), is included for comparison. The black line indicates the time the UV lights surrounding the chamber were turned on. The freshly emitted particles have a  $D_f'$  of  $2.15 \pm 0.05$  indicating a non-spherical particle, similar to the  $D_f'$  of 2.20 of Xue et al. (2009) for freshly emitted diesel particles. After 4 h of photochemical exposure, the  $D_f'$  rises nearly linearly to 2.60.

The  $D_f'$  analysis here is a bulk measurement of all the particles selected in the APM which increased from 50 nm to 250 nm equivalent sizes. While it should be theoretically possible to link the  $D_f'$  reported here to the dynamic shape factor,  $\chi$  (Kaspar, 1982; DeCarlo et al., 2004), we do not do so here because of the limitations an ensemble method (the APM-SMPS system) has when expanded to a size-dependent individual particle analysis.

The APM-SMPS data set agrees with the TEM analysis of the previous section – that freshly emitted biomass burning aerosol are not spherical. Because the TEM operates at near-vacuum pressures, there is the possibility that volatile material may evaporate from the fractal backbone. This would skew the perceived importance of the particle's morphology. However, this second confirmation of morphology from the APM-SMPS indicates that volatile evaporation in the TEM is not completely the reason for shape factors greater than one being reported.

### Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As  $D'_f$  increases with time, the aerosol becomes more spherical with photochemical exposure. This is likely due to a coating of secondary organic aerosol (SOA) forming on the primary emissions (Nakao et al., 2011). The SOA fills in the void space inside the fractal agglomerates. This aged fractal-like dimension is smaller than the aged biomass burning plume aerosol measured by China et al. (2013) but the  $D'_f$  reported here is not directly comparable to their  $D_f$ . In addition to the different derivations, China et al. (2013) focused on larger particles than those investigated here, on a shorter period of atmospheric aging (1–2 h), and during the smoldering phase of the burn. However, both of these results confirm, through two different methods, that applying the electrical mobility diameter as the volume equivalent diameter will result in an over-estimation of particle size and volume.

### 4.3 Applying a true volume equivalent diameter to the CCN analysis of biomass burning aerosol

The assumption that the electrical mobility diameter cubed is equal to the true volume of the particle will result in a skewed CCN prediction for aerosol from combustion sources. The previous work of Dusek et al. (2011) has investigated conceptually the importance of fractal-morphology in CCN analysis. We present experimental data for both the hygroscopicity and morphology of the particle and apply it to  $\kappa$ -Köhler analysis.

Figure 6 shows an example of the CCN to CN activation ratio as a function of mobility diameter for biomass burning at a supersaturation of 0.37 %. Both the raw electrical mobility activation diameters and the volume equivalent diameters are shown on the figure. The volume equivalent diameters are determined on a point-by-point basis by using a logarithmic regression of the data in Fig. 4 and Eq. (2). The shape factor is assumed to be equal to unity for particles smaller than 30 nm where the regression falls below one. The activation ratio of ammonium sulfate at the same supersaturation is included for comparison. Figure 6 shows the shape factor of a particle affects its perceived mobility diameter. Since  $\chi$  is greater than unity for particles larger than 30 nm, the activation curve shifts to smaller electrical mobility diameters. In this example, the



diameter where half of all particles activate ( $D_{p50}$ ) is measured at 75 nm electrical mobility but the true volume equivalent diameter for that particle is at 60 nm.

Figure 7 shows the total effect that accounting for the volume equivalent diameter can have on the calculation of  $\kappa$ . The data from a non-photochemically exposed chamise biomass burning injection into the chamber is presented. The markers in Fig. 7 are the activation diameters for each scan of the SMPS-CCNC at a supersaturation. The red line is the  $\kappa$  fit between the points as shown where the lines of constant  $\kappa$  are forced such that critical supersaturation scales with  $d_p^{-3/2}$ . The implications of this are discussed later. The black line is the fit of the points if the mobility diameter is converted to the volume equivalent diameter via the process described for Fig. 6. The hygroscopicity of manzanita biomass, without morphological corrections, has a  $\kappa$  of  $0.38 \pm 0.09$ . This value agrees with Engelhart et al.'s (2012) hygroscopicity measurements of primary (non-aged) chamise emissions. When the shape factors of the particles is taken into account using the TEM data collected concurrently, the calculated  $\kappa$  increases to 0.52. Since the particles are not spherical, their volume equivalent diameters are smaller than their mobility diameters. Therefore, using only the mobility diameter when calculating  $\kappa$  can substantially underestimate hygroscopicity.

This analysis of particle morphology may help reconcile the difference in calculated hygroscopicity between filter reconstituted biomass burning aerosol and in-situ sampled aerosol. Carrico et al. (2008) reported HTDMA and CCN derived  $\kappa$  values for biomass burning filter samples extracted in water and methanol. The CCN derived  $\kappa$  from Carrico et al.'s study of Alaskan Duff core aerosol extracted in water was 0.148. This value is significantly higher than the on-line sampling of Alaskan Duff core done by both Petters et al. (2009) and Engelhart et al. (2012) which reported a CCN derived  $\kappa$  value of 0.09 and 0.07, respectively. The 50 % increase in filter reconstituted hygroscopicity may be due to the use of mobility diameter as the volume equivalent diameter for non-spherical aerosols. In filter reconstitution, the liquid solution is first atomized and dried before flowing into the downstream instruments. The process of drying may fill in void space in a non-spherical backbone thus producing a particle whose volume

## Measured morphology of biomass burning CCN

M. Giordano et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



equivalent and electrical mobility diameters are identical. It is important to note that Carrico et al. (2008) reported a lower hygroscopicity value for reconstituted sagebrush than both Petters et al.'s (2009) and Engelhart et al.'s (2012) in-situ sampled aerosol. This discrepancy may be due to different burning or feedstock conditions that may influence particle shape factor. However, since reconstitution of aerosols can increase measured hygroscopicity, this also suggests that cloud processing of fractal aerosols should be considered for biomass burning aerosol's climate impacts.

#### 4.4 Sensitivity of $\kappa$ to particle shape factor

Because the shape factor of particles derived from biomass burning can range from 1 to greater than 1.5, the importance of particle morphology can be negligible to substantial. While the current literature on the morphology of biomass burning aerosols is sparse, the work done on diesel particles suggests that burn conditions may play a part in particle morphology (Park et al., 2004; Nakao et al., 2011). Along with the affects that particle aging have on morphology shown in this study, it is prudent to analyze the potential range of hygroscopicities that biomass burning aerosols can have.

For any non-spherical, fractal particle generated from combustion, the larger the activation diameter in a CCNC, the greater the difference between the mobility diameter and volume equivalent diameter will be. Figure 8 shows how sensitive  $\kappa$  can be when applying a volume equivalent diameter calculation. In Fig. 8, hygroscopicity data from biomass burning tests by Petters et al. (2009) is used. The grey shaded area is the sensitivity of black spruce ( $\kappa = 0.07$ ) when applying a dynamic shape factor of 1 (right-most boundary of the shaded area) and of 1.5 (left-most boundary of the shaded area). Here, black spruce ranges from a  $\kappa$  of 0.07 to 0.24. The blue shaded area is the same treatment for sage brush ( $\kappa = 0.33$  to 1.1 after applying  $\chi$ ). Douglas fir, oak, titi, and needlegrass, two relatively low-hygroscopic and two moderately hygroscopic biomass burning species, are included for comparison. In this analysis we apply a constant shape factor across all diameters as opposed to the mobility diameter-dependent treatment used earlier. This sensitivity analysis therefore shows the area that the true

$\kappa$  line falls in but not necessarily its slope. It is also a large range of  $\chi$  which captures the effects that burn conditions and the species itself may have on particle morphology. This range is indicative of the effects that photochemical aging can have on morphology.

Figure 9 shows the sensitivity of the slope of the  $\kappa$ -Köhler isolines (heretofore referred to as the experimental exponent or exponent) to two different methods of applying shape factor to CCN analysis. Shown in the figure are an application of a constant shape factor ( $\chi \neq f(d_{\text{dry}})$ ) and a variable shape factor ( $\chi = f(d_{\text{dry}})$ ). The data shown is data from the chamise experiments shown in Fig. 7. The constant application of  $\chi$  does not exhibit any change in the experimental exponent of the  $\kappa$  lines.  $S_c$  scales with  $d_d^{-3/2}$  as predicted by Köhler theory. This suggests that the solute dissolution process is thermodynamically ideal, the particle completely dissolves and surface tension effects are negligible. When a variable shape factor is applied,  $s_c$  then scales with  $d_d^{-1.8}$ . Kumar et al. (2011) observed a similar effect on the  $\kappa$  exponent in wet processed soil and dust samples. Their study observed an unusually low experimental exponent of  $-2.16$  for the wet processed mineral sample, Arizona Test Dust (ATD). This deviation from Köhler theory could be caused by sparingly soluble species, an externally mixed sample (composition as a function of particle size), or activation physics. The presence of black carbon (BC) in biomass burning may act as the sparingly soluble species for biomass burning generated aerosol. The presence of BC could explain the deviation from Köhler theory for the corrected data set from combustion aerosol and further investigation into this phenomenon is required. The complexity and non-ideal behavior of the aerosol solute can be masked if electrical mobility diameter is used in CCN analysis.

## 5 Conclusions

This work investigated the morphology and hygroscopicity of biomass burning aerosol. We show, through two independent measurements that particles emitted from biomass combustion are not spherical and that the degree of non-sphericity can change with

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



photochemical exposure. Thus, for biomass burning aerosol sampled near source with electrical mobility techniques, size and volume measurements are likely overestimated. As the ageing process occurs, SOA condenses on the fractal particle and the volume changes (measured through electrical mobility) are small compared to changes in particle mass. We also measure the dynamic shape factor of freshly emitted particles and apply it to  $\kappa$ -Köhler theory to ensure a mobility diameter derived volume. We find that particle morphology is a function of a particle's electrical mobility diameter, which is consistent with literature, and a particle's dynamic shape factor can range from 1.06 to 1.42 for electrical mobility equivalent diameters of 50 to 100 nm. These mobility diameters are relevant for CCN activation and therefore represent an underestimation of  $\kappa$  since the volume of particle that is being activated is smaller than assumed. Alone, this suggests an underestimation of up to almost 50 %. The uncertainty induced by excluding particle morphology in CCN data sets is consistent with the range of biomass burning  $\kappa$  values presented in literature data. The issue is further complicated by changes in particle morphology that occur with photochemical aging and time. Biomass burning derived aerosols experience a decrease in hygroscopicity with aging. The results presented here suggest that morphological changes as well as chemical compositional changes may contribute to that phenomenon. A number of questions still remain to be investigated about the true nature of biomass hygroscopicity. Since shape factor,  $\chi$ , is a function of particle size, in practice this means that the  $\kappa$ -isolines do not follow those of an ideal solution, where critical supersaturation scales with  $d_d^{-3/2}$ . Scaling with ideal theory suggests that critical solubility and surface tension limits are negligible and thus non-ideal effects are partially hidden from electrical mobility-CCNC observation. The corrected data suggests that  $s_c$  scales with  $d_d^{-1.8}$  and does not follow ideal thermodynamic behavior. Non-ideal behavior is observed because the process of chemical transformation (secondary aerosol formation) occurs concurrently with physical rounding (decrease in the fractal nature) of the particles. The importance of timescales for chemical vs. physical evolution of ambient biomass burning aerosols needs to be further explored.

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Acknowledgements.* Many thanks to David Weise and Joey Chong of the USDA Forest Service for their advice and pivotal role in the acquisition of fuels. The authors would also like to thank David Cocker and the Keck Foundation for use of the environmental chamber. MG would also like to thank Bibiana Lopez for her work supporting this paper. MG would like to thank California NASA Space Grant Consortium for partially supporting this work. MG and AA would also like to thank the US Environmental Protection Agency for funding this work made possible by EPA grant number 83504001. Any opinions, findings, and conclusions expressed in this material are those of the authors and do not necessarily reflect the views of NSF, NASA, or EPA.

## References

- Andreae, M. O., Rosenfeld, D., Artaxo, P., Costa, A. A., Frank, G. P., Longo, K. M., and Silva-Dias, M. A. F.: Smoking rain clouds over the Amazon, *Science*, 303, 1337–1342, doi:10.1126/science.1092779, 2004.
- Baron, P. A., Sorensen, C. M., and Brockmann, J. E.: Nonspherical particle measurements: shape factors, fractals, and fibers, in: *Aerosol Measurement: Principles, Techniques, and Applications*, edited by: Baron, P. A. and Willeke, K., John Wiley, New York, 705–749, 2001.
- Carrico, C. M., Petters, M. D., Kreidenweis, S. M., Collett Jr., J. L., Engling, G., and Malm, W. C.: Aerosol hygroscopicity and cloud droplet activation of extracts of filters from biomass burning experiments, *J. Geophys. Res.*, 113, D08206, doi:10.1029/2007JD009274, 2008.
- China, S., Mazzoleni, C., Gorkowski, K., Aiken, A. C., and Dubey, M. K.: Morphology and mixing state of individual freshly emitted wildfire carbonaceous particles, *Nature Communications*, 4, 2122, doi:10.1038/ncomms3122, 2013.
- Crutzen, P. J. and Andreae, M. O.: Biomass burning in the tropics: impact on atmospheric chemistry and biogeochemical cycles, *Science*, 250, 1669–1678, doi:10.1126/science.250.4988.1669, 1990.

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Dahneke, B.: Slip correction factors for nonspherical bodies – I introduction and continuum flow, *J. Aerosol Sci.*, 4, 139–145, 1973a.
- Dahneke, B.: Slip correction factors for nonspherical bodies – II free molecule flow, *J. Aerosol Sci.*, 4, 147–161, 1973b.
- 5 Dusek, U., Frank, G. P., Massling, A., Zeromskiene, K., Iinuma, Y., Schmid, O., Helas, G., Hennig, T., Wiedensohler, A., and Andreae, M. O.: Water uptake by biomass burning aerosol at sub- and supersaturated conditions: closure studies and implications for the role of organics, *Atmos. Chem. Phys.*, 11, 9519–9532, doi:10.5194/acp-11-9519-2011, 2011.
- Engelhart, G. J., Hennigan, C. J., Miracolo, M. A., Robinson, A. L., and Pandis, S. N.: Cloud  
10 condensation nuclei activity of fresh primary and aged biomass burning aerosol, *Atmos. Chem. Phys.*, 12, 7285–7293, doi:10.5194/acp-12-7285-2012, 2012.
- Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in atmospheric constituents and in radiative forcing, in: *Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and New York, NY, USA, 131–215, 2007.
- 15 Fuchs, N. A.: *The Mechanics of Aerosols*, Pergamon, New York, 1964.
- Giordano, M. R., Short, D. Z., Hosseini, S., Lichtenberg, W., and Asa-Awuku, A.: Changes in droplet surface tension affect the observed hygroscopicity of photochemically aged biomass burning aerosol, *Environ. Sci. Technol.*, 47, 10980–10986, doi:10.1021/es401867j, 2013.
- Haywood, J. and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: a review, *Rev. Geophys.*, 38, 513–543, 2000.
- 25 Hennigan, C. J., Miracolo, M. A., Engelhart, G. J., May, A. A., Presto, A. A., Lee, T., Sullivan, A. P., McMeeking, G. R., Coe, H., Wold, C. E., Hao, W.-M., Gilman, J. B., Kuster, W. C., de Gouw, J., Schichtel, B. A., J. L. Collett Jr., Kreidenweis, S. M., and Robinson, A. L.: Chemical and physical transformations of organic aerosol from the photo-oxidation of open biomass burning emissions in an environmental chamber, *Atmos. Chem. Phys.*, 11, 7669–7686, doi:10.5194/acp-11-7669-2011, 2011.
- 30 Kaspar, G.: Dynamics and measurement of smokes, I. Size characterization of non-spherical particles, *Aerosol Sci. Tech.*, 1, 187–199, 1982.

# Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Khalizov, A. F., Zhang, R., Zhang, D., Xue, H., Pagels, J., and McMurry, P. H.: Formation of highly hygroscopic soot aerosols upon internal mixing with sulfuric acid vapor, *J. Geophys. Res.*, 114, D05208, doi:10.1029/2008JD010595, 2009.
- Knutson, E. O. and Whitby, K. T.: Aerosol classification by electric mobility: apparatus, theory, and applications, *J. Aerosol Sci.*, 6, 443–451, doi:10.1016/0021-8502(75)90060-9, 1975.
- Kohler, H.: The nucleus in and the growth of hygroscopic droplets, *Trans. Farad. Soc.*, 32, 1152–1161, 1936.
- Kousaka, Y., Endo, Y., Ichitsubo, H., and Alonso, M.: Orientation-specific dynamic shape factors for doublets and triplets of spheres in the transition regime, *Aerosol Sci. Tech.*, 24, 36–44, 1996.
- Koylu, U. O. and Faeth, G. M.: Structure of overfire soot in buoyant turbulent diffusion flames at long residence times, *Comb. Flame*, 89, 140–156, 1992.
- Koylu, U. O., Faeth, G. M., Farias, T. L., and Carvalho, M. G.: Fractal and projected structure properties of soot aggregates, *Comb. Flame*, 100, 621–633, 1995.
- Kumar, P., Sokolik, I. N., and Nenes, A.: Cloud condensation nuclei activity and droplet activation kinetics of wet processed regional dust samples and minerals, *Atmos. Chem. Phys.*, 11, 8661–8676, doi:10.5194/acp-11-8661-2011, 2011.
- Kuwata, M. and Kondo, Y.: Measurements of particle masses of inorganic salt particles for calibration of cloud condensation nuclei counters, *Atmos. Chem. Phys.*, 9, 5921–5932, doi:10.5194/acp-9-5921-2009, 2009.
- Latham, T. L., Beyersdorf, A. J., Thornhill, K. L., Winstead, E. L., Cubison, M. J., Hecobian, A., Jimenez, J. L., Weber, R. J., Anderson, B. E., and Nenes, A.: Analysis of CCN activity of Arctic aerosol and Canadian biomass burning during summer 2008, *Atmos. Chem. Phys.*, 13, 2735–2756, doi:10.5194/acp-13-2735-2013, 2013.
- Lee, K. O., Cole, R., Sekar, R., Choi, M. Y., Zhu, J., Kang, J., and Bae, C.: Detailed Characterization of Morphology and Dimensions of Diesel Particulates via Thermophoretic Sampling, *SAE Technical Paper Series 2001-01-3572*, available at: <http://papers.sae.org/2001-01-3572/> (last access: 10 January 2014), 2001.
- Liu, X. and Wang, J.: How important is organic aerosol hygroscopicity to aerosol indirect forcing?, *Environ. Res. Lett.*, 5, 044010, doi:10.1088/1748-9326/5/4/044010, 2010.
- Malloy, Q. G. J., Nakao, S., Qi, L., Austin, R., Stothers, C., Hagino, H., and Cocker, D. R. III: Real-time aerosol density determination utilizing a modified scanning mobility par-

ticle sizer – aerosol particle mass analyzer system, *Aerosol Sci. Tech.*, 43, 673–678, doi:10.1080/02786820902832960, 2009.

Martins, J. A., Gonçalves, F. L. T., Morales, C. A., Fisch, G. F., Pinheiro, F. G. M., Júnior, J. B. V. L., and Dias, M. A. F. S.: Cloud condensation nuclei from biomass burning during the Amazonian dry-to-wet transition season, *Meteorol. Atmos. Phys.*, 104, 83–93, 2009.

McMurry, P. H., Wang, X., Park, K., and Ebara, K.: The relationship between mass and mobility for atmospheric particles: a new technique for measuring particle density, *Aerosol Sci. Tech.*, 36, 227–238, 2002.

Meakin, P., Donn, B., and Mulholland, G. W.: Collisions between point masses and fractal aggregates, *Langmuir*, 5, 510–518, 1989.

Moore, R. H., Nenes, A., and Medina, J.: Scanning mobility CCN analysis – a method for fast measurements of size-resolved CCN distributions and activation kinetics, *Aerosol Sci. Tech.*, 44, 861–871, doi:10.1080/02786826.2010.498715, 2010.

Nakao, S., Shrivastava, M., Nguyen, A., Jung, H., and Cocker III, D.: Interpretation of secondary organic aerosol formation from diesel exhaust photooxidation in an environmental chamber, *Aerosol Sci. Tech.*, 45, 8, 964–972, doi:10.1080/02786826.2011.573510, 2011.

Novakov, T. and Corrigan, C. E.: Cloud condensation nucleus activity of the organic component of biomass smoke particles, *Geophys. Res. Lett.*, 23, 2141–2144, 1996.

Oh, C. and Sorensen, C. M.: The Effect of overlap between monomers on the determination of fractal cluster morphology, *J. Colloid Interface Sci.*, 193, 17–25, 1997.

Park, K., Cao, F., Kittelson, D. B., and McMurry, P. H.: Relationship between particle mass and mobility for diesel exhaust particles, *Environ. Sci. Technol.*, 37, 577–583, 2003.

Park, K., Kittelson, D. B., and McMurry, P. H.: Structural properties of diesel exhaust particles measured by transmission electron microscopy (TEM): relationships to particle mass and mobility, *Aerosol Sci. Tech.*, 38, 881–889, doi:10.1080/027868290505189, 2004.

Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.

Petters, M. D., Carrico, C. M., Kreidenweis, S. M., Prenni, A. J., DeMott, P. J., Collett, J. L., and Moosmüller, H.: Cloud condensation nucleation activity of biomass burning aerosol, *J. Geophys. Res.*, 114, D22205, doi:10.1029/2009JD012353, 2009.

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Pöschl, U.: Atmospheric aerosols: composition, transformation, climate and health effects, *Angew. Chem. Int. Edit.*, 44, 7520–7540, 2005.
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124, 2001.
- 5 Raymond, T. M. and Pandis, S. N.: Cloud activation of single-component organic aerosol particles, *J. Geophys. Res.*, 107, 4787, doi:10.1029/2002JD002159, 2002.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmos. Chem. Phys.*, 5, 799–825, doi:10.5194/acp-5-799-2005, 2005.
- 10 Reutter, P., Su, H., Trentmann, J., Simmel, M., Rose, D., Gunthe, S. S., Wernli, H., Andreae, M. O., and Pöschl, U.: Aerosol- and updraft-limited regimes of cloud droplet formation: influence of particle number, size and hygroscopicity on the activation of cloud condensation nuclei (CCN), *Atmos. Chem. Phys.*, 9, 7067–7080, doi:10.5194/acp-9-7067-2009, 2009.
- Roberts, G. C. and Nenes, A.: A continuous-flow streamwise thermal-gradient
- 15 CCN chamber for atmospheric measurements, *Aerosol Sci. Tech.*, 39, 206–221, doi:10.1080/027868290913988, 2005.
- Rose, D., Gunthe, S. S., Mikhailov, E., Frank, G. P., Dusek, U., Andreae, M. O., and Pöschl, U.: Calibration and measurement uncertainties of a continuous-flow cloud condensation nuclei counter (DMT-CCNC): CCN activation of ammonium sulfate and sodium chloride aerosol
- 20 particles in theory and experiment, *Atmos. Chem. Phys.*, 8, 1153–1179, doi:10.5194/acp-8-1153-2008, 2008.
- Rose, D., Nowak, A., Achtert, P., Wiedensohler, A., Hu, M., Shao, M., Zhang, Y., Andreae, M. O., and Pöschl, U.: Cloud condensation nuclei in polluted air and biomass burning smoke near the mega-city Guangzhou, China – Part 1: Size-resolved measurements and implications for the modeling of aerosol particle hygroscopicity and CCN activity, *Atmos. Chem. Phys.*, 10, 3365–3383, doi:10.5194/acp-10-3365-2010, 2010.
- 25 Rose, D., Gunthe, S. S., Su, H., Garland, R. M., Yang, H., Berghof, M., Cheng, Y. F., Wehner, B., Achtert, P., Nowak, A., Wiedensohler, A., Takegawa, N., Kondo, Y., Hu, M., Zhang, Y., Andreae, M. O., and Pöschl, U.: Cloud condensation nuclei in polluted air and biomass burning smoke near the mega-city Guangzhou, China – Part 2: Size-resolved aerosol chemical
- 30 composition, diurnal cycles, and externally mixed weakly CCN-active soot particles, *Atmos. Chem. Phys.*, 11, 2817–2836, doi:10.5194/acp-11-2817-2011, 2011.



**Measured  
morphology of  
biomass burning  
CCN**

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schmidt-Ott, A., Baltensperger, U., Gaggeler, H. W., and Jost, D. T.: Scaling behavior of physical parameters describing agglomerates, *J. Aerosol Sci.*, 21, 711–717, 1990.

Seaton, A., Godden, D., MacNee, W., and Donaldson, K.: Particulate air pollution and acute health effects, *Lancet*, 345, 176–178, 1995.

- 5 Shulman, M. L., Jacobson, M. C., Carlson, R. J., Synovec, R. E., and Young, T. E.: Dissolution behavior and surface tension effects of organic compounds in nucleating cloud droplets, *Geophys. Res. Lett.*, 23, 277–280, 1996.

- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arel-  
lano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004,  
10 *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.


Ward, D. E. and Hao, W. M.: Projections of emissions from burning of biomass for use in studies of global climate and atmospheric chemistry, in: *Proceedings of the 84th Annual Meeting and Exhibition Air & Waste Management Association*, Volume 19, 16–21 June 1991, Vancouver, British Colombia, Canada, 1991.

- 15 Xue, H., Khalizov, A. F., Wang, L., Zheng, J., and Zhang, R.: Effects of coating of dicarboxylic acids on the mass-mobility relationship of soot particles, *Environ. Sci. Technol.*, 43, 2787–2792, 2009.


- Yokelson, R. J., Crounse, J. D., DeCarlo, P. F., Karl, T., Urbanski, S., Atlas, E., Campos, T., Shinozuka, Y., Kapustin, V., Clarke, A. D., Weinheimer, A., Knapp, D. J., Montzka, D. D.,  
20 Holloway, J., Weibring, P., Flocke, F., Zheng, W., Toohey, D., Wennberg, P. O., Wiedinmyer, C., Mauldin, L., Fried, A., Richter, D., Walega, J., Jimenez, J. L., Adachi, K., Buseck, P. R., Hall, S. R., and Shetter, R.: Emissions from biomass burning in the Yucatan, *Atmos. Chem. Phys.*, 9, 5785–5812, doi:10.5194/acp-9-5785-2009, 2009.

- 25 Zhang, R., Khalizov, A. F., Pagels, J., Zhang, D., Xue, H., and McMurry, P. H.: Variability in morphology, hygroscopicity, and optical properties of soot aerosols during atmospheric processing, *P. Natl. Acad. Sci. USA*, 105, 10291–10296, doi:10.1073/pnas.0804860105, 2008.







**Fig. 1.** Schematic of the APM-DMA-TEM setup.



**Fig. 2.** The three measured parameters from the TEM pictures used in this study: the primary particle projected area ( $A_p$ ), primary particle diameter ( $d_p$ ), and the projected area of the particle ( $A_a$ ).



**Fig. 3.** TEM analysis of primary particle diameters of agglomerates generated from biomass burning of chamise. Data is fit with a normal Gaussian distribution. The mean of the distribution is  $25.7 \pm 2.1$  nm.



**Fig. 4.** Shape factor ( $\chi$ ) vs. the APM selected mobility diameter equivalent for freshly emitted biomass burning aerosol.

## Measured morphology of biomass burning CCN

M. Giordano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Fig. 5.** APM-SMPS derived fractal-like dimension data vs. time of experiment. A fractal-like dimension of 3 indicates a spherical particle while  $D_f < 3$  is non-spherical. Black line indicates time of UV lights being turned on. Ammonium sulfate calibration data is included for comparison.




**Fig. 6.** Activation curves for manzanita using the mobility diameter and the volume equivalent diameter calculated using the shape factor data from Fig. 4. The activation curve of ammonium sulfate is included for comparison. The black squares represent the raw electrical mobility activation diameters and the red crosses are the volume equivalent diameters.






**Fig. 7.** Supersaturation vs. the dry activation diameter for one chamise experiment. Markers indicate the data from the SMPS-CCNC and the red line is the best-fit  $\kappa$  between the points. The black line indicates the  $\kappa$  when the volume equivalent diameter is used instead of the mobility diameter.



**Fig. 8.** Sensitivity analysis of  $\kappa$  to  $\chi$ .  $\kappa$  data for each biomass burning species is taken from Petters et al. (2009). The sensitivity of  $\kappa$  to  $\chi$  is analyzed using two constant dynamic shape factors of 1 to 1.5. The region between the two  $\chi$  values is shaded in grey for black spruce and blue for sage brush. The  $\kappa$  lines for Douglas fir, oak, titi, and rush are shown for comparison.



**Fig. 9.** Variation of  $\kappa$ -Köhler isolines when applying a constant or no shape factor and applying a shape factor as a function of particle size. The blue triangles have a critical supersaturation that scales with  $d_d^{-3/2}$  due to a constant shape factor being applied while the red circles scale with  $d_d^{-1.8}$  due to the  $\chi - d_{\text{mobility}}$  relationship from Fig. 4 being applied.