

Discussions for “Cirrus and water vapour transport
in the tropical tropopause layer – Part 2: Roles of
ice nucleation and sedimentation, cloud dynamics,
and moisture conditions” – Reviewer A

Tra Dinh, Stephan Fueglistaler, Dale Durran, Thomas Ackerman

We would like to thank the reviewer for the encouraging comments for the manuscript. Please find below our responses to the reviewer’s minor revision requests. We agree with all the reviewer’s comments and have revised the manuscript accordingly.

1. **Reviewer** — Suggest adding either “cold cirrus clouds” or “Tropical Tropopause Layer” to the title because these simulations/conclusions do not necessarily hold true for warm cirrus clouds.

Authors — The title has been changed to “Cirrus and water vapour transport in the tropical tropopause layer – Part 2: Roles of ice nucleation and sedimentation, cloud dynamics, and moisture conditions.” Please see also our responses to Reviewer B for further explanation of this change in title of the manuscript.

2. **Reviewer** — P. 13304, Line 19: intepretations → interpretations.

Authors — Thanks for spotting this spelling mistake. We have fixed it accordingly.

3. **Reviewer** — P. 13311, Lines 7-10 (Rather, of interest ...): What observations are you referring to in this sentence? Are there any observations to support your model simulations and conclusions? Although this is a theoretical study, quite a bit of observational work has been done to study these TTL processes. Can you link your results to these studies?

Authors — The reference to observations here is indeed confusing. We have deleted the second half of the sentence so that it now reads: “Rather, of interest here is the question to what extent the strongly simplified inf-sed calculation captures the dehydration induced by the cloud.”

Please refer to our response to Comment 1 of Reviewer B for discussions comparing between model simulations and observations.

4. **Reviewer** — Figure 3: The q_v is averaged over the domain (correct?). What is the average cloud base/top height relative to the location of the hydrated/dehydrated layer for each scenario? In the text you mention that air passes through the cloud base is hydrated and the air that passes through the top is dehydrated. But in Fig. 3b for the all-phys dry case, the layer is hydrated at the top of the domain. I would be interested to see where the cloud top is relative to these simulations.

Authors — The difference between the Eulerian and Lagrangian interpretations is quite interesting (but may be confusing). Figure 4 (originally Fig. 3) is indeed averaged over the domain (the Eulerian sense), and the change in water vapour here (Δq_v) includes the advective tendencies. Conversely, the advective tendencies do not contribute to the change (δq_v) that the air parcels experience (because the parcels move with the air flow). Hence, we may have situations when, in the Eulerian sense hydration occurs ($\Delta q_v > 0$), but in the Lagrangian sense dehydration occurs ($\delta q_v < 0$).

In Figure 5 of the revised manuscript, we show the domain average change in water vapour due to the source/sink terms associated with ice growth/sublimation only (i.e. the advective tendencies are excluded). Figure 5 shows that the domain average ice-to-vapour conversion is consistent with the change in water vapour along air parcels (the Lagrangian interpretation), with dehydration occurs always in the upper half of the cloud layer.

We have added the following explanation for the differences between Fig. 4 and Fig. 5: “To separate the advective tendencies from the impacts on the moisture profiles of microphysical processes, we compute the accumulative mass of water associated with ice-to-vapour exchange during the model integration (Fig. 5). The exchange mass between vapour and condensates is recorded at the time and location when/where (de)hydration occurs (not at the end of the model integration). With the advective tendencies excluded, Fig. 5 shows that microphysical processes consistently lead to dehydration in the upper half of the cloud layer (in contrast to Fig. 4b where the advective

tendencies result in hydration above 17.1 km in the all-phys simulation for the dry scenario).”

The cloud moves about both vertically and horizontally during the simulations. Please see the supplemental animations for illustration. We have added Fig. 6 which shows the location of the cloud at 3.5 d.

5. **Reviewer** — Figure 4: Is the conversion from ice to vapor or vapor to ice? Please clarify in the figure caption.

Authors — The conversion is from ice to vapour. The caption (now Fig. 5) is revised as: “The profiles of accumulative mass exchange from ice to vapour over the model integration.”

6. **Reviewer** — P. 13316, Lines 10–11 (“Of more interest ...”): I don’t think I understand how you drew this conclusion. From your discussion, I thought that the inf-sed scenario did not produce much nucleation. Can you refer to the figure(s) to help make this statement more clear?

Authors — This paragraph is indeed not clear. We rewrote it as: “The left panels in Fig. 8 compare the changes in specific humidity of air parcels in the inf-sed runs versus those in the all-phys runs for the dry (top panel) and moist (bottom panel) scenarios. The joint histograms (Fig. 8a and Fig. 8c) show a large number of air parcels for which the specific humidity is essentially unchanged in the inf-sed runs, but significant dehydration or hydration occurs in the all-phys runs. In addition, for dehydrated air parcels, δq_v is more negative in the all-phys than inf-sed calculations in both the dry and moist scenarios (the data for $\delta q_v < 0$ lie above the one-to-one line in Fig. 8a and Fig. 8c). In other words, dehydrated parcels dehydrate more in the all-phys calculation. In the inf-sed runs, the immediate fallout of ice crystals limits dehydration to the saturation mixing ratio q_s at the nucleation time. Further dehydration does not occur in the same air parcels because the temperature never drops sufficiently low to bring the relative humidity above the nucleation threshold again. In the all-phys runs, dehydration continues after nucleation due to growth of ice crystals, and dehydration up to q_s at the minimum temperature of the Kelvin wave passage can be obtained.”

Please also note that we have rewritten a large part of Sect. 3.3, which is hopefully clearer and easier to read now.

7. **Reviewer** — P. 13316, Line 20: Figure 6c refers to the moist scenario. I

think you mean Fig 6b (not 6c).

Authors — Thanks! It is panel b indeed (the original Fig. 6 is now Fig. 8).

8. **Reviewer** — Figure 6: The scale is not defined for the histogram color bar, but I imagine it is number or counts. Please define in the caption.

Authors — We added: “The colour bar shows the number of air parcel counts.”

Discussions for “Cirrus and water vapour transport in the tropical tropopause layer – Part 2: Roles of ice nucleation and sedimentation, cloud dynamics, and moisture conditions” – Reviewer B

Tra Dinh, Stephan Fueglistaler, Dale Durran, Thomas Ackerman

We would like to thank the reviewer for the thoughtful comments that have helped us revise and improve the manuscript. Accordingly, additional descriptions for model configurations, ice number concentrations and in-cloud relative humidities have been added to the revised manuscript. Please see the highlighted text and the new Fig. 3 in the revised manuscript for the changes.

We have changed the title of the manuscript (now “Cirrus and water vapour transport in the tropical tropopause layer – Part 2: Roles of ice nucleation and sedimentation, cloud dynamics, and moisture conditions”). This work emerged from Dinh et al. (2012), i.e. Part 1, but it took us longer than expected to publish it. Hence, we originally did not call it Part 2, but in light of the reviewer’s comments that wish for more thorough discussions of model setups and several important results that have been addressed in Dinh et al. (2012), we reconsidered our position, and label it as Part 2. References in the text to Dinh et al. (2012) are provided, including:

- “This work follows up on a case modelling study by Dinh et al. (2012), who simulated a TTL cirrus cloud with characteristics similar to observations.”
- “The simulations build on previous model development by Dinh et al. (2012), to which the readers may wish to refer for technical details omitted here.”
- “The base-state (unperturbed) temperature profile $\tilde{T}(z)$ is taken from a sounding typical for the tropics (see Dinh et al., 2012, Fig. 1).”

- “The 2D profiles of q_v and RH_i for the dry scenario are illustrated in Dinh et al. (2012, their Fig. 4).”
- “These initial conditions, in combination with the Kelvin wave forcing, produce thin cirrus clouds around the tropopause that are both similar to observations (see Dinh et al. (2012) for detailed comparison between model simulations and observations) and confined within the model domain for the entire duration of simulation.”
- “The cloud evolution in the 2D domain in the all-phys simulation for the dry scenario is illustrated in the supplemental animation of Dinh et al. (2012).”

Below please find our detailed responses to the reviewer’ comments.

1. **Reviewer** — The entry of water vapour into the stratosphere or equivalently the issue of dehydration is closely related to the microphysical issue of low ice number concentrations in the TTL. Low concentrations would lead to less effective relaxation to equilibrium of the clouds; thus, high water vapour concentrations will remain and may be transported into the stratosphere. This correlation should be emphasised in the study in more details. Actually, ice crystal number concentrations and relative humidities obtained during the long-term simulation should be evaluated. The number concentrations should be compared with measurements as reported by Krämer et al. (2009) and Jensen et al. (2010). The simulations should mostly produce low concentrations, since the large-scale forcing is quite small. Thus, I would expect that the cloud should contain (high) ice supersaturation; in terms of the analysis of the upward water vapour transport, this might be an issue since for downward moisture transport during cloud lifting, less supersaturation outside the cloud is required. Thus, the distribution of relative humidity inside/outside clouds should be checked carefully.

Authors — Detailed comparison of cloud characteristics (cloud lifetime, ice number concentration, ice water content, ice crystal size distribution, cloud radiative heating, ...) between model simulations and observations (by Krämer et al. (2009); Jensen et al. (2010) as well as many others) have been provided in Dinh et al. (2012, Sects. 4.2 and 4.3). In the revised manuscript, we mentioned that the readers may find these comparisons in Dinh et al. (2012). The comparison indicates that the characteristics of the simulated clouds agree with observations.

We have shown the ice number concentration in Fig. 1b. In addition, the analysis for in-cloud relative humidities is now included and presented in Fig. 3. Please see the last paragraph of Sect. 3.1 for the relevant discussion. The results indicate that, on average over the cloudy area, the ice number concentration generated in the model is sufficiently high (but is within observations) such that the distribution of in-cloud relative humidities maximises at 100 %; that is, the largest fraction of the cloudy air is maintained close to saturation throughout the cloud lifetime. There is a realistic spread of RH_i to values above/below 100 % in regions subject to mixing with environmental air and/or where the ice number concentration is low.

2. **Reviewer** — Although the model might be explained in details in former publications, please repeat the key details of the microphysics parameterisation and the dynamics (e.g. underlying equations). The resolution of the model seems to be quite unbalanced, the horizontal grid spacing is quite large compared to the vertical spacing. The size distribution seems to be quite narrow, I would expect ice crystals larger than $50\mu\text{m}$ in the TTL. The radiation parameterisation is not clear to me. Do you use an explicit radiation transfer model for calculating heating rates of ice crystals online? Please clarify these issues.

Authors — The equations describing the dynamical core of the model are quite complex and have been provided in Dinh et al. (2012, their Sects. 3 and 4). Please see the descriptions for the microphysical processes in Sect. 2.2 of the revised manuscript. The radiation scheme is mentioned as in: “Additionally, we compute the cloud radiative heating that results from the absorption of radiation by ice crystals using the radiative transfer scheme described in Durran et al. (2009, their Sect. 3a).”

The spatial resolution has been chosen specifically for TTL cirrus, which are very thin (~ 1 km in thickness) but very wide (hundreds up to a few thousand kilometres in width). Our model calculations (not shown) indicate that the results are more sensitive to the vertical resolution than the horizontal resolution. Also, Jensen et al. (2010) show that the relative humidities and cloud properties observed in TTL cirrus change sharply in very thin layers (of a few metres). In addition, Murphy (2014) suggests that high vertical resolution (again, on order of a few metres) is required to accurately compute ice sedimentation for TTL cirrus.

Please see Dinh et al. (2012, Fig. 7), where we showed that the size distribution of ice crystals is consistent between model and observations.

3. **Reviewer** — The setup of the scenarios is not clear at all. Vertical profiles of relative humidity and temperature would help; it is not clear how the humidity profiles look like for the different cases (dry versus moist). In addition, the stratification is of high interest, since during the simulation the stable layers are destabilised by radiation feedback. As remarked above, it would be interesting if a change in stratification would crucially change the results (i.e. the moisture transport). Please be aware of that the absolute values of the RH_i profile in Jensen et al. (2005) are much too high and can be explained by measurement errors (see recent comparison and evaluation by Fahey et al. (2014)).

Authors — The vertical profile of temperature can be found in Dinh et al. (2012): “The base-state (unperturbed) temperature profile $\bar{T}(z)$ is taken from a sounding typical for the tropics (see Dinh et al., 2012, Fig. 1).” We also added that “the 2D profiles of q_v and RH_i for the dry scenario are illustrated in Dinh et al. (2012, their Fig. 4).”

The cloud radiative heating results in (i) a cloud-scale circulation and (ii) small-scale convection in the destabilized layer at the cloud top. Both of these have impacts on the cloud evolution and the water vapour transport. We have not specifically separated the impacts of the cloud-scale circulation from that of convection, and think the discussion provided is sufficient for this paper, but agree that in a next step this question could be addressed.

We added a footnote: “Note that the recent comprehensive instrumentation evaluation by Fahey et al. (2014) suggests that measurement errors may have caused overestimation in the relative humidity reported in Jensen et al. (2005).”

4. **Reviewer** — The cloud evolution (as shown in Fig. 2) should be explained in more details, since this is the major result leading to differences between dry and moist profiles. For instance, you have to explain why the ice crystals become larger in the moist scenario. Which processes lead to larger crystals, is it just enhanced diffusional growth or do other processes play a role? 2D plots of relative humidity and ice mass/number concentrations might also help for clarification.

Authors — The reason for larger ice crystals in the moist scenario has been provided in the manuscript, and as below:

“The radiatively induced horizontal motions widen the cloud tops and narrow the cloud bases. As the clouds deform into trapezoidal shapes (Fig. 2), ice

crystals fall into initially clear air at the (tilted) lateral sides of the clouds (and also at the bases of the clouds). At the sides of the clouds in regions subject to horizontal inflow of environmental air, ice crystals grow to larger sizes in the moist scenario (where the ambient air is supersaturated) than in the dry scenario (where the ambient air is subsaturated)."

Please see Dinh et al. (2012, Fig. 4) for the 2D profile of RH_i .

5. **Reviewer** — It seems that the forcing of the whole scenario is exclusively given by large-scale Kelvin waves. Although this might be appropriate for an idealised investigation, the role of gravity waves in the TTL and their impact on cirrus clouds should not be neglected. Former studies addressed this issue from a microphysical point of view (e.g. Spichtinger and Krämer (2013); Jensen and Pfister (2004) and it should be mentioned at least.

Authors — The cloud radiative heating generates gravity waves, and these are resolved in the all-phys simulations. We have added the following discussions to Sect. 2.2 of the manuscript: "As further discussed in Sect. 3.1, the cloud radiative heating induces (i) a cloud-scale circulation (itself a gravity wave signal) and (ii) small-scale convective cells at the cloud top, whose buoyancy forces generate small-scale gravity waves which propagate vertically outwards from the convective layer (see also Dinh et al. (2010)). The all-phys simulations fully resolve the perturbations (in both temperature and velocities) of these gravity waves generated by the cloud radiative heating. The importance of gravity waves to the microphysical processes in TTL cirrus has also been suggested by Jensen and Pfister (2004); Spichtinger and Krämer (2013)." We hope to address the roles of gravity waves more thoroughly in a follow-up paper by adding gravity waves generated by other sources outside clouds.

6. **Reviewer** — Figure 6 is very hard to understand; maybe you could try to make a simple version of it to explain the main features in a simpler way.

Authors — We have not come up with a simpler way to present this information. However, we have rewritten a large part of Sect. 3.3, which is hopefully clearer now. Please also see our responses to Comments 6 and 8 of Reviewer A.

7. **Reviewer** — The supplement is not very user-friendly; it would be much better to upload the short movies in a common format instead of embedding

them into a pdf file. Please change the format of the supplement accordingly.

Authors — The movies will be also provided in GIF format. Also, we have used a frame at $t = 3.5$ d of the movies as Fig. 6 of the revised manuscript.

References

- Dinh, T., D. R. Durran, and T. Ackerman, 2010: Maintenance of tropical tropopause layer cirrus. *J. Geophys. Res.*, **115** (D02104), 1–15, doi:10.1029/2009JD012735.
- Dinh, T., D. R. Durran, and T. Ackerman, 2012: Cirrus and water vapor transport in the tropical tropopause layer - Part 1: A specific case modeling study. *Atmos. Chem. Phys.*, **12** (20), 9799–9815, doi:10.5194/acp-12-9799-2012.
- Durran, D. R., T. Dinh, M. Ammerman, and T. Ackerman, 2009: The mesoscale dynamics of thin tropical tropopause cirrus. *J. Atmos. Sci.*, **66** (9), 2859–2873, doi:10.1175/2009JAS3046.1.
- Fahey, D. W., et al., 2014: The AquaVIT-1 intercomparison of atmospheric water vapor measurement techniques. *Atmos. Meas. Tech. Discuss.*, **7** (4), 3159–3251, doi:10.5194/amtd-7-3159-2014.
- Jensen, E. and L. Pfister, 2004: Transport and freeze-drying in the tropical tropopause layer. *J. Geophys. Res.*, **109** (D02207), 1–16.
- Jensen, E. J., L. Pfister, T.-P. Bui, P. Lawson, and D. Baumgardner, 2010: Ice nucleation and cloud microphysical properties in tropical tropopause layer cirrus. *Atmos. Chem. Phys.*, **10** (3), 1369–1384, doi:10.5194/acp-10-1369-2010.
- Jensen, E. J., et al., 2005: Ice supersaturations exceeding 100% at the cold tropical tropopause : implications for cirrus formation and dehydration. *Atmos. Chem. Phys.*, **5**, 851–862.
- Krämer, M., et al., 2009: Ice supersaturations and cirrus cloud crystal numbers. *Atmos. Chem. Phys.*, **9** (11), 3505–3522, doi:10.5194/acp-9-3505-2009.

- Murphy, D. M., 2014: Rare temperature histories and cirrus ice number density in a parcel and one-dimensional model. *Atmos. Chem. Phys. Discuss.*, **14**, 10 701–10 723, doi:10.5194/acpd-14-10701-2014.
- Spichtinger, P. and M. Krämer, 2013: Tropical tropopause ice clouds: a dynamic approach to the mystery of low crystal numbers. *Atmos. Chem. Phys.*, **13** (19), 9801–9818, doi:10.5194/acp-13-9801-2013.