



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Simulating black carbon and dust and their radiative forcing in seasonal snow: a case study over North China with field campaign measurements

C. Zhao¹, Z. Hu^{2,1}, Y. Qian¹, L. R. Leung¹, J. Huang², M. Huang¹, J. Jin³, M. Flanner⁴, R. Zhang^{2,1}, H. Wang¹, H. Yan^{2,1}, Z. Lu⁵, and D. G. Streets⁵

¹Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA

²Key Laboratory for Semi-Arid Climate Change of the Ministry of Education, Lanzhou University, Gansu, China

³Departments of Watershed Sciences and Plants, Soils, and Climate, Utah State University, Logan, UT, USA

⁴University of Michigan, Ann Arbor, MI, USA

⁵Argonne National Laboratory Argonne, IL, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 9 May 2014 – Accepted: 15 May 2014 – Published: 23 May 2014

Correspondence to: C. Zhao (chun.zhao@pnnl.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

14, 13331–13379, 2014

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

A state-of-the-art regional model, WRF-Chem, is coupled with the SNICAR model that includes the sophisticated representation of snow metamorphism processes available for climate study. The coupled model is used to simulate the black carbon (BC) and dust concentrations and their radiative forcing in seasonal snow over North China in January–February of 2010, with extensive field measurements used to evaluate the model performance. In general, the model simulated spatial variability of BC and dust mass concentrations in the top snow layer (hereafter BCS and DSTS, respectively) are quantitatively or qualitatively consistent with observations. The model generally moderately underestimates BCS in the clean regions but significantly overestimates BCS in some polluted regions. Most model results fall into the uncertainty ranges of observations. The simulated BCS and DSTS are highest with $> 5000 \text{ ng g}^{-1}$ and up to 5 mg g^{-1} , respectively, over the source regions and reduce to $< 50 \text{ ng g}^{-1}$ and $< 1 \mu\text{g g}^{-1}$, respectively, in the remote regions. BCS and DSTS introduce a similar magnitude of radiative warming ($\sim 10 \text{ W m}^{-2}$) in the snowpack, which is comparable to the magnitude of surface radiative cooling due to BC and dust in the atmosphere. This study represents a significant effort in using a regional modeling framework to simulate BC and dust and their direct radiative forcing in snowpack. Although a variety of observational datasets have been used to attribute model biases, some uncertainties in the results remain, which highlights the need for more observations, particularly concurrent measurements of atmospheric and snow aerosols and the deposition fluxes of aerosols, in future campaigns.

1 Introduction

Snow is an important component in the Earth's climate system. Water from snowmelt generates runoff to fill rivers and reservoirs in many regions of the world. Snow cover can change the surface albedo of grassland by a factor of 3–4 and forested regions by

ACPD

14, 13331–13379, 2014

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a factor of 2–3 (Thomas and Rowntree, 1992; Betts and Ball, 1997). Therefore, snow, at a larger scale, regulates the temperature of the Earth’s surface and alters the general circulation, and, at a smaller scale, affects regional climate (e.g., Barnett et al., 1988; Walland and Simmonds, 1996). Small changes in snow albedo can have large impacts on surface warming due to the rapid feedbacks involving changes to snow morphology, sublimation and melt rates (Bond et al., 2013). Snow albedo can be influenced by many factors including snow depth, snow grain size, and snow impurity. Snow impurity is mainly due to the light absorbing aerosols (LAA) deposited on/in the snowpack, including primarily black carbon (BC), brown carbon (BrC), and dust (Warren and Wiscombe, 1980, 1985; Painter et al., 2007; Flanner et al., 2007, 2009, 2012). For example, the visible snowpack albedo can be reduced from 0.95 for pure snow to 0.1 for dirty snow with $100 \mu\text{g g}^{-1}$ BC (Warren and Wiscombe, 1980). A recent estimate for global effective BC forcing in snow and ice is $+0.13 \text{ W m}^{-2}$ (Bond et al., 2013). Dust residing in snow can also reduce snow albedo, particularly for the visible bands. In the upper Colorado river basin, a total end-of-year dust concentration of $0.23\text{--}4.16 \text{ mg g}^{-1}$ in the top 3 cm of the snow column can result in mean springtime dust radiative forcing in snow from 31 to 49 W m^{-2} at the alpine site and 45 to 75 W m^{-2} at the subalpine site (Painter et al., 2012; Skiles et al., 2012).

The LAA effects in snow could significantly affect the atmospheric general circulation and hydrological cycle at global and regional scales in multiple ways (Qian et al., 2009, 2011; Painter et al., 2010; Skiles et al., 2012; Bond et al., 2013). For example, over Asia, BC-in-snow could increase the surface air temperature and reduce spring snowpack over the Tibetan Plateau (TP) (Qian et al., 2011). Earlier snowmelt can shift the timing of peak runoff and its duration and modulate the soil moisture and surface fluxes. These impacts may result in an earlier onset of the South Asian Monsoon and an increase of moisture, cloudiness, and convective precipitation over northern India (Qian et al., 2011). They may also affect the East Asian Monsoon, resulting in changes in summer precipitation pattern (Qian et al., 2011). In the western US, Qian et al. (2009) found that BC deposition on the mountain snowpack could also cause a reduction in

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



snow accumulation and an earlier snowmelt that impacts water resources during the drier summer months. In the upper Colorado river basin, dust radiative forcing in snow could shorten the snow cover duration by 21 to 51 days, cause the peak runoff to occur 3 weeks earlier on average, and reduce the annual runoff by more than $\sim 5\%$ (Painter et al., 2010; Skiles et al., 2012).

Although there have been some modeling studies estimating the LAA concentrations in snow and their impact on snow albedo, solar absorption in snow, and climate and hydrological cycle, most previous studies used global climate models (e.g., Flanner et al., 2007, 2009, 2012; Qian et al., 2011, 2014) or snow energy balance models coupled with simplified radiation schemes with observed LAA concentrations in snow as inputs (e.g., Painter et al., 2010, 2012; Skiles et al., 2012). Both amounts and properties of snow and aerosols are extremely heterogeneous, particularly in regions with complex topography (e.g., Hansen and Nazarenko, 2004; Painter et al., 2010; Zhao et al., 2013a). Global climate models with relatively coarse horizontal resolutions (typically $1 \sim 2^\circ$) cannot fully resolve the spatial and temporal variability of aerosols in snow. In addition, global climate models with some simplified treatments of physics and chemistry processes may introduce additional uncertainties in simulating snow and aerosols (Zhao et al., 2013b, c).

Few studies of the climatic impacts of LAA in snowpack used regional climate models. Although regional models are more computationally demanding, the ability to resolve complex terrain using high spatial resolution is particularly important for simulating mountain precipitation and snowpack (e.g., Leung et al., 2003; Leung and Qian, 2003; Rasmussen et al., 2011; Yoon et al., 2012). More sophisticated physics and chemistry treatments used in some regional models can provide additional benefits. Although Qian et al. (2009) used a regional modeling framework to investigate the impact of BC in snow over the Western US, they estimated the BC content in snow based on a single year simulation with a coupled chemistry component and applied the same BC deposition to simulations of multiple years to estimate BC-snow effects. This method limits the interactions between BC deposition and climate that could influence

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the BC-induced snow albedo perturbations. Furthermore, snow albedo perturbations from snow aging were ignored, which limits the potential for accelerated snow aging induced by BC-in-snow effect. Lastly, they did not simulate the dust-in-snow effect, which likely had a much greater impact than BC in some regions (e.g., remote mountains) in specific seasons (Painter et al., 2007).

A recent version (v3.5.1) of WRF-Chem, the Weather Research and Forecasting (WRF) model (Skamarock et al., 2008) coupled with a chemistry component (Grell et al., 2005), was publicly released with the Community Land Model (CLM) v4.0 (Lawrence et al., 2011) as one of the options for land surface model (Jin et al., 2012). The Snow, Ice, and Aerosol Radiative Model (SNICAR) (Flanner and Zender, 2005) as part of the CLMv4.0 (Flanner et al., 2007, 2009, 2012) provides an opportunity to use the regional climate modeling framework with fully coupled aerosol and snow to further investigate the impacts of LAA in snowpack on regional climate and hydrological cycle. WRF-Chem is a state-of-the-art regional model that has been used extensively to study aerosols and their impacts on air quality and climate (e.g., Grell et al., 2005; Fast et al., 2006, 2009; Gustafson et al., 2007; Gao et al., 2011, 2014; Shrivastava et al., 2011; Chen et al., 2013, 2014; Zhao et al., 2010, 2011, 2012, 2013a, c). SNICAR has a sophisticated representation of snow metamorphism processes. Coupled to CLM within the Community Earth System Model (CESM) (Flanner et al., 2007), SNICAR has been widely used to simulate snow albedo as well as the radiative impact of LAA in snow on climate (e.g., Flanner et al., 2007, 2009, 2012; Qian et al., 2011, 2014).

Although SNICAR is released with the CLMv4.0 in WRF-Chem (v3.5.1), it is not connected with the atmospheric aerosol deposition and, therefore, aerosol radiative effect in snowpack does not function. This study couples the WRF-Chem simulated aerosols with SNICAR and makes aerosol radiative effect in snowpack function. The modification of SNICAR by Flanner et al. (2012) is also implemented into the model. The coupled model is used to simulate the BC and dust concentrations and their radiative forcing in snowpack during a field campaign that collected seasonal snow on a road trip at 46 sites over North China in January–February of 2010. Brown carbon is

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

excluded from the SNICAR simulations in this study because its optical properties are poorly constrained (Flanner et al., 2007, 2009). This study represents the first effort on evaluating WRF-Chem for simulating BC and dust in snowpack at a relatively high spatial resolution against the field campaign measurements of BC content in snow, and on estimating the radiative forcing of BC and dust in snow over North China. The rest of the paper is organized as follows. Sections 2 and 3 provide detailed description of the WRF-Chem and SNICAR models and the observations used in this study. In Sect. 4, the simulated spatial and temporal variation of snow and aerosol content in snow are evaluated against measurements and characterized. Then, the radiative forcing of BC and dust in snow over North China are estimated. The findings are summarized and discussed in Sect. 5.

2 Model

In this study, WRF-Chem (v3.5.1), briefly described in Sect. 2.1, is used to couple the SNICAR model with interactive aerosols as described in Sect. 2.2 below. Section 2.3 discusses the setup of model simulations. The emissions used in the simulations including anthropogenic and biomass burning emissions and online calculated emissions of mineral dust and sea-salt are described in Sect. 2.4.

2.1 WRF-Chem

In this study, the MOSAIC (Model for Simulating Aerosol Interactions and Chemistry) aerosol model (Zaveri et al., 2008) coupled with the CBM-Z (Carbon Bond Mechanism) photochemical mechanism (Zaveri and Peters, 1999), one of the chemistry options in WRF-Chem, is used and coupled with the SNICAR model. The MOSAIC aerosol scheme uses the sectional approach to represent aerosol size distributions with a number of discrete size bins, either four or eight bins in the current version of WRF-Chem (Fast et al., 2006). All major aerosol components including sulfate (SO_4^{2-}),

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

nitrate (NO_3^-), ammonium (NH_4^+), black carbon (BC), organic matter (OM), sea salt, and mineral dust are simulated in the model. The MOSAIC aerosol scheme includes physical and chemical processes of nucleation, condensation, coagulation, aqueous phase chemistry, and water uptake by aerosols. Aerosol dry deposition is simulated following the approach of Binkowski and Shankar (1995). Wet removal of aerosols by grid-resolved stratiform clouds/precipitation includes in-cloud removal (rainout) and below-cloud removal (washout) by impaction and interception, following Easter et al. (2004) and Chapman et al. (2009). Convective transport and wet removal of aerosols by cumulus clouds follow Zhao et al. (2013c).

Aerosol optical properties such as extinction, single-scattering albedo, and asymmetry factor for scattering are computed as a function of wavelength for each model grid box. Aerosols are assumed internally mixed in each bin, i.e., a complex refractive index is calculated by volume averaging for each bin for each chemical constituent of aerosols. The Optical Properties of Aerosols and Clouds (OPAC) dataset (Hess et al., 1998) is used for the SW and LW refractive indices of aerosols, except that a constant value of $1.53 + 0.003i$ is used for the SW refractive index of dust following Zhao et al. (2010, 2011). A detailed description of the computation of aerosol optical properties in WRF-Chem can be found in Fast et al. (2006) and Barnard et al. (2010). Aerosol radiative feedback is coupled with the Rapid Radiative Transfer Model (RRTMG) (Mlawer et al., 1997; Iacono et al., 2000) for both shortwave (SW) and longwave (LW) radiation as implemented by Zhao et al. (2011). The optical properties and direct radiative forcing of individual aerosol species in the atmosphere are diagnosed following the methodology described in Zhao et al. (2013a). Aerosol-cloud interactions were included in the model by Gustafson et al. (2007) for calculating the activation and re-suspension between dry aerosols and cloud droplets.

2.2 SNICAR

The Snow, Ice, and Aerosol Radiation model (SNICAR) is a multilayer model that accounts for vertically heterogeneous snow properties and heating and influence of the

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ground underlying snow (Flanner and Zender, 2005; Flanner et al., 2007, 2009, 2012). The model uses the theory from Wiscombe and Warren (1980) and the two-stream, multilayer radiative approximation of Toon et al. (1989). SNICAR can well simulate snow surface albedo as well as the radiative absorption within each snow layer. It can also simulate aerosol content and radiative effect in snow, and was first used to study the aerosol heating and snow aging in a global climate model by Flanner et al. (2007). The radiative influence of aerosol on snow albedo simulated by SNICAR has been validated with recent laboratory and field measurements (Brandt et al., 2011; Hadley and Kirchstetter, 2012). More detailed description of the SNICAR model can be found in Flanner and Zender (2005) and Flanner et al. (2007, 2012). In WRF-Chem with CLM chosen as the land surface scheme, SNICAR defines radiative layers that match the five thermal layers in CLM that vertically resolve snow thermal processes, densification, and melt-water transport (e.g., Oleson et al., 2010). Here, we focus on the description of processes associated with modeling aerosols and their optical properties in snow-pack.

2.2.1 Aerosol deposition and scavenging

BC and dust can deposit on/in snow through dry and wet deposition processes as discussed above. In this study, atmospheric aerosol wet deposition is calculated with the prognostic precipitation, thus avoiding inconsistencies associated with prescribed aerosol deposition and meteorology. Following Flanner et al. (2012), BC can mix externally and internally with precipitation hydrometeors while dust can only mix externally with precipitation hydrometeors. BC and dust removed through dry deposition processes (sedimentation and turbulent mix-out), often the dominant sink of large particles near emission sources, are mixed externally with snow grains. Dust removed through wet deposition is also treated as being externally mixed with snow grains. BC removed through below-cloud scavenging, where particles are collected by falling hydrometeors through Brownian motion, electrostatic forces, collision and/or impaction, are also assumed to remain attached to the outside of the hydrometeors rather than

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



becoming internalized. Cloud-borne BC collected by stratiform precipitation is treated as being internally mixed with snow grains. Interstitial and cloud-borne BC removed through convective scavenging processes is assumed to mix externally and internally, respectively, with snow grains. Although determining the mixing state of BC deposited through convective scavenging is uncertain, only a few percent of precipitation falling on snow surfaces comes from convection (Flanner et al., 2007; Sect. 4.2.1). Following Flanner et al. (2012), we do not simulate the distribution of aerosol number concentrations within the precipitating hydrometeors, which may influence total absorption.

In this study, SNICAR in WRF-Chem simulates four tracers of dust and two tracers of BC in snow following Flanner et al. (2012). The four tracers of dust represent four sizes of dust with diameters of $0.1 \sim 1 \mu\text{m}$, $1 \sim 2.5 \mu\text{m}$, $2.5 \sim 5 \mu\text{m}$, and $5 \sim 10 \mu\text{m}$. All four dust tracers are assumed to mix externally with snow grains. With the MO-SAIC aerosol model, WRF-Chem can simulate dust in the atmosphere with four size bins (i.e., $0.039\text{--}0.156 \mu\text{m}$, $0.156\text{--}0.625 \mu\text{m}$, $0.625\text{--}2.5 \mu\text{m}$, and $2.5\text{--}10.0 \mu\text{m}$ in dry diameter) or eight size bins ($0.039\text{--}0.078 \mu\text{m}$, $0.078\text{--}0.156 \mu\text{m}$, $0.156\text{--}0.312 \mu\text{m}$, $0.312\text{--}0.625 \mu\text{m}$, $0.625\text{--}1.25 \mu\text{m}$, $1.25\text{--}2.5 \mu\text{m}$, $2.5\text{--}5.0 \mu\text{m}$, and $5.0\text{--}10.0 \mu\text{m}$ in dry diameter). Both of them are coupled with the dust tracers in SNICAR. The coupling of dust size distributions between WRF-Chem and SNICAR are summarized in Table 1. Although both four- and eight-bin size representations in MOSAIC are coupled with SNICAR, the analysis presented in this paper focuses on the simulations using the eight-bin size representation because it is more accurate in representing aerosol size distributions (Zhao et al., 2013c). The two BC tracers in SNICAR represent interstitial and within-ice BC, respectively, in order to account for the light absorption by snowpack containing BC particles residing within snow grains (will be discussed in Sect. 2.2.2). Following Flanner et al. (2012), this study fixes two BC tracers in snow with $0.1 \mu\text{m}$ dry effective radius, although size distributions of BC may influence the BC absorption in snow, which deserves further investigation.

The scavenging of aerosols in snow by melt-water is simulated by SNICAR. CLM accounts for melt-water drainage by adding excess water to the layer beneath when

the liquid content exceeds the layer's holding capacity, defined by snow porosity and irreducible water saturation. SNICAR assumes aerosol inclusion in melt-water is proportional to its mass mixing ratio multiplied by a scavenging factor. Thus the change rate of aerosol mass in each layer i is:

$$\frac{dm_i}{dt} = k(q_{i+1}c_{i+1} - q_i c_i) + D \quad (1)$$

where m_i is the absolute mass of aerosol in layer i , k is the scavenging ratio, q_i is the mass flux of water out of layer i , c_i is the mass mixing ratio of aerosol in layer i (aerosol mass divided by liquid and solid water mass), and D is the sum of wet and dry deposition of atmospheric aerosols, added only to the surface layer of snowpack. After deposition, aerosols are mixed instantly and uniformly in the model surface layer, which never exceeds 3 cm thick. In this study, following Flanner et al. (2012), k for interstitial BC (k_{pho}) and within-ice BC (k_{phi}) are assumed to be 0.03 and 0.20, respectively, and k for four dust tracers are 0.02, 0.02, 0.01, and 0.01, respectively. These scavenging ratios are highly uncertain and need to be constrained by observations in the future (Flanner et al., 2012; Qian et al., 2014), but the values used here are generally reasonable compared to observations in the high latitudes (Doherty et al., 2013). In addition, as fresh snowfall is added, CLM continuously divides the model surface snow layer and prevents excessive aerosol accumulation. Therefore, the concentrations of each tracer within each snow layer depend on atmospheric deposition rates, new snowfall, ice sublimation, meltwater flushing, and layer combinations and divisions (Flanner et al., 2007, 2012; Oleson et al., 2010).

2.2.2 Optical properties

Optical properties of BC mixed internally and externally with snow grains are treated separately in SNICAR (Flanner et al., 2012). In this study, SNICAR accounts for the light absorption by snowpack containing BC particles residing within snow grains, which could significantly increase the solar absorption compared to interstitial BC. The refrac-

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tive indices of BC from Chang and Charalampopoulos (1990) and the effective radius of $0.1\ \mu\text{m}$ are used for offline calculated Mie parameters for BC (i.e., extinction, single scattering albedo, and scattering asymmetry parameters) (Flanner et al., 2007, 2012). Building on Chylek et al. (1984), the dynamic effective medium approximation method is applied to derive absorption enhancement ratio over broad size ranges of snow grains and BC expected for snowpack. Then, the mass absorption cross-section of BC internally mixed with snow grains is scaled by the derived enhancement ratio, which is determined as a function of BC and snow effective radius online via a lookup table. Application of the enhancement ratio was justified as a reasonable approach to account for absorption enhancement by BC internally mixed with snow grains (Flanner et al., 2012). Dust optical properties in snowpack are obtained using the Maxwell–Garnett mixing approximation along with the Mie Theory. Dust in snowpack is assumed to be a blend of SiO_2 (47.6 % by volume), limestone (2 %), montmorillonite (25 %), illite (25 %), and hematite (0.4 %) with four size bins following the lognormal distribution with a number median radius of $0.414\ \mu\text{m}$ and standard deviation of 2 (Flanner et al., 2009). Since dust is radiatively active, its co-existence could decrease the absorption by BC in snowpack (Flanner et al., 2009). Accounting for enhanced absorption by dust mixed internally with snow grains could further decrease BC absorption, but this is not considered in this study following Flanner et al. (2012).

SNICAR treats the snow particle as a collection of ice spheres, with effective number median radius of $30\text{--}1500\ \mu\text{m}$. Mie parameters for one visible ($0.3\text{--}0.7\ \mu\text{m}$) and four near-infrared (NIR) ($0.7\text{--}1.0$, $1.0\text{--}1.2$, $1.2\text{--}1.5$, and $1.5\text{--}5.0\ \mu\text{m}$) wave bands are computed offline for lognormal distributions of this wide size range of snow grains for computational efficiency. For radiative transfer calculations, the snow grains and six aerosol tracers are treated as external mixtures (with absorption enhancements applied to within-ice BC), by summing the extinction optical depths of each component, weighting the individual single-scattering albedos by optical depths, and weighting the asymmetry parameters by the product of optical depths and single-scattering albedos

(Flanner et al., 2007, 2012). A detailed description of the computation of optical properties of snow with aerosols in SNICAR can be found in Flanner et al. (2012).

2.3 Numerical experiments

In this study, the WRF-Chem simulation is performed at $36\text{ km} \times 36\text{ km}$ horizontal resolution with 120×90 grid cells ($95\text{--}140^\circ\text{ E}$, $30\text{--}60^\circ\text{ N}$) (Fig. 1) and 35 vertical layers up to 100 hPa. Although only the results for January–February of 2010 are analyzed, the simulation is conducted from 1 October 2009 to 25 February 2010 to allow accumulation of snowpack and aerosols in snowpack. The meteorological initial and lateral boundary conditions are derived from the National Center for Environmental Prediction final analysis (NCEP/FNL) data at 1° horizontal resolution and 6 h temporal intervals. The modeled u-component and v-component wind and atmospheric temperature are nudged towards the NCEP/FNL reanalysis data with a nudging time scale of 6 h (Stauf-fer and Seaman, 1990). The chemical initial and boundary conditions are provided by the quasi-global WRF-Chem simulations for the same time period (Zhao et al., 2013c). The MYJ (Mellor–Yamada–Janjic) planetary boundary layer scheme, CLM land surface scheme, Morrison 2-moment microphysics scheme, Kain–Fritsch cumulus scheme, and RRTMG longwave and shortwave radiation schemes are used in this study.

2.4 Emissions

Anthropogenic emissions are obtained from the Asian emission inventory described by Zhang et al. (2009) at $0.5^\circ \times 0.5^\circ$ horizontal resolution for 2006 except that BC, OM, and sulfate are from the China emission inventory described by Lu et al. (2011) at a $0.1^\circ \times 0.1^\circ$ horizontal spatial resolution and a monthly temporal resolution for the simulation period. Biomass burning emissions for January and February of 2010 are obtained from the Global Fire Emissions Database, Version 3 (GFEDv3) with a monthly temporal resolution and a $0.5^\circ \times 0.5^\circ$ horizontal resolution (van der Werf et al., 2010) and vertically distributed following the injection heights suggested by Dentener et al. (2006)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

for the Aerosol InterComparison project (AeroCom). Sea salt and dust emissions follow Zhao et al. (2013c). Vertical dust fluxes are calculated with the GOCART dust emission scheme (Ginoux et al., 2001), and the emitted dust particles are distributed into the MOSAIC aerosol size bins following a theoretical expression based on the physics of scale-invariant fragmentation of brittle materials derived by Kok (2011). For MOSAIC 8-bin, dust particles are emitted into eight size bins with mass fractions of $10^{-6}\%$, $10^{-4}\%$, 0.02 %, 0.2 %, 1.5 %, 6 %, 26 %, and 45 %, respectively. For MOSAIC 4-bin, dust particles are emitted into four size bins with mass fractions of $10^{-4}\%$, 0.22 %, 7.5 %, and 71 %, respectively. WRF-Chem has been widely used to simulate the dust life cycle and climatic impact at a global scale (Zhao et al., 2013a) and regional scales over West Africa (Zhao et al., 2010, 2011), Saudi Arabia (Kalenderski et al., 2013), North America (Zhao et al., 2012), and East Asia (Chen et al., 2013, 2014). The BC and dust emissions within the model domain are indicated in Fig. 1.

3 Observation

3.1 North China field campaign (NCC)

The primary observational dataset used in this study to evaluate the model simulations of snow and BC content in snow is from a field campaign that collected seasonal snow in January/February 2010 on a road trip at 46 sites in North China. The sampling locations are shown in Fig. 1. At each site, samples were gathered from individual snow pits at several depths to examine snow deposited at different times during the winter. The sampling sites were chosen to be far from local sources of pollution. About 300 snow samples were collected. Processing and initial analyses were carried out at Lanzhou University in China and at three temporary laboratories, using the filtering techniques pioneered by Clarke and Noone (1985) and also used for Arctic snow by Doherty et al. (2010). Each sample was melted rapidly in a microwave oven and then immediately drawn through a $0.4\ \mu\text{m}$ Nuclepore filter to extract the particulates.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The BC masses in snow samples are estimated using the wavelength dependence of the measured absorption from the Integrating Sandwich Sphere (ISSW) spectrophotometer (Grenfell et al., 2011). The spectrally-resolved total light absorption is divided into BC and non-BC fractions based on the absorption Angstrom exponent of the material on the filter, and by assigning different absorption Angstrom exponents to BC and non-BC light absorbing aerosols (Wang et al., 2013). The measured BC mass is an equivalent mass assuming a 550 nm BC mass absorption efficiency of $6.3 \text{ m}^2 \text{ g}^{-1}$ (Wang et al., 2013). More details about the measurements in the NCC can be found in Huang et al. (2011), Wang et al. (2013), and Zhang et al. (2013). For comparison between simulations and observations, model results are sampled at the observation sampling time at each site in January and February of 2010. The simulated BC mass content in the top snow layer (which never exceeds 3 cm thickness) is compared with the observational values averaged at the top layer (2–5 cm depending on sites) snow samples.

3.2 CMC reanalysis

The Canadian Meteorological Center (CMC) reanalysis dataset consists of Northern Hemisphere snow depth analysis data processed by the CMC. Snow depth data are obtained from surface synoptic observations, meteorological aviation reports, and special aviation reports that are acquired from the World Meteorological Organization (WMO) information system. The CMC reanalysis dataset includes daily observations from 1998 at a spatial resolution of $24 \text{ km} \times 24 \text{ km}$. Monthly averages of snow depth and estimated snow water equivalent (SWE) are provided, where SWE is estimated using a density look-up table. More descriptions and the data can be found from the CMC website (<http://nsidc.org/data/nsidc-0447.html>).

4 Results

4.1 Snowpack simulations

Before characterizing the distributions of BC and dust in snow, it is necessary to evaluate and understand potential biases of the WRF-Chem simulated snow distributions with available observations or reanalysis over North China. Figure 2 shows the spatial distributions of snow depth and SWE from the CMC reanalysis data and the WRF-Chem simulation averaged for January–February of 2010, with the NCC observations of snow depth at specific locations embedded in each panel. In general, the simulation and CMC reanalysis data are consistent with spatial correlation coefficient of 0.6, both showing snow depth increases northward and with increasing topography, although the model simulates much larger snow depth near the northern boundary and snow cover extends further south. Although both model and reanalysis data capture reasonably the campaign observed spatial variation of snow depth (Fig. 2), the model overestimates the observed snow depth, while reanalysis underestimates it (Fig. 3). Overall, the reanalysis better captures the observed spatial variability with a correlation coefficient of 0.73 compared to 0.51 for the simulation, but the model matches the observed average snow depth over the 46 observation sites better, as indicated by the mean values of 13.6 cm and 9.7 cm from the simulation and reanalysis, respectively, compared to 13.2 cm from observations. In addition, the spatial distributions of SWE from the CMC reanalysis data and the WRF-Chem simulation averaged for January–February of 2010 are also shown in Fig. 2. Again, the model results and CMC reanalysis data show consistent spatial distribution of SWE with a correlation coefficient of 0.66. Model simulates a much larger SWE than the CMC reanalysis near the north boundary. The comparison is consistent with that for snow depth.

One key factor affecting the snow simulation is surface temperature. Figure 4 shows the spatial distribution of 2 m temperature from the observations (colored circle) and the corresponding WRF-Chem simulation over North China in January–February of 2010. Daily observations of 2 m temperature in January–February of 2010 at 225 sites

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in North China are obtained from the Chinese National Meteorological Center (CNMC). The simulation can capture well the observed 2 m temperature with spatial correlation coefficient of 0.95 over the region of interest. The 2 m temperature decreases dramatically from around freezing level to about -30°C with increasing latitude. The simulated 2 m temperature averaged over the observation sites is -13°C , which is 2°C colder than that from observations, but the caveats in comparing grid mean simulated values with station data should be noted. The model cold bias is particularly noticeable over areas north of 40°N . Another key factor affecting the snow simulation is surface precipitation of snow. However, there is no publicly available observation for evaluation.

4.2 Simulations of BC and dust concentrations in snow

4.2.1 BC concentrations in snow

Figure 5 shows the spatial distribution of BC mass concentration in the top snow layer from the WRF-Chem simulation over North China in January–February of 2010. The campaign observed BC mass concentrations in the top snow layer (BCS) are also shown as the overlaid colored circles. The model simulates the highest BCS with $> 5000\text{ ng g}^{-1}$ over Northeast China ($110\text{--}130^{\circ}\text{E}$ and $38\text{--}48^{\circ}\text{N}$), where both BC deposition and snow depth are high (Figs. 2 and 6). BCS reduces away from Northeast China and reaches $< 50\text{ ng g}^{-1}$ near the north boundary of the domain. This is consistent with the NCC campaign data that show high BCS in heavily industrialized regions and smaller concentrations ($30\text{--}50\text{ ng g}^{-1}$) farther north in North China (51°N), indicating that BCS is determined by both snow coverage and BC deposition. Figure 6 shows the spatial distribution of dry and wet deposition fluxes of BC on snow from the WRF-Chem simulation in January–February of 2010. The deposition fluxes are averaged spatially and temporally over only snow-covered surface, representing the deposition on snow. Dry and wet deposition fluxes of BC are more comparable over Central China ($105\text{--}120^{\circ}\text{E}$ and $30\text{--}40^{\circ}\text{N}$), but dry deposition dominates the total BC deposition in Northeast China, where the highest BCS is found. The reason is that, over the surface

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



that is frequently covered by snow, dry deposition is more efficient than wet deposition that is limited to rainy or snow days (figures not shown). Wet deposition mainly results from resolved stratiform precipitation that accounts for more than 95 % of the total precipitation during the study period (figures not shown).

BCS are compared more explicitly with observations at each site in Fig. 7, following the sequence of each stop in the campaign from left to right on the x-axis. During the campaign, the road trip started from Inner Mongolia (sites 1–15), which is relatively clean compared to Northeast China. Most of the observed BCS over this region are around hundreds of ng g^{-1} . The cleanest snow (tens of ng g^{-1}) was sampled near the north boundary of China (sites 21–24). Polluted snow was sampled in the industrial regions of Northeast China (sites 26–40). There was no data available for sites 41–46.

The WRF-Chem simulation captures the observed sharp increase of BCS towards more polluted sites, from 30–50 ng g^{-1} at 51° N to over thousands of ng g^{-1} at 42° N. The simulated BCS agrees well with the observations at sites outside of industrial source regions (sites 1–32), except for site 6. The model generally has negative biases (within a factor of two) in these relatively clean sites. It is noteworthy that the simulation exhibits large daily and diurnal variation of BCS as denoted by the box and the whisker bar, respectively, in Fig. 7. Although the simulated BCS at the sampling time at site 6 significantly overestimates the observed BCS, the simulated mean value of January–February is closer to the observation and has a negative bias. Besides site 6, differences between the simulated values at the snow sampling time and the monthly average exist in many other sites to different degrees. This indicates the caveats in comparing model simulated monthly mean values with observations from seasonal snow sampling, a common practice in global modeling studies (e.g., Huang et al., 2011; Qian et al., 2014), because temporal variations of seasonal snow can be significant. The relative biases increase at sites 32–40 located within the industrial source regions. The model generally overestimates the BCS in these sites. It is interesting to see from Fig. 7b that the model generally has smaller biases in regions outside of the industrial source regions but large positive biases are common near or

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

MISR over land. Part of the difference between the MODIS and MISR retrievals may be due to their different overpass time, as aerosol concentrations may build up from morning to early afternoon. Ge et al. (2010) found that the MISR retrieved AOD has more reasonable agreements with ground-based observations over China compared to the MODIS “Deep-Blue” products. Overall, the model captures well both satellite retrievals, showing high AOD over the industrial regions of Central and Northeast China. The model results are more consistent with the MISR retrievals. The region with high BC deposition fluxes (Fig. 6) also has large AOD. Although satellite AOD can generally evaluate the model performance in simulating aerosols, more explicit measurements are needed to evaluate the simulated BC in the atmosphere. In addition, some biases of simulated BCS may also come from uncertainties of BC deposition rate, particularly for dry deposition rate that dominates the BC deposition over Northeast China where the simulated BCS show relatively large biases. In order to identify the uncertainty sources of BCS, concurrent measurements of atmospheric and snow BC and the deposition fluxes of BC are needed in future campaign.

Besides uncertainties in the model, the difference between observations and simulations could also result from uncertainties in the BCS retrievals (Wang et al., 2013). The measured BC mass is an equivalent mass with assumptions of BC optical properties in snowpack, such as assuming a 550 nm BC mass absorption efficiency of $6.3 \text{ m}^2 \text{ g}^{-1}$ (Wang et al., 2013). Therefore, the observations also show large uncertainty ranges (Fig. 7). In general, most model results fall into the uncertainty ranges of the observations. Furthermore, the observations at each site are assumed to be representative of the average over a $36 \text{ km} \times 36 \text{ km}$ model grid cell, which may not be true in some cases due to the inhomogeneous spatial distribution of snow and BCS.

4.2.2 Dust concentrations in snow

Figure 9 shows the spatial distribution of dust mass concentrations in the top snow layer from the WRF-Chem simulations over North China in January–February of 2010. The dust mass concentrations in the top snow layer (DSTS) are highest over Northwest

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



China and Mongolia (90–110° E and 40–45° N) and Liaoning province, which generally follow the dust source distribution (Fig. 1). DSTS can reach $\sim 5 \text{ mg g}^{-1}$ near the dust source regions, which is comparable to the magnitudes of the observed dust mass concentrations in snow after dust storms in the Upper Colorado River Basin of the US (Painter et al., 2012). DSTS gradually reduces with distance away from the dust source regions and reaches $\sim 10 \mu\text{g g}^{-1}$ over Northeast and Central China and $\sim 100 \text{ ng g}^{-1}$ near the northern boundary of the domain. The spatial distribution of DSTS follows the coincidence of dust deposition (Fig. 10) and snow coverage (Fig. 2). Both the MODIS and MISR retrievals show a band of moderate AOD in the northwest of the domain (Fig. 8), which is captured by the model. This band of moderate AOD is mainly contributed by dust (figures not shown) emitted locally from the Gobi Desert or transported eastward from the Taklimakan Desert to the west (Chen et al., 2014). The region with high dust deposition fluxes (Fig. 10) coincides well with this AOD band.

Although, there is no direct measurement of DSTS in the NCC campaign, the simulated DSTS spatial distribution is consistent with the campaign finding that DSTS was the dominant absorbing aerosol in snow over the grasslands of Northwest China and Mongolia, but is reduced significantly eastward. In addition, the NCC campaign also found a significant amount of dust along with BC in snow in Liaoning province (Huang et al., 2011). The DSTS was detected in the campaign through visual comparison of the sampling filters with a set of standard samples (Huang et al., 2011). DSTS is much higher than BCS near the dust source regions but comparable to, if not lower than, BCS over Northeast China and Central China, where BC emissions dominate. Huang et al. (2011) also found that the estimated absorbing aerosol concentrations in snow-pack range from 100 to 4300 ng g^{-1} BC equivalent in the dust-dominated regions (e.g., Inner Mongolia), higher than those in the BC-dominated regions (e.g., Northeast China) that ranged from 40 to 1600 ng g^{-1} .

4.3 Direct radiative forcing of BC and dust in snow

The single scattering albedo (SSA) complement (1-SSA) of BC, dust, and snow grains are calculated offline following the methods described in Sect. 2.2.2 and shown in Fig. 11. BC has the strongest absorption at all the five wavebands compared to dust and snow grains. BC absorption at the five wavebands decreases with increasing size from 0.05 μm to 0.5 μm (figures not shown). On the contrary, absorption of dust and snow grains increases with increasing size, except for the waveband of 1.5–5.0 μm , at which dust absorption is strongest at the smallest size (submicron particles). Dust and snow grains also have larger variation of SSA complement among the five wave bands than BC. Generally, absorption of snow grains increases with increasing wavelength, while that of dust decreases with increasing wavelength except for submicron dust particles. Although dust is much more absorbing than snow grains at wavelengths shorter than 1.0 μm , it is noteworthy that dust SSA complements could be smaller than that of snow grains, depending on the snow and dust effective radius and wavelengths. This indicates that dust in snow could increase snow albedo at some specific cases, unlike BC that always reduces snow albedo.

Figure 12 shows the absorption enhancement ratio of BC (at 0.1 μm effective radius) mixed internally with snow grains, which is also calculated offline (Sect. 2.2.2) at five spectral wavebands as a function of snow grain effective radius. BC absorption is enhanced when BC is mixed internally with snow grains compared to mixed externally, as indicated by the enhancement ratios larger than one, except for the waveband of 1.5–5.0 μm , at which BC absorption could be reduced when mixed internally with snow grains larger than $\sim 250 \mu\text{m}$ effective radius. In general, the BC absorption enhancement effect increases with decreasing wavelength and decreases with increasing snow grain effective radius. In most cases, BC absorption could be enhanced by a factor of 1.5–2 dependent on the snow grain effective radius and wavelength.

Figure 13 shows the spatial distribution of BC and dust direct radiative forcing in snowpack from the WRF-Chem simulation over North China in January–February of

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2010. The forcing is averaged when snow is present, representing the mean increase in energy absorption by snowpack. It is shown that both BC and dust introduce radiative warming in snow. The warming patterns are consistent with the spatial distributions of BCS and DSTS (Figs. 5 and 9). The warming effect reaches $\sim 10 \text{ W m}^{-2}$ for both BC and dust in snow. Although, dust is much less absorbing than BC (Fig. 11), it results in similar magnitude of warming in snow as BC due to its much larger mass content in snow than BC. As a reference, the direct radiative forcing of atmospheric BC and dust at the surface is also shown. The patterns of direct radiative forcing of atmospheric BC and dust follow the distributions of their concentrations in the atmosphere (figures not shown). Atmospheric BC and dust reduce radiation reaching surface and have a cooling effect of $\sim -10 \text{ W m}^{-2}$ near the source regions. The pattern of BC warming in snow shows higher magnitude in south than in north, while that of BC cooling at the surface is reversed. The cooling pattern at the surface of atmospheric dust roughly coincides with the warming pattern induced by dust in snow. It is interesting to note that BC and dust in snow result in comparable magnitudes of surface radiative forcing as in the atmosphere but with an opposite sign, which may indicate that BC and dust radiative forcing in snow and at the surface could offset each other in winter over some regions of North China, particularly for dust.

5 Conclusions

In this study, the WRF-Chem model is coupled with the SNICAR model to simulate the BC and dust concentrations and their radiative forcing in seasonal snow over North China in January–February of 2010, consistent with a field campaign region and period. This study represents the first effort in evaluating WRF-Chem for simulating seasonal snowpack and its impurities against various observations made over North China. The direct radiative forcing of BC and dust in snow is also estimated for the first time at a relatively high spatial resolution over North China. In general, the model captures well the observed spatial distributions of surface temperature and snow properties including

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



snow depth and snow equivalent water, which are the basis for reasonably estimating aerosol contents in snowpack.

The simulation shows that BCS is the highest ($> 5000 \text{ ng g}^{-1}$) over Northeast China and reduces away from Northeast China to $< 50 \text{ ng g}^{-1}$ near the northern boundary of the domain. DSTS is highest (up to 5 mg g^{-1}) over Northwest China and Mongolia and Liaoning province and reduces away from the dust source regions to $< 1 \mu\text{g g}^{-1}$ over Northeast and Central China and $< 100 \text{ ng g}^{-1}$ near the north boundary of domain. The spatial variability of BCS and DSTS follows that of dust deposition fluxes and snow coverage. In this study case, the dry deposition fluxes of BC and dust are generally larger than the wet deposition fluxes. The wet deposition results mainly from the resolved stratiform precipitation. DSTS is much higher than BCS near the dust source regions but comparable to, if not lower than, BCS over Northeast China and Central China with large BC emissions.

The model simulated spatial variability of BCS and DSTS are consistent with the field observations. Without direct measurements of dust content in snow, the simulated DSTS can only be evaluated qualitatively. However, the simulated DSTS reaches several mg g^{-1} near the dust source regions, which is comparable to the magnitude of the observed dust mass concentrations in snow after dust storms in the Upper Colorado River Basin of the United States (Painter et al., 2012). The simulated BCS can be evaluated quantitatively with direct measurements. The agreement between the simulated and observed BCS is generally better (within a factor of two) at the sites outside of the industrial source regions than at the sites within the industrial source regions. The difference between observations and simulations may be partly because the observations at each site are assumed to be representative of the $36 \text{ km} \times 36 \text{ km}$ model grid cell, which may not be true in some cases due to the inhomogeneous spatial distribution of snow and BCS. It may be difficult for model to resolve the sub-grid variability, which is typically larger over the source regions with large emissions than relatively clean areas. This is partly reflected by the fact that the model generally overestimates the BCS at sites inside the industrial regions. There are also uncertainties from the BCS retrievals

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



near-infrared wavebands depending on the sizes. In one of our sensitivity simulations with the modal approach to represent dust size distributions as in Zhao et al. (2013c), dust could result in cooling effect in snow in some specific cases. In addition, following Flanner et al. (2012), the size-resolved atmospheric BC is lumped together in snow and assumed a fixed effective radius. More work is needed to resolve the BC size distribution in snow. Furthermore, the atmospheric dust, if mixed internally with other aerosols, can be activated as cloud droplet. However, the effect of particles mixed internally with snow grains is only considered for BC in snowpack. Dust particles are all assumed to mix externally with snow grains. BC amounts in snow grains may also be underestimated. Although this study accounts for the ice nucleation of liquid droplets, which may contain BC through cloud droplet activation, ice nucleation of aerosols for ice- and mixed-phase clouds are not included. Therefore, the deposition of ice-borne aerosols is not considered. Finally, the inconsistency between the size distributions of deposited dust and that used for dust optical properties calculation in snowpack may also introduce uncertainties. Future study should define consistent optical properties for the atmospheric and snow dust.

In this study, although the simulated spatial distributions of BC and dust content in snowpack are generally evaluated with observations, it is still difficult to fully quantify and attribute the model biases in simulating snow impurities, which could result from uncertainties in many model processes, including emissions, deposition processes, and snow melting scavenging. To appropriately represent BC and dust content in snow, the model needs to simulate reasonably well the aerosol lifecycle in snow as discussed by Flanner et al. (2007, 2012) and Qian et al. (2014). The parametric uncertainties of the SNICAR model need to be quantified and constrained with observations. Furthermore, the aerosol lifecycle in the atmosphere should also be reasonably well simulated. The model needs to reproduce the atmospheric aerosol concentrations and treat the deposition processes correctly. Although the WRF-Chem simulated AOD is generally consistent with satellite retrievals, no direct measurement of BC or dust in the atmosphere is available for comparison. In addition, there is also no measurement to eval-

uate the simulated deposition fluxes. In order to fully understand model biases in simulating snow impurities, concurrent measurements of atmospheric and snow aerosols and the deposition fluxes of aerosols are needed in future field campaigns.

Finally, this study only estimates the direct radiative forcing by BC and dust in snowpack, which results from the enhanced absorption of solar radiation by BC and dust. It should be noted that BC and dust can indirectly enhance snow absorption by increasing the grain size due to acceleration of grain growth from the direct effect (first indirect effect) and by the earlier exposure of a darker substrate (second indirect effect) (Hansen and Nazarenko, 2004), which cannot be diagnosed in a single experiment. Further studies are needed to examine BC and dust impact with both direct and indirect effects in snow and hence the impact on climate and hydrological cycle, which can be quantified with numerical experiments by including and excluding aerosols in snow. The WRF-Chem model coupled with SNICAR can also be applied to other regions of the globe to understand the impact of BC and dust in snowpack on the regional climate and water supplies.

Acknowledgements. This research was supported by the Office of Science of the US Department of Energy (DOE) as part of the Regional & Global Climate Modeling (RGCM) program. J. Huang acknowledges support from the National Basic Research Program of China (2012CB955301). H. Wang acknowledges support from the DOE Earth System Modeling program. R. Zhang was supported by the China Scholarship Fund. This study used computing resources from the PNNL Institutional Computing. Pacific Northwest National Laboratory is operated by Battelle Memorial Institute for the DOE under contract DE-AC05-76RL01830.

References

Barnard, J. C., Fast, J. D., Paredes-Miranda, G., Arnott, W. P., and Laskin, A.: Technical Note: Evaluation of the WRF-Chem “Aerosol Chemical to Aerosol Optical Properties” Module using data from the MILAGRO campaign, *Atmos. Chem. Phys.*, 10, 7325–7340, doi:10.5194/acp-10-7325-2010, 2010.

ACPD

14, 13331–13379, 2014

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Barnett, T. P., Dumenil, L., Schlese, U., and Roeckner, E.: The effect of Eurasian snow cover on global climate, *Science*, 239, 504–507, doi:10.1126/science.239.4839.504, 1988.
- Betts, A. K. and Ball, J. H.: Albedo over the boreal forest, *J. Geophys. Res.*, 102, 28901–28909, doi:10.1029/96JD03876, 1997.
- 5 Binkowski, F. S. and Shankar, U.: The regional particulate matter model: 1. Model description and preliminary results, *J. Geophys. Res.*, 100, 26191–26209, 1995.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: a scientific assessment, *J. Geophys. Res.*, 118 5380–5552, 2013.
- Brandt, R. E., Warren, S. G., and Clarke, A. D.: A controlled snowmaking experiment testing the relation between black carbon content and reduction of snow albedo, *J. Geophys. Res.*, 116, D08109, doi:10.1029/2010JD015330, 2011.
- Chang, H. and Charalampopoulos, T. T.: Determination of the wavelength dependence of refractive indices of flame soot, *P. Roy. Soc. Lond. A*, 430, 577–591, 1990.
- Chapman, E. G., Gustafson Jr., W. I., Easter, R. C., Barnard, J. C., Ghan, S. J., Pekour, M. S., and Fast, J. D.: Coupling aerosol-cloud-radiative processes in the WRF-Chem model: Investigating the radiative impact of elevated point sources, *Atmos. Chem. Phys.*, 9, 945–964, doi:10.5194/acp-9-945-2009, 2009.
- 20 Chen, S., Huang, J., Zhao, C., Qian, Y., Leung, L. R., and Yang, B.: Modeling the Transport and Radiative Forcing of Taklimakan Dust over the Tibetan Plateau in Summer, *J. Geophys. Res.*, 118, 797–812, doi:10.1002/jgrd.50122, 2013.
- Chen, S., Zhao, C., Qian, Y., Leung, L. R., Huang, J., Huang, Z., Bi, J., Zhang, W., Shi, J., Yang, L., Li, D., and Li, J.: Regional modeling of dust mass balance and radiative forcing over East Asia using WRF-Chem, *Aeolian Res.*, in press, 2014.
- Chylek, P., Ramaswamy, V., and Cheng, R. J.: Effect of graphitic carbon on the albedo of clouds, *J. Atmos. Sci.*, 41, 3076–3084, doi:10.1175/1520-0469(1984)041<3076:EOGCOT>2.0.CO;2, 1984.
- 30 Clarke, A. D. and Noone, K. J.: Soot in the Arctic snow-pack: a cause for perturbations in radiative transfer, *Atmos. Environ.*, 19, 2045–2053, 1985.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmos. Chem. Phys.*, 6, 4321–4344, doi:10.5194/acp-6-4321-2006, 2006.

Diner, D. J., Beckert, J., Reilly, T., Bruegge, C., Conel, J., Kahn, R., Martonchik, J., Ackerman, T., Davies, R., Gerstl, S., Gordon, H., Muller, J., Myneni, R., Sellers, P. J., Pinty, B., and Verstraete, M.: Multi-angle Imaging SpectroRadiometer (MISR) instrument description and experiment overview, *IEEE T., Geosci. Remote*, 36, 1072–1087, 1998.

Diner, D. J., Abdou, W. A., Bruegge, C. J., Conel, J. E., Crean, K. A., Gaitley, B. J., Helmlinger, M. C., Kahn, R. A., Martonchik, J. V., and Pilorz, S. H.: MISR aerosol optical depth retrievals over southern Africa during the SAFARI-2000 dry season campaign, *Geophys. Res. Lett.*, 28, 3127–3130, 2001.

Doherty, S. J., Warren, S. G., Grenfell, T. C., Clarke, A. D., and Brandt, R. E.: Light-absorbing impurities in Arctic snow, *Atmos. Chem. Phys.*, 10, 11647–11680, doi:10.5194/acp-10-11647-2010, 2010.

Doherty, S. J., Grenfell, T. C., Forsstrom, S., Hegg, D. L., Brandt, R. E., and Warren, S. G.: Observed vertical redistribution of black carbon and other insoluble light-absorbing particles in melting snow, *J. Geophys. Res. Atmos.*, 118, 5553–5569, doi:10.1002/jgrd.50235, 2013.

Easter, R. C., Ghan, S. J., Zhang, Y., Saylor, R. D., Chapman, E. G., Laulainen, N. S., Abdul-Razzak, H., Leung, L. R., Bian, X., and Zaveri, R. A.: MIRAGE: model description and evaluation of aerosols and trace gases, *J. Geophys. Res.*, 109, D20210, doi:10.1029/2004JD004571, 2004.

Fast, J., Aiken, A. C., Allan, J., Alexander, L., Campos, T., Canagaratna, M. R., Chapman, E., DeCarlo, P. F., de Foy, B., Gaffney, J., de Gouw, J., Doran, J. C., Emmons, L., Hodzic, A., Herndon, S. C., Huey, G., Jayne, J. T., Jimenez, J. L., Kleinman, L., Kuster, W., Marley, N., Russell, L., Ochoa, C., Onasch, T. B., Pekour, M., Song, C., Ulbrich, I. M., Warneke, C., Welsh-Bon, D., Wiedinmyer, C., Worsnop, D. R., Yu, X.-Y., and Zaveri, R.: Evaluating simulated primary anthropogenic and biomass burning organic aerosols during MILAGRO: implications for assessing treatments of secondary organic aerosols, *Atmos. Chem. Phys.*, 9, 6191–6215, doi:10.5194/acp-9-6191-2009, 2009.

Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., and Grell, G. A.: Evolution of ozone, particulates, and aerosol direct forcing in an urban area

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

using a new fully-coupled meteorology, chemistry, and aerosol model, *J. Geophys. Res.*, 111, D21305, doi:10.1029/2005JD006721, 2006.

Flanner, M. G. and Zender, C. S.: Snowpack radiative heating: influence on Tibetan Plateau climate, *Geophys. Res. Lett.*, 32, L06501, doi:10.1029/2004GL022076, 2005.

5 Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day climate forcing and response from black carbon in snow, *J. Geophys. Res.*, 112, D11202, doi:10.1029/2006JD008003, 2007.

Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan, V., and Rasch, P. J.: Springtime warming and reduced snow cover from carbonaceous particles, *Atmos. Chem. Phys.*, 9, 2481–2497, doi:10.5194/acp-9-2481-2009, 2009.

10 Flanner, M. G., Liu, X., Zhou, C., Penner, J. E., and Jiao, C.: Enhanced solar energy absorption by internally-mixed black carbon in snow grains, *Atmos. Chem. Phys.*, 12, 4699–4721, doi:10.5194/acp-12-4699-2012, 2012.

15 Gao, Y., Liu, X., Zhao, C., and Zhang, M.: Emission controls versus meteorological conditions in determining aerosol concentrations in Beijing during the 2008 Olympic Games, *Atmos. Chem. Phys.*, 11, 12437–12451, doi:10.5194/acp-11-12437-2011, 2011.

Gao, Y., Zhao, C., Liu, X., Zhang, M., and Leung, L. R.: Regional modeling of aerosol and its radiative forcing over East Asia using WRF-Chem, *Atmos. Environ.*, 92, 250–266, 2014.

20 Ge, J., Su, J., Ackerman, T. P., Fu, Q., Huang, J., and Shi, J.: Dust aerosol optical properties retrieval and radiative forcing over northwestern China during the 2008 China – US joint field experiment, *J. Geophys. Res.*, 115, D00K12, doi:10.1029/1009JD013263, 2010.

Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.*, 106, 20225–20273, 2001.

25 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39, 6957–6976, 2005.

Grenfell, T. C., Doherty, S. J., Clarke, A. D., and Warren, S. G.: Light absorption from particulate impurities in snow and ice determined by spectrophotometric analysis of filters, *Appl. Optics*, 50, 2037–2048, 2011.

30 Gustafson, W. I., Chapman, E. G., Ghan, S. J., Easter, R. C., and Fast, J. D.: Impact on modeled cloud characteristics due to simplified treatment of uniform cloud condensation nuclei during NEAQS 2004, *Geophys. Res. Lett.*, 34, L19809, doi:10.1029/2007GL030021, 2007.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Hadley, O. L. and Kirchstetter, T. W.: Black-carbon reduction of snow albedo, *Nature Climate Change*, 2, 437–440, 2012.
- Hansen, J. and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *P. Natl. Acad. Sci. USA*, 101, 423–428, 2004.
- 5 Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: the Software Package OPAC, *B. Am. Meteorol. Soc.*, 79, 831–844, doi:10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2, 1998.
- Hsu, N. C., Tsay, S., King, M., and Herman, J. R.: Deep blue retrievals of Asian Aerosol Properties during ACE-Asia, *IEEE T. Geosci. Remote*, 44, 3180, doi:10.1109/TGRS.2006.879540, 10 2006.
- Huang, J., Fu, Q., Zhang, W., Wang, X., Zhang, R., Ye, H., and Warren, S.: Dust and black carbon in seasonal snow across northern China, *B. Am. Meteorol. Soc.*, 92, 175–181, doi:10.1175/2010BAMS3064.1, 2011.
- Iacono, M. J., Mlawer, E. J., S. A. Clough and Morcrette, J.-J.: Impact of an improved longwave radiation model, RRT M. on the energy budget and thermodynamic properties of the NCAR community climate mode, CCM3, *J. Geophys. Res.*, 105, 14873–14890, 2000.
- 15 Jin, J. and Wen, L.: Evaluation of snowmelt simulation in the Weather Research and Forecasting model, *J. Geophys. Res.*, 117, D10110, doi:10.1029/2011JD016980, 2012.
- Kalenderski, S., Stenchikov, G., and Zhao, C.: Modeling a typical winter-time dust event over the Arabian Peninsula and the Red Sea, *Atmos. Chem. Phys.*, 13, 1999–2014, doi:10.5194/acp-13-1999-2013, 2013.
- 20 Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate the size of the global dust cycle, *P. Natl. Acad. Sci. USA*, 108, 1016–1021, 2011.
- 25 Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in version 4 of the Community Land Model, *J. Adv. Model. Earth Sys.*, 3, M03001, doi:10.1029/2011MS000045, 2011.
- 30 Leung, L. R. and Qian, Y.: The sensitivity of precipitation and snowpack simulations to model resolution via nesting in regions of complex terrain, *J. Hydrometeorol.*, 4, 1025–1043, 2003.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Leung, L. R., Qian, Y., and Bian, X. D.: Hydroclimate of the western United States based on observations and regional climate simulation of 1981–2000, Part I: Seasonal statistics, *J. Climate*, 16, 1892–1911, doi:10.1175/1520-0442(2003)016<1892:HOTWUS>2.0.CO;2, 2003.
- Lu, Z., Zhang, Q., and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, *Atmos. Chem. Phys.*, 11, 9839–9864, doi:10.5194/acp-11-9839-2011, 2011.
- Martonchik, J. V., Diner, D. J., Kahn, R., and Gaitley, B.: Comparison of MISR and AERONET aerosol optical depths over desert sites, *Geophys. Res. Lett.*, 31, L16102, doi:10.1029/2004GL019807, 2004.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., M. J. Iacono and Clough, S. A.: RRTM, a validated correlated- k model for the longwave, *J. Geophys. Res.*, 102, 16663–16682, 1997.
- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddes, J., Heald, C. L., Hoffman, F., Lamarque, J.-F., Mahowald, N., Niu, G.-Y., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stöckli, R., Wang, A., Yang, Z.-L., Zeng, X., and Zeng, X.: Technical Description of version 4.0 of the Community Land Model (CLM), Tech. Rep. NCAR/TN-478 + STR, National Center for Atmospheric Research, 2010.
- Painter, T. H., Barrett, A. P., Landry, C. C., Neff, J. C., Cassidy, M. P., Lawrence, C. R., McBride, K. E., and Farmer, G. L.: Impact of disturbed desert soils on duration of mountain snow cover, *Geophys. Res. Lett.*, 34, L12502, doi:10.1029/2007GL030284, 2007.
- Painter, T. H., Deems, J. S., Belnap, J., Hamlet, A. F., Landry, C. C., and Udall, B.: Response of Colorado River runoff to dust radiative forcing in snow, *P. Natl. Acad. Sci. USA*, 107, 17125–17130, doi:10.1073/pnas.0913139107, 2010.
- Painter, T. H., Skiles, S. M., Deems, J. S., Bryant, A. C., and Landry, C. C.: Dust radiative forcing in snow of the Upper Colorado River Basin: 1. A 6 year record of energy balance, radiation, and dust concentrations, *Water Resour. Res.*, 48, W07521, doi:10.1029/2012WR011985, 2012.
- Qian, Y., Gustafson Jr., W. L., Leung, L. R., and Ghan, S. J.: Effects of soot-induced snow albedo change on snowpack and hydrological cycle in western United States based on Weather Research and Forecasting chemistry and regional climate simulations, *J. Geophys. Res.*, 114, D03108, doi:10.1029/2008JD011039, 2009.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Qian, Y., Flanner, M. G., Leung, L. R., and Wang, W.: Sensitivity studies on the impacts of Tibetan Plateau snowpack pollution on the Asian hydrological cycle and monsoon climate, *Atmos. Chem. Phys.*, 11, 1929–1948, doi:10.5194/acp-11-1929-2011, 2011.

Qian, Y., Wang, H., Zhang, R., Flanner, M. G., and Rasch, P. J.: Black carbon in snow and its radiative forcing over the Arctic and Northern China: uncertainty associated with deposition and in-snow processes, *Environ. Res. Lett.*, in press, 2014.

Rasmussen, R. J., Liu, C., Ikeda, K., Gochis, D., Yates, D., Chen, F., Tewari, M., Barlage, M., Dudhia, J., Yu, W., and Miller, K.: High-resolution coupled climate runoff simulations of seasonal snowfall over Colorado: a process study of current and warmer climate, *J. Climate*, 24, 3015–3048, 2011.

Shrivastava, M., Fast, J., Easter, R., Gustafson Jr., W. I., Zaveri, R. A., Jimenez, J. L., Saide, P., and Hodzic, A.: Modeling organic aerosols in a megacity: comparison of simple and complex representations of the volatility basis set approach, *Atmos. Chem. Phys.*, 11, 6639–6662, doi:10.5194/acp-11-6639-2011, 2011.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X., Wang, W., and Powers, J. G.: A description of the advanced research WRF version 3, NCAR Tech. Note, NCAR/TN-475+STR, 8 pp., Natl. Cent. for Atmos. Res., Boulder, Colo., available at: http://www.mmm.ucar.edu/wrf/users/docs/arw_v3.pdf (last access: May 2014), 2008.

Skiles, S. M., Painter, T. H., Deems, J. S., Bryant, A. C., and Landry, C. C.: Dust radiative forcing in snow of the Upper Colorado River Basin: 2. Interannual variability in radiative forcing and snowmelt rates, *Water Resour. Res.*, 48, W07522, doi:10.1029/2012WR011986, 2012.

Stauffer, D. R. and Seaman, N. L.: Use of four-dimensional data assimilation in a limited-area mesoscale model, Part I: Experiments with synoptic-scale data, *Mon. Weather Rev.*, 118, 1250–1277, 1990.

Thomas, G. and Rowntree, P. R.: The boreal forests and climate, *Q. J. Roy. Meteor. Soc.*, 118, 469–497, doi:10.1002/qj.49711850505, 1992.

Toon, O. B., McKay, C. P., Ackerman, T. P., and Santhanam, K.: Rapid calculation of radiative heating rates and photodissociation rates in inhomogeneous multiple scattering atmospheres, *J. Geophys. Res.*, 94, 16,287–16,301, 1989.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Walland, D. J. and Simmonds, I.: Modelled atmospheric response to changes in Northern Hemisphere snow over, *Clim. Dynam.*, 13, 25–34, doi:10.1007/s003820050150, 1996.
- Wang, X., Doherty, S. J., and Huang, J.: Black carbon and other light-absorbing impurities in snow across Northern China, *J. Geophys. Res.-Atmos.*, 118, 1471–1492, doi:10.1029/2012JD018291, 2013.
- Warren, S. G. and Wiscombe, W. J.: A model for the spectral albedo of snow, II. Snow containing atmospheric aerosols, *J. Atmos. Sci.*, 37, 2734–2745, doi:10.1175/1520-0469(1980)037<2734:AMFTSA>2.0.CO;2, 1980.
- Warren, S. G. and Wiscombe, W. J.: Dirty snow after nuclear war, *Nature*, 313, 467–470, doi:10.1038/313467a0, 1985.
- Wiscombe, W. J. and Warren, S. G.: A model for the spectral albedo of snow, I: Pure snow, *J. Atmos. Sci.*, 37, 2712–2733, 1980.
- Yoon, J.-H., Ruby Leung, L., and Correia Jr., J.: Comparison of dynamically and statistically downscaled seasonal climate forecasts for the cold season over the United States, *J. Geophys. Res.*, 117, D21109, doi:10.1029/2012JD017650, 2012.
- Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *J. Geophys. Res.*, 104, 30387–30415, 1999.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for simulating aerosol interactions and chemistry (MOSAIC), *J. Geophys. Res.*, 113, D13204, doi:10.1029/2007JD008792, 2008.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, doi:10.5194/acp-9-5131-2009, 2009.
- Zhang, R., Hegg, D. A., Huang, J., and Fu, Q.: Source attribution of insoluble light-absorbing particles in seasonal snow across northern China, *Atmos. Chem. Phys.*, 13, 6091–6099, doi:10.5194/acp-13-6091-2013, 2013.
- Zhao, C., Liu, X., Leung, L. R., Johnson, B., McFarlane, S. A., Gustafson Jr., W. I., Fast, J. D., and Easter, R.: The spatial distribution of mineral dust and its shortwave radiative forcing over North Africa: modeling sensitivities to dust emissions and aerosol size treatments, *Atmos. Chem. Phys.*, 10, 8821–8838, doi:10.5194/acp-10-8821-2010, 2010.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Zhao, C., Liu, X., Ruby Leung, L., and Hagos, S.: Radiative impact of mineral dust on monsoon precipitation variability over West Africa, *Atmos. Chem. Phys.*, 11, 1879–1893, doi:10.5194/acp-11-1879-2011, 2011.
- 5 Zhao, C., Liu, X., and Leung, L. R.: Impact of the Desert dust on the summer monsoon system over Southwestern North America, *Atmos. Chem. Phys.*, 12, 3717–3731, doi:10.5194/acp-12-3717-2012, 2012.
- Zhao, C., Leung, L. R., Easter, R., Hand, J., and Avise, J.: Characterization of speciated aerosol direct radiative forcing over California, *J. Geophys. Res.*, 118, 2372–2388, doi:10.1029/2012JD018364, 2013a.
- 10 Zhao, C., Liu, X., Qian, Y., Yoon, J., Hou, Z., Lin, G., McFarlane, S., Wang, H., Yang, B., Ma, P.-L., Yan, H., and Bao, J.: A sensitivity study of radiative fluxes at the top of atmosphere to cloud-microphysics and aerosol parameters in the community atmosphere model CAM5, *Atmos. Chem. Phys.*, 13, 10969–10987, doi:10.5194/acp-13-10969-2013, 2013b.
- 15 Zhao, C., Chen, S., Leung, L. R., Qian, Y., Kok, J. F., Zaveri, R. A., and Huang, J.: Uncertainty in modeling dust mass balance and radiative forcing from size parameterization, *Atmos. Chem. Phys.*, 13, 10733–10753, doi:10.5194/acp-13-10733-2013, 2013c.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Table 1. Matching of dust size distributions between WRF-Chem and SNICAR. The values represent mass fractions of deposited atmospheric dust from WRF-Chem into each size of dust in snow in SNICAR.

		SNICAR 4 bin (diameter in μm)				
			0.1–1	1–2.5	2.5–5	5–10
WRF-Chem	4 bin (Diameter in μm)	0.039–0.156	1	0	0	0
		0.156–0.625	1	0	0	0
		0.625–2.5	0	1	0	0
		2.5–10	0	0	0.5	0.5
WRF-Chem	8 bin (Diameter in μm)	0.039–0.078	1	0	0	0
		0.078–0.156	1	0	0	0
		0.156–0.312	1	0	0	0
		0.312–0.625	1	0	0	0
		0.625–1.25	0	1	0	0
		1.25–2.5	0	1	0	0
		2.5–5.0	0	0	1	0
		5.0–10.0	0	0	0	1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 1. Model domain with BC and dust emissions in January–February of 2010. The 46 observation sites in the North China campaign are denoted by pentagrams in black.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 2. Spatial distribution of snow depth (cm) and snow water equivalent (cm) from the WRF-Chem simulations and CMC reanalysis over North China in January–February of 2010. Snow depth (cm) from the campaign observations (colored circle) is also shown.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 3. The campaign observations of snow depth (cm) vs. the corresponding WRF-Chem simulations (black dots) and CMC reanalysis (red dots).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Spatial distribution of 2 m temperature from the observations (colored circle) and the corresponding WRF-Chem simulations over North China in January–February of 2010.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 5. Spatial distribution of BC mass concentration in the top snow layer from the WRF-Chem simulations over North China in January–February of 2010. The campaign observed BC mass concentrations in the top snow layer (colored circle) are also shown.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 6. Spatial distribution of dry and wet deposition fluxes of BC on snow from the WRF-Chem simulations over North China in January–February of 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 7. (a) BC mass concentrations in the top snow layer from the campaign observations (black) and the corresponding WRF-Chem simulations (red). The box plot of observations shows the minimum and maximum possible values from retrievals and the bar within the box shows the most likely value; the whisker bar of model results shows the minimum and maximum simulated values at 24 h of January–February. The asterisk represents the 24 h average value for January–February. The box of model results shows the minimum and maximum simulated values at observation sampling hour of January–February. The bar within the box represents the simulated value at the observation sampling hour and day. (b) Ratio (red circles) of BC mass concentration in the top snow layer from the model at the observation sampling time and observation along with latitudes (green line).

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 8. Spatial distribution of AOD over North China in January–February of 2010 from the MISR and Aqua-MODIS retrievals and the corresponding WRF-Chem simulations. The model results are sampled at the MISR (~ 10:45 LT) and Aqua-MODIS (~ 13:30 LT) overpass time, respectively, at the locations where retrievals are available.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 9. Spatial distribution of dust mass concentration in the top snow layer from the WRF-Chem simulations over North China in January–February of 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 10. Spatial distribution of dry and wet deposition fluxes of dust on snow from the WRF-Chem simulations over North China in January–February of 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 11. Single scattering albedo (SSA) complement ($1-SSA$) at the five spectral wavebands of snow grains as a function of effective radius (colored solid lines), BC with an effective radius of $0.1 \mu\text{m}$ (asterisk), and dust with four different sizes (Dust1 (circle), Dust2 (square), Dust3 (triangle), and Dust4 (diamond)).

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 12. Absorption enhancement ratio of BC ($0.1 \mu\text{m}$ effective radius) internally mixed with snow grains at the five spectral wavebands as a function of snow grain effective radius.

Simulating black carbon and dust and their radiative forcing in seasonal snow

C. Zhao et al.




Figure 13. Spatial distribution of BC and dust direct radiative forcing at the surface and in the snow from the WRF-Chem simulations over North China in January–February of 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)