Atmos. Chem. Phys. Discuss., 14, 14471–14518, 2014 www.atmos-chem-phys-discuss.net/14/14471/2014/ doi:10.5194/acpd-14-14471-2014 © Author(s) 2014. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Development of a 10 year (2001–2010) 0.1° dataset of land-surface energy balance for mainland China

X. Chen^{1,2}, Z. Su¹, Y. Ma², S. Liu³, Q. Yu⁴, and Z. Xu³

¹Faculty of Geo-Information Science and Earth Observation, University of Twente, Enschede, the Netherlands

²Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, China

³State Key Laboratory of Remote Sensing Science, School of Geography, Beijing Normal University, Beijing, China

⁴Plant Functional Biology and Climate Change Cluster, University of Technology, Sydney, P.O. Box 123, Broadway, NSW 2007, Australia

Received: 20 February 2014 - Accepted: 14 May 2014 - Published: 5 June 2014

Correspondence to: X. Chen (x.chen@utwente.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.



Abstract

In the absence of high resolution estimates of the components of surface energy balance for China, we developed an algorithm based on the surface energy balance system (SEBS) to generate a dataset of land-surface energy and water fluxes on a monthly time scale from 2001 to 2010 at a 0.1° × 0.1° spatial resolution by using multi-satellite and meteorological forcing data. A remote-sensing-based method was developed to estimate canopy height, which was used to calculate roughness length and flux dynamics. The land-surface flux dataset was validated against "ground-truth" observations from 11 flux tower stations in China. The estimated fluxes correlate well with the stations' measurements for different vegetation types and climatic conditions (av-10 erage bias = 15.3 W m^{-2} , RMSE = 26.4 W m^{-2}). The quality of the data product was also assessed against the GLDAS dataset. The results show that our method is efficient for producing a high-resolution dataset of surface energy flux for the Chinese landmass from satellite data. The validation results demonstrate that more accurate downward long-wave radiation datasets are needed to be able to accurately estimate turbulent fluxes and evapotranspiration when using the surface energy balance model. Trend analysis of land-surface radiation and energy exchange fluxes revealed that the Tibetan Plateau has undergone relatively stronger climatic change than other parts

of China during the last 10 years. The capability of the dataset to provide spatial and temporal information on water-cycle and land–atmosphere interactions for the Chinese landmass is examined. The product is free to download for studies of the water cycle and environmental change in China.

1 Introduction

As China is one of the fastest growing and urbanizing economies in the world, changes in land cover and land use can significantly influence the environment by altering land– atmosphere energy and water exchanges (Suh and Lee, 2004; Lin et al., 2009). For



instance, rapid urban expansion has substantially changed surface heat fluxes in the Pearl River delta (PRD) (Lin et al., 2009) and has increased sensible heat fluxes in the Beijing metropolitan area (Zhang et al., 2009a). The variability of surface energy balance and its partitioning may also have an important impact on climate variability

in China (Sun and Wu, 2001). Similarly, changes in surface energy fluxes have been shown to alter the intensity of the East Asian monsoon (Zhou and Huang, 2008; Qiu, 2013; Hsu and Liu, 2003). In short, understanding variation in energy fluxes is important for the study of climate change in China (Brauman et al., 2007). Nevertheless, the spatial and temporal variability of China's land-surface energy balance, and the magnitude of each, are still unknown.

While it is of critical importance to understand the partitioning of water and energy distribution across China's terrestrial surface, accurate monitoring of their spatial and temporal variation is notoriously difficult (Ma et al., 2011). Several field experiments are being carried out to monitor turbulent fluxes over selected land cover in China by using ground-based eddy covariance devices (Liu et al., 2012; Wang et al., 2010;

Yu et al., 2006; Ma et al., 2008b; Li et al., 2009). However, these measurements are only representative of small areas around the locations where the measurements are being made. For this reason, establishment of an eddy-covariance flux network cannot provide a complete land-surface heat flux picture for the entire Chinese landmass.

15

A number of products can be derived from land-surface energy fluxes. Jung et al. (2009), for example, generated global spatial flux fields by using a network upscaling method. However their flux network included only a limited number of flux stations in China. The Global Soil Wetness Project 2 (GSWP-2) (Dirmeyer et al., 2006) produced a global land surface product on a 1° × 1° grid for the period 1986 to 1995.

²⁵ The Global Land Data Assimilation System (GLDAS) (Rodell et al., 2004) can provide a global coverage in the form of 3 hourly, 0.25° data. Furthermore, products from the European Centre for Medium-Range Weather Forecasts (ECMWF) interim reanalysis (ERA-Interim) (Dee et al., 2011), the National Centers for Environmental Prediction (NCEP) (Kalnay et al., 1996), Modern-Era Retrospective Analysis for Research and



Applications (MERRA) (Rienecker et al., 2011) and other reanalysis data can also provide temporally continuous – but coarse – spatial resolution datasets of land surface fluxes. Jiménez et al. (2011) made an inter-comparison of different land-surface heat flux products. When these products were applied at continental scales, the different approaches resulted in large differences (Vinukollu et al., 2011a; Jiménez et al., 2011; Mueller et al., 2011).

The problems met by using currently available flux data in climate studies of China have been reported by Zhou and Huang (2010). Zhu et al. (2012) have also reported that summer sensible heat flux derived from eight datasets (including NCEP, ERA, and GLDAS) of China's Tibetan Plateau region differ from each other in their spatial distribution. In addition, all the flux datasets mentioned above are based on model simulations, which have deficiencies for studying changes in water-cycle and land–air interactions in China (Chen et al., 2013c; Su et al., 2013; Wang and Zeng, 2012; Ma et al., 2008a).

10

A spatially and temporally explicit estimate of surface energy fluxes is of considerable interest for hydrological assessments and meteorological and climatological investigations (Norman et al., 2003). Satellite-sensed data of surface variables can be used to produce maps of heat and water fluxes at different scales (Wang and Liang, 2008; X. Lui et al., 2012; Liu et al., 2010; Vinukollu et al., 2011b). Remote sensing approaches

- to estimate surface heat and water fluxes have been largely used on regional scales (Fan et al., 2007; Ma et al., 2011; Jia et al., 2012; Zhang et al., 2009b; Z. Li et al., 2012; Shu et al., 2011), but there is no analysis of satellite-derived data currently underway to produce a complete, physically-consistent, decadal land-surface heat flux dataset (Jiménez et al., 2009) for the Chinese landmass. The use of remotely-sensed data of-
- fers the potential of acquiring observations of variables such as albedo, land surface temperature, and NDVI at a continental scale for China. Figure 1 shows an example of an NDVI map for China.

Since surface fluxes cannot be directly detected by satellite-borne sensors, an alternative for estimating continental water and energy fluxes can be derived by applying



the aerodynamic theory of turbulent flux transfer (Ma et al., 2011) or by establishing statistical relationships between related satellite observations and land surface fluxes (Jiménez et al., 2009; Wang et al., 2007). Most remotely-sensed latent heat flux or evapotranspiration products have null values in urban, water, snow, barren and desert areas (Mu et al., 2007; Wang et al., 2007; Jiménez et al., 2009). This is due to the lack

- of a uniform representation of turbulent exchange processes over different types of land cover in their method. Meanwhile, the aerodynamic turbulent transfer method can describe the flux exchange through changes in surface roughness length over different land covers. Statistical methods establish relationships between satellite-sensed
- ¹⁰ observations (e.g. NDVI, LST, albedo) and land surface fluxes through various fitting techniques (Wang et al., 2007). The simple relationships established cannot give a reasonable approximation for extreme conditions such as bare soil or other types of non-canopy land cover (e.g. lakes, deserts) because land covers behave significantly differently in land-surface energy flux partitioning. Fortunately, turbulent flux transfer parameterization can overcome the shortcomings of statistical methods and produce
- spatially continuous distributions of land-surface energy fluxes with prepared meteorological forcing data. For this reason we chose a more physically-based method – turbulent flux parameterization – to produce the dataset.

The challenge in using turbulent flux parameterization lies in the transition from regional to continental and global scales, because meteorological data of high resolution (i.e. 1–10 km) are not easily obtained for a large region. Recently, Chinese scientists have produced high resolution meteorological forcing data that can be used in our study. Another issue is the complexity met with the method when combining different spatial and temporal sampling input variables. This is discussed in detail in Sect. 3.1.

²⁵ The last difficulty that has surrounded application of turbulent flux parameterization at continental scales is the acquisition of roughness length. To address this difficulty, we have developed a remote-sensing-based mixing technique to estimate canopy heights at a continental scale and use the resulting canopy height dataset to derive, for the very first time, the dynamic variation of surface roughness length for the Chinese landmass.



In our study we set out to estimate turbulent heat fluxes simulated with energy balance and aerodynamic parameterization formulas that are based on a revised model of the surface energy balance system (SEBS) (Chen et al., 2013a, b; Su, 2002; Timmermans, 2011); tests show that the revised model delivers better performance and ⁵ improvements in cases where the type of land cover in China is bare soil, short canopy or snow (Chen et al., 2013a, b). Sensible heat flux in SEBS was derived from the difference between surface temperature and air temperature by using Monin–Obukhov sim-

ilarity theory and bulk atmospheric boundary layer similarity (Brutsaert, 1999), which parameterizes ground surface momentum and heat-transfer coefficient maps to take
into account surface roughness, canopy height, vegetation cover, and meteorological stability (Su et al., 2001; Su, 2002; Chen et al., 2013b). The latent heat flux can then be estimated from an energy balance model, assuming surface net radiation and ground flux are known (Ma et al., 2002; Allen et al., 2011; Vinukollu et al., 2011b).

Complex topography and climatic conditions in China make it very difficult to obtain a clear picture of the distribution of energy and water fluxes with a high spatial resolution over a relatively long period for such a large area. To derive the surface energy balance terms for the Chinese landmass, we used high resolution reanalysis data, which merges model outputs, remote sensing observations, and in-situ measurements. In addition, we also assessed the accuracy of the surface energy balance terms (net rediction complete latent heat, and ground heat fluxes) and their alignatic terms (net

radiation, sensible heat, latent heat, and ground heat fluxes) and their climatic trends in the preceding decade (2001–2010).

After defining the equations of the SEBS model (Sect. 2), we describe (in Sect. 3) the input data and ground-truth measurements used in the study. Further, we assess the capacity of the remote-sensing-based product to reproduce the range and variability of

²⁵ measured fluxes by comparing them with in-situ flux tower measurements, followed by trend analysis of the spatial patterns of the fluxes (Sect. 4). Concluding remarks are found in Sect. 5.



2 Model description and development

The surface energy balance system model known as SEBS (Su, 2002) uses aerodynamic resistance to create a spatially coherent estimate of land heat fluxes. Some model inputs can be obtained from remote sensing data, while others can be obtained

from meteorological forcing data (e.g. GLDAS, ERA and NCEP reanalysis data). The model's equations and the required forcing variables are described in the remainder of this section.

The surface energy balance equation can be expressed as:

¹⁰ Rn = $G_0 + H + LE$,

25

where Rn is the net radiation flux; G_0 is the ground heat flux, which is parameterized by its relationship with Rn (Su et al., 2001); *H* is the sensible heat flux; and LE is the latent heat flux.

LE is computed by using the evaporative fraction after deriving the other three vari-¹⁵ ables in Eq. (1) and taking into consideration energy and water limits (Su, 2002). As these fluxes were produced with a monthly average temporal resolution, energy storage in vegetation is not considered.

Net radiation flux is:

Rn = $(1 - \alpha) \times SWD + LWD - LWU$,

where α is broadband albedo; SWD is downward surface short-wave radiation; and LWD and LWU are downward and upward surface long-wave radiation, respectively. Here satellite observed albedo is used. LWU is derived from land surface temperature (LST) using the Stefan–Boltzmann law. Land surface emissivity is derived as described in Chen et al. (2013a). LWD and SWD values are obtained from meteorological

forcing data.



(1)

(2)

Sensible heat flux (H) is computed according to the Monin–Obukhov similarity theory (MOST):

$$H = k u_* \rho c_p (\theta_0 - \theta_a) \left[\ln \left(\frac{z - d}{z_{0h}} \right) - \Psi_h \left(\frac{z - d}{L} \right) + \Psi_h \left(\frac{z_{0h}}{L} \right) \right]^{-1}, \tag{3}$$

⁵ where *k* is the von Karman constant; u_* is friction velocity; ρ is air density; c_p is specific heat for moist air; θ_0 is the potential temperature at the ground surface; θ_a is the potential air temperature at height *z*; *d* is the zero plane displacement height; Ψ_h is the stability correction function for sensible heat transfer (Brutsaert, 1999); and *L* is the Obukhov length. In our study θ_a was obtained from meteorological forcing data and θ_0 was derived from Moderate Resolution Imaging Spectroradiometer (MODIS) LST data. For more detailed information about u_* and the calculation of *L*, see Su (2002) and Chen et al. (2013b).

The roughness height for heat transfer (z_{0h}) in Eq. (3) is calculated as follows:

$$z_{\rm 0h} = \frac{z_{\rm 0m}}{\exp(kB^{-1})}.$$
 (4)

15

Using the fractional canopy coverage, kB^{-1} at each pixel can be derived according to the following modification of the equation described by Su et al. (2001):

$$kB^{-1} = f_{\rm c}^2 \times kB_{\rm c}^{-1} + f_{\rm s}^2 \times kB_{\rm s}^{-1} + 2 \times f_{\rm c} \times f_{\rm s} \times kB_{\rm m}^{-1},$$
(5)

where f_c is fractional canopy coverage and f_s is the fraction of bare soil in one pixel; kB_c^{-1} is the kB^{-1} of the canopy; kB_s^{-1} is the kB^{-1} of bare soil; and kB_m^{-1} is kB^{-1} for mixed bare soil and canopy. As kB^{-1} is the most important parameter in a MOST-based calculation of sensible heat flux, kB^{-1} has been updated by Chen et al. (2013b). The momentum roughness length used to calculate kB_s^{-1} was given a value of 0.004 (Chen et al., 2013b), and the heat roughness length of bare soil was calculated according to



Yang et al. (2002). The new kB^{-1} gives a better performance than the previous version of kB^{-1} (Chen et al., 2013b, a). Detailed evaluations of the new parameterization of kB^{-1} can be found in Chen et al. (2013b).

The roughness height for momentum transfer z_{om} in Eq. (4) is derived from canopy ⁵ height (HC), leaf area index (LAI) and the canopy momentum transfer model (Massman, 1997):

$$z_{om} = HC \times (1 - d/HC) \times exp(-k \times \beta),$$

$$\beta = C_1 - C_2 \times exp(-C_3 \times C_d \times LAI),$$
(6)
(7)

¹⁰ where $C_1 = 0.32$, $C_2 = 0.26$, and $C_3 = 15.1$ are model constants related to the bulk surface drag coefficient (Massman, 1997). The three constants have been tested for several canopies (Chen et al., 2013b; Cammalleri et al., 2010) and evaluated as one of the best solutions for canopy turbulent-flux parameterization (Cammalleri et al., 2010). C_d is the drag coefficient, which typically equals 0.2 (Goudriaan, 1977); *d* is displacement height, which is derived from HC and the wind speed extinction coefficient (Su, 2002; Su et al., 2001).

As Chen et al. (2013b) have pointed out, HC is vital for turbulent heat simulations, which makes accurate estimation of HC for the Chinese landmass important for this study. A remote-sensing-based canopy height method (Chen et al., 2013b) was further developed to estimate canopy height distribution for the whole China. Simard et al. (2011) produced a global forest canopy-height map using data from the Geoscience Laser Altimeter System (GLAS) aboard ICESat (Ice, Cloud, and land Elevation Satellite). However, short-canopy (e.g. maize, rice, wheat) height information cannot be acquired by laser techniques. Since short-canopy height usually varies by season
throughout the year – crops are planted in spring and harvested in autumn – we calculated short-canopy height using an NDVI-based equation from Chen et al. (2013b):

14479

$$HC = HC_{min} + \frac{HC_{max} - HC_{min}}{(NDVI_{max} - NDVI_{min})} \times (NDVI - NDVI_{min})$$



(8)

where HC_{max} and HC_{min} are the maximum and minimum short-canopy height; HC_{min} is set to 0.002 m; and HC_{max} is set to 2.5 m, corresponding to the greatest height of seasonal crops in China. NDVI_{min} and NDVI_{max} are minimum and maximum NDVI values during our 10 year study period. Each short-canopy pixel was given an NDVI_{min} and NDVI_{max} value to calculate the canopy height. The NDVI-based short-canopy height method above was used to fill relevant pixels with forest canopy heights of less than 10 m. Higher canopy heights (greater than 10 m) were assumed to be constant, i.e. with no seasonal change. By merging canopy heights greater than 10 m and variable short-canopy data, we constructed dynamic monthly maps of canopy heights for the Chinese
landmass for the period of 2001–2010. These maps were then used to calculate heat fluxes.

3 Data and validation

Our modeling approach makes use of a variety of satellite-based sensor data and meteorological forcing data to estimate monthly energy and water fluxes across China.

The forcing data can come from satellite-based or reanalysis datasets. Due to the influence of weather, satellite-sensed visible and thermal band data (e.g. NDVI, albedo, LST) often have spatial and temporal gaps in daily data. Various temporal and spatial gap-filling algorithms have been developed to produce continuous monthly data for satellite-sensed variables (Chen et al., 2004; Moody et al., 2005). In order to avoid both spatial and temporal gaps in the final product, we selected some specific satellite-sensed datasets for this study (see Table 1). Detailed information about each input

variable is described in following subsections.

The longest period covered by the forcing dataset is approximately 31 years; the shortest is about 10 years. Spatial resolution of the dataset varies from 0.01 to 0.25° and its sample frequency from 3 h to 1 month. The meteorological forcing data developed by the Institute of Tibetan Plateau Research, Chinese Academy of Sciences (hereafter referred to as ITPCAS forcing data) (He, 2010) was constructed to study



meteorological variation in China. ITPCAS forcing data covers the entire landmass of China and has the highest temporal resolution among the input datasets used. Other variables such as LST and albedo, for example, have coarser temporal resolutions (monthly) and global coverage. When combining data of different spatial and temporal scaling issues need to be addressed.

Estimates of land-surface energy flux can be subject to large errors, due to bias in the meteorological forcing input data. The spatial distribution of meteorological variables is closely related to topography (Li et al., 2013). When interpolating meteorological input variables to finer scales, these effects have to be accounted for (Sheffield et al., 2006),

- which goes beyond the scope of our study. Therefore we chose to resample the satellite product of high spatial resolution to a lower spatial resolution that matches the resolution of the meteorological input data. Also, the meteorological data were averaged to monthly values that have the same temporal resolution as the remotely-sensed input variables. ITPCAS forcing data provides us data of the highest spatial resolution among
- the meteorological forcing data currently available (e.g. ERA-interim, NCEP, GLDAS, MERRA). Taking into account of all these items, our aim was to produce a monthly product of 0.1° × 0.1° resolution land-surface heat fluxes that contains neither spatial nor temporal gaps and can be used to study seasonal and inter-annual variability in the hydrological and energy cycles of China.

20 3.1 Input datasets

3.1.1 Meteorological forcing data

In studies previous to ours, reanalysis data have been applied in many different ways, for example to construct land-surface forcing data (Sheffield et al., 2006), to detect climate trends (Taniguchi and Koike, 2008), and to investigate water and energy cycles at regional and continental scales (Roads and Betts, 2000). Reanalysis data has also been applied by the remote sensing community to derive estimates of global terrestrial evapotranspiration and gross primary production (Mu et al., 2007; Yuan et al.,



2010). Few studies, however, have used reanalysis data together with remotely-sensed ground data to derive global land-energy fluxes (sensible heat flux, latent heat flux, net radiation, etc.).

- Researchers have developed several kinds of reanalysis data. Comparisons and evaluations of these reanalysis products with in-situ observations have been performed for individual sites, specific regions, and the entire globe (Wang and Zeng, 2012; Decker et al., 2011). It is well known that inaccuracies existing in reanalysis forcing data may have substantial impacts on the simulation of land-surface energy partitioning. It is difficult to choose which reanalysis data is better for use as forcing data. Additionally,
- the spatial resolution of all of the above reanalysis/forcing datasets is not as high as that of remote sensing data. The ITPCAS forcing dataset was produced by merging a variety of data sources. This dataset benefits in particular from the merging of information from 740 weather stations operated by the China Meteorological Administration that have not been used in other forcing data. The dataset has already been used to
- ¹⁵ run land surface models and has been shown to be more accurate than other forcing datasets (Chen et al., 2011; Liu and Xie, 2013). ITPCAS meteorological forcing data include variables such as instantaneous near-surface air temperature (T_a), near-surface air pressure (P), near-surface air specific humidity (Q), near-surface wind speed (Ws) at a temporal resolution of 3 h, 3 hourly mean downward surface short-wave (SWD) ²⁰ and downward surface long-wave (LWD) radiation. The time period covered is from
- 1979 to 2010; the spatial resolution has a grid size of $0.1^{\circ} \times 0.1^{\circ}$.

3.1.2 MOD11C3 land surface temperature

MODIS (Moderate-resolution Imaging Spectroradiometer) sensors have been used to produce several global and continental scale LST datasets. MOD11C3 V5 products
 ²⁵ (Wan, 2009) are validated over a range of representative conditions with an average bias of less than 1 Kelvin (Coll et al., 2009; Wan and Li, 2008). The MOD11C3 V5 LST product has a 0.05° grid size, a monthly temporal resolution without gaps and covers the period March 2000 to October 2012. It provides monthly daytime and night-time



LST values. In our study we averaged daytime and night-time values to represent monthly means.

After interpolating MOD11C3 V5 LST to a $0.1^{\circ} \times 0.1^{\circ}$ resolution, we picked out LST values of pixels that included the 11 flux tower stations from which in-situ measurements were gathered. The pixel values were validated against the in-situ LST measurements. Detailed information about each station is given in Sect. 3.2. The linear correlation (R = 1.0), RMSE (= 1.9 K) and MB (mean value of the satellite data minus in-situ observation = 0.5 K) indicate that the quality of remotely-sensed LST data in China is high. They also show that MOD11C3 V5 LST captures the in-situ LST vari-

¹⁰ ability of different elevations and land surfaces, which is described in Sect. 4.1.

3.1.3 Albedo

Land surface albedo determines the fraction of short-wave radiation absorbed by the ground, thus influencing the surface energy budget. Studies of land-surface energy balance require temporal and spatial albedo input data without gaps. Several research ¹⁵ projects have been devoted to producing long-term time series of surface albedo from various satellite-borne sensors (Riihel et al., 2013; Muller et al., 2012; Liu et al., 2012). However most of the albedo products do not provide gap-filled time-series albedo maps. Taking MODIS MCD43B albedo product as an example, 20 to 40 % of the pixels of global landmass miss valid albedo values every year (Liu et al., 2012). Twenty ²⁰ percent invalid values in albedo input data will result in the same amount of empty val-

- ²⁰ percent invalid values in albedo input data will result in the same amount of empty values in heat flux output, an issue that limits albedo data that can be used in our study. After checking several albedo products (including GlobAlbedo (Muller et al., 2012), CMSAF cLouds, Albedo and RAdiation Surface Albedo (CLARA-SAL albedo) (Riihel et al., 2013), and MCD43B), we decided to use GlobAlbedo as its data does not contain
- spatial or temporal gaps. This albedo dataset is based on a monthly sample and has a spatial resolution of 0.05°, which we interpolated to a 0.1° resolution for our study.



3.1.4 NDVI

The Normalized Difference Vegetation Index (NDVI) is regarded as a reliable indicator of vegetation parameters. NDVI has been widely used to explore vegetation dynamics and their relationships with environmental factors (Piao et al., 2006). NDVI data from

the Systeme Pour l'Observation de la Terre (SPOT) VEGETATION sensor, distributed by Vito, have a spatial resolution of 1 km × 1 km and a temporal resolution of 10 days (synthesized on days 1, 11 and 21 of each month). In order to reduce noise resulting from clouds, the maximum NDVI value in a month for each pixel is selected to represent the canopy status of that month.

10 3.1.5 Canopy fraction

Canopy fraction (f_c) is defined as the fraction of ground surface covered by the vegetation canopy (varying from 0 to 1). f_c in SEBS is used to distinguish the contributions of vegetation and soil to the roughness parameterization. Here f_c was derived from NDVI data using the following equation:

15
$$f_{\rm c} = \frac{\rm NDVI - \rm NDVI_{min}}{\rm NDVI_{max} - \rm NDVI_{min}}.$$

3.2 Validation data

The product generated by our model needed to be validated by comparing it with an independent observational dataset. The energy balance measurement system (eddy covariance, four component radiation and ground heat flux) at flux sites is widely ac-

²⁰ covariance, four component radiation and ground heat flux) at flux sites is widely accepted as a method for direct measurement of energy and fluxes and is widely applied for assessing global evapotranspiration products (Zhang et al., 2010; Jung et al., 2011; Yan et al., 2012; Fisher et al., 2008).

To validate the product, we compiled a dataset from 11 flux stations in China with land ²⁵ cover types including bare soil, alpine meadow, forest, cropland, orchard, grassland,

(9)

and wetlands. Elevations of these stations range from 5 to 4800 m. The observational dataset includes data from Maqu (MQ) (Chen et al., 2013b; Wang et al., 2013), Wenjiang (WJ) (Zhang et al., 2012), Bijie (BJ) (Ma et al., 2006), Miyun (MY) (S. M. Liu et al., 2013), Daxing (DX) (S. M. Liu et al., 2013), Guantao (GT) (S. M. Liu et al., 2011,

⁵ 2013), Yucheng (YC) (Flerchinger et al., 2009), Dongtan (DT) (Zhao et al., 2009), SC (Semi-Arid Climate and Environment Observatory of Lanzhou University) (Huang et al., 2008; Wang et al., 2010; Guan et al., 2009), and Weishan (WS) stations (Lei and Yang, 2010a, b). Detailed information about each site is listed in Table 2.

Half-hourly fluxes were processed using standardized quality control procedures,

¹⁰ which are described in the literature references for each station. The half-hourly *H*, LE, and four component radiation were then averaged to monthly values. Monthly average values derived from less than 70 % of the flux data in each month were not used in the validations. Gap filling was not used for the flux measurement data.

4 Results

4.1 Validation against flux tower measurements

The accuracy of remote-sensing-based land-surface heat fluxes is questionable without validation against ground-based measurements (Meir and Woodward, 2010). This subsection describes the validation of the SEBS model against heat flux measurements from a diverse range of climates.

In order to analyze the source of flux calculation errors, variables related to surface radiation fluxes were all validated against flux station observations. Table 3 shows that H and LE have RMSE values slightly less than 22 W m⁻², which is lower than the RMSE values of products of other statistical methods (see Table 7 in Wang et al., 2007 and Table 5 in Jiménez et al., 2009). Indeed, Kalma et al. (2008) assessed 30 published LE validation results obtained by using ground flux measurements and reported an



average RMSE value of about 50 W m⁻² and relative errors of 15–30 %. The RMSE of our LE dataset is significantly lower than their averaged RMSE value.

We also compared our validation results with that of other, similar products produced by a previous version of SEBS. Vinukollu et al. (2011b), for instance, produced global

- ⁵ land surface fluxes with RMSE values of 40.5 W m⁻² (sensible flux) and 26.1 W m⁻² (latent flux) (calculated from Table 4 in Vinukollu et al., 2011b), which are larger than those in our study. Table 3 lists the values of the statistical parameters for the validation of a data product produced by GLDAS (which has the highest spatial resolution compared with other available terrestrial energy-flux datasets) against the same measurements
- from the Chinese flux stations as used in our study. According to the mean values of the statistical variables, the quality of our flux dataset is comparable to GLDAS' model and data assimilation results. These comparisons of accuracy demonstrate that our revised model is efficient for producing a high-resolution dataset of land-surface energy fluxes for China.
- ¹⁵ Net radiation has relatively higher RMSE and MB values than *H*, LE and G_0 in the dataset because its accuracy is dependent on the accuracy of the other variable estimates (albedo, LST, SWD, LWD, LWU, etc.). Any errors in these variables can cause bias in net radiation. LWD, for example, has a linear-fitting slope value of 0.9, with most points located around the fitting line. The correlation coefficient is as high as 0.98, thus
- ²⁰ demonstrating that there is still room for improvement of the LWD algorithms. LWD in ITPCAS was calculated with algorithms developed from measurements from across the Tibetan Plateau. The LWD algorithms may not, therefore, be accurate for other parts of China (K. Yang, personal communication, 2013). This underlines the need for more accurate LWD radiation fluxes in order to improve the accuracy of turbulent fluxes and evapotranspiration.
 - In addition to the statistical evaluation of model results against observations, seasonal and inter-annual changes in the model results also need to be checked. Yucheng station, which is an agricultural experimental station with winter wheat and summer maize as dominant crops was taken as an example (Fig. 2). Crops at Yucheng station



mature twice per year, which is representative of warm temperate farming cropland, typical for the North China Plain. A two-year flux dataset was used to compare against values extracted from our model-derived product. The inter-annual and seasonal LST and LWU data closely match the in-situ observations. The SWD term also successfully

- ⁵ captures seasonal variations. LWD is systematically lower than observations. The LE produced at Yucheng station not only captures seasonal variation, but also responds at step stages, which occur when the wheat is harvested or maize seeds have just been sown (from June to August). The increased sensible heat and decreased latent heat flux observed in July 2003 were caused by the wheat harvest, however this signal
- change is not captured by the model result. The simulated sensible and latent heat produced by SEBS has a one-month lag when compared to reality. This phenomenon is caused by adopting a maximum monthly NDVI value, resulting in faulty representation of canopy status changes in the month of June.
- The Semi-Arid Climate and Environment Observatory of Lanzhou University (SC station) is situated on China's Loess Plateau, at 1965.8 m above sea level. Annual mean precipitation there is 381.1 mm and annual evapotranspiration is 1528.5 mm (Huang et al., 2008). Being typical of stations operating under arid conditions, its flux measurements were compared with the grid point values extracted from the model product (Fig. 3). In 2008 the land surface around the station was covered by snow from 19
 January to 20 February. Consequently the GlobAlbedo value was high for February.
- Unexpectedly, albedo was relatively low for January, which could be caused by the coarse temporal sampling of the station pixel by the satellite sensor. The calculated monthly sensible heat and latent heat in January 2008 have biases of -11.7 (with an observed monthly mean sensible heat = 15 W m^{-2}) and -7.6 W m^{-2} (with an observed monthly mean latent heat = 4.8 W m^{-2}), respectively. The relatively large bias for SC
- station when covered with snow may be caused by the mixed pixel around the station.

The results of other stations have been included in Supplement submitted with this paper. Comparison with the results of these other stations shows that model estimates of surface energy balance variables match the magnitude and seasonal variation



observed at stations in several contrasting ecosystems. Comparisons between the fluxtower-measured and the modeled fluxes show that latent fluxes were more accurate than sensible fluxes. Comparisons with other studies, which are presented in Table 4, show that the accuracy of our dataset is one of the best among high-resolution datasets of land surface fluxes.

4.2 Spatial distribution of land-surface energy fluxes

Using maps of average annual land-surface radiation and energy fluxes, we analyzed the spatial patterns of radiation and energy fluxes for the Chinese landmass and compared them with other products, such as GLDAS. The highest values of downward surface solar radiation (Fig. 4a) are located in the southwest of the Tibetan Plateau, while the lowest values occur in the Sichuan Basin (SB). The highest levels of upward short-wave radiation (Fig. 4c) occur around the snow-covered peaks of the Himalaya (HM), Karakorum (KRM) and Kunlun (KLM), and the Qilian (QLM) and Nyainqentanglha (NQM) mountain ranges. The strongest net solar radiation (SWD minus SWU) on

the Chinese landmass occurs in the southern part of the Tibetan Plateau (see Supplement). The downward and upward long-wave radiation (Fig. 4b and c) on the Tibetan Plateau are the lowest for the entire Chinese landmass. Southern China has the highest levels of upward and downward long-wave radiation. The highest values of net long-wave radiation (LWU minus LWD) occur in the southern and western parts of the Tibetan Plateau (see Supplement).

Figure 5 shows that northwestern China (NWC), the western Tibetan Plateau (TP), the inner Mongolian Plateau (MP) and the Loess Plateau (LP) have the highest yearly average values for surface sensible-heat flux. Croplands of the northern China Plain (NCP, including the lowlands of Shandong, Henan, and Hebei provinces) and the north-

eastern China Plain (NEP, including the lowlands of Liaoning, Jilin, and Heilongjiang provinces) have low average yearly values for sensible heat flux. The Pearl River delta (PRD) and Tarim (TRB) and Sichuan (SCB) basins also have low levels of sensible heat



flux, as do the Yinchuan (YCB) and the inner Mongolian basins (IMB) along the Yellow River. This spatial distribution is consistent with GLDAS results (see Supplement).

Simulated annual latent heat fluxes (Fig. 5b) exhibit a southeast to northwest decreasing gradient, which is consistent with other studies (Y. Liu et al., 2013b). The

- ⁵ southeastern Tibetan Plateau has high levels of annual latent heat flux. The Gobi desert, in the northwest of China (NWC), has the lowest annual latent heat flux, followed by the western Tibetan Plateau and the inner Mongolian Plateau (MP). Lake regions along the Yangtze River and the region of basins along the Yellow River have relatively high levels of latent heat flux.
- ¹⁰ The highest levels of annual average surface net radiation (Fig. 5c) can be found in southwestern China and the Lasha Basin (LB); the lowest levels occur in the Sichuan (SCB) and Junggar Basins (JB). The highest levels of annual average ground-heat flux (Fig. 5c) are to be found in western China, due to large amounts of incoming solar radiation that occur under dry conditions. The monthly average of G_0 is negligible when ¹⁵ compared with other fluxes.
- The role of plateau heating on Asia's monsoons is being discussed vigorously (Qiu, 2013; Wu et al., 2012; Boos and Kuang, 2010). Figure 6 shows seasonal comparisons of *H* between boreal winter (DJF), spring (MAM), summer (JJA) and autumn (SON). The largest area of positive sensible heating occurs in spring. Lee et al. (2011) have
 ²⁰ shown that contrasting sensible heat fluxes between the Chinese landmass and the seas surrounding it during the pre-monsoon period (April–May) affect monsoon development in East Asia. Figure 6a shows that sources of sensible heating in spring occur over the Tibetan and several other plateaus in China. During summer, the highest sensible heat fluxes are to be found on the western Tibetan Plateau, the eastern Loess
 ²⁵ Plateau (LP) and in northwestern China (NWC).

LE in summer has the largest area of high latent heating, followed by that in spring, autumn and winter (Fig. 7). Latent heat in summer is highest in southeastern and southern China as a result of abundant rainfall in these regions. Similarly on irrigated land, such as that found in Yinchuan (YB), the inner Mongolian basin (IMB) and the



downstream Fbasins of the Tianshan (TM) and Kunlun (KLM) mountains, latent heat and evapotranspiration are high due to the ample supply of water in summer. Latent heat fluxes in autumn and winter are significantly lower than those of the other two seasons. The magnitudes and spatial patterns of LE in China of our product are generally consistent with other reports (Yao et al., 2013; Mu et al., 2007; Jung et al., 2010).

Net radiation in summer has the highest values of the four seasons. Most of the Chinese landmass acts as a source of surface energy for the atmosphere (Fig. 8).

4.3 Trend analysis

The ability to capture the inter- and intra-annual variation for each land-surface energy variable is of interest to researchers of monsoon phenomena and climate change (Zhu et al., 2012). Indeed, understanding these variations is essential for studies on climate change and water-resource-related issues. We have calculated annual average values for each flux variable. The nonparametric Mann–Kendall test (MK) is one of the most widely used methods for hydro-meteorological time series analysis (Z. Liu et al.,

¹⁵ 2013; Gan, 1998). The MK method was applied to the series of annual average fluxes to check variations during the period 2001–2010. The resulting slope indicates that downward surface short-wave radiation increased during that decade over the majority of the Tibetan Plateau (Fig. 9).

The ground solar measurements at China Meteorological Administration (CMA) sta-

- tions during 2003–2006, as shown in Fig. 1b of Yang et al. (2012), confirms the increasing trend of downward surface short-wave radiation found in our study. The annual mean visibility measured at these stations also displays an increasing trend (Fig. 2a of Yang et al., 2012), while ERA-40 reanalyzed precipitable-water and station-observed specific humidity show a decreasing trend from 2000 to 2006 (Fig. 3a of Yang
- et al., 2012). These results indicate that the atmosphere over the plateau is becoming drier, which would explain why SWD has increased during the decade.

The upward short-wave radiation over the Himalaya (HM), the Ganges (GM), the Karakorum (KRM), and the Qilian (QLM) and Nyainqentanglha (NQM) mountain ranges



has also increased over the last 10 years, which may be caused by the glacial retreat that has occurred in these areas (Scherler et al., 2011; Yao et al., 2004). Lasha basin (LB) has the steepest rising trend in LWU, perhaps because of the relatively greater degree of anthropogenic (e.g. urbanization) activity occurring in this area. The trend analysis did not reveal any clear spatial pattern in downward long-wave radiation. Net 5 radiation over several high mountain ranges (including the Himalaya, the Ganges, the Karakorum and the Qilian and Nyaingentanglha mountain ranges) increased by approximately 5 W m⁻² between 2001 and 2010 (Fig. 10). The strongest increase in net radiation occurred in the central part of the Tibetan Plateau. As Matthew (2010) has pointed out, soil moisture in the central Tibetan Plateau showed an increasing trend 10 from 1987 to 2008. Wetter soil can cause the ground surface to absorb more net radiation and thus increase latent heat flux. Moreover, wetter soil can increase soil heating capacity (Guan et al., 2009) and so further increase ground heat flux. The increases in net radiation and soil moisture may also explain a rising trend in latent heat in the central Tibetan Plateau. Clearly, the plateau is experiencing accelerated environmental 15 changes (Zhong et al., 2011; Salama et al., 2012). Indeed, land-surface radiation and energy trend analyses also show that the Tibetan Plateau is experiencing a relatively stronger change in land-surface radiation (verified by Tang et al., 2011) and energy exchange than other parts of China.

20 5 Conclusions and discussion

In view of China's highly fragmented landscape, high-resolution land-surface heat flux maps are necessary for hydrological studies. As China includes arid, semi-arid, humid, and semi-humid regions, quantifying its water and energy budgets is a challenge. We have developed the surface energy balance system (SEBS) further to produce

a land-surface heat flux dataset at a continental scale of higher resolution than datasets derived using other methods. In summary, using remote sensing data and surface meteorological information, a data product of monthly resolution has been developed for



land-surface heat flux analysis. We have validated our remote-sensing-based approach with in-situ observations from 11 flux stations in China. Taking into account the limitations of available spatial data and computing resources, we applied the model to the entire Chinese landmass using a 0.1° resolution meteorological dataset, MODIS LST, vegetation indices and other variables to generate a climatological dataset of land-

- surface energy balance for a 10 year period. The modeling results for both pixel-point and spatial distribution demonstrate that this approach meets our aims in terms of (a) being robust across a variety of land cover and climate types and (b) performing well for the temporal and spatial scales of interest. The spatial distribution maps generated
- for each variable of surface energy balance give important background information on the terrestrial hydrology and energy cycles. This product also demonstrates the impact of topography and climatic conditions on land–air energy and moisture exchanges in China.
- The applicability of remote-sensing-based estimates of land surface fluxes is ham-¹⁵ pered by limited temporal coverage of satellite sensors (Ryu et al., 2012). Remote sensing data are snapshots of the land surface status at a particular point in space and time (Ryu et al., 2011). It is challenging to compare remote-sensing-based monthly flux data with ground measurements that are made on time scales ranging from half-hourly through to monthly.
- The energy flux product we have developed has a spatial resolution of approximately 10 km, while flux towers have a footprint of tens to hundreds of meters. The tower footprint may not be representative of the larger pixel of the product, and this mismatch will result in errors if the mean of the satellite pixel is different from that of the flux tower footprint. Remote-sensing-based studies stress that direct comparison is a challenge
- ²⁵ because scale mismatch (Norman et al., 2003) and heterogeneity of the land surface reduce the spatial representativeness of ground-site measurements (Mi et al., 2006). Another challenge is validating the grid-box-based simulation results on the scale of the Chinese landmass, since reliable observations of flux data are only available from a few sites in the simulated region.



Potential effects of changes in turbulent heat fluxes on the monsoon over East Asia (Lee et al., 2011) as a result of China's recent urbanization can be studied further using our product. As an independent satellite-based product, it can also be used as a data source for evaluating land surface models. The product can be downloaded from the URL upon the request to the contact author: https://drive.google.com/folderview?id= B7yGrB1U9eDec2JFbnA5eldlVHc&usp=sharing.

The Supplement related to this article is available online at doi:10.5194/acpd-14-14471-2014-supplement.

Acknowledgements. This study was supported by the Chinese National Key Program for De veloping Basic Sciences (2010CB951701), the Chinese National Natural Science Foundation (41275010), CAS-KNAW joint Ph.D. researcher project, the ESA WACMOS project and the FP7 CORE-CLIMAX project. The forcing dataset used in our study was developed at the Data Assimilation and Modeling Center for Tibetan Multi-spheres, Institute of Tibetan Plateau Research, Chinese Academy of Sciences. We thank Y. Kun for his comments during the writing of this paper.

this paper. For our study we used eddy covariance data acquired from the scientific community and networks. We acknowledge Wenjiang, Ali, Yucheng, and Weishan stations, Nagqu Station of Plateau Climate and Environment, Magqu Zoige Plateau Wetlands Ecosystem Research Station, and the Semi-Arid Climate and Environment Observatory of Lanzhou University for providing their in-situ measurement datasets. We also acknowledge X. Xu (CMA), L. Tian (ITP, CAS), Y. Zhang (CAREERI, CAS), S. Xu (IGSNNR, CAS), B. Zhao (Fudan University), and H. Lei (Tsinghua University) for providing us with their flux datasets.

References

Allen, R., Irmak, A., Trezza, R., Hendrickx, J. M. H., Bastiaanssen, W., and Kjaersgaard, J.: Satellite-based ET estimation in agriculture using SEBAL and METRIC, Hydrol. Process., 25, 4011–4027, doi:10.1002/hyp.8408, 2011.



25

- 14494
- Collection-5 MODIS Land Surface Temperature Products Users' Guide: available at: http://www. icess.ucsb.edu/modis/LstUsrGuide/usrguide.html (last access: July 2013), 2009.

of the V5 MODIS land surface temperature product, J. Geophys. Res., 114, D20102,

- ²⁵ Chen, Y., Yang, K., Qin, J., Zhao, L., Tang, W., and Han, M.: Evaluation of AMSR-E retrievals and GLDAS simulations against observations of a soil moisture network on the central Tibetan Plateau, J. Geophys. Res.-Atmos., 118, 4466–4475, doi:10.1002/jgrd.50301, 2013c. Coll, C., Wan, Z., and Galve, J. M.: Temperature-based and radiance-based validations
- Chen, Y., Yang, K., He, J., Qin, J., Shi, J., Du, J., and He, Q.: Improving land surface temperature modeling for dry land of China, J. Geophys. Res., 116, D20104, doi:10.1029/2011jd015921, 2011.
- 1607–1618, doi:10.5194/hess-17-1607-2013, 2013a.
 Chen, X., Su, Z., Ma, Y., Yang, K., Wen, J., and Zhang, Y.: An improvement of roughness height parameterization of the Surface Energy Balance System (SEBS) over the Tibetan Plateau, J. Appl. Meteorol. Clim., 52, 607–622, doi:10.1175/jamc-d-12-056.1, 2013b.
- Remote Sens. Environ., 91, 332–344, doi:10.1016/j.rse.2004.03.014, 2004.
 Chen, X., Su, Z., Ma, Y., Yang, K., and Wang, B.: Estimation of surface energy fluxes under complex terrain of Mt. Qomolangma over the Tibetan Plateau, Hydrol. Earth Syst. Sci., 17,
- Sci., 14, 2643–2659, doi:10.5194/hess-14-2643-2010, 2010.
 Chen, J., Jönsson, P., Tamura, M., Gu, Z., Matsushita, B., and Eklundh, L.: A simple method for reconstructing a high-quality NDVI time-series data set based on the Savitzky–Golay filter, Density 2004 10 2014.
- Minacapilli, M.: The impact of in-canopy wind profile formulations on heat flux estimation in an open orchard using the remote sensing-based two-source model, Hydrol. Earth Syst.
- Brutsaert, W.: Aspects of bulk atmospheric boundary layer similarity under free-convective conditions, Rev. Geophys., 37, 439–451, doi:10.1029/1999rg900013, 1999.
 Cammalleri, C., Anderson, M. C., Ciraolo, G., D'Urso, G., Kustas, W. P., La Loggia, G., and
- lation versus plateau heating, Nature, 463, 218–222, doi:10.1038/nature08707, 2010. Brauman, K., Daily, G., Duarte, T., and Mooney, H.: The nature and value of ecosystem services: an overview highlighting hydrologic services, in: Annual Review of Environment and

5

10

30

98, 2007.

doi:10.1029/2009jd012038, 2009.

Resources, Annual Review of Environment and Resources, Annual Reviews, Palo Alto, 67-

Boos, W. R. and Kuang, Z.: Dominant control of the South Asian monsoon by orographic insu-



Discussion

Paper

Discussion

Paper

Discussion Paper

Discussion Paper

10 years China land-surface energy balance dataset

X. Chen et al.





Decker, M., Brunke, M. A., Wang, Z., Sakaguchi, K., Zeng, X., and Bosilovich, M. G.: Evaluation of the reanalysis products from GSFC, NCEP, and ECMWF using flux tower observations, J. Climate, 25, 1916–1944, doi:10.1175/jcli-d-11-00004.1, 2011.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,

- Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration
- and performance of the data assimilation system, Q. J. Roy. Meteor. Soc., 137, 553–597, doi:10.1002/qj.828, 2011.
 - Dirmeyer, P. A., Gao, X., Zhao, M., Guo, Z., Oki, T., and Hanasaki, N.: GSWP-2: multimodel analysis and implications for our perception of the land surface, B. Am. Meteorol. Soc., 87, 1381–1397, doi:10.1175/bams-87-10-1381, 2006.
- Fan, L., Liu, S., Bernhofer, C., Liu, H., and Berger, F. H.: Regional land surface energy fluxes by satellite remote sensing in the Upper Xilin River Watershed (Inner Mongolia, China), Theor. Appl. Climatol., 88, 231–245, doi:10.1007/s00704-006-0241-9, 2007.
 - Fisher, J. B., Tu, K. P., and Baldocchi, D. D.: Global estimates of the land–atmosphere water flux based on monthly AVHRR and ISLSCP-II data, validated at 16 FLUXNET sites, Remote Sens. Environ., 112, 901–919, doi:10.1016/j.rse.2007.06.025, 2008.
 - Flerchinger, G., Xaio, W., Marks, D., Sauer, T., and Yu, Q.: Comparison of algorithms for incoming atmospheric long-wave radiation, Water Resour. Res., 45, W03423, doi:10.1029/2008wr007394, 2009.
 - Gan, T. Y.: Hydroclimatic trends and possible climatic warming in the Canadian Prairies, Water Resour. Res., 34, 3009–3015, doi:10.1029/98wr01265, 1998.
- Goudriaan, J.: Crop Micrometeorology: a Simulation Study, Ph.D. thesis, Wageningen University, Wageningen, 249 pp., 1977.
 - Guan, X., Huang, J., Guo, N., Bi, J., and Wang, G.: Variability of soil moisture and its relationship with surface albedo and soil thermal parameters over the Loess Plateau, Adv. Atmos. Sci.,
- ³⁰ 26, 692–700, doi:10.1007/s00376-009-8198-0, 2009.

20

25

He, J.: Development of Surface Meteorological Dataset of China with High Temporal and Spatial Resolution, M.S., Inst. of Tibetan Plateau Res., Chin. Acad. of Sci., Beijing, China, 2010.



- Hsu, H.-H., and Liu, X.: Relationship between the Tibetan Plateau heating and East Asian summer monsoon rainfall, Geophys. Res. Lett., 30, 2066, doi:10.1029/2003gl017909, 2003.
- Huang, J., Zhang, W., Zuo, J., Bi, J., Shi, J., Wang, X., Chang, Z., Huang, Z., Yang, S., Zhang, B., Wang, G., Feng, G., Yuan, J., Zhang, L., Zuo, H., Wang, S., Fu, C., and Jifan, C.:
- ⁵ An overview of the semi-arid climate and environment research observatory over the Loess Plateau, Adv. Atmos. Sci., 25, 906–921, 2008.
 - Jia, Z., Liu, S., Xu, Z., Chen, Y., and Zhu, M.: Validation of remotely sensed evapotranspiration over the Hai River Basin, China, J. Geophys. Res., 117, D13113, doi:10.1029/2011jd017037, 2012.
- Jiménez, C., Prigent, C., and Aires, F.: Toward an estimation of global land surface heat fluxes from multisatellite observations, J. Geophys. Res., 114, D06305, doi:10.1029/2008jd011392, 2009.
 - Jiménez, C., Prigent, C., Mueller, B., Seneviratne, S. I., McCabe, M. F., Wood, E. F., Rossow, W. B., Balsamo, G., Betts, A. K., Dirmeyer, P. A., Fisher, J. B., Jung, M., Kana-
- ¹⁵ mitsu, M., Reichle, R. H., Reichstein, M., Rodell, M., Sheffield, J., Tu, K., and Wang, K.: Global intercomparison of 12 land surface heat flux estimates, J. Geophys. Res., 116, D02102, doi:10.1029/2010jd014545, 2011.
 - Jung, M., Reichstein, M., and Bondeau, A.: Towards global empirical upscaling of FLUXNET eddy covariance observations: validation of a model tree ensemble approach using a bio-sphere model, Biogeosciences, 6, 2001–2013, doi:10.5194/bg-6-2001-2009, 2009.
- sphere model, Biogeosciences, 6, 2001–2013, doi:10.5194/bg-6-2001-2009, 2009. Jung, M., Reichstein, M., Ciais, P., Seneviratne, S. I., Sheffield, J., Goulden, M. L., Bonan, G., Cescatti, A., Chen, J., de Jeu, R., Dolman, A. J., Eugster, W., Gerten, D., Gianelle, D., Gobron, N., Heinke, J., Kimball, J., Law, B. E., Montagnani, L., Mu, Q., Mueller, B., Oleson, K., Papale, D., Richardson, A. D., Roupsard, O., Running, S., Tomelleri, E., Viovy, N., Weber, U.,
- ²⁵ Williams, C., Wood, E., Zaehle, S., and Zhang, K.: Recent decline in the global land evapotranspiration trend due to limited moisture supply, Nature, 467, 951–954, 2010.
 - Jung, M., Reichstein, M., Margolis, H. A., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D.,
- Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, J. Geophys. Res., 116, G00J07, doi:10.1029/2010jg001566, 2011.



Kalma, J., McVicar, T., and McCabe, M.: Estimating land surface evaporation: a review of methods using remotely sensed surface temperature data, Surv. Geophys., 29, 421–469, doi:10.1007/s10712-008-9037-z, 2008.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M.,

Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40 year reanalysis project, B. Am. Meteorol. Soc., 77, 437–471, doi:10.1175/1520-0477(1996)077<0437:tnyrp>2.0.co;2, 1996.

Lee, E., Barford, C., Kucharik, C., Felzer, B., and Foley, J.: Role of turbulent heat fluxes over

- ¹⁰ land in the monsoon over East Asia, International Journal of Geosciences, 2 420–431, doi:10.4236/ijg.2011.24046a, 2011.
 - Lei, H. and Yang, D.: Seasonal and interannual variations in carbon dioxide exchange over a cropland in the North China Plain, Glob. Change Biol., 16, 2944–2957, doi:10.1111/j.1365-2486.2009.02136.x, 2010a.
- Lei, H. and Yang, D.: Interannual and seasonal variability in evapotranspiration and energy partitioning over an irrigated cropland in the North China Plain, Agr. Forest Meteorol., 150, 581–589, doi:10.1016/j.agrformet.2010.01.022, 2010b.
 - Li, X., Li, X., Li, Z., Ma, M., Wang, J., Xiao, Q., Liu, Q., Che, T., Chen, E., Yan, G., Hu, Z., Zhang, L., Chu, R., Su, P., Liu, Q., Liu, S., Wang, J., Niu, Z., Chen, Y., Jin, R., Wang, W.,
- Ran, Y., Xin, X., and Ren, H.: Watershed allied telemetry experimental research, J. Geophys. Res.-Atmos., 114, D22103, doi:10.1029/2008jd011590, 2009.
 - Li, X., Liang, S., Yuan, W., Yu, G., Cheng, X., Chen, Y., Zhao, T., Feng, J., Ma, Z., Ma, M., Liu, S., Chen, J., Shao, C., Li, S., Zhang, X., Zhang, Z., Sun, G., Chen, S., Ohta, T., Varlagin, A., Miyata, A., Takagi, K., Saiqusa, N., and Kato, T.: Estimation of evapotranspiration over the terrestrial ecosystems in China, Ecohydrology, 7, 139–149, doi:10.1002/eco.1341, 2012.
 - Li, X., Wang, L., Chen, D., Yang, K., Xue, B., and Sun, L.: Near-surface air temperature lapse rates in the mainland China during 1962–2011, J. Geophys. Res.-Atmos., 118, 7505–7515, doi:10.1002/jgrd.50553, 2013.

25

Li, Z., Zheng, F.-L., and Liu, W.-Z.: Spatiotemporal characteristics of reference evapotranspira-

tion during 1961–2009 and its projected changes during 2011–2099 on the Loess Plateau of China, Agr. Forest Meteorol., 154–155, 147–155, doi:10.1016/j.agrformet.2011.10.019, 2012.



- Lin, W., Zhang, L., Du, D., Yang, L., Lin, H., Zhang, Y., and Li, J.: Quantification of land use/land cover changes in Pearl River Delta and its impact on regional climate in summer using numerical modeling, Reg. Environ. Change, 9, 75–82, doi:10.1007/s10113-008-0057-5, 2009.
- Liu, R., Wen, J., Wang, X., Wang, L., Tian, H., Zhang, T. T., Shi, X. K., Zhang, J. H., and
 Lv, SH. N.: Actual daily evapotranspiration estimated from MERIS and AATSR data over the Chinese Loess Plateau, Hydrol. Earth Syst. Sci., 14, 47–58, doi:10.5194/hess-14-47-2010, 2010.
 - Liu, H. Z., Feng, J. W., Järvi, L., and Vesala, T.: Four-year (2006–2009) eddy covariance measurements of CO₂ flux over an urban area in Beijing, Atmos. Chem. Phys., 12, 7881–7892, doi:10.5194/acp-12-7881-2012, 2012.
- Liu, J.-G. and Xie, Z.-H.: Improving simulation of soil moisture in China using a multiple meteorological forcing ensemble approach, Hydrol. Earth Syst. Sci., 17, 3355–3369, doi:10.5194/hess-17-3355-2013, 2013.

10

20

- Liu, N. F., Liu, Q., Wang, L. Z., Liang, S. L., Wen, J. G., Qu, Y., and Liu, S. H.: A statistics-based temporal filter algorithm to map spatiotemporally continuous shortwave albedo from MODIS data, Hydrol. Earth Syst. Sci., 17, 2121–2129, doi:10.5194/hess-17-2121-2013, 2013.
 - Liu, S. M., Xu, Z. W., Wang, W. Z., Jia, Z. Z., Zhu, M. J., Bai, J., and Wang, J. M.: A comparison of eddy-covariance and large aperture scintillometer measurements with respect to the energy balance closure problem, Hydrol. Earth Syst. Sci., 15, 1291–1306, doi:10.5194/hess-15-1291-2011, 2011.
 - Liu, S. M., Xu, Z. W., Zhu, Z. L., Jia, Z. Z., and Zhu, M. J.: Measurements of evapotranspiration from eddy-covariance systems and large aperture scintillometers in the Hai River Basin, China, J. Hydrol., 487, 24–38, doi:10.1016/j.jhydrol.2013.02.025, 2013.

Ma, L., Zhang, T., Li, Q., Frauenfeld, O. W., and Qin, D.: Evaluation of ERA-40, NCEP-1, and

- NCEP-2 reanalysis air temperatures with ground-based measurements in China, J. Geophys. Res., 113, D15115, doi:10.1029/2007jd009549, 2008a.
 - Ma, Y., Su, Z., Li, Z., Koike, T., and Menenti, M.: Determination of regional net radiation and soil heat flux over a heterogeneous landscape of the Tibetan Plateau, Hydrol. Process., 16, 2963–2971, 2002.
- Ma, Y., Zhong, L., Su, Z., Ishikawa, H., Menenti, M., and Koike, T.: Determination of regional distributions and seasonal variations of land surface heat fluxes from Landsat-7 Enhanced Thematic Mapper data over the central Tibetan Plateau area, J. Geophys. Res., 111, D10305, doi:10.1029/2005jd006742, 2006.



14498

14499

- Ma, Y., Kang, S., Zhu, L., Xu, B., Tian, L., and Yao, T.: Tibetan Observation and Research Platform – atmosphere–land interaction over a heterogeneous landscape, B. Am. Meteorol. Soc., 89, 1487–1492, doi:10.1175/2008BAMS2545.1, 2008.
- Ma, Y., Zhong, L., Wang, B., Ma, W., Chen, X., and Li, M.: Determination of land surface heat fluxes over heterogeneous landscape of the Tibetan Plateau by using the MODIS and in situ
 - data, Atmos. Chem. Phys., 11, 10461–10469, doi:10.5194/acp-11-10461-2011, 2011. Massman, W. J.: An analytical one-dimensional second-order closure model of turbulence statistics and the lagrangian time scale within and above plant canopies of arbitrary structure,
 - Bound.-Lay. Meteorol., 83, 407–421, 1997.

15

- Matthew, O.: Characterization of the Effects of Climate Variation on Land Surface Temperature and Soil Moisture Through Stochastic Analysis of Long Term SSM/I Observations Over the Tibetan Plateau, Master, International Institute for Geo-information Science and Earth Observation, University of Twente, Enschede, the Netherlands, 1–67, 2010.
 - Meir, P. and Woodward, F. I.: Amazonian rain forests and drought: response and vulnerability, New Phytol., 187, 553–557, doi:10.1111/j.1469-8137.2010.03390.x, 2010.
 - Mi, N., Yu, G. R., Wang, P. X., Wen, X. F., and Sun, X. M.: A preliminary study for spatial representiveness of flux observation at ChinaFLUX sites, Sci. China Ser. D, 49, 24–35, doi:10.1007/s11430-006-8024-9, 2006.

Moody, E. G., King, M. D., Platnick, S., Schaaf, C. B., and Feng, G.: Spatially complete global

- spectral surface albedos: value-added datasets derived from Terra MODIS land products, IEEE T. Geosci. Remote, 43, 144–158, doi:10.1109/tgrs.2004.838359, 2005.
 - Mu, Q., Heinsch, F. A., Zhao, M., and Running, S. W.: Development of a global evapotranspiration algorithm based on MODIS and global meteorology data, Remote Sens. Environ., 111, 519–536, doi:10.1016/j.rse.2007.04.015, 2007.
- ²⁵ Mueller, B., Seneviratne, S. I., Jimenez, C., Corti, T., Hirschi, M., Balsamo, G., Ciais, P., Dirmeyer, P., Fisher, J. B., Guo, Z., Jung, M., Maignan, F., McCabe, M. F., Reichle, R., Reichstein, M., Rodell, M., Sheffield, J., Teuling, A. J., Wang, K., Wood, E. F., and Zhang, Y.: Evaluation of global observations-based evapotranspiration datasets and IPCC AR4 simulations, Geophys. Res. Lett., 38, L06402, doi:10.1029/2010gl046230, 2011.
- Muller, J.-P., López, G., Watson, G., Shane, N., Kennedy, T., Yuen, P., Lewis, P., Fischer, J., Guanter, L., Domench, C., Preusker, R., North, P., Heckel, A., Danne, O., and Krämer, U.: The ESA GlobAlbedo Project for mapping the Earth's land surface albedo for 15 years from



European Sensors, IEEE Geoscience and Remote Sensing Symposium (IGARSS), IEEE, Munich, Germany, 22–27.7.12, 2012.

- Norman, J. M., Anderson, M. C., Kustas, W. P., French, A. N., Mecikalski, J., Torn, R., Diak, G. R., Schmugge, T. J., and Tanner, B. C. W.: Remote sensing of surface energy fluxes
- at 101 m pixel resolutions, Water Resour. Res., 39, 1221, doi:10.1029/2002wr001775, 2003.
 Piao, S., Mohammat, A., Fang, J., Cai, Q., and Feng, J.: NDVI-based increase in growth of temperate grasslands and its responses to climate changes in China, Global Environ. Chang., 16, 340–348, doi:10.1016/j.gloenvcha.2006.02.002, 2006.

Qiu, J.: Monsoon Melee, Science, 340, 1400–1401, doi:10.1126/science.340.6139.1400, 2013.

- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's modern-era retrospective analysis for research and applications, J. Climate. 24, 3624–3648. doi:10.1175/icli-d-11-00015.1.2011.
- Riihelä, A., Manninen, T., Laine, V., Andersson, K., and Kaspar, F.: CLARA-SAL: a global 28 yr timeseries of Earth's black-sky surface albedo, Atmos. Chem. Phys., 13, 3743–3762, doi:10.5194/acp-13-3743-2013, 2013.

Roads, J. and Betts, A.: NCEP-NCAR and ECMWF reanalysis surface water and energy

- ²⁰ budgets for the Mississippi River Basin, J. Hydrometeorol., 1, 88–94, doi:10.1175/1525-7541(2000)001<0088:nnaers>2.0.co;2, 2000.
 - Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C. J., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D., and Toll, D.: The Global Land Data Assimilation System, B. Am. Meteorol. Soc., 85, 381–394, doi:10.1175/bams-85-3-381. 2004.
- Ryu, Y., Baldocchi, D. D., Kobayashi, H., van Ingen, C., Li, J., Black, T. A., Beringer, J., van Gorsel, E., Knohl, A., Law, B. E., and Roupsard, O.: Integration of MODIS land and atmosphere products with a coupled-process model to estimate gross primary productivity and evapotranspiration from 1 km to global scales, Global Biogeochem. Cy., 25, GB4017, doi:10.1029/2011gb004053, 2011.

25

Ryu, Y., Baldocchi, D. D., Black, T. A., Detto, M., Law, B. E., Leuning, R., Miyata, A., Reichstein, M., Vargas, R., Ammann, C., Beringer, J., Flanagan, L. B., Gu, L., Hutley, L. B., Kim, J., McCaughey, H., Moors, E. J., Rambal, S., and Vesala, T.: On the temporal upscaling of evap-



otranspiration from instantaneous remote sensing measurements to 8 day mean daily-sums, Agr. Forest Meteorol., 152, 212–222, doi:10.1016/j.agrformet.2011.09.010, 2012.

- Salama, M. S., Velde, R., Zhong, L., Ma, Y., Ofwono, M., and Su, Z.: Decadal variations of land surface temperature anomalies observed over the Tibetan Plateau by the Special
- ⁵ Sensor Microwave Imager (SSM/I) from 1987 to 2008, Climatic Change, 114, 769–781, doi:10.1007/s10584-012-0427-3, 2012.

Scherler, D., Bookhagen, B., and Strecker, M. R.: Spatially variable response of Himalayan glaciers to climate change affected by debris cover, Nat. Geosci., 4, 156–159, 2011.

Sheffield, J., Goteti, G., and Wood, E. F.: Development of a 50 year high-resolution global

- dataset of meteorological forcings for land surface modeling, J. Climate, 19, 3088–3111, doi:10.1175/jcli3790.1, 2006.
 - Shu, Y., Stisen, S., Jensen, K. H., and Sandholt, I.: Estimation of regional evapotranspiration over the North China Plain using geostationary satellite data, Int. J. Appl. Earth Obs., 13, 192–206, doi:10.1016/j.jag.2010.11.002, 2011.
- ¹⁵ Simard, M., Pinto, N., Fisher, J. B., and Baccini, A.: Mapping forest canopy height globally with spaceborne lidar, J. Geophys. Res., 116, G04021, doi:10.1029/2011jg001708, 2011.
 - Su, Z.: The Surface Energy Balance System (SEBS) for estimation of turbulent heat fluxes, Hydrol. Earth Syst. Sci., 6, 85–100, doi:10.5194/hess-6-85-2002, 2002.

Su, Z., Schmugge, T., Kustas, W. P., and Massman, W. J.: An evaluation of two models for

- estimation of the roughness height for heat transfer between the land surface and the atmosphere, J. Appl. Meteorol., 40, 1933–1951, 2001.
 - Su, Z., de Rosnay, P., Wen, J., Wang, L., and Zeng, Y.: Evaluation of ECMWF's soil moisture analyses using observations on the Tibetan Plateau, J. Geophys. Res.-Atmos., 118, 5304– 5318, doi:10.1002/jgrd.50468, 2013.
- Suh, M.-S. and Lee, D.-K.: Impacts of land use/cover changes on surface climate over east Asia for extreme climate cases using RegCM2, J. Geophys. Res.-Atmos., 109, D02108, doi:10.1029/2003jd003681, 2004.

Sun, L. and Wu, G.: Influence of land evapotranspiration on climate variations, Sci. China Ser. D, 44, 838–846, doi:10.1007/bf02907096, 2001.

Tang, W.-J., Yang, K., Qin, J., Cheng, C. C. K., and He, J.: Solar radiation trend across China in recent decades: a revisit with quality-controlled data, Atmos. Chem. Phys., 11, 393–406, doi:10.5194/acp-11-393-2011, 2011.



- Taniguchi, K. and Koike, T.: Seasonal variation of cloud activity and atmospheric profiles over the eastern part of the Tibetan Plateau, J. Geophys. Res.-Atmos., 113, 10104–10104, 2008.
 Timmermans, J.: Coupling Optical and Thermal Directional Radiative Transfer to Biophysical
- Processes in Vegetated Canopies, Ph.D., Faculty of Geo-information Science and Earth Observation, University of Twente, Enchede, the Netherlands, 1–157, 2011.
- servation, University of Twente, Enchede, the Netherlands, 1–157, 2011.
 Vinukollu, R. K., Meynadier, R., Sheffield, J., and Wood, E. F.: Multi-model, multi-sensor estimates of global evapotranspiration: climatology, uncertainties and trends, Hydrol. Process., 25, 3993–4010, doi:10.1002/hyp.8393, 2011a.
- Vinukollu, R. K., Wood, E. F., Ferguson, C. R., and Fisher, J. B.: Global estimates of
 evapotranspiration for climate studies using multi-sensor remote sensing data: evaluation of three process-based approaches, Remote Sens. Environ., 115, 801–823, doi:10.1016/j.rse.2010.11.006. 2011b.
 - Wan, Z. and Li, Z. L.: Radiance-based validation of the V5 MODIS land–surface temperature product, Int. J. Remote Sens., 29, 5373–5395, doi:10.1080/01431160802036565, 2008.
- ¹⁵ Wang, A. and Zeng, X.: Evaluation of multireanalysis products with in situ observations over the Tibetan Plateau, J. Geophys. Res., 117, D05102, doi:10.1029/2011jd016553, 2012.
 Wang, G., Huang, J., Guo, W., Zuo, J., Wang, J., Bi, J., Huang, Z., and Shi, J.: Observation analysis of land–atmosphere interactions over the Loess Plateau of northwest China, J. Geophys. Res., 115, D00K17, doi:10.1029/2009jd013372, 2010.
- ²⁰ Wang, K. and Liang, S.: An improved method for estimating global evapotranspiration based on satellite determination of surface net radiation, vegetation index, temperature, and soil moisture, J. Hydrometeorol., 9, 712–727, doi:10.1175/2007jhm911.1, 2008.
- Wang, K., Wang, P., Li, Z., Cribb, M., and Sparrow, M.: A simple method to estimate actual evapotranspiration from a combination of net radiation, vegetation index, and temperature, J.
 Geophys. Res.-Atmos., 112, D15107, doi:10.1029/2006jd008351, 2007.
 - Wang, S., Zhang, Y., Lv, S., Liu, H., and Shang, L.: Estimation of turbulent fluxes using the flux-variance method over alpine meadows surface in eastern Tibetan Plateau, Adv. Atmos. Sci., 30, 411–424, doi:10.1007/s00376-012-2056-1, 2013.
- Wu, G., Liu, Y., He, B., Bao, Q., Duan, A., and Jin, F.-F.: Thermal controls on the Asian summer monsoon, Sci. Rep., 2, 404, doi:10.1038/srep00404, 2012.
- Yan, H., Wang, S. Q., Billesbach, D., Oechel, W., Zhang, J. H., Meyers, T., Martin, T. A., Matamala, R., Baldocchi, D., Bohrer, G., Dragoni, D., and Scott, R.: Global estimation of evapo-



transpiration using a leaf area index-based surface energy and water balance model, Remote Sens. Environ., 124, 581–595, 2012.

Yang, K., Koike, T., Fujii, H., Tamagawa, K., and Hirose, N.: Improvement of surface flux parametrizations with a turbulence-related length, Q. J. Roy. Meteor. Soc., 128, 2073–2087, 2002.

5

25

- Yang, K., Ding, B., Qin, J., Tang, W., Lu, N., and Lin, C.: Can aerosol loading explain the solar dimming over the Tibetan Plateau?, Geophys. Res. Lett., 39, L20710, doi:10.1029/2012gl053733, 2012.
- Yao, T., Wang, Y., Liu, S., Pu, J., Shen, Y., and Lu, A.: Recent glacial retreat in High Asia in China and its impact on water resource in Northwest China, Sci. China Ser. D, 47, 1065– 1075, doi:10.1360/03yd0256, 2004.
 - Yao, Y., Liang, S., Cheng, J., Liu, S., Fisher, J. B., Zhang, X., Jia, K., Zhao, X., Qin, Q., Zhao, B., Han, S., Zhou, G., Zhou, G., Li, Y., and Zhao, S.: MODIS-driven estimation of terrestrial latent heat flux in China based on a modified Priestley–Taylor algorithm, Agr. Forest Meteorol., 171–172, 187–202, doi:10.1016/j.agrformet.2012.11.016, 2013.
- 1/1–1/2, 18/–202, doi:10.1016/j.agrformet.2012.11.016, 2013.
 Yu, G.-R., Wen, X.-F., Sun, X.-M., Tanner, B. D., Lee, X., and Chen, J.-Y.: Overview of ChinaFLUX and evaluation of its eddy covariance measurement, Agr. Forest Meteorol., 137, 125–137, doi:10.1016/j.agrformet.2006.02.011, 2006.

Yuan, W., Liu, S., Yu, G., Bonnefond, J.-M., Chen, J., Davis, K., Desai, A. R., Goldstein, A. H.,

- ²⁰ Gianelle, D., Rossi, F., Suyker, A. E., and Verma, S. B.: Global estimates of evapotranspiration and gross primary production based on MODIS and global meteorology data, Remote Sens. Environ., 114, 1416–1431, doi:10.1016/j.rse.2010.01.022, 2010.
 - Zhang, C., Chen, F., Miao, S., Li, Q., Xia, X., and Xuan, C.: Impacts of urban expansion and future green planting on summer precipitation in the Beijing metropolitan area, J. Geophys. Res.-Atmos., 114, D02116, doi:10.1029/2008jd010328, 2009a.
 - Zhang, K., Kimball, J. S., Nemani, R. R., and Running, S. W.: A continuous satellite-derived global record of land surface evapotranspiration from 1983 to 2006, Water Resour. Res., 46, W09522, doi:10.1029/2009wr008800, 2010.

Zhang, L., Li, Y., Li, Y., and Zhao, X.: Seasonal changes of turbulent fluxes at a typical agricul-

tural site in the Chengdu Plain based on quality-controlled data, J. Meteorol. Soc. Jpn., 90, 195–202, 2012.



Zhong, L., Su, Z., Ma, Y., Salama, M. S., and Sobrino, J. A.: Accelerated changes of environmental conditions on the Tibetan Plateau caused by climate change, J. Climate, 24, 6540-6550, doi:10.1175/jcli-d-10-05000.1, 2011.

Atmos. Sci., 27, 500-512, doi:10.1007/s00376-009-9081-8, 2010.

- 15
 - Zhu, X., Liu, Y., and Wu, G.: An assessment of summer sensible heat flux on the Tibetan Plateau from eight data sets, Science China Earth Sciences, 55, 779-786, doi:10.1007/s11430-012-4379-2, 2012.

Zhang, X., Ren, Y., Yin, Z.-Y., Lin, Z., and Zheng, D.: Spatial and temporal variation patterns of reference evapotranspiration across the Qinghai-Tibetan Plateau during 1971-2004, J. Geophys. Res., 114, D15105, doi:10.1029/2009jd011753, 2009b.

Zhao, B., Yan, Y., Guo, H., He, M., Gu, Y., and Li, B.: Monitoring rapid vegetation succession in

- estuarine wetland using time series MODIS-based indicators: an application in the Yangtze 5 River Delta area, Ecol. Indic., 9, 346–356, doi:10.1016/j.ecolind.2008.05.009, 2009.
 - Zhou, L. and Huang, R.: Interdecadal variability of sensible heat flux in arid and semi-arid region of Northwest China and its relationship to summer precipitation in China, Chinese J. Atmos. Sci., 32, 1276–1288, 2008 (in Chinese).
- balance dataset **Discussion** Paper ¹⁰ Zhou, L. and Huang, R.: An assessment of the guality of surface sensible heat flux derived X. Chen et al. from reanalysis data through comparison with station observations in Northwest China, Adv. **Title Page** Abstract Introduction Conclusions References **Discussion** Paper **Tables** < Back Full Screen / Esc **Discussion** Paper

Discussion Paper



ACPD

14, 14471–14518, 2014

10 years China

land-surface energy

Table 1. Input datasets used for	calculating land	d surface	fluxes for	or China	(see	Sects.	2 and	5 b
for an explanation of abbreviation	າຣ).							

Variables	Data source	Temporal resolution	Availability	Domain	Spatial resolution (degrees)	Method
SWD	ITPCAS	3h	1979–2010	China land	0.1	Reanalysis
SWU	ITPCAS and GlobAlbedo	3h	1982–2009	China land	0.1	Satellite
						and Reanalysis
LWD	ITPCAS	3 h	1979–2010	China land	0.1	Reanalysis
LWU	MOD11C3	1 month	2000–2012	China land	0.05	Satellite
Та	ITPCAS	3 h	1979–2010	China land	0.1	Reanalysis
Q	ITPCAS	3 h	1979–2010	China land	0.1	Reanalysis
Ws	ITPCAS	3h	1979–2010	China land	0.1	Reanalysis
Ρ	ITPCAS	3 h	1979–2010	China land	0.1	Reanalysis
LST	MOD11C3 V5	1 month	2000–2012	Global	0.05	Satellite
HC	GLAS and SPOT VEGETATION	1 month	2000–2012	China land	0.01	Satellite
α	GlobAlbedo	1 month	2000–2010	Global	0.05	Satellite
NDVI	SPOT VEGETATION	10 days	1998–2012	Global	0.01	Satellite



ACPD 14, 14471–14518, 2014								
10 years China land-surface energy balance dataset								
X. Che	en et al.							
Title	Page							
Abstract	Introduction							
Conclusions	References							
Tables	Figures							
14	►I							
Back	Close							
Full Scre	een / Esc							
Printer-friendly Version								
Interactive	Discussion							
C	B Y							

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

Table 2. Flux tower sites supplying measurement data for product validation.

	Lat[deg]/Lon[deg]	Land cover	Eddy covariance	Radiometer	Measurement period	Site elevation (m)	Reference
WJ	30.4200° N/103.5000° E	Crop	CSAT3, Licor7500 (10 HZ)	CNR-1	Mar 2008–Aug 2009	539	Zhang et al. (2012)
MQ	33.8872° N/102.1406° E	Alpine meadow	CSAT3, Licor7500(10 HZ)	CNR-1	Apr 2009–May 2010	3439	Wang et al. (2013)
AL	33.3905° N/79.7035° E	Bare soil	CSAT3, Licor7500(10 HZ)	CNR-1	Jul 2010–Dec 2010	4700	Ma et al. (2008b)
BJ	31.3686° N/91.8986° E	Alpine grass	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2008–Dec 2010	4520	Ma et al. (2011)
MY	40.6038° N/117.3233° E	Orchard	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2008–Dec 2010	350	Liu et al. (2013a)
DX	39.6213° N/116.4270° E	Crop	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2008–Dec 2010	100	Liu et al. (2013a)
GT	36.5150° N/115.1274° E	Crop	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2008–Dec 2010	30	Liu et al. (2013a)
YC	36.9500° N/116.600° E	Crop	CSAT3, Licor7500(10 HZ)	CNR-1	Oct 2002–Oct 2004	13	Flerchinger et al. (2009)
DT	31.5169° N/121.9717° E	Wetland	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2005–Dec 2007	5	Zhao et al. (2009)
SC	35.95° N/104.133° E	Dry land	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2007–Dec 2008	1965	Huang et al. (2008)
WS	36.6488° N/116.0543° E	Winter wheat/ summer maize	CSAT3, Licor7500(10 HZ)	CNR-1	Jan 2006–Dec 2008	30	Lei and Yang (2010b)

ACPD 14, 14471–14518, 2014								
10 year land-surfa balance	10 years China land-surface energy balance dataset							
X. Che	en et al.							
Title	Page							
Abstract	Introduction							
Conclusions	References							
Tables	Figures							
14	۶I							
	•							
Back	Close							
Full Scre	een / Esc							
Printer-friendly Version								
Interactive	Discussion							
C	B Y							

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

Table 3. Comparison of accuracy of our flux data product and GLDAS against in-situ measurements from 11 Chinese flux towers.

			Energy flux				Radiation flux				
		H (W m ⁻²)	$LE (W m^{-2})$	$G_0 (W m^{-2})$	Rn (W m ⁻²)	Mean	SWD (W m ⁻²)	SWU (W m ⁻²)	$LWD (Wm^{-2})$	$LWU (W m^{-2})$	Mean
Our flux data product	Slope	0.39	0.9	0.87	0.92	0.77	0.95	0.68	0.91	0.95	0.87
	Intercept	-0.5	-6.1	6.1	-20.2	-8.2	13.6	10.9	-0.66	16.6	9.9
	RMSE	21.5	21.9	11.7	36.2	22.8	28.3	10.2	32.8	9.6	20.2
	MB	14.7	10.1	-5.7	26.3	11.4	-5.7	-0.65	28.9	2.4	6.2
	R	0.41	0.85	0.50	0.86	0.66	0.89	0.78	0.98	0.99	0.91
	Sample	280	284	197	313	270	310	307	307	307	308
GLDAS	Slope	0.77	0.87	0.58	1.0	0.81	0.99	0.75	0.87	1.0	0.90
	Intercept	20.83	5.1	-1.34	8.0	8.2	34.9	13.1	27.7	-4.5	17.8
	RMSE	26.6	20.6	6.7	17.9	17.9	45.6	15.9	19.2	11.1	23.0
	MB	-15.8	0.75	3.0	-10.4	-5.6	-32.87	-4.6	13.5	-3.2	-6.8
	R	0.46	0.80	0.61	0.95	0.71	0.87	0.65	0.99	0.98	0.87
	Sample	249	250	162	281	236	275	272	272	275	274



Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

Table 4. Comparison of statistical values reported in similar studies.

Reference	Research area	Method	Statistical parameters	H (W m ⁻²)	LE (W m ⁻²)	Flux network	Note
This study	Chinese landmass	SEBS	RMSE MB	23.1 16.8	21.9 8.3	flux towers in China	
			R	0.6	0.8		
Wang et al. (2007)	Southern Great Plains, USA	Regression method	RMSE	×	29.8	flux towers in Southern Great Plains, USA	calculated from Table 9
			MB	×	12.17		
			R	×	0.91		
Jiménez et al. (2009)	global	Statistical method	RMSE	×	×	AmeriFlux	calculated from Tables 5 and 7
			MB	-5.23	7.9		
			R	0.68	0.76		
Vinukollu et al. (2011b)	global	SEBS	RMSE	40.5	26.1	AmeriFlux	calculated from Table 4
			MB	27.98	-7.74		
			R	0.53	0.51		



Figure 1. A NDVI map of the Chinese landmass based on SPOT satellite data. The symbols indicate major physical phenomena: Tibetan Plateau (TP), northwestern China (NWC), inner Mongolian Plateau (MP), Loess Plateau (LP), North China Plain (NP), northeastern China Plain (NEP); Pearl River delta (PRD), Sichuan (SCB), Yinchuan (YCB), the inner Mongolian (IMB), and Lasha (LB), Tarim (TRB), Junggar (JB) basins; the Himalaya (HM), Ganges (GM), Kunlun (KL), Karakorum (KRM), Tianshan (TM), Nyainqentanglha (NQM) and Qilian mountain (QLM) ranges. The plateau and plain letter symbols are in red type. The basins letter symbols are in green type. The flux station letter symbols are in yellow type. Blue lines show several of the major rivers in China. Black lines indicate the borders of provinces.





Figure 2. Time-series comparison of SEBS input and output variables against measurements at Yucheng station. Black lines are SEBS results; red lines are measured values.





Figure 3. Time-series comparison of SEBS input and output variables against measurements at SC station. Black lines are SEBS results; red lines are measured values.





Figure 4. Maps of annual average **(a)** downward short-wave radiation (SWD), **(b)** downward long-wave radiation (LWD), **(c)** upward short-wave radiation (SWU), and **(d)** upward long-wave radiation (LWU) from 2001 to 2010. Black lines show several major rivers in China.





Figure 5. Maps of multiyear (2001–2010) means of retrieved fluxes: (a) sensible heat flux (H), (b) latent heat flux (LE), (c) net radiation (Rn), and (d) ground heat flux (G_0). White lines show several major rivers in China.





Figure 6. Maps of seasonal average sensible heat flux for **(a)** March–May (MAM), **(b)** June–August (JJA), **(c)** September–November (SON), and **(d)** December–February (DJF) from 2001 to 2010. Black lines show several major rivers in China.





Figure 7. Maps of seasonal average latent heat flux for **(a)** March–May (MAM), **(b)** June–August (JJA), **(c)** September–November (SON), and **(d)** December–February (DJF) from 2001 to 2010. White lines show several major rivers in China.





Figure 8. Maps of seasonal average net radiation for **(a)** March–May (MAM), **(b)** June–August (JJA), **(c)** September–November (SON), and **(d)** December–February (DJF) from 2001 to 2010. White lines show several major rivers in China.





Figure 9. Spatial trends of **(a)** SWD (downward short-wave), **(b)** LWD (downward long-wave), **(c)** SWU (upward short-wave), and **(d)** LWU (upward long-wave radiation) for the Chinese land-mass from 2001 to 2010.





Figure 10. Spatial trends of (a) sensible heat flux (H), (b) latent heat flux (LE), (c) net radiation (Rn), and (d) ground heat flux (G_0) on the Chinese landmass from 2001 to 2010.

