

1 Emission of iodine containing volatiles by selected microalgae species

2

3 U. R. Thorenz^{1*}, L. J. Carpenter², R.-J. Huang^{1,3,4}, M. Kundel¹, J. Bosle¹ and T. Hoffmann¹

4 [1] Institute of Inorganic and Analytical Chemistry, Johannes Gutenberg-University of Mainz,
5 55128 Mainz, Germany

6 [2] Department of Chemistry, University of York, Heslington, York YO10 5DD, UK

7 [3] School of Physics & Centre for Climate and Air Pollution Studies, Ryan Institute, National
8 University of Ireland Galway, University Road, Galway, Ireland

9 [4] Laboratory of Atmospheric Chemistry, Paul Scherrer Institute (PSI), 5232 Villigen, Switzerland

10 *Now at the Max Planck Institute for Chemistry, 55128 Mainz, Germany

11 Correspondence to: T. Hoffmann (t.hoffmann@uni-mainz.de)

12

13 Abstract

14 In this study we present the results of an emission study of different phytoplankton samples in
15 aqueous media treated with elevated ozone levels. Halocarbon measurements show that the
16 samples tested released bromoform and different iodocarbons including iodomethane,
17 iodochloromethane and diiodomethane. Iodide and iodate levels in the liquid phase were
18 representative of concentrations of surface water in a natural environment. Measurement of volatile
19 iodine (I₂) emissions from two diatom samples (*Mediopyxis helysia* and *Porosira glacialis*) and the
20 background sample (F/2-medium from filtered natural seawater), showed that the quantity of I₂
21 evolved depends on the ozone concentration in the air. This behaviour was assumed to be caused by
22 the oxidation reaction mechanism of iodide with ozone. The I₂ emission flux agrees with model
23 calculations at different iodide concentrations. The I₂ emission of a natural plankton concentrate
24 sample was, however, very low compared to other samples and showed no dependence on ozone.
25 The reason for this was shown to be the low iodide concentration in the algal suspension, which
26 seems to be the limiting factor in the oxidative formation of I₂.

27

28 Introduction

29 Iodine chemistry plays an essential role in the marine boundary layer (MBL) due to its effect on the
30 destruction of tropospheric ozone, perturbation of the HO_x/NO_x cycle and the formation of new
31 particles and cloud condensation nuclei, thereby leading to changes in the global radiative forcing
32 (Hoffmann et al., 2001; von Glasow and Crutzen, 2003; O'Dowd and Hoffmann, 2005; Bloss et al.,
33 2005; Huang et al., 2010a, b). This essential role of iodine and of other activated halogens is shown in
34 field measurements in the marine boundary layer (MBL), laboratory chamber experiments or
35 incubation experiments of different algae and in atmospheric models (Carpenter, 2003; Küpper et al.,

1 2008; Kundel et al., 2012; McFiggans et al., 2000). The biogeochemical cycle of iodine is controlled by
2 large iodine exchanges from the oceans to the atmosphere, driven by marine biotic and abiotic
3 production (Schall et al., 1997). Volatilized species are photolabile iodocarbons like CH_2I_2 , CH_3I , $\text{C}_2\text{H}_5\text{I}$,
4 CH_2ICl , CH_2IBr and molecular iodine (I_2). Marine species like macroalgae and microalgae play a
5 dominant role in the emission of these compounds (Carpenter et al., 1999, Huang et al., 2013, Saiz-
6 Lopez and Plane, 2004).

7 Since molecular iodine and iodocarbons are photochemically unstable (lifetimes between about
8 some tens of seconds for I_2 and a few days for CH_3I) they are photolysed under UV-visible light to
9 form $\text{I}\cdot$ atoms, which are then instantly oxidised by ozone to form the iodine monoxide radical $\text{IO}(\text{g})$
10 (Hoffmann et al., 2001; Saiz-Lopez et al., 2006). Further oxidation reactions of IO in the gas phase
11 then can form low volatile iodine oxides (I_xO_y) which may nucleate under certain conditions and form
12 new particles.

13 Recently it was proposed that the ozone loss over the tropical Atlantic Ocean was higher than
14 calculated from global atmospheric models, and that this additional ozone destruction is induced by
15 halogens such as bromine and iodine (Read et al., 2008). Biogenic emissions, such as the already
16 studied iodocarbon emissions by phytoplankton species, e.g. coccolithophorids, diatoms and
17 chlorophytes, (Colomb et al. 2008) are too low to explain the differences in model calculations and
18 observations (Mahajan et al. 2010), therefore additional sources of the reactive iodine species are
19 discussed, one of them being the surface reaction of ozone with seawater.

20 Garland and Curtis first discovered that the emission of molecular iodine from the surface of artificial
21 and natural seawater is proportional to the ozone concentration at the air/water interface (Garland
22 and Curtis, 1981). Sakamoto and co-workers examined the reaction mechanism of the iodide
23 oxidation by ozone at the air/water interface, resulting in the formation of the intermediates IOOO^-
24 and HOI and the emission products IO and I_2 (Sakamoto et al., 2009). Further laboratory experiments
25 show that different organics affect the reaction of iodide with ozone, e.g. fulvic acid enhances the I_2
26 formation, but not the formation of IO (Hayase et al., 2010, 2012).

27 Since the formation of I_2 and IO from the air/water interface is dependent on the iodide
28 concentration in seawater, the reaction path found by Garland and Curtis may explain elevated
29 iodine emissions in areas of higher phytoplankton activity (Garland and Curtis, 1981). The ability of
30 different phytoplankton, e.g. diatoms, to reduce iodate, which is ubiquitous in the open ocean, to
31 iodide was shown for natural and elevated iodate concentrations (Wong et al. 2002; Chance et al.
32 2007) and for the different growth states (Bluhm et al. 2010) of the phytoplankton cultures. A
33 correlation of iodine species in the particle phase and average chlorophyll exposure of air masses
34 along back trajectories was found by Lai et al, 2011, indicating the link between phytoplankton
35 activity and emission of atmospheric iodine.

36 Since the formation of I_2 and IO is correlated to the iodide concentration (Sakamoto et al. 2009) and
37 the iodide concentration of surface waters is correlated to phytoplankton (Bluhm et al. 2010), this
38 study investigates links between iodide concentrations in microalgae-containing seawater and abiotic
39 formation and emission of I_2 , utilising laboratory experiments of the reaction of the seawater surface
40 with ozone.

1 Materials and Methods

2 Experimental set-up

3 Two diatom cultures (*M. helysia*, *Porosira glacialis* both from the Alfred Wegener Institut/Sylt), were
4 kept in F/2 seawater medium for growing. These media were prepared from filtered natural
5 seawater from the shores of Sylt and additional nutrients which the diatoms need to grow (0.88
6 mmol NO₃⁻, 0.04 mmol PO₄³⁻ and 0.01 mmol SiO₃²⁻) and which is a common used medium as
7 described by Guillard et al., 1975 and Kraberg et al. 2012. Both cultures were incubated in the F/2
8 medium at 16°C with 12-h-light-12-h-dark cycling (LUMILUX Plus Eco daylight lamp; approx.. 40 µmol
9 PAR) for 4 weeks prior the experiment. Just before the emission experiment, the algal suspensions
10 were diluted in a 2:1 ratio in F/2 medium and homogenised by stirring. In addition to the diatom
11 cultures, a plankton concentrate was collected from the North Sea (55°01.562N; 8°27.113E) on May
12 24th 2012 using a 80 µm and 200 µm Apstein plankton net and diluted using the same medium as for
13 the diatom cultures. Microscopic observations showed that the plankton concentrate sample was
14 dominated by colonies of the haptophyte *Phaeocystis sp.* and only a low amount of diatoms were
15 present.

16 For each experiment 1.5 L of the sample (i.e. diatom suspension, natural plankton concentrate or
17 background (F/2 medium)) was introduced into a glass chamber tube (10 L), shown in Figure 1, and
18 three magnetic stirrers were switched on immediately. A continuous flow of synthetic air (3.4 L min⁻¹)
19 was channelled over the stirred algae suspension (surface area 2250 cm²) in the first experiment with
20 no ozone and in the second experiment with elevated ozone levels of 100 ppb. The ozone was
21 generated using an UV radiation source and the resulting ozone levels were measured using an
22 ozone analyzer (Dasibi Environmental Corp. Model 1008-RS, Glendale, USA). To measure the
23 emission of I₂ and halocarbons, α-cyclodextrin-coated denuders (Huang and Hoffmann, 2009; Huang
24 et al., 2010c) and adsorption tubes (Kundel et al. 2012) were mounted at the other end of the tube
25 chamber together with the ozone monitor. The chamber outflow was sampled using two membrane
26 pumps, one with 0.50 L min⁻¹ for the denuders and the other using 0.15 L min⁻¹ for the adsorption
27 tubes. To assure an overpressure over the sampling time a U-shaped tube filled with ultra-pure water
28 was mounted in the centre exit of the glass chamber to measure the overpressure hydrostatically.
29 The whole set-up was wrapped with aluminium foil to prevent photolysis of I₂ and halocarbon
30 compounds. Potential wall losses of I₂ and halocarbons were investigated using diffusion (I₂) and
31 permeation (halocarbons) test gas sources with a dilution chamber; no wall losses were observed
32 within the precision of the measurements for the three concentrations tested: 70 ppt, 270 ppt and
33 700 ppt for iodine and 0.2 ppt, 2.4 ppt and 8.3 ppt for the halocarbons and within using the stated
34 gas flows.

35 To monitor the emissions of I₂ and halocarbons from the liquid samples, an evaporation standard
36 was added to the microalgal suspension in order to highlight any problems related to air sampling.
37 This standard was 1,3-dibromopropane diluted in ultrapure water (500 µl of 0.94 µg L⁻¹ which was
38 then diluted with the sample to 1.5 L). The standard was chosen given the results from a first set of
39 experiments with *M. helysia* and *Coscinodiscus wailesii* which show no detectable traces of this
40 compound. We decided not to add any iodine containing compounds to prevent interferences with
41 the I₂ emission.

1 Halocarbon measurements

2 Air samples of 6.75 L sampling volume were pre-concentrated at a flow rate of 150 ml min⁻¹ on
3 thermal desorption tubes filled with 100 mg Tenax TA 60/80 and 150 mg CarbotrapTM 20/40 both
4 provided by Supelco (Bellefonte, PA, USA). The samples were analysed using a self-made thermal
5 desorption device mounted on a gas chromatograph (TraceGC, Thermo Scientific, Dreieich Germany)
6 - mass spectrometer (PolarisQ, Thermo Scientific, Dreieich, Germany). During the desorption period
7 of 6 minutes the cryotrap was cooled to -160 °C. Afterwards the cryotrap was rapidly heated to 270
8 °C for injection. The analytes were separated on a DB624 Durabond column (60 m; 0,32 mm; 1,8 µm
9 FT) using helium as carrier gas with a constant pre-column pressure of 0.5 bar. The temperature
10 program was: 55 °C (4 min), ramp with 5 °C min⁻¹ to 120 °C (4 min) and ramp with 8 °C min⁻¹ to 200 °C
11 (4 min). Halocarbons were detected using a mass spectrometer in NCI mode with methane as
12 reagent gas (2.5 ml min⁻¹), the primary electron energy was set to 120 eV and an emission current of
13 50 mA in single ion monitoring mode (SIM) was used. Iodinated compounds (CH₃I, C₂H₅I, CH₂ICl,
14 CH₂I₂, 1-C₃H₇I, 2-C₃H₇I, 1-nC₄H₉I, 2-nC₄H₉I, 1-iso-C₄H₉I) were quantified using *m/z* 127 and
15 brominated compounds (CH₂Br₂, CH₃Br, 1,3-C₃H₆Br₂) were quantified using *m/z* 79 and 81 at a 1:1
16 ratio. A five point calibration was done in the range between 0.01 ng and 1 ng using the continuously
17 diluted output of a permeation test gas source (Thorenz et al. 2012). The detection limits for the
18 individual iodocarbons were 0.003-0.088 ppt and for the bromocarbons were 0.004 – 0.009 ppt. For
19 each series of measurements, the calibration was done in triplicate (precision of method 3-13%).

20 I₂, Iodide and Iodate measurements

21 Sampling of gaseous I₂ was performed using the denuder technique described by Huang and
22 Hoffmann, 2009. Brown glass denuder tubes (6 mm i.d., 50 cm length) were coated using a α-
23 cyclodextrin suspension (2.5 mg mL⁻¹ in methanol) and sealed with polypropylene caps. Before
24 sampling the denuders were stored in a fridge. For sampling the denuders were mounted vertically
25 with a glass tube of 15 cm upstream to achieve laminar flow. The sampling flow was 500 mL min⁻¹ for
26 45 min. After sampling the denuders were sealed and stored in a fridge until derivatization. For
27 derivatization the α-cyclodextrine coating was eluted with ultrapure water (20 mL), then 25 µL N,N-
28 dimethylaniline (1 µg mL⁻¹ in methanol), 500 µL phosphate buffer (pH 6.4) and 500 µL 2-
29 iodosobenzoate (4 mg mL⁻¹) were added, the mixture was shaken for 2 hours. After adding 3 ml
30 sodium acetate the sample was extracted with 100 µL cyclohexane and 100µL 2,4,6-tribromoaniline
31 (internal standard: IS) in cyclohexane (250ppb).

32 Iodide and iodate were derivatized from seawater to form the same product as described for I₂.
33 Iodide was oxidized to form I₂ by using iodosobenzoate and iodate was reduced first to iodide and
34 then oxidized to form I₂. 10 mL aliquots of seawater were analysed for iodide and for total iodine,
35 iodate was calculated by difference. The method for iodide derivatization was slightly changed from
36 the one described by Mishra et al. (2000). The use of sodium hydrogen sulfite as an agent to reduce
37 iodate to iodide is described by Schwehr and Santschi (2003).

$$38 \quad I_{\text{seawater}} = I^- + IO_3^-$$

39 To measure iodide, 10 mL seawater were mixed with 1 ml ethylenediaminetetraacetic acid solution
40 (0.5%), 500 µL phosphate buffer, 500 µL N,N-dimethylaniline, 500 µL iodosobenzoate and shaken.

1 After adding 3 ml sodium acetate the sample was extracted with 100 μ L cyclohexane and 100 μ L
2 2,4,6-tribromoaniline (IS) in cyclohexane (250 ppb).

3 To measure iodate an aliquot of 10 mL seawater was mixed with 1 mL ethylenediaminetetraacetic
4 acid solution (0.5%), 1 mL hydrochloric acid (3.7%) and 500 μ L sodium hydrogen sulfite solution
5 ($283.9 \mu\text{mol L}^{-1}$) to reduce the iodate. Afterwards 500 μ L sodium acetate, 4 mL phosphate buffer, 500
6 μ L N,N-dimethylaniline, 500 μ L iodosobenzoate were added. After shaking the sample was again
7 extracted with 100 μ L cyclohexane and 100 μ L 2,4,6-tribromoaniline (IS) in cyclohexane (250 ppb).

8 1 μ L of the cyclohexane extract was injected to the GC-MS System (6850 GC & 5973 MS, Agilent
9 Technologies, Waldbronn, Germany) at a constant flow of 1 mL min^{-1} of helium (99.999%) and, the
10 chromatographic separation was performed using a capillary column FS Supreme 5 MS with a length
11 of 30 m, inner diameter of 0.25 mm and film thickness of 0.25 μ m (CS Chromatographie Servieve,
12 Langenwehe, Germany) with a temperature program starting at 50°C (for 3 min), then heating up at
13 30 °C min^{-1} to 220°C (for 3 min). The mass spectrometer measured in electron ionisation mode at 70
14 eV, the specific fragments of the product 4-iodo-N,N-dimethylaniline was extracted at m/z 247 (M+)
15 and of the internal standard 2,4,6-tribromoaniline at m/z 329 (M+).

16 Chlorophyll measurements

17 The analytical method for chlorophyll α (chl α) measurements is described by Edler et al 1979. An
18 aliquot of 50-100 mL water samples were filtered on glass fibre filters (GF/F-Whatman). The dry
19 filters were put in polypropylene vials and extracted with 7.5 ml acetone. The extract was stored
20 together with the filter in a dark fridge at 3°C overnight and centrifuged the next day (5500 rpm, 7
21 min) at 5°C. The absorption of the supernatant was measured against acetone using an Uvikon XL
22 double beam spectrophotometer at $\lambda = 750 \text{ nm}$, 663 nm, 645 nm and 630 nm. To calculate the
23 concentration of chl α the equation of Jeffrey and Humphrey, 1975 was used. Chl α can be a good
24 indicator for microalgae biomass (Roy 2010; Bluhm et al. 2010; Colomb et al. 2008), and has been
25 used to calculated emission rates of iodine-containing volatiles from phytoplankton. This calculation
26 was not used here, since the mechanisms of synthesis and release of these iodine containing gases is
27 still unclear. All gaseous compounds in this study are therefore given as measured mixing ratio and
28 the chl α value of the corresponding algae suspension is also given.

29 Results and Discussion

30 Halocarbons

31 The emission rates of the natural halocarbons and the evaporation standard, given in Table 1, were
32 calculated by the amount measured in the adsorption tubes divided by the emission time and the
33 surface area of the suspension sample ($\text{pmol min}^{-1} \text{m}^{-2}$). The halocarbon emission rates showed no
34 effect on the different ozone levels; therefore the data for each sample are summarized for high and
35 low ozone conditions. An evaporation standard was added to the different samples to recognize
36 differences in emission rates of the organic compounds from the aqueous phase. The standard was
37 added in a 10 to 100-fold excess compared to natural concentrations of bromocarbons in Atlantic
38 seawater (Carpenter et al. 2000) to reduce the effect of natural 1,3-dibromopropane which may alter
39 the mixing ratio of the evaporation standard measured. In the chosen concentration a natural

1 abundance would change the result only by 1-10% compared to the spike solution. The results of the
2 measurements of 1,3-dibromopropane showed very constant values, as can be seen from the low
3 standard deviation between the different samples and replicates. This result indicates a stable and
4 reliable experimental setup in terms of evaporation of volatile compounds from the water surface
5 and of the mixing of the bulk water.

6 The measured emission rates of the natural halocarbons show that the brominated compound,
7 CHBr_3 , is elevated compared to the iodocarbons emission rates. This result fits to observations of the
8 natural abundance of halocarbons in seawater as described in earlier studies (Roy et al. 2011). The
9 emission rate of CHBr_3 is higher for the two diatom cultures (*M. helysia* and *P. glacialis*) than for the
10 plankton samples containing *Phaeocystis sp.* and the background. Again, this result matches field and
11 laboratory data showing a link between elevated CHBr_3 concentrations in seawater and the
12 simultaneous occurrence of diatoms (Colomb et al. 2008, Quack et al. 2007, Moore et al. 1996).

13 The iodocarbon emissions in experiments using the background (F/2 medium) and the plankton
14 concentrate were dominated by CH_3I , followed by CH_2ICl and CH_2I_2 . However, for the diatom
15 cultures, CH_2I_2 was the dominant iodocarbon emitted with CH_3I and CH_2ICl both showing lower
16 emission rates. The emission of iodocarbons from the F/2 background is not surprising for two
17 reasons; first, the medium was produced from natural shoreline filtered water, which already may
18 contain iodocarbons (Wong and Cheng, 1998). The second reason may be related to iodocarbon-
19 producing bacteria (Amachi et al. 2001; Amachi et al. 2003). These bacteria could have been present
20 and active in the natural seawater water used to produce the F/2 medium, since it was not sterilized
21 prior to use. Additionally, the emission rates of CH_2I_2 , CH_2ICl and CH_3I in the diatom samples *P.*
22 *glacialis* and *M. helysia* were significant higher compared to the background (Wilcoxon rank sum test
23 $p=0.00032$ and $p=0.00007$, respectively). This increase in emission can be explained by the capability
24 of the diatoms to produce iodocarbons which had already been reported by Moore et al. (1996). To
25 compare the natural plankton concentrate with the cultivated diatom cultures and the background
26 one must keep in mind that chl α concentrations are biomass tracers reflecting the abundance of
27 phytoplankton. The results for the chl α measurement, given in Table 1, clearly show that the natural
28 plankton concentrate contains less biomass than the cultured diatoms. Therefore, we conclude that
29 the lower iodocarbon emissions of the plankton concentrate compared to the diatom cultures is
30 partly due to lower biomass density. The lower iodocarbon emission rates in the natural plankton
31 concentrate could also be related to iodine uptake of natural occurring micro algae (van Bergeijk et
32 al. 2013). The emission flux summed for the three iodocarbons in the four samples' background,
33 plankton concentrate, *P. glacialis* and *M. helysia*, was in the range $0.21 - 1.02 \text{ pmol min}^{-1} \text{ m}^{-2}$, $0.14 -$
34 $0.58 \text{ pmol min}^{-1} \text{ m}^{-2}$, $0.50 - 1.35 \text{ pmol min}^{-1} \text{ m}^{-2}$ and $0.57 - 1.53 \text{ pmol min}^{-1} \text{ m}^{-2}$, respectively. We are
35 not aware of emission studies investigating the flux of iodocarbons from micro algae suspensions to
36 directly compare these results. To establish a connection to other experimental observations the
37 results listed above are compared to incubation studies of marine aggregates producing iodocarbons
38 and calculated emission fluxes in coastal, seaweed-rich regions. Hughes et al. (2008) measured the
39 iodocarbon production of different marine aggregates to be within 0.71 to $6.90 \text{ pmol h}^{-1} \text{ L}^{-1}$. The
40 production rate is difficult to compare to the presented results, since the flux in our study is based on
41 the production by the microalgae species and evaporation from the surface, whereas Hughes et al.
42 (2008) measured the production in the aqueous phase. Jones et al. (2009) calculated iodocarbon
43 emissions at a sampling site surrounded by fields of macro algae in open sea water at Roscoff,

1 France. The flux of iodocarbons was estimated to $85.28 \text{ pmol min}^{-1} \text{ m}^{-2}$, two orders of magnitude
2 higher than the flux obtained in the present study. Thus it appears that on an areal basis, the
3 natural populations of microalgae studied here are much less prevalent emitters of iodocarbons than
4 seaweeds and marine aggregates.

5

6 Iodide and iodate

7 The concentrations of iodide and iodate in the different samples are shown in Table 1. For each
8 sample, the mean and range for six replicates are shown; no differences in iodide and iodate
9 concentrations were observed under elevated (100 ppb O_3) and low ozone (0 ppb O_3) conditions.
10 The iodate concentrations in the background and in the three plankton samples were in the same
11 range, with mean concentrations between 438 and 448 nmol L^{-1} . These iodate concentrations are in
12 the range measured for the open ocean of 400 to 500 nmol L^{-1} iodate in most oceanic regions (Bluhm
13 et al. 2011). The ubiquity of iodate suggests that its concentration is not a limiting factor.

14 The iodide concentrations in the two diatom cultures, *P. glacialis* and *M. helysia*, are slightly elevated
15 with mean values of $12.70 \text{ nmol L}^{-1}$ and $16.84 \text{ nmol L}^{-1}$, respectively, compared to the background
16 iodide concentration of $10.35 \text{ nmol L}^{-1}$ and the plankton concentrate iodide concentration of 6.47
17 nmol L^{-1} . This enhanced iodide concentration indicates the reduction of iodate by the two diatom
18 cultures, which was also found by Bluhm et al. 2010 and Wong et al. 2002 for different
19 phytoplankton species. Such a reduction of iodate to iodide will result in a decrease in the iodate
20 concentration, however, for the measured iodate concentration in this study the expected decrease
21 falls within the analytical precision of the measurement. The iodide concentrations in all samples are
22 comparable with oceanic surface water concentrations, for example around 10 - 30 nmol L^{-1} in the
23 Weddel Sea surface water (Bluhm et al. 2011).

24 The low iodide concentration of the plankton concentrate sample compared to the background
25 sample is surprising, but may be assigned to an overall low level of different nutrients, like phosphate
26 and silicate, in the Wadden Sea of Sylt at springtime (Weisse et al. 1986), although the level of iodate
27 was consistent. Another possible reason for the low iodide concentration in the plankton
28 concentrate could be iodine uptake by microalgae present in the natural plankton sample (van
29 Bergeijk et al. 2013).

30 Ozone measurements

31 The results of the ozone measurement for the samples: background, *P. glacialis*, *M. helysia* and the
32 plankton concentrate were normalized against a background measurement obtained using ultra-pure
33 water in the chamber. This was performed in order to account for losses of ozone through wall
34 reactions, losses on the water surface, and losses due to droplet formation from stirring. The ozone
35 consumption was calculated using a Continuous Stirred-Tank Reactor (CSTR) approach with 668 ng
36 min^{-1} ozone (100 ppb) introduced into the chamber (total volume: 10 L , flow: 3.4 L min^{-1} and
37 residence time: 2.94 min). The difference between the introduced ozone flow and measured ozone
38 flow is considered as consumed ozone, due to the oxidation of iodide and other ozone depleting
39 reactions in the samples. To calculate the consumed ozone, the flow rate was summarized over 45
40 min of the experiment. Ozone consumption was clearly observed for all samples. The background

1 sample showed the weakest ozone consumption of 58 nmol, followed by the sample of *P. glacialis*
2 with 186 nmol and the plankton concentrate with 253 nmol. The highest ozone consumption was
3 shown by *M. helysia* with 335 nmol.

4 I₂ emissions

5 The I₂ emission rate was calculated by dividing the amount of I₂ by the sampling time and the
6 suspension surface area. The results for the four samples are shown in Figure 2. The background and
7 the two diatom samples, *M. helysia* and *P. glacialis* show significant higher emission rates when the
8 ozone level is elevated (100 ppb O₃) compared to conditions where no ozone is present (0 ppb O₃).
9 The difference between the high and low ozone conditions is small for the background, increases for
10 the *P. glacialis* sample and is highest for the *M. helysia* sample. The plankton concentrate does not
11 show a significant dependence of the I₂ emission rate on the ozone level. The ozone-dependent
12 increase in the I₂ emission rate of the other samples indicates that iodide, which is present at the
13 air/water interface, is oxidised by ozone to form I₂, which is consistent with the results from artificial
14 and natural seawater (Garland and Curtis, 1981, Sakamoto et al. 2009).

15 Figure 3 shows the change in I₂ emission rate ([I₂ at 100 ppb ozone] – [I₂ at 0 ppb ozone]) of the
16 different samples as a function of the iodide concentration measured in the bulk water. A linear
17 correlation fits the data well with a Pearson coefficient of R² = 0.998. This behaviour indicates a
18 direct proportional relationship, which was also seen by Sakamoto et al. 2009 for small iodide
19 concentrations (0 – 5 mmol L⁻¹). Carpenter et al. (2013) also observed that the I₂ emission is
20 dependent on the aqueous iodide concentration. The proposed reaction sequence, as shown in
21 equations (1)-(5), explains the relationship between the iodide concentration in the aqueous phase
22 and the I₂ emissions (Sakamoto et al. 2009).



28 The plankton sample does not show an elevated I₂ emission at 100 ppb ozone compared to zero
29 ozone. This observation indicates that in the plankton sample an additional I₂ loss process takes
30 place. Reactions or partitioning of I₂ in an organic surface layer, which was discussed in Carpenter et
31 al. (2013), would be one possibility to explain these results. In fact the specific microalgae found in
32 the plankton concentrate, *Phaeocystis sp.*, is known to produce high amounts of organic matter
33 (Eberlein et al. 1985). An alternative explanation is the low iodide concentration in the plankton
34 concentrate, which may be related to iodide uptake by the natural occurring plankton communities.
35 The iodide concentrations and ozone mixing ratios in this study represent more likely natural
36 conditions compared to the study of Sakamoto et al. (iodide concentration between 0.01 – 50 mmol
37 L⁻¹ and ozone mixing ratio from 2 – 298 ppm). However, the results presented here demonstrate that

1 even under low iodide concentrations, representative of natural conditions of the MBL, a significant
2 formation of I₂ by the ozone driven oxidation of iodide at the air/water interface takes place, until
3 the iodide concentration gets too low.

4 Comparing the I₂ and iodocarbon emission rates, it is clear that the volatile iodine emissions are
5 dominated by I₂. Therefore I₂ emissions from natural seawater surfaces are more relevant for
6 atmospheric processes than the emission of iodocarbons. At the same time the experiments
7 presented here show that the emission of iodocarbons is not linked to the formation of I₂ at the
8 air/water interface (Martino et al., 2009), since no correlation between I₂ emissions or O₃ mixing
9 ratio and iodocarbon emissions was observed.

10 Calculated emissions for the background, *P. glacialis* and *M. helysia* were 8.32×10^5 , 1.47×10^6 and
11 2.40×10^6 molecules cm⁻² s⁻¹, respectively. Modelled emissions calculated using the kinetic model of
12 the aqueous interfacial layer by Carpenter et al. (2013) for the iodide concentration measured were
13 1.16×10^6 , 1.67×10^6 and 2.91×10^6 molecules cm⁻² s⁻¹, respectively. The measured and modelled
14 values agree well, showing that the model is able to predict emissions for natural iodide
15 concentrations.

16 Figure 4 shows the change in the I₂ emission rate plotted versus the consumed ozone for the four
17 different samples. This was done to see whether ozone depletion in the flow chamber is mainly
18 driven by the iodide or if other factors are important. The graph shows that the ozone depletion
19 correlates with the enhancement in the I₂ emission rate for the two diatom samples and for the
20 background. Therefore, the ratio of the formation of I₂ to the amount of O₃ consumed, calculated as
21 $R(I_2) = n(I_2) / n(O_3)$, with $n(I_2)$ = amount of I₂ formed and $n(O_3)$ = amount of O₃ consumed during the
22 experiment, was used to determine the dependence of I₂ formation on O₃. $R(I_2)$ has a maximum value
23 of 1, which, referring to eqs.1-5, indicates that every molecule of ozone which is consumed produces
24 one molecule of I₂. The formation ratio for the background sample was the highest with $R(I_2) =$
25 0.14‰, followed by the samples of *M. helysia* with $R(I_2) = 0.08‰$ and *P. glacialis* $R(I_2) = 0.07‰$. This
26 means that a higher degree of biological activity of the sample decreases the formation ratio. The
27 decrease of I₂ emission in the surface reaction of ozone with iodide was also seen by Carpenter et al.
28 when turning from iodide solutions to sea water, which contains more organic substances (Carpenter
29 et al. 2013).

30 The plankton concentrate also depletes ozone, although there is no enhancement in I₂ emission.
31 Therefore, another mechanism in ozone depletion must be taking place, possibly induced by other
32 ozone reactive substances formed or excreted from *Phaeocystis sp.*. Another explanation is a
33 reduced release of I₂ and a higher release of HOI, which was not measured in this study. In fact,
34 Carpenter and coworkers observed HOI as the main iodine compound released in their experiments,
35 followed by I₂ (Carpenter et al. 2013).

36

37 Conclusions

38 Different phytoplankton suspensions were treated with high and low ozone levels. Halocarbons
39 including bromoform, iodomethane, iodochloromethane and diiodomethane, were released from
40 the suspensions independent of the ozone level. The use of an evaporation standard in the aqueous

1 phase indicated that the emission rates of all gaseous organics were quite stable. The iodide and
2 iodate concentration in the liquid phase also showed no dependence on the ozone level in the gas
3 phase and were comparable to concentrations in surface water in the open ocean. The emission flux
4 of the iodocarbons was lower compared to the calculated flux at a coastal, kelp-rich site in Roscoff,
5 France, an observation which emphasizes the higher emission of iodocarbons from macroalgae
6 compared to microalgae. The emission rates of iodocarbons were also lower than the emission of I₂,
7 confirming that I₂ emissions from the remote ocean dominate over organic iodine sources for the
8 MBL (Jones et al., 2010; Lawler 2012; Carpenter et al., 2013). The emission of I₂ showed a
9 dependency on the ozone level in the air as well as on the iodide concentration in the sample
10 suspension, as has been found previously (Carpenter et al. 2013 and other refs). For the two diatom
11 samples, *M. helysia* and *P. glacialis*, and the background sample, a correlation was found for the I₂
12 emission and the ozone consumption during the experiment. The I₂ emissions from the plankton
13 concentrate, taken in the Wadden Sea of Sylt, were lower than the other samples and showed no
14 dependence on the ozone levels. An explanation could be the lower iodide concentration in the
15 plankton sample, since iodide is the limiting factor for the oxidative reaction. Another explanation
16 may be the preferred formation and emission of HOI when organic compounds are present in the
17 liquid phase. The experiments showed that different algae suspensions (*M. helysia* and *P. glaciales*)
18 are capable of emitting I₂ by the reaction of ozone with dissolved iodide at the air/water interface
19 under natural conditions. However, it remains unclear whether iodine emissions from aquatic
20 systems can be fully understood without the simultaneous measurement of HOI.

21

22 Acknowledgements

23 The authors gratefully acknowledge the financial support of the German Research Foundation (DFG:
24 Deutsche Forschungsgemeinschaft) within the Research Training Group 826: "Trace Analysis of
25 Elemental Species: Development of Methods and Applications."

26

1 References

- 2 Amachi S., Kamagata ., Kanagawa T., Muramatsu Y.: Bacteria mediate methylation of iodine in
3 marine and terrestrial environments. *Appl. Environ. Microbiol.* 67, 2718-2722, 2001.
- 4 Amachi S., Kasahara M., Hanada S., Kamagata Y., Shinoyama H., Fujii T., Muramatsu Y.: Microbial
5 participation in iodine volatilization from soils, *Environ. Sci. Tech.* 37, 3885-3890 2003.
- 6 Bloss W.J., Lee J.D., Johnson G.P., Sommariva R., de Heard, Saiz-Lopez A., Plane J.M., McFiggans G.,
7 Coe H., Flynn M., Williams P., Rickard A.R., Fleming Z.L.: Impact of halogen monoxide chemistry
8 upon boundary layer OH and HO₂ concentrations at a coastal site, *Geophys. Res. Lett.*, 32,
9 doi:10.1029/2004GL022084, 2005.
- 10 Bluhm K., Croot P.L., Huhn O.R.G.: Distribution of iodide and iodate in the Atlantic sector of the
11 southern ocean during austral summer, *Deep-sea research part II- topical studies in*
12 *oceanography* 58, 2733-2748, 2011.
- 13 Bluhm K., Croot P., Wuttig K., Lochte K.: Transformation of iodate to iodide in marine phytoplankton
14 driven by cell senescence, *Aquat. Biol.* 11, 1-15, 2010.
- 15 Carpenter L.J.: Iodine in the marine boundary layer, *Chem. Rev.* 103, 4953-4962, 2003.
- 16 Carpenter L.J., MacDonald S.M., Shaw M.D., Kumar R., Saunders R.W., Parthipan R., Wilson J., Plane
17 J.M.C.: Atmospheric iodine levels influenced by sea surface emissions of inorganic iodine, *Nature*
18 *Geosci.* 6, 108-111, 2013.
- 19 Carpenter L.J., Malin G., Liss P.S., Kupper F.C.: Novel biogenic iodine-containing trihalomethanes and
20 other short-lived halocarbons in the coastal East Atlantic, *Global. Biogeochem. Cy.*, 14, 1191-
21 1204, 2000.
- 22 Carpenter L.J., Sturges W.T., Penkett S.A., Liss P.S., Alicke B., Hebestreit K., Platt U.: Short-lived alkyl
23 iodides and bromides at Mace Head, Ireland: Links to biogenic sources and halogen oxide
24 production, *J. Geophys. Res.-Atmos.*, 104, 1679-1689, 1999.
- 25 Chance R., Malin G., Jickells T., Baker A.R.: Reduction of iodate to iodide by cold water diatom
26 cultures. *Mar. Chem.*, 105, 169-180, 2007.
- 27 Colomb A., Yassaa N., Williams J., Peeken I., Lochte K.: Screening volatile organic compounds (VOCs)
28 emissions from five marine phytoplankton species by head space gas chromatography/mass
29 spectrometry (HS-GC/MS). *J. Environ. Monit.*, 10, 325-330, 2008.
- 30 Eberlein K., Leal M.T., Hammer K.D., Hickel W. Dissolved organic substances during a *Phaeocystis*
31 *pouchetii* bloom in the German Bight (North Sea), *Mar. Biol.*, 89, 311-316, 1985.
- 32 Edler L.: *Phytoplankton and chlorophyll.* Baltic Marine Biologists, vol 5. Univ. of Stockholm,
33 Stockholm, 1979
- 34 Garland J.A., Curtis H.: Emission of iodine from the sea-surface in the presence of ozone, *J. Geophys.*
35 *Res.-Oc., ATM* 86, 3183-3186, 1981.
- 36 Hayase S., Yabushita A., Kawasaki M.: Iodine Emission in the Presence of Humic Substances at the
37 Water's Surface, *J. Phys. Chem.*, 116, 5779-5783, 2012.

- 1 Hayase S., Yabushita A., Kawasaki M., Enami S., Hoffmann M.R., Colussi A.J.: Heterogeneous reaction
2 of gaseous ozone with aqueous iodide in the presence of aqueous organic species, *J. Phys.*
3 *Chem. A* 114,6016–6021, 2010.
- 4 Hoffmann T., O'Dowd C.D., Seinfeld J.H.: Iodine oxide homogeneous nucleation: An explanation for
5 coastal new particle production, *Geophys. Res. Lett.*, 28, 1949–1952, 2001.
- 6 Huang R. J., Hoffmann T.: Development of a coupled diffusion denuder system combined with gas
7 chromatography/Mass spectrometry for the separation and quantification of molecular iodine
8 and the activated iodine compounds iodine monochloride and hypoiodous acid in the marine
9 atmosphere, *Anal. Chem.* 81,1777–178, 2009.
- 10 Huang R. J., Seitz K., Buxmann J., Poehler D., Hornsby K.E., Carpenter L.J., Platt U., Hoffmann T.: In
11 situ measurements of molecular iodine in the marine boundary layer: the link to macroalgae and
12 the implications for O₃, IO, OIO and NO_x, *Atmos. Chem. Phys.*, 10, 4823–4833, 2010a.
- 13 Huang, R. J., Seitz, K., Neary, T., O'Dowd, C. D., Platt, U., Hoffmann T.: Observations of high
14 concentrations of I₂ and IO in coastal air supporting iodine-oxide driven coastal new particle
15 formation, *Geophys. Res. Lett.*, 37, L03803, doi:10.1029/2009GL041467, 2010b.
- 16 Huang, R. J., Hou, X. L., Hoffmann, T.: Extensive evaluation of a diffusion denuder for the
17 quantification of atmospheric stable and radioactive molecular iodine, *Environ. Sci. Technol.*, 44,
18 5061–5066, 2010c.
- 19 Huang, R. J., Thorenz, U. R., Kundel, M., Venables, D. S., Ceburnis, D., Ho. K. F., Chen, J., Vogel, A. L.,
20 Kupper, F. C., Smyth, P. P. A., Nitschke, U., Stengel, D. B., Berresheim, H., O'Dowd, C. D.,
21 Hoffmann, T.: The seaweeds *Fucus vesiculosus* and *Ascophyllum nodosum* are significant
22 contributors to coastal iodine emissions. *Atmos. Chem. Phys.*, 13, 5255-5264, 2013.
- 23 Hughes, C., Malin, G., Turley, C., Keely, B., Nightingale, P., Liss, P.: The production of volatile
24 iodocarbons by biogenic marine aggregates. *Limnol. Oceanogr.*, 53,867-872.2008.
- 25 Jeffrey S.W., Humphrey G.F.: New Spectrophotometric equations for determining chlorophylls A, B,
26 C1 and C2 in higher plants, algae and natural phytoplankton. *Biochem. Physiol. Pfl.*, 167, 191–
27 194, 1975.
- 28 Jones C.E., Hornsby K.E., Dunk R.M., Leigh R.J., Carpenter L.J. Coastal measurements of short-lived
29 reactive iodocarbons and bromocarbons at Roscoff, Brittany during the RHaMBLe campaign,
30 *Atmos. Chem. and Phys.*, 9, 8757–8769, 2009.
- 31 Kraberg, A., Carstens, K., Peters, S., Tilly, K., & Wiltshire, K. (2012). The diatom *Mediopyxis helysia*
32 Kühn, Hargreaves & Halliger 2006 at Helgoland Roads: a success story? *Helgoland Marine*
33 *Research*, 66(3), 463–468. doi:10.1007/s10152-011-0277-9Kundel M., Thorenz U.R., Petersen
34 J.H., Huang R., Bings N.H., Hoffmann T.: Application of mass spectrometric techniques for the
35 trace analysis of short-lived iodine-containing volatiles emitted by seaweed, *Anal. Bioanal.*
36 *Chem.*, 402, 3345–3357, 2012.
- 37 Küpper F.C., Carpenter L.J., McFiggans G.B., Palmer C.J., Waite T.J., Boneberg E., Woitsch S., Weiller
38 M., Abela R., Grolimund D., Potin P., Butler A., Luther G.W., Kroneck P.M.H., Meyer-Klaucke W.,
39 Feiters M.C.: Iodide accumulation provides kelp with an inorganic antioxidant impacting
40 atmospheric chemistry, *Proc. Natl. Acad. Sci. U.S.A.*, 105, 6954–6958, 2008.
- 41 Lai S.C., Williams J., Arnold S.R., Atlas E.L., Gebhardt S., Hoffmann T.: Iodine containing species in the
42 remote marine boundary layer: A link to oceanic phytoplankton, *Geophys Res. Lett.*, 38, 2011.

- 1 Mahajan A.S., Plane J.M.C., Oetjen H., Mendes L., Saunders R.W., Saiz-Lopez A., Jones C.E., Carpenter
2 L.J., McFiggans G.B.: Measurement and modelling of tropospheric reactive halogen species over
3 the tropical Atlantic Ocean. *Atmos. Chem. Phys.*, 10, 4611–4624, 2010.
- 4 McFiggans G., Plane J.M., Allan B.J., Carpenter L.J., Coe H., O'Dowd C.: A modeling study of iodine
5 chemistry in the marine boundary layer. *J. Geophys. Res.-Atm.*, 105, 14371–14385, 2000.
- 6 Mishra S., Singh V., Jain A., Verma K.K.: Determination of iodide by derivatization to 4-iodo-N,N-
7 dimethylaniline and gas chromatography-mass spectrometry, *Analyst*, 125, 459–464, 2000.
- 8 Moore R.M., Webb M., Tokarczyk R., Wever R.: Bromoperoxidase and iodoperoxidase enzymes and
9 production of halogenated methanes in marine diatom cultures. *J. Geophys. Res.-Oc.*, 101,
10 20899–20908, 1996.
- 11 O'Dowd C.D., Hoffmann T.: Coastal new particle formation: A review of the current state-of-the-art,
12 *Environ. Chem.* 2, 245–255, 2005.
- 13 Quack B., Peeken I., Petrick G., Nachtigall K.: Oceanic distribution and sources of bromoform and
14 dibromomethane in the Mauritanian upwelling, *J. Geophys. Res.-Oc.*, 112, 2007.
- 15 van Bergeijk S.A., Hernandez Javier L., Heyland A., Manchado M., Pedro Canavate J.: Uptake of
16 iodide in the marine haptophyte *Isochrysis* sp. (T.ISO) driven by iodine oxidation, *J. Phycol.*, 49,
17 640–647, 2013.
- 18 von Glasow R. and Crutzen P.J. Tropospheric Halogen Chemistry. In: *Treatise on Geochemistry*
19 *Update1*, vol 4.02, pp 1–67, 2003.
- 20 Read K.A., Mahajan A.S., Carpenter L.J., Evans M.J., Faria B.V.E., Heard D.E., Hopkins J.R., Lee J.D.,
21 Moller S.J., Lewis A.C., Mendes L., McQuaid J.B., Oetjen H., Saiz-Lopez A., Pilling M.J., Plane
22 J.M.C.: Extensive halogen-mediated ozone destruction over the tropical Atlantic Ocean, *Nature*,
23 453, 1232–1235, 2008.
- 24 Roy R.: Short-term variability in halocarbons in relation to phytoplankton pigments in coastal waters
25 of the central eastern Arabian Sea, *Estuar Coast. Shelf. S.*, 88, 311–321, 2010.
- 26 Roy R., Pratihary A., Narvenkar G., Mochemadkar S., Gauns M., Naqvi S.W.A.: The relationship
27 between volatile halocarbons and phytoplankton pigments during a *Trichodesmium* bloom in
28 the coastal eastern Arabian Sea. *Estuar Coast. Shelf. S.*, 95, 110–118, 2011.
- 29 Saiz-Lopez A., Plane J.M.: Novel iodine chemistry in the marine boundary layer, *Geophys. Res. Lett.*,
30 31, 2004.
- 31 Saiz-Lopez A., Plane J.M., McFiggans G., Williams P.I., Ball S.M., Bitter M., Jones R.L., Hongwei C.,
32 Hoffmann T.: Modelling molecular iodine emissions in a coastal marine environment: the link to
33 new particle formation. *Atmos. Chem. Phys.* 6, 883–895, 2006.
- 34 Sakamoto Y., Yabushita A., Kawasaki M., Enami S.: Direct emission of I₂ molecule and IO radical from
35 the heterogeneous reactions of gaseous ozone with aqueous potassium iodide solution, *J. Phys.*
36 *Chem. A*, 113, 7707–7713, 2009.
- 37 Schall C., Heumann K.G., Kirst G.O.: Biogenic volatile organoiodine and organobromine hydrocarbons
38 in the Atlantic Ocean from 42 degrees N to 72 degrees, *S. Fresen. J. Anal. Chem.*, 359, 298–305,
39 1997.

- 1 Schwehr K.A., Santschi P.H.: Sensitive determination of iodine species, including organo-iodine for
2 freshwater and seawater samples using high performance liquid chromatography and
3 spectrophotometric detection, *Anal. Chim. Acta*, 482, 59–71, 2003.
- 4 Thorenz U.R., Kundel M., Müller L., Hoffmann T.: Generation of standard gas mixtures of
5 halogenated, aliphatic, and aromatic compounds and prediction of the individual output rates
6 based on molecular formula and boiling point, *Anal. Bioanal. chem.*, 2012.
- 7 Weisse T., Grimm N., Hickel W., Martens P.: Dynamics of *Phaeocystis pouchetii* blooms in the
8 Wadden Sea of Sylt (German Bight, North Sea), *Estuar. Coast. Shelf. S.*, 171–182, 1986
- 9 Wong G.T., Piumsomboon A.U., Dunstan W.M.: The transformation of iodate to iodide in marine
10 phytoplankton cultures. *Mar. Ecol.-Prog Ser.* 237, 27–39, 2002.
- 11 Wong G.T.F., Cheng X.-H.: Dissolved organic iodine in marine waters: Determination, occurrence and
12 analytical implications. *Mar.Chem.* 59, 271-281, 1998.
- 13

1 Table 1: Halocarbon emission rates, concentrations of chlorophyll α , iodide and iodate in the
 2 four different sample suspensions


Sample		F/2 medium background	<i>P. glacialis</i>	<i>M. helysia</i>	plankton concentrate
		Range	Range	Range	Range
		(Mean)	(Mean)	(Mean)	(Mean)
CH ₃ I	pmol min ⁻¹ m ⁻²	0.17 – 0.72 (0.35)	0.21 – 0.69 (0.45)	0.32 – 0.82 (0.53)	0.08 – 0.37 (0.19)
CH ₂ lCl	pmol min ⁻¹ m ⁻²	0.02 – 0.22 (0.11)	0.02 – 0.22 (0.16)	0.04 – 0.22 (0.18)	0.02 – 0.12 (0.07)
CH ₂ l ₂	pmol min ⁻¹ m ⁻²	0.02 – 0.08 (0.07)	0.27 – 0.44 (0,36)	0.21 – 0.50 (0.37)	0.04 – 0.09 (0.07)
CHBr ₃	pmol min ⁻¹ m ⁻²	1.76 – 1.90(1.81)	1.99 – 2.17 (2.09)	1.75 – 2.17 (2.09)	1.75 – 2.33 (1.82)
chl α	$\mu\text{g L}^{-1}$	n.d.	257	927	2.5
Iodide	nmol L ⁻¹	6.6 - 15.6 (10.4)	7.3 - 19.7 (12.7)	9.9 - 21.9 (16.8)	3.5 - 9.5 (6.5)
Iodate	nmol L ⁻¹	402 - 538 (428)	408 - 478 (448)	397 - 499 (446)	424 - 478 (442)
1,3-C ₃ H ₆ Br ₂ *	pmol min ⁻¹ m ⁻²	7.77 \pm 0.04	7.78 \pm 0.59	7.77 \pm 0.99	7.69 \pm 0.07
$\Sigma_{\text{Iodocarbon/chl } \alpha}$	pmol/g	n.d.	19.75	6.06	694.88

* evaporation standard given as

mean \pm standard deviation

chl α was measured for each sample once halocarbons, iodide, iodate mean values and ranges are calculated from 6 replicates


$\Sigma_{\text{Iodocarbon/chl } \alpha}$ Iodocarbon emissions were summed for the experimental conditions (time and surface area) and normalized to chl α in the watery phase



1

2 Figure 1: Experimental setup of the chamber with the phytoplankton suspension


3



4


5 Figure 2: Iodine emission rates normalized for the surface area of the different samples at 0 ppb and
 6 100 ppb ozone. The error bars represent the standard deviation of the three replicates of each
 7 experiment.

8



1

2 Figure 3: Correlation of the change in the I₂ emission and the iodide concentration in the mikro algae
 3 suspension



1
2 Figure 4: Function of the change in the total I₂ emissions in relation to the amount of consumed
3 ozone

4

5