

1 **Authors response to referee and marked-up manuscript version (page 12)**

2
3 **Anonymous Referee #1**

4
5 **1. In my opinion, the paper titled "Emission of iodine containing volatiles by selected microalgae**
6 **species" by Thorenz et al., is not suitable for publication in ACPD without major revisions.**

7
8 We revised the paper taken the points raised by the referee into account. We have also clarified the
9 connection to atmospheric science and highlighted the importance of I₂ emission from the air /water
10 interface.

11
12 Changes in the text:

13 Paragraph added in result and discussion:

14 "However, the results presented here demonstrate that even under low iodide concentrations,
15 representative of natural conditions of the MBL, a significant formation of I₂ by the ozone driven
16 oxidation of iodide at the air/water interface takes place, until the iodide concentration gets too low"

17
18 Paragraph added in Conclusion:

19 "The experiments showed that different algae suspensions (*M. herysia* and *P. glaciales*) are capable
20 of emitting I₂ by the reaction of ozone with dissolved iodide at the air/water interface under natural
21 conditions. "

22
23 **2. This is a biological incubation study of various phytoplanktons for the detection of many**
24 **iodocarbons produced under conditions that are not normal in seawater.**

25
26 We revised the paper to make clear that this study is not a classical incubation study, which
27 investigates the production of trace gases from phytoplankton as function of e.g. their growth
28 rate/phase/nutrient limitation, but that it is an (gaseous) emission study performed under conditions
29 mimicking the natural environment. We investigated the emission of different iodine species (organic
30 = iodocarbons and inorganic = I₂) from natural sea water and micro algae suspensions at the
31 air/water interface under conditions where the iodide and iodate content in the water phase are
32 representative of those in seawater. Molecular iodine is an important contributor to reactive
33 atmospheric iodine, and to date emissions studies have focused either on coastal macroalgae, or on
34 the abiotic source. This is the first study that we are aware of that has investigated I₂ emissions from
35 microalgae, which are widespread across the global oceans.

36
37 Changes in the text can be found in the corrected manuscript.

38
39 **3. The link of this work to either atmospheric physics or atmospheric chemistry is weak.**

40
41 We disagree that iodine emission from seawater at the air/water interface is not relevant for
42 atmospheric chemistry. Modelling and observational studies (e.g. Read et al., 2008; McFiggans et al.,
43 2010; Saiz-Lopez et al., 2012) show that iodine significantly reduces tropospheric ozone and in
44 certain regions constitutes an important particle nucleation mechanism. We have modified the
45 abstract, introduction and conclusion to more clearly and explicitly state how this study is relevant
46 for atmospheric research.

47
48 Changes in the text can be seen in the corrected manuscript.

49
50 **4. There is specialist jargon used in the paper that is not defined, like "F/2 aqueous media". What**
51 **are the advantages and disadvantages of using this type of media?**

1
2 "F/2 medium" is now defined and explained in the experimental section. It has been removed from
3 the abstract.

4
5 Changes in the text:

6 "Two diatom cultures (*M. helysia*, *Porosyra glacialis* both from the Alfred Wegener Institut/Sylt),
7 were kept in F/2 seawater medium for growing. These media were prepared from filtered natural
8 seawater from the shores of Sylt and additional nutrients which the diatoms need to grow (0.88
9 mmol NO₃⁻, 0.04 mmol PO₄³⁻ and 0.01 mmol SiO₃²⁻) and which is a common used medium as
10 described by Guillard et al., 1975 and Kraberg et al. 2012."

11
12
13 **5. As the authors point out, it is extremely difficult to compare the emissions measured in the**
14 **incubation studies to those in the real world. Usually, emissions in incubation studies are greater**
15 **than those measured in the real world, but here the study yields emissions two orders of**
16 **magnitude lower than measured in the real world.**

17
18 The previous studies referred to here are iodocarbon emissions from a coastal region strongly
19 influenced by macroalgae and from marine aggregates. The latter was not a "real world" study but
20 an incubation study (Hughes et al., 2008). Macroalgae are the strongest known emitters of halogens
21 and it was expected to find much lower (per area) emissions from microalgae.

22 Changes in the text:

23 "Jones et al. (2009) calculated iodocarbon emissions at a sampling site surrounded by fields of macro
24 algae in open sea water at Roscoff, France. The flux of iodocarbons was estimated to 85.28 pmol min⁻¹
25 m⁻², two orders of magnitude higher than the flux obtained in the present study. Thus it appears
26 that on an areal basis, the natural populations of microalgae studied here are much less prevalent
27 emitters of iodocarbons than seaweeds and marine aggregates."

28 **6. I would recommend that the authors submit the paper to a specialist journal in microbiology,**
29 **because it is hard to interpret this work for the atmospheric science community.**

30
31 We disagree with this statement; the emission of I₂ from seawater under natural conditions (O₃
32 mixing ratio in air and iodide concentration in seawater) is interesting for the atmospheric science
33 community (e.g. Lawler et al., 2014). In particular, the change in emission of the natural plankton
34 concentrate compared with the cultures is interesting, since the latter samples clearly show the
35 abiotic formation of I₂ from iodide and O₃ and the natural plankton sample does not. In this case the
36 formation of HOI instead of I₂ is indeed interesting for the atmospheric science community, as
37 pointed out in Carpenter et al., 2013.

38
39 **7. I think that further work is needed to understand why their observed emissions are so low.**

40
41 As we point out in lines 21-27 on page 14587, the emissions of I₂ not unexpectedly low, rather, they
42 are in good agreement with the model for the background seawater and the diatom cultures. The I₂
43 emission of the plankton concentrate is low, but also the iodide is low, therefore it is in agreement
44 with the model. We have expanded the discussion to include iodide uptake by the natural occurring
45 plankton.

46 The emissions of iodocarbons are low compared to shoreline water in a kelp field. Kelps are known as
47 big producers of iodocarbons, especially during tidal dryness. We agree with the referee that there

1 should be further studies to understand iodocarbon emission rates from different kind of
2 seawater/planktons, but the iodocarbon emission of phytoplankton was not the main goal of this
3 study (See statements two and five).

4

5

6

1 **Anonymous Referee #2**

2 Received and published: 13 August 2014

3
4 **1) The emission of halogenated compounds strongly depends on algal species (e.g. Tokarczyk and**
5 **Moore 1994), growth stage (e.g. Moore et al. 1996) and, at least for macroalgae, on environmental**
6 **factors such as irradiance (e.g. Laturus et al. 2004) and temperature (e.g. Nitschke et al. 2013).**
7 **Thus, the authors should provide essential and detailed information about the strains examined**
8 **(strain numbers), the culture/ maintenance and experimental conditions (irradiance: PAR,**
9 **probably UV [type of bulbs, fluorescent tubes], temperature, photoperiod) and the growth phase**
10 **which cultures had reached at the time of the experiment (lag, log, stationary?). Since growth of**
11 **diatoms is silica-dependent, they should also mention that this micronutrient was present (or**
12 **absent?) in the f/2 growth medium used (concentration?). I am further interested in the fact why**
13 **species from different geographical regions were chosen (temperate: *M. helysia*, Antarctic/Arctic:**
14 ***P. glacialis*). Are these species key components in their natural habitats (bloom-forming, high**
15 **abundance, high biomass)? Was the habitat temperature regime considered during maintenance**
16 **and experiments?**

17
18 We agree that the emission is strongly dependent on algal species, growth stage and environmental
19 factors. Therefore we give detailed information about the growth of the two diatom cultures, which
20 were cultured before the actual emission experiments stated in our study. Both diatom cultures were
21 personal isolates from the AWI colleagues at Sylt and were cultured in a cooling room at 16 °C in a 12
22 h light cycle using a Lumilux natural daylight lamp. The culture maintenance is now described with all
23 these details in the experimental section. The different concentrations of the nutrients added to the
24 filtered natural seawater to prepare the F/2 medium are added to the text. We had no possibility to
25 check the growth phase, therefore we do not state it. The two diatoms were chosen in cooperation
26 with our colleagues in Sylt and Helgoland. In an earlier study we investigated *Mediopyxis helysia*,
27 which is an intensively studied diatom in Helgoland (Kraberg et al. 2012), with promising results.
28 *Porosira glacialis* was chosen, because we wanted another diatom originating from a different
29 habitat and we were dependent on the isolates present in Sylt. We kept both species using the same
30 growing conditions, although their originating habitat is different, as we had no experimental
31 opportunity to culture them individually. We added the reasons for choosing the species for our
32 experiments to the manuscript in the experimental part.

33
34 Changes in the text where done according to the answer to the referee in the experimental part.

35
36 **2) Many algal species do not only release iodide into seawater (e.g. Chance et al. 2009, Nitschke et**
37 **al. 2013), where it can undergo transformation (e.g. Chance et al. 2009, Bluhm et al. 2010), but**
38 **they also efficiently absorb it (van Bergeijk et al. 2013). Such iodide uptake may explain the low**
39 **concentration in the medium as measured by the authors. Iodine uptake is a topic that the authors**
40 **should address in their discussion to place their findings into a biogeochemical context.**

41
42 The iodide uptake of algal species is important and therefore added to the discussion, especially for
43 the natural algae sample this leads to an interesting point. The uptake itself was not measured but is
44 now mentioned in the discussion of the low iodocarbon emission and low iodide concentration in the
45 natural plankton concentrate.

46
47 Changes in the text:

48 Added section: "Another possible reason for the low iodide concentration in the plankton
49 concentrate could be iodine uptake by microalgae present in the natural plankton sample (van
50 Bergeijk et al. 2013). "

51 From: "An alternative explanation is the low iodide concentration in the plankton concentrate."

1 To:” An alternative explanation is the low iodide concentration in the plankton concentrate, which
2 may be related to iodide uptake by the natural occurring plankton communities.”
3

4 **3) Regarding the halocarbon emission rates (Table 1), although mean values for CH₃I and CH₂ICl
5 differ slightly between “background” (which may be called “blank measurement”), *P. glacialis*, *M.*
6 *helysia* and the natural plankton sample, the actual range of values is quite similar. In order to
7 attribute the emission of these compounds to algal cells present, the authors may back up their
8 data by statistical analyses, meaning they should provide proof that emissions rates were
9 significantly different/higher when algae were present. I would recommend to perform either a
10 simple t-test (“background” against “culture”) or, for more information, a one-way ANOVA for
11 each compound. If the assumptions for the t-test and the one-way ANOVA are violated, the
12 authors can perform Mann-Whitney U-tests or Kruskal-Wallis tests, respectively. The term
13 “significant” can only be used when a statistical test revealed a significance. Also, the authors
14 observed the emission of CH₂I₂ from *P. glacialis*; this finding contradicts Moore et al.(1996). Any
15 explanation (see the following comments)?
16**

17 A Mann-Whitney U test was performed (which I know as Wilcoxon rank sum test) to investigate
18 whether or not the increases in CH₃I, CH₂I₂ and CH₂ICl are significant. To investigate the significance
19 the whole sample dataset was used, all 6 replicates for the iodocarbons. The test proves the
20 significant increase in iodocarbons when diatoms are present. This discussion is now added to the
21 manuscript. In our discussion we still include other probable formation pathways as considered by
22 the reviewer in the following comments. Regarding that Moore et al. did not observe that *P. glacialis*
23 produces CH₂I₂ we refer to the publication when Moore et al state: ” *P. glacialis* produced CH₂I₂
24 either in insignificant quantities or it was not possible to say that the quantities were substantially
25 different from the control treatment. This does not mean that this organism is unable to produce
26 CH₂I₂.” Therefore, we do not see a contradiction with Moore et al.. The sampling and experimental
27 conditions are very different for the presented study and the study of Moore et al, as they measured
28 production using a purge and trap system, while we measure emissions using a chamber system with
29 a constant flow and adsorption tubes which are analyzed later. One reason why we measured CH₂I₂
30 when Moore et al. did not, aside from the experimental differences, could be the different strains of
31 the micro algae, different environmental conditions in the medium or other environmental factors.
32 All these points are interesting, but are not the primary scope of the presented study (abiotic
33 formation and emission of I₂ from natural seawater to the atmosphere) However, we feel that these
34 points make a good case for investigating the production of iodocarbons by different micro algae
35 strains in more detail and with the presented method again.
36

37 Changes in the text:

38 Section added:” Additionally, the emission rates of CH₂I₂, CH₂ICl and CH₃I in the diatom samples *P.*
39 *glacialis* and *M. helysia* were significant higher compared to the background (Wilcoxon rank sum test
40 $p=0.00032$ and $p=0.00007$, respectively).”
41

42
43 **4) Were halocarbons potentially present in the growth medium? They may not originate from algal
44 cells. Where did the seawater used for media preparation originate from? For example,
45 concentrations of organic iodine species can be high in coastal or nearshore waters (Wong and
46 Cheng 1998); a fact that is neglected by the authors (see p 14581, l 10) and should be addressed.**
47

48 To be aware of emission from the growth medium we used the medium in our emission experiment
49 as we used the diatom cultures, which we explain now in more detail in the experimental part. The
50 halocarbons emitted from the growth medium were stated in Table 1 in the row f/2 medium

1 background. To emphasize the emission from the background we add the discussion about emission
2 of halocarbons from natural seawater as suggested by the referee.

3
4 Changes in the text: (paragraphs added)

5 “The emission of iodocarbons from the F/2 background is not surprising for two reasons; first, the
6 medium was produced from natural shoreline filtered water, which already may contain iodocarbons
7 (Wong and Cheng, 1998). The second reason may be related to iodocarbon-producing bacteria
8 (Amachi et al. 2001; Amachi et al. 2003). These bacteria could have been present and active in the
9 natural seawater water used to produce the F/2 medium, since it was not sterilized prior to use.”

10
11 **5) Also, some bacteria are known to take up, release and emit iodine species (Amachi et al. 2001,**
12 **Amachi et al. 2003). The presence of bacteria in the growth medium and/or in association with**
13 **algae might have influenced the results presented. For example, the authors show that**
14 **halocarbons were also emitted from pure growth medium without algal cells (Table 1: f/2 medium**
15 **“background” range). Any explanation? Was the f/2 medium sterilised before usage? Were the**
16 **algal strains axenic?**

17
18 Bacteria that are able to take up, release and emit iodine cannot be excluded in our emission
19 experiment; therefore we add them to our discussion. The emission from the medium without algal
20 cells is discussed now for both emissions due to halocarbons in the seawater used to produce the
21 medium and for bacteria.

22
23 Changes in the text:

24 see above

25
26 **6) Regarding the determination of I₂ emission rates, was the experimental set-up (Fig. 1)**
27 **characterised for potential wall losses? I₂ is quite “sticky” and large surfaces can potentially act as**
28 **efficient sink for I₂, implying that I₂ emission rates are probably underestimated. How was this**
29 **issue addressed?**

30
31 Wall losses have been evaluated for the set up described in Fig. 1 in the lab using a I₂ diffusion
32 source. We measured the I₂ mixing ratios after diluting the source flow (500ppt I₂) and compared it
33 to the mixing ratios at the end of the glass chamber described in Fig. 1 using the same flows as
34 described in the micro algae experiment. We could not observe wall losses larger than our analytical
35 precision (RSD =5%). We also checked losses to the water surface by adding ultra pure water, and
36 again we did not observe any losses to the water surface. We include our results on wall losses for I₂
37 in the experimental description.

38
39 Changes in the text: (paragraph added)

40 “Potential wall losses of I₂ and halocarbons were investigated using diffusion (I₂) and permeation
41 (halocarbons) test gas sources; no wall losses were observed within the precision of the
42 measurements using the stated gas flows.”

43
44 **7) Chlorophyll concentration can vary with environmental factors and under stress conditions; it**
45 **represents therefore not the best proxy for biomass. Chlorophyll a data may be supported by cell**
46 **numbers.**

47
48 We agree that Chlorophyll a concentrations are not the best proxy for biomass. We did not have the
49 opportunity to count the cells at the AWI in List, Sylt or to measure the dry weight or anything else,
50 therefore we cannot state cell numbers or other proxys for the biomass. Since chlorophyll a is the

1 only possibility to relate the emissions of volatile iodine compounds to the biomass, we use this
2 proxy in our manuscript.

3

4 **8) The study is placed into the field of biology and the link to atmospheric processes is relatively**
5 **weak. Thus, I am not sure that ACP is a suitable journal for the work presented. In any case, before**
6 **publication, the authors should address the above points.**

7

8 We disagree with the referee in this point. The study is placed in the field of atmospheric processes,
9 since the emission of molecular iodine and iodocarbons from the hydrosphere to the atmosphere is
10 important for many atmospheric processes (ozone depletion, particle formation, perturbation of
11 oxidative cycles in the atmosphere, (Carpenter et al., 2013)). We tried to clarify the atmospheric
12 relation of the study in the text, by clarifying that the study is not an incubation study but an
13 emission study. Additionally we clarified the importance of the emission of I₂ (which is formed
14 abiotically) versus the emission of iodocarbons (which are related to the field of biology). We also
15 tried to reduce the biological language (by reducing the use of biology related vocabulary like F/2
16 medium) to clarify that the study is not related to biology in the first place, but we still kept all the
17 explanations of micro algal processes, since one goal of the study was to relate the biological
18 formation of iodide from iodate by microalgae and the atmospheric relevant process of I₂ formation
19 at the air/water interface.

20

21 Changes in the text:

22 There are several changes in the text, which can be seen in the corrected manuscript, as example just
23 a few are mentioned here.

24 Paragraph added in result and discussion:

25 “Comparing the I₂ and iodocarbon emission rates, it is clear that the volatile iodine emissions are
26 dominated by I₂. Therefore I₂ emissions from natural seawater surfaces are more relevant for
27 atmospheric processes than the emission of iodocarbons.”

28 Paragraph added in Conclusion:

29 “The emission rates of iodocarbons were also lower than the emission of I₂, confirming that I₂
30 emissions from the remote ocean dominate over organic iodine sources for the MBL (Jones et al.,
31 2010; Lawler 2012; Carpenter et al., 2013).”

32

33

1 **Anonymous Referee #3**

2

3 **- For stoichiometrically comparing production rates, it would be desirable to compute**
 4 **the rates for halocarbons and iodine (not only for iodide and iodate) in moles (Table 1).**

5

6 To make it easier to stoichiometrically compare the halocarbon emission rates, they were computed
 7 to pico moles. Table 1 was changed accordingly. The summed emission rates in the text were also
 8 changed. The emission rate of iodine, shown in the Figures 2 and 3, were also changed to pico moles.
 9 Therefore, it is now straightforward to compare the emission rates of iodocarbons and I₂ as reported
 10 in this study with emission or production rates from other studies.

11

12 Changes in the text:

13 From:

14 "The emission flux summed for the three iodocarbons in the four samples F/2-medium, plankton concentrate,
 15 *P. glacialis* and *M. helysia* was in the range of 0.034 – 0.163 ng min⁻¹ m⁻², 0.025 – 0.098 ng min⁻¹ m⁻², 0.106 –
 16 0.264 ng min⁻¹ m⁻² and 0.153 – 0.288 ng min⁻¹ m⁻², respectively."

17

18 To:

19 "The emission flux summed for the three iodocarbons in the four samples' background, plankton concentrate,
 20 *P. glacialis* and *M. helysia*, was in the range 0.21 – 1.02 pmol min⁻¹ m⁻², 0.14 – 0.58 pmol min⁻¹ m⁻², 0.50 – 1.35
 21 pmol min⁻¹ m⁻² and 0.57 – 1,53 pmol min⁻¹ m⁻², respectively."

22


23

24 Changes in Table 1:


Sample		F/2 medium background	<i>P. glacialis</i>	<i>M. helysia</i>	plankton concentrate
		Range (Mean)	Range (Mean)	Range (Mean)	Range (Mean)
CH ₃ I	pmol min ⁻¹ m ⁻²	0.17 – 0.72 (0.35)	0.21 – 0.69 (0.45)	0.32 – 0.82 (0.53)	0.08 – 0.37 (0.19)
CH ₂ I ₂	pmol min ⁻¹ m ⁻²	0.02 – 0.22 (0.11)	0.02 – 0.22 (0.16)	0.04 – 0.22 (0.18)	0.02 – 0.12 (0.07)
CH ₂ I ₂	pmol min ⁻¹ m ⁻²	0.02 – 0.08 (0.07)	0.27 – 0.44 (0,36)	0.21 – 0.50 (0.37)	0.04 – 0.09 (0.07)
CHBr ₃	pmol min ⁻¹ m ⁻²	1.76 – 1.90(1.81)	1.99 – 2.17 (2.09)	1.75 – 2.17 (2.09)	1.75 – 2.33 (1.82)
chl α	µg L ⁻¹	n.d.	257.27	926.59	2.53
Iodide	nmol L ⁻¹	6.60 - 15.69 (10.35)	7.32 - 19.71 (12.70)	9.90 - 21.94 (16.84)	3.52 - 9.45 (6.47)
Iodate	nmol L ⁻¹	402 - 538 (428)	408 - 478 (448)	397 - 499 (446)	424 - 478 (442)
1,3-C ₃ H ₆ Br ₂ *	pmol min ⁻¹ m ⁻²	7.77 ± 0.04	7.78 ± 0.59	7.77 ± 0.99	7.69 ± 0.07
Σ _{Iodocarbon} /chl α	pmol/g	n.d.	19.75	6.06	694.88

25

26 Figures changed:



1



2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

- Also, for comparing rates between species, it would be desirable to express them on a basis per g (or mg) chlorophyll a or maybe even g fresh weight (or dry weight) – if the conversion factor between chlorophyll and biomass is known (Table 1).

Since the conversion factor is not known the emission rates cannot be expressed on fresh or dry weight basis. The summed iodocarbon emission rates of the three algae suspensions were expressed on the basis per g chlorophyll a, this information was added to table 1.(see above) We think it is inappropriate to discuss the iodocarbon emission expressed on the basis of chlorophyll a in the text, since the reader could misinterpret emission as formation. Chl a is a concentration measured for the algae suspension, the iodocarbons were measured in the gas phase passing above the surface of the algae suspensions. This study is not a classical incubation study to investigate the formation of halocarbons by microalgae and to measure the concentrations built up in the water. The study presented here is an emission study to investigate the emission of iodocarbons by aqueous suspensions containing different microalgae. As a consequence we prefer not to discuss emission rates based on chlorophyll a in the text, however, for comparison to other studies the sum of iodocarbon emission based on chlorophyll a is stated in Table 1.

- There seems to be a discrepancy between the text and Table 1. Is the iodate concentration actually going up or down over time in a batch culture? And, if it does go down (in case it is reduced as the text suggests) can the differential amount of moles

1 **iodine be traced – in other words, does that iodate become iodide, molecular iodine,**
2 **iodocarbons, or a combination of all these?**

3

4 Since we did not measure the iodate and iodide concentrations during the course of the experiment
5 we are unable to comment this issue. The concentrations were measured after the cultures had 4
6 weeks for growing. The F/2 medium was treated the same way, however, without any micro algae
7 cultures in it. We therefore assume that the background is a representative background for the
8 iodide and iodate concentrations in the plankton samples and is comparable to the samples before
9 the micro algae were grown.

10

11 When we compared the iodide and iodate concentrations of the different samples and stated that
12 the iodate concentrations were in the same range for all samples, we presume that the amount of
13 iodate which the micro algae are able to reduce to iodide is not measurable, since the concentration
14 differences are too low to be measured with the analytical precision of the methods used. When we
15 discuss the slightly elevated iodide concentrations in the micro algae cultures compared to the
16 background, we are aware that this observation is not statistically significant, however, at least an
17 indication that the microalgae indeed reduced iodate to iodide, as expected from previous studies.

18

19 The iodate concentrations are between 397 and 538 nmol L⁻¹ for the different samples, about two
20 orders of magnitude higher than the iodide concentrations. Therefore, the reduction of iodate falls
21 within our analytical precision. For the iodide measurements the analytical precision is much higher
22 and the natural variability of the concentrations lower, therefore, we believe that the discussion of
23 the formation of iodide is scientifically sound.

24

25 To improve this part of the manuscript, we introduced changes in the individual sections on iodide
26 and iodate in the results and discussion part.

27

28 Changes in the text:(the following paragraph was added)

29 “Such a reduction of iodate to iodide will result in a decrease in the iodate concentration, however, for the
30 measured iodate concentration in this study the expected decrease falls within the analytical precision of the
31 measurement. The iodide concentrations in all samples are comparable with oceanic surface water
32 concentrations, for example around 10-30 nmol L⁻¹ in the Weddel Sea surface water (Bluhm et al. 2011).”

33

34 **- A particularly interesting question which the manuscript does not address or even**
35 **raise at all: Actually do the data tell us or suggest anything, which is the precursor**
36 **(iodine source) for the formation of iodocarbons – iodide or iodate?**

37

38 This is indeed an interesting question, however, we think it is not possible to answer this question
39 based on the results and experimental set up chosen for this study. We measured the iodocarbons in
40 the gas phase and it is obvious that they are released from the different algae solutions. The emission
41 rates of iodocarbons measured for the background sample suggests that iodocarbons were already
42 present in the seawater which was used to prepare the media to grow the diatoms. The emissions
43 rate of iodocarbons in the diatom samples are elevated compared to the background, therefore we
44 assume that diatoms are capable of producing iodomethanes, however, we cannot judge based on
45 our experiments if they use iodate or iodide. We know that iodide is favoured for the biotic (SAM,
46 haloperoxidase) and abiotic (photochemical formation with DOM (Moore and Zafirou, 1994))
47 formation reactions, since iodine has the same oxidation state in iodide as it has in iodocarbons (-1),
48 however, whether the micro algae reduce iodate to iodide to form the halocarbons or if they use the
49 iodide directly which is already present in the water cannot be resolved. What we can assume is the
50 discrepancy in the emission of iodocarbons and I₂. The iodocarbon emissions are not related to the

1 formation of I₂ at the air/water interface, since the formation of iodocarbons is not different for high
2 and low ozone conditions. This conclusion was added to the I₂ emission section in the manuscript.

3

4 Changes in the text: (the following paragraph was added):

5 “Comparing the I₂ and iodocarbon emission rates, it is clear that the volatile iodine emissions are dominated by
6 I₂. Therefore I₂ emissions from natural seawater surfaces are more relevant for atmospheric processes than
7 the emission of iodocarbons. At the same time the experiments presented here show that the emission of
8 iodocarbons is not linked to the formation of I₂ at the air/water interface (c.f. Martino et al., 2009), since no
9 correlation between I₂ emissions or O₃ mixing ratio and iodocarbon emissions was observed.”

10

1 Emission of iodine containing volatiles by selected microalgae species

2

3 U. R. Thorenz^{1*}, L. J. Carpenter², R.-J. Huang^{1,3,4}, M. Kundel¹, J. Bosle¹ and T. Hoffmann¹

4 [1] Institute of Inorganic and Analytical Chemistry, Johannes Gutenberg-University of Mainz,
5 55128 Mainz, Germany

6 [2] Department of Chemistry, University of York, Heslington, York YO10 5DD, UK

7 [3] School of Physics & Centre for Climate and Air Pollution Studies, Ryan Institute, National
8 University of Ireland Galway, University Road, Galway, Ireland

9 [4] Laboratory of Atmospheric Chemistry, Paul Scherrer Institute (PSI), 5232 Villigen, Switzerland

10 *Now at the Max Planck Institute for Chemistry, 55128 Mainz, Germany

11 Correspondence to: T. Hoffmann (t.hoffmann@uni-mainz.de)

12

13 Abstract

14 | In this study we present the results of an incubation-emission study of different phytoplankton
15 | samples in F/2-aqueous media treated with elevated ozone levels. Halocarbon measurements show
16 | that the samples tested released bromoform and different iodocarbons including iodomethane,
17 | iodochloromethane and diiodomethane. Iodide and iodate levels in the liquid phase were
18 | representative of concentrations of surface water in a natural environment. Measurement of volatile
19 | iodine (I₂) emissions from two diatom samples (*Mediopyxis helysia* and *Porosira glacialis*) and the
20 | background sample (F/2-medium from locally-seawater-filtered natural seawater), showed that the
21 | quantity of I₂ evolved depends on the ozone concentration in the air. This behaviour was assumed to
22 | be caused by the oxidation reaction mechanism of iodide with ozone. The I₂ emission flux agrees
23 | with model calculations at different iodide concentrations. The I₂ emission of a natural plankton
24 | concentrate sample was, however, very low compared to other samples and showed no dependence
25 | on ozone. The reason for this was shown to be the low iodide concentration in the algae suspension,
26 | which seems to be the limiting factor in the oxidative formation of I₂.

27

28 Introduction

29 Iodine chemistry plays an essential role in the marine boundary layer (MBL) due to its effect on the
30 destruction of tropospheric ozone, perturbation of the HO_x/NO_x cycle and the formation of new
31 particles and cloud condensation nuclei, thereby leading to changes in the global radiative forcing
32 (Hoffmann et al., 2001; von Glasow and Crutzen, 2003; O'Dowd and Hoffmann, 2005; Bloss et al.,
33 2005; Huang et al., 2010a, b). This essential role of iodine and of other activated halogens is shown in
34 field measurements in the marine boundary layer (MBL), laboratory chamber experiments or
35 incubation experiments of different algae and in atmospheric models (Carpenter, 2003; Küpper et al.,

1 2008; Kundel et al., 2012; McFiggans et al., 2000). The biogeochemical cycle of iodine is controlled by
2 large iodine exchanges from the oceans to the atmosphere, driven by marine biotic and abiotic
3 production (Schall et al., 1997). Volatilized species are photolabile iodocarbons like CH_2I_2 , CH_3I , $\text{C}_2\text{H}_5\text{I}$,
4 CH_2ICl , CH_2IBr and molecular iodine (I_2). Marine species like macroalgae and microalgae play a
5 dominant role in the emission of these compounds (Carpenter et al., 1999, Huang et al., 2013, Saiz-
6 Lopez and Plane, 2004).

7 Since molecular iodine and iodocarbons are photochemically unstable (lifetimes between about
8 some tens of seconds for I_2 and a few days for CH_3I) they are photolysed under UV-visible light to
9 form $\text{I}\cdot$ atoms, which are then instantly oxidised by ozone to form the iodine monoxide radical $\text{IO}(\text{g})$
10 (Hoffmann et al., 2001; Saiz-Lopez et al., 2006). Further oxidation reactions of IO in the gas phase
11 then can form low volatile iodine oxides (I_xO_y) which may nucleate under certain conditions and form
12 new particles.

13 Recently it was proposed that the ozone loss over the tropical Atlantic Ocean was higher than
14 calculated from global atmospheric models, and that this additional ozone destruction is induced by
15 halogens such as bromine and iodine (Read et al., 2008). Biogenic emissions, such as the already
16 studied iodocarbon emissions by phytoplankton species, e.g. coccolithophorids, diatoms and
17 chlorophytes, (Colomb et al. 2008) are too low to explain the differences in model calculations and
18 observations (Mahajan et al. 2010), therefore additional sources of the reactive iodine species are
19 discussed, one of them being the surface reaction of ozone with seawater.

20 Garland and Curtis first discovered that the emission of molecular iodine from the surface of artificial
21 and natural seawater is proportional to the ozone concentration at the air/water interface (Garland
22 and Curtis, 1981). Sakamoto and co-workers examined the reaction mechanism of the iodide
23 oxidation by ozone at the air/water interface, resulting in the formation of the intermediates IOOO^-
24 and HOI and the emission products IO and I_2 (Sakamoto et al., 2009). Further laboratory experiments
25 show that different organics affect the reaction of iodide with ozone, e.g. fulvic acid enhances the I_2
26 formation, but not the formation of IO (Hayase et al., 2010, 2012).

27 Since the formation of I_2 and IO from the air/water interface is dependent on the iodide
28 concentration in seawater, the reaction path found by Garland and Curtis may explain elevated
29 iodine emissions in areas of higher phytoplankton activity (Garland and Curtis, 1981). The ability of
30 different phytoplankton, e.g. diatoms, to reduce iodate, which is ubiquitous in the open ocean, to
31 iodide was shown for natural and elevated iodate concentrations (Wong et al. 2002; Chance et al.
32 2007) and for the different growth states (Bluhm et al. 2010) of the phytoplankton cultures. A
33 correlation of iodine species in the particle phase and average chlorophyll exposure of air masses
34 along back trajectories was found by Lai et al, 2011, indicating the link between phytoplankton
35 activity and emission of atmospheric iodine.

36 Since the formation of I_2 and IO is correlated to the iodide concentration (Sakamoto et al. 2009) and
37 the iodide concentration of surface waters is correlated to phytoplankton (Bluhm et al. 2010), this
38 study investigates links between phytoplankton, iodide concentrations in microalgae-containing
39 seawater and abiotic formation and emission of I_2 emission, utilising laboratory experiments of the
40 reaction of the seawater surface with ozone.

1 Materials and Methods

2 Experimental set-up

3 Two diatom cultures (*M. helysia*, *Porosira glacialis*) both from the Alfred Wegener Institut/Sylt,
4 were kept in F/2 seawater medium for growing. These media were prepared from filtered natural
5 seawater from the shores of Sylt and additional nutrients which the diatoms need to grow (0.88
6 mmol NO₃⁻, 0.04 mmol PO₄³⁻ and 0.01 mmol SiO₃²⁻) and which is a common used medium as
7 described by (Guillard, et al., 1975) and Kraberg et al. 2012. Both cultures were kept incubated in the
8 F/2 medium at 16°C with 12-h-light-12-h-dark cycling (LUMILUX Plus Eco daylight lamp; approx.. 40
9 μmol PAR) for at least 4 weeks prior the experiment. Just before the emission experiment, the algal
10 suspensions were then diluted in a 2:1 ratio in F/2 medium and homogenised by stirring. In addition
11 to the diatom cultures, a plankton concentrate was collected from the North Sea (55°01.562N;
12 8°27.113E) on May 24th 2012 using a 80 μm and 200 μm Apstein plankton net and diluted using the
13 same F/2 medium as for the diatom cultures. Microscopic observations showed that the plankton
14 concentrate sample was dominated by colonies of the haptophyte *Phaeocystis sp.* and only a low
15 amount of diatoms was were present in the sample, as determined by a microscopic analyses.

16 For each experiment, 1.5 L of the sample (i.e. diatom suspension, natural plankton concentrate or
17 background (F/2 medium)) was introduced into a glass chamber tube (10 L), shown in Figure 1, and
18 three magnetic stirrers were switched on immediately. A continuous flow of synthetic air (3.4 L min⁻¹)
19 was channelled over the stirred algae suspension in the first experiment with no ozone and in the
20 second experiment with elevated ozone levels of 100 ppb. The ozone was generated using an UV
21 radiation source and the resulting ozone levels were measured using an ozone analyzer (Dasibi
22 Environmental Corp. Model 1008-RS, Glendale, USA). To measure the emission of I₂ and halocarbons,
23 α-cyclodextrin-coated denuders (Huang and Hoffmann, 2009; Huang et al., 2010c) and adsorption
24 tubes (Kundel et al. 2012) were mounted at the other end of the tube chamber together with the
25 ozone monitor. The chamber outflow was sampled using two membrane pumps, one with 0.50 L min⁻¹
26 for the denuders and the other using 0.15 L min⁻¹ for the adsorption tubes. To assure an
27 overpressure over the sampling time a U-shaped tube filled with ultra-pure water was mounted in
28 the centre exit of the glass chamber to measure the overpressure hydrostatically. The whole set-up
29 was wrapped with aluminium foil to prevent photolysis of I₂ and halocarbon compounds. Potential
30 wall losses of I₂ and halocarbons were investigated using diffusion (I₂) and permeation (halocarbons)
31 test gas sources; no wall losses were observed within the precision of the measurements using the
32 stated gas flows.

33 To monitor the emissions of I₂ and halocarbons from the liquid samples, an evaporation standard
34 was added to the microalgaemicroalgal-suspension in order to highlight any problems related to air
35 sampling. This standard was 1,3-dibromopropane diluted in ultrapure water (500 μl of 0.94 μg L⁻¹
36 which was then diluted with the sample to 1.5 L). The standard was chosen given the results from a
37 first set of experiments with *M. helysia* and *Coscinodiscus wailesii* which show no detectable traces of
38 this compound. We decided not to add any iodine containing compounds to prevent interferences
39 with the I₂ emission.

40 Halocarbon measurements

1 | Air samples of 6.75 L sampling volume were pre-concentrated at a flow rate of 150 ml min⁻¹ on
2 thermal desorption tubes filled with 100 mg Tenax TA 60/80 and 150 mg Carbotrap™ 20/40 both
3 provided by Supelco (Bellefonte, PA, USA). The samples were analysed using a self-made thermal
4 desorption device mounted on a gas chromatograph (TraceGC, Thermo Scientific, Dreieich Germany)
5 - mass spectrometer (PolarisQ, Thermo Scientific, Dreieich, Germany). During the desorption period
6 of 6 minutes the cryotrap was cooled to -160 °C. Afterwards the cryotrap was rapidly heated to 270
7 °C for injection. The analytes were separated on a DB624 Durabond column (60 m; 0,32 mm; 1,8 µm
8 FT) using helium as carrier gas with a constant pre-column pressure of 0.5 bar. The temperature
9 program was: 55 °C (4 min), ramp with 5 °C min⁻¹ to 120 °C (4 min) and ramp with 8 °C min⁻¹ to 200 °C
10 (4 min). Halocarbons were detected using a mass spectrometer in NCI mode with methane as
11 reagent gas (2.5 ml min⁻¹), the primary electron energy was set to 120 eV and an emission current of
12 50 mA in single ion monitoring mode (SIM) was used. Iodinated compounds (CH₃I, C₂H₅I, CH₂ICl,
13 CH₂I₂, 1-C₃H₇I, 2-C₃H₇I, 1-nC₄H₉I, 2-nC₄H₉I, 1-iso-C₄H₉I) were quantified using *m/z* 127 and
14 brominated compounds (CH₂Br₂, CH₃Br, 1,3-C₃H₆Br₂) were quantified using *m/z* 79 and 81 at a 1:1
15 ratio. A five point calibration was done in the range between 0.01 ng and 1 ng using the continuously
16 diluted output of a permeation test gas source (Thorenz et al. 2012). The detection limits for the
17 individual iodocarbons were 0.003-0.088 ppt and for the bromocarbons were 0.004 – 0.009 ppt. For
18 each series of measurements, the calibration was done in triplicate (precision of method 3-13%).

19 I₂, Iodide and Iodate measurements

20 Sampling of gaseous I₂ was performed using the denuder technique described by Huang and
21 Hoffmann, 2009. Brown glass denuder tubes (6 mm i.d., 50 cm length) were coated using a α-
22 cyclodextrin suspension (2.5 mg mL⁻¹ in methanol) and sealed with polypropylene caps. Before
23 sampling the denuders were stored in a fridge. For sampling the denuders were mounted vertically
24 with a glass tube of 15 cm upstream to achieve laminar flow. The sampling flow was 500 mL min⁻¹ for
25 45 min. After sampling the denuders were sealed and stored in a fridge until derivatization. For
26 derivatization the α-cyclodextrin coating was eluted with ultrapure water (20 mL), then 25 µL N,N-
27 dimethylaniline (1 µg mL⁻¹ in methanol), 500 µL phosphate buffer (pH 6.4) and 500 µL 2-
28 iodosobenzoate (4 mg mL⁻¹) were added, the mixture was shaken for 2 hours. After adding 3 ml
29 sodium acetate the sample was extracted with 100 µL cyclohexane and 100µL 2,4,6-tribromoaniline
30 (internal standard: IS) in cyclohexane (250ppb).

31 Iodide and iodate were derivatized from seawater to form the same product as described for I₂.
32 Iodide was oxidized to form I₂ by using iodosobenzoate and iodate was reduced first to iodide and
33 then oxidized to form I₂. 10 mL aliquots of seawater were analysed for iodide and for total iodine,
34 iodate was calculated by difference. The method for iodide derivatization was slightly changed from
35 the one described by Mishra et al. (2000). The use of sodium hydrogen sulfite as an agent to reduce
36 iodate to iodide is described by Schwehr and Santschi (2003).

$$37 \quad I_{\text{seawater}} = I^- + IO_3^-$$

38 To measure iodide, 10 mL seawater were mixed with 1 ml ethylenediaminetetraacetic acid solution
39 (0.5%), 500 µL phosphate buffer, 500 µL N,N-dimethylaniline, 500 µL iodosobenzoate and shaken.
40 After adding 3 ml sodium acetate the sample was extracted with 100 µL cyclohexane and 100 µL
41 2,4,6-tribromoaniline (IS) in cyclohexane (250 ppb).

1 To measure iodate an aliquot of 10 mL seawater was mixed with 1 mL ethylenediaminetetraacetic
2 acid solution (0.5%), 1 mL hydrochloric acid (3.7%) and 500 µL sodium hydrogen sulfite solution
3 (283.9 µmol L⁻¹) to reduce the iodate. Afterwards 500µL sodium acetate, 4 mL phosphate buffer, 500
4 µL N,N-dimethylaniline, 500 µL iodosobenzoate were added. After shaking the sample was again
5 extracted with 100 µL cyclohexane and 100µL 2,4,6-tribromoaniline (IS) in cyclohexane (250 ppb).

6 1 µL of the cyclohexane extract was injected to the GC-MS System (6850 GC & 5973 MS, Agilent
7 Technologies, Waldbronn, Germany) at a constant flow of 1 mL min⁻¹ of helium (99.999%) and, the
8 chromatographic separation was performed using a capillary column FS Supreme 5 MS with a length
9 of 30 m, inner diameter of 0.25 mm and film thickness of 0.25µm (CS Chromatographie Service,
10 Langenwehe, Germany) with a temperature program starting at 50°C (for 3 min), then heating up at
11 30 °C min⁻¹ to 220°C (for 3 min). The mass spectrometer measured in electron ionisation mode at 70
12 eV, the specific fragments of the product 4-iodo-N,N-dimethylaniline was extracted at *m/z* 247 (M+)
13 and of the internal standard 2,4,6-tribromoaniline at *m/z* 329 (M+).

14 Chlorophyll measurements

15 The analytical method for chlorophyll α (chl α) measurements is described by Edler et al 1979. An
16 aliquot of 50-100 mL water samples were filtered on glass fibre filters (GF/F-Whatman). The dry
17 filters were put in polypropylene vials and extracted with 7.5 ml acetone. The extract was stored
18 together with the filter in a dark fridge at 3°C overnight and centrifuged the next day (5500 rpm, 7
19 min) at 5°C. The absorption of the supernatant was measured against acetone using an Uvikon XL
20 double beam spectrophotometer at $\lambda = 750$ nm, 663 nm, 645 nm and 630 nm. To calculate the
21 concentration of chl α the equation of Jeffrey and Humphrey, 1975 was used-. Chl α can be a good
22 indicator for microalgae biomass (Roy 2010; Bluhm et al. 2010; Colomb et al. 2008), and has been
23 used to calculate emission rates of iodine-containing volatiles from phytoplankton. This calculation
24 was not used here, since the mechanisms of synthesis and release of these iodine containing gases is
25 still unclear. All gaseous compounds in this study are therefore given as measured mixing ratio and
26 the chl α value of the corresponding algae suspension is listed also given aside.

27 Results and Discussion

28 Halocarbons

29 The emission rates of the natural halocarbons and the evaporation standard, given in Table 1 **Fehler!**
30 **Verweisquelle konnte nicht gefunden werden.**, were calculated by the amount measured in the
31 adsorption tubes divided by the emission time and the surface area of the suspension sample
32 (pmol ng min⁻¹ m⁻²). The halocarbon emission rates showed no effect on the different ozone levels;
33 therefore the data for each sample are summarized for high and low ozone conditions. An
34 evaporation standard was added to the different samples to recognize differences in emission rates
35 of the organic compounds from the aqueous phase. The standard was added in a 10 to 100-fold
36 excess compared to natural concentrations of bromocarbons in Atlantic seawater (Carpenter et al.
37 2000) to reduce the effect of natural 1,3-dibromopropane which may alter the mixing ratio of the
38 evaporation standard measured. In the chosen concentration a natural abundance would change the
39 result only by 1-10% compared to the spike solution. The results of the measurements of 1,3-
40 dibromopropane showed very constant values, as can be seen from the low standard deviation

1 between the different samples and replicates. This result indicates a stable and reliable experimental
2 setup in terms of evaporation of volatile compounds from the water surface and of the mixing of the
3 bulk water.

4 The measured emission rates of the natural halocarbons show that the brominated compound,
5 CHBr_3 , is elevated compared to the iodocarbons emission rates. This result ~~actually~~ fits to
6 observations of the natural abundance of halocarbons in seawater as described in earlier studies
7 (Roy et al. 2011). The emission rate of CHBr_3 is higher for the two diatom cultures (*M. helysia* and *P.*
8 *glacialis*) than for the plankton samples containing *Phaeocystis sp.* and the background ~~F/2 medium~~.
9 Again, this result matches field and laboratory data showing a link between elevated CHBr_3
10 concentrations in seawater and the simultaneous occurrence of diatoms (Colomb et al. 2008, Quack
11 et al. 2007, Moore et al. 1996).

12 The iodocarbon emissions in experiments using the background (F/2 medium) and the plankton
13 concentrate were dominated by CH_3I , followed by CH_2ICl and CH_2I_2 . However, for the diatom
14 cultures, CH_2I_2 was the dominant iodocarbon emitted with CH_3I and CH_2ICl both showing lower
15 emission rates. The emission of iodocarbons from the F/2 background is not surprising for two
16 reasons; first, the medium was produced from natural shoreline filtered water, which already may
17 contain iodocarbons (Wong and Cheng, 1998). The second reason may be related to iodocarbon-
18 producing bacteria (Amachi et al. 2001; Amachi et al. 2003). These bacteria could have been present
19 and active in the natural seawater water used to produce the F/2 medium, since it was not sterilized
20 prior to use. Additionally, the emission rates of CH_2I_2 , CH_2ICl and CH_3I in the diatom samples *P.*
21 *glacialis* and *M. helysia* were significant higher compared to the background (Wilcoxon rank sum test
22 $p=0.00032$ and $p=0.00007$, respectively). This increase in emission can be explained by the capability
23 of the diatoms to produce iodocarbons which had already been reported by Moore et al. (1996). To
24 compare the natural plankton concentrate with the cultivated diatom cultures and the background
25 one must keep in mind that chl_a concentrations are biomass tracers reflecting the abundance of
26 phytoplankton. The results for the chl_a measurement, given in Table 1, clearly show that the
27 natural plankton concentrate contains less biomass than the cultured diatoms. Therefore, we
28 conclude that the lower iodocarbon emissions of the plankton concentrate compared to the diatom
29 cultures is partly due to lower biomass density. The lower iodocarbon emission rates in the natural
30 plankton concentrate could also be related to iodine uptake of natural occurring micro algae (van
31 Bergeijk et al. 2013).

32 The emission flux summed for the three iodocarbons in the four samples' ~~F/2-mediumbackground~~,
33 plankton concentrate, *P. glacialis* and *M. helysia*, was in the range 0.21 – 1.02 $\text{pmol min}^{-1} \text{m}^{-2}$, 0.14 –
34 0.58 $\text{pmol min}^{-1} \text{m}^{-2}$, 0.50 – 1.35 $\text{pmol min}^{-1} \text{m}^{-2}$ and 0.57 – 1.53 $\text{pmol min}^{-1} \text{m}^{-2}$ of 0.034 – 0.163 ng
35 $\text{min}^{-1} \text{m}^{-2}$, 0.025 – 0.098 $\text{ng min}^{-1} \text{m}^{-2}$, 0.106 – 0.264 $\text{ng min}^{-1} \text{m}^{-2}$ and 0.153 – 0.288 $\text{ng min}^{-1} \text{m}^{-2}$,
36 respectively. We are not aware of ~~emission incubation~~ studies investigating the flux of iodocarbons
37 from micro algae suspensions to directly compare these results. To establish a connection to other
38 experimental observations the results listed above are compared to incubation studies of marine
39 aggregates producing iodocarbons and calculated emission fluxes ~~for open in coastal, seaweed-rich~~
40 regions-sea water. Hughes et al. (2008) measured the iodocarbon production of different marine
41 aggregates to be within 0.71 to 6.90 $\text{pmol h}^{-1} \text{L}^{-1}$ to 66 $\text{ng min}^{-1} \text{L}^{-1}$. The production rate is difficult to
42 compare to the presented results, since the flux in our study is based on the production by the

1 microalgae species and evaporation from the surface, whereas Hughes et al. (2008) measured the
2 production in the aqueous phase. Jones et al. (2009) calculated iodocarbon emissions at a sampling
3 site surrounded by fields of macro algae in open sea water at Roscoff, France. The flux of
4 iodocarbons was estimated to 85.28 pmol min⁻¹ m⁻², two orders of magnitude higher than the flux
5 obtained in the present study. Thus it appears that on an areal basis, the natural populations of
6 microalgae studied here are much less prevalent emitters of iodocarbons than seaweeds and marine
7 aggregates.

9 Iodide and iodate

10 The concentrations of iodide and iodate in the different samples are shown in Table 1. For each
11 sample, the mean and range for six replicates are shown; no differences in iodide and iodate
12 concentrations were observed under elevated (100 ppb O₃) and low ozone (0 ppb O₃) conditions.

13 The iodate concentrations in the background ~~F/2 medium~~ and in the three plankton samples were in
14 the same range, with mean concentrations between 438 and 448 nmol L⁻¹. These iodate
15 concentrations are in the range measured for the open ocean of 400 to 500 nmol L⁻¹ iodate in most
16 oceanic regions (Bluhm et al. 2011). The ~~ubiquitous~~ubiquity of iodate suggests that its concentration
17 is not a limiting factor.

18 The iodide concentrations in the two diatom cultures, *P. glacialis* and *M. helysia*, are slightly elevated
19 with mean values of 12.70 nmol L⁻¹ and 16.84 nmol L⁻¹, respectively, compared to the ~~F/2~~
20 ~~medium~~background iodide concentration of 10.35 nmol L⁻¹ and the plankton concentrate iodide
21 concentration of 6.47 nmol L⁻¹. This enhanced iodide concentration indicates the reduction of iodate
22 by the two diatom cultures, which was also found by Bluhm et al. 2010 and Wong et al. 2002 for
23 different phytoplankton species. Such a reduction of iodate to iodide will result in a decrease in the
24 iodate concentration, however, for the measured iodate concentration in this study the expected
25 decrease falls within the analytical precision of the measurement. The iodide concentrations in all
26 samples are comparable with oceanic surface water concentrations, for example around 10-30 nmol
27 L⁻¹ in the Weddel Sea surface water (Bluhm et al. 2011).

28 The low iodide concentration of the plankton concentrate sample compared to the ~~F/2~~ background
29 sample is surprising, but may be assigned to an overall low level of different nutrients, like phosphate
30 and silicate, in the Wadden Sea of Sylt at springtime (Weisse et al. 1986), although the level of iodate
31 was consistent. Another possible reason for the low iodide concentration in the plankton
32 concentrate could be iodine uptake by microalgae present in the natural plankton sample (van
33 Bergeijk et al. 2013).

34 Ozone measurements


35 The results of the ozone measurement for the samples: ~~F/2 medium~~background, *P. glacialis*, *M.*
36 *helysia* and the plankton concentrate were normalized against a background measurement obtained
37 using ultra-pure water in the chamber. This was performed in order to account for losses of ozone
38 through wall reactions, losses on the water surface, and losses due to droplet formation from
39 stirring. The ozone consumption was calculated using a Continuous Stirred-Tank Reactor (CSTR)
40 approach with 668 ng min⁻¹ ozone (100 ppb) introduced into the chamber (total volume: 10 L, flow:

1 3.4 L min⁻¹ and residence time: 2.94 min). The difference between the introduced ozone flow and
 2 measured ozone flow is considered as consumed ozone, due to the oxidation of iodide and other
 3 ozone depleting reactions in the samples. To calculate the consumed ozone, the flow rate was
 4 summarized over 45 min of the experiment. Ozone consumption was clearly observed for all
 5 samples. The F/2-background sample showed the weakest ozone consumption of 58 nmol, followed
 6 by the sample of *P. glacialis* with 186 nmol and the plankton concentrate with 253 nmol. The highest
 7 ozone consumption was shown by *M. helysia* with 335 nmol.

8 I₂ emissions

9 The I₂ emission rate was calculated by dividing the amount of I₂ by the sampling time and the
 10 suspension surface area. The results for the four samples are shown in Figure 2. The F/2
 11 mediumbackground and the two diatom samples, *M. helysia* and *P. glacialis* show significant higher
 12 emission rates when the ozone level is elevated (100 ppb O₃) compared to conditions where no
 13 ozone is present (0 ppb O₃). The difference between the high and low ozone conditions is small for
 14 the F/2 mediumbackground, increases for the *P. glacialis* sample and is highest for the *M. helysia*
 15 sample. The plankton concentrate does not show a significant dependence of the I₂ emission rate on
 16 the ozone level. The ozone-dependent increase in the I₂ emission rate of the other samples indicates
 17 that iodide, which is present at the air/water interface, is oxidised by ozone to form I₂, which is
 18 consistent with the results from artificial and natural seawater (Garland and Curtis, 1981, Sakamoto
 19 et al. 2009).

20 Figure 3 shows the change in I₂ emission rate ([I₂ at 100 ppb ozone] – [I₂ at 0 ppb ozone]) of the
 21 different samples as a function of the iodide concentration measured in the bulk water. A linear
 22 correlation fits the data well with a Pearson coefficient of R² = 0.998. This behaviour indicates a
 23 direct proportional relationship, which was also seen by Sakamoto et al. 2009 for small iodide
 24 concentrations (0 – 5 mmol L⁻¹). Carpenter et al. (2013) also observed that the I₂ emission is
 25 dependent on the aqueous iodide concentration. The proposed reaction sequence, as shown in
 26 equations (1)-(5), explains the relationship between the iodide concentration in the aqueous phase
 27 and the I₂ emissions (Sakamoto et al. 2009).



33 The plankton sample does not show an elevated I₂ emission at 100 ppb ozone compared to zero
 34 ozone. This observation indicates that in the plankton sample an additional I₂ loss process takes
 35 place. Reactions or partitioning of I₂ in an organic surface layer, which was discussed in Carpenter et
 36 al. (2013), would be one possibility to explain these results. In fact the specific microalgae found in
 37 the plankton concentrate, *Phaeocystis sp.*, is known to produce high amounts of organic matter

1 (Eberlein et al. 1985). An alternative explanation is the low iodide concentration in the plankton
2 concentrate, which may be related to iodide uptake by the natural occurring plankton communities.
3 The iodide concentrations and ozone mixing ratios in this study represent more likely natural
4 conditions compared to the study of Sakamoto et al. (iodide concentration between 0.01 – 50 mmol
5 L⁻¹ and ozone mixing ratio from 2 – 298 ppm). However, the results presented here demonstrate that
6 even under low iodide concentrations, representative of natural conditions of the MBL, a significant
7 formation of I₂ by the ozone driven oxidation of iodide at the air/water interface takes place, until
8 the iodide concentration gets too low.

9 Comparing the I₂ and iodocarbon emission rates, it is clear that the volatile iodine emissions are
10 dominated by I₂. Therefore I₂ emissions from natural seawater surfaces are more relevant for
11 atmospheric processes than the emission of iodocarbons. At the same time the experiments
12 presented here show that the emission of iodocarbons is not linked to the formation of I₂ at the
13 air/water interface (c.f. Martino et al., 2009), since no correlation between I₂ emissions or O₃ mixing
14 ratio and iodocarbon emissions was observed.

15 Calculated emissions for the F/2-mediumbackground, *P. glacialis* and *M. helysia* were 8.32 x 10⁵, 1.47
16 x10⁶ and 2.40 x 10⁶ molecules cm⁻² s⁻¹, respectively. Modelled emissions calculated using the kinetic
17 model of the aqueous interfacial layer by Carpenter et al. (2013) for the iodide concentration
18 measured were 1.16 x 10⁶, 1.67 x 10⁶ and 2.91 x 10⁶ molecules cm⁻² s⁻¹, respectively. The measured
19 and modelled values agree well, showing that the model is able to predict emissions for natural
20 iodide concentrations.

21 Figure 4 shows the change in the I₂ emission rate plotted versus the consumed ozone for the four
22 different samples. This was done to see whether ozone depletion in the flow chamber is mainly
23 driven by the iodide or if other factors are important. The graph shows that the ozone depletion
24 correlates with the enhancement in the I₂ emission rate for the two diatom samples and for the F/2
25 mediumbackground. Therefore, the ratio of the formation ratio of I₂ in dependence on to the
26 amount of O₃ consumed, was calculated by as $R(I_2) = n(I_2) / n(O_3)$, with $n(I_2)$ = amount of I₂ formed
27 and $n(O_3)$ = amount of O₃ consumed during the experiment, was used to determine the dependence
28 of I₂ formation on O₃. R(I₂) has a maximum value of 1, regarding which, referring to eqs.1-5, indicates
29 that if every molecule of ozone which is consumed produces one molecule of I₂. The formation ratio
30 for the F/2-background sample was the highest with R(I₂) = 0.14‰, followed by the samples of M.
31 helysia with R(I₂) = 0.08‰ and P. glacialis R(I₂) = 0.07‰. This means that a higher degree of biological
32 activity of the sample decreases the formation ratio. The decrease of I₂ emission in the surface
33 reaction of ozone with iodide was also seen by Carpenter et al. when turning from iodide solutions to
34 sea water, which contains more organic substances (Carpenter et al. 2013).

35 The plankton concentrate also depletes ozone, although there is no enhancement in I₂ emission.
36 Therefore, another mechanism in ozone depletion obviously takes must be taking place, possibly
37 induced by other ozone reactive substances formed or excreted from *Phaeocystis sp.*. Another
38 explanation is a reduced release of I₂ and a higher release of HOI, which was not measured in this
39 study. In factdeed, Carpenter and coworkers observed HOI as the main iodine compound released in
40 their experiments, followed by I₂ (Carpenter et al. 2013).

41

1 Conclusions

2 Different phytoplankton suspensions were treated with high and low ozone levels. Halocarbons
3 including bromoform, iodomethane, iodochloromethane and diiodomethane, were released from
4 the suspensions independent of the ozone level. The use of an evaporation standard in the aqueous
5 phase indicated that the emission rates of all gaseous organics were quite stable. The iodide and
6 iodate concentration in the liquid phase also showed no dependence on the ozone level in the gas
7 phase and were comparable to concentrations in surface water in the open ocean. The emission flux
8 of the iodocarbons was lower compared to the calculated flux at a natural site in Roscoff, France, an
9 observation which emphasizes the higher emission of iodocarbons from macroalgae compared to
10 microalgae. The emission flux of the iodocarbons was lower compared to the calculated flux at a
11 coastal, kelp-rich site in Roscoff, France, an observation which emphasizes the higher emission of
12 iodocarbons from macroalgae compared to microalgae. The emission rates of iodocarbons were also
13 lower than the emission of I₂, confirming that I₂ emissions from the remote ocean dominate over
14 organic iodine sources for the MBL (Jones et al., 2010; Lawler 2012; Carpenter et al., 2013). The
15 emission of I₂ showed a dependency on the ozone level in the air as well as on the iodide
16 concentration in the sample suspension, as has been found previously (Carpenter et al. 2013 and
17 other refs). For the two diatom samples, *M. helysia* and *P. glacialis*, and the ~~F/2-medium~~-background
18 sample, a correlation was found for the I₂ emission and the ozone consumption during the
19 experiment. The I₂ emissions from the plankton concentrate, taken in the Wadden Sea of Sylt, were
20 as-lower than the other samples and showed no dependence on the ozone levels. An explanation
21 could be the lower iodide concentration in the plankton sample, since iodide is the limiting factor for
22 the oxidative reaction. Another explanation may be the preferred formation and emission of HOI
23 when organic compounds are present in the liquid phase. The experiments showed that different
24 algae suspensions (*M. helysia* and *P. glacialis*) are capable of emitting I₂ by the reaction of ozone
25 with dissolved iodide at the air/water interface under natural conditions. However, it remains
26 unclear whether iodine emissions from aquatic systems can be fully understood without the
27 simultaneous measurement of HOI.

28

29 Acknowledgements

30 The authors gratefully acknowledge the financial support of the German Research Foundation (DFG:
31 Deutsche Forschungsgemeinschaft) within the Research Training Group 826: "Trace Analysis of
32 Elemental Species: Development of Methods and Applications."

33

1 References

- 2 [Amachi S., Kamagata T., Kanagawa T., Muramatsu Y.: Bacteria mediate methylation of iodine in](#)
3 [marine and terrestrial environments. *Appl. Environ. Microbiol.* 67, 2718-2722, 2001.](#)
- 4 [Amachi S., Kasahara M., Hanada S., Kamagata Y., Shinoyama H., Fujii T., Muramatsu Y.: Microbial](#)
5 [participation in iodine volatilization from soils, *Environ. Sci. Tech.* 37, 3885-3890 2003.](#)
- 6 Bloss W.J., Lee J.D., Johnson G.P., Sommariva R., de Heard, Saiz-Lopez A., Plane J.M., McFiggans G.,
7 Coe H., Flynn M., Williams P., Rickard A.R., Fleming Z.L.: Impact of halogen monoxide chemistry
8 upon boundary layer OH and HO₂ concentrations at a coastal site, *Geophys. Res. Lett.*, 32,
9 doi:10.1029/2004GL022084, 2005.
- 10 Bluhm K., Croot P.L., Huhn O.R.G.: Distribution of iodide and iodate in the Atlantic sector of the
11 southern ocean during austral summer, Deep-sea research part II- topical studies in
12 oceanography 58, 2733–2748, 2011.
- 13 Bluhm K., Croot P., Wuttig K., Lochte K.: Transformation of iodate to iodide in marine phytoplankton
14 driven by cell senescence, *Aquat. Biol.* 11, 1–15, 2010.
- 15 Carpenter L.J.: Iodine in the marine boundary layer, *Chem. Rev.* 103, 4953–4962, 2003.
- 16 Carpenter L.J., MacDonald S.M., Shaw M.D., Kumar R., Saunders R.W., Parthipan R., Wilson J., Plane
17 J.M.C.: Atmospheric iodine levels influenced by sea surface emissions of inorganic iodine, *Nature*
18 *Geosci.* 6, 108–111, 2013.
- 19 Carpenter L.J., Malin G., Liss P.S., Kupper F.C.: Novel biogenic iodine-containing trihalomethanes and
20 other short-lived halocarbons in the coastal East Atlantic, *Global. Biogeochem. Cy.*, 14, 1191–
21 1204, 2000.
- 22 Carpenter L.J., Sturges W.T., Penkett S.A., Liss P.S., Alicke B., Hebestreit K., Platt U.: Short-lived alkyl
23 iodides and bromides at Mace Head, Ireland: Links to biogenic sources and halogen oxide
24 production, *J. Geophys. Res.-Atmos.*, 104, 1679–1689, 1999.
- 25 Chance R., Malin G., Jickells T., Baker A.R.: Reduction of iodate to iodide by cold water diatom
26 cultures. *Mar. Chem.*, 105, 169–180, 2007.
- 27 Colomb A., Yassaa N., Williams J., Peeken I., Lochte K.: Screening volatile organic compounds (VOCs)
28 emissions from five marine phytoplankton species by head space gas chromatography/mass
29 spectrometry (HS-GC/MS). *J. Environ. Monit.*, 10, 325–330, 2008.
- 30 Eberlein K., Leal M.T., Hammer K.D., Hickel W. Dissolved organic substances during a *Phaeocystis*
31 *pouchetii* bloom in the German Bight (North Sea), *Mar. Biol.*, 89, 311-316, 1985.
- 32 Edler L.: Phytoplankton and chlorophyll. *Baltic Marine Biologists*, vol 5. Univ. of Stockholm,
33 Stockholm, 1979
- 34 Garland J.A., Curtis H.: Emission of iodine from the sea-surface in the presence of ozone, *J. Geophys.*
35 *Res.-Oc., ATM* 86, 3183–3186, 1981.
- 36 Hayase S., Yabushita A., Kawasaki M.: Iodine Emission in the Presence of Humic Substances at the
37 Water's Surface, *J. Phys. Chem.*, 116, 5779–5783, 2012.

- 1 Hayase S., Yabushita A., Kawasaki M., Enami S., Hoffmann M.R., Colussi A.J.: Heterogeneous reaction
2 of gaseous ozone with aqueous iodide in the presence of aqueous organic species, *J. Phys.*
3 *Chem. A* 114,6016–6021, 2010.
- 4 Hoffmann T., O'Dowd C.D., Seinfeld J.H.: Iodine oxide homogeneous nucleation: An explanation for
5 coastal new particle production, *Geophys. Res. Lett.*, 28, 1949–1952, 2001.
- 6 Huang R. J., Hoffmann T.: Development of a coupled diffusion denuder system combined with gas
7 chromatography/Mass spectrometry for the separation and quantification of molecular iodine
8 and the activated iodine compounds iodine monochloride and hypoiodous acid in the marine
9 atmosphere, *Anal. Chem.* 81,1777–178, 2009.
- 10 Huang R. J., Seitz K., Buxmann J., Poehler D., Hornsby K.E., Carpenter L.J., Platt U., Hoffmann T.: In
11 situ measurements of molecular iodine in the marine boundary layer: the link to macroalgae and
12 the implications for O₃, IO, OIO and NO_x, *Atmos. Chem. Phys.*, 10, 4823–4833, 2010a.
- 13 Huang, R. J., Seitz, K., Neary, T., O'Dowd, C. D., Platt, U., Hoffmann T.: Observations of high
14 concentrations of I₂ and IO in coastal air supporting iodine-oxide driven coastal new particle
15 formation, *Geophys. Res. Lett.*, 37, L03803, doi:10.1029/2009GL041467, 2010b.
- 16 Huang, R. J., Hou, X. L., Hoffmann, T.: Extensive evaluation of a diffusion denuder for the
17 quantification of atmospheric stable and radioactive molecular iodine, *Environ. Sci. Technol.*, 44,
18 5061–5066, 2010c.
- 19 Huang, R. J., Thorenz, U. R., Kundel, M., Venables, D. S., Ceburnis, D., Ho, K. F., Chen, J., Vogel, A. L.,
20 Kupper, F. C., Smyth, P. P. A., Nitschke, U., Stengel, D. B., Berresheim, H., O'Dowd, C. D.,
21 Hoffmann, T.: The seaweeds *Fucus vesiculosus* and *Ascophyllum nodosum* are significant
22 contributors to coastal iodine emissions. *Atmos. Chem. Phys.*, 13, 5255-5264, 2013.
- 23 Hughes, C., Malin, G., Turley, C., Keely, B., Nightingale, P., Liss, P.: The production of volatile
24 iodocarbons by biogenic marine aggregates. *Limnol. Oceanogr.*, 53,867-872.2008.
- 25 Jeffrey S.W., Humphrey G.F.: New Spectrophotometric equations for determining chlorophylls A, B,
26 C1 and C2 in higher plants, algae and natural phytoplankton. *Biochem. Physiol. Pfl.*, 167, 191–
27 194, 1975.
- 28 Jones C.E., Hornsby K.E., Dunk R.M., Leigh R.J., Carpenter L.J. Coastal measurements of short-lived
29 reactive iodocarbons and bromocarbons at Roscoff, Brittany during the RHaMBLe campaign,
30 *Atmos. Chem. and Phys.*, 9, 8757–8769, 2009.
- 31 [Kraberg, A., Carstens, K., Peters, S., Tilly, K., & Wiltshire, K. \(2012\). The diatom *Mediopyxis*
32 *helysia* Kühn, Hargreaves & Halliger 2006 at Helgoland Roads: a success story?
33 *Helgoland Marine Research*, 66\(3\), 463–468. doi:10.1007/s10152-011-0277-9](#)
- 34 Kundel M., Thorenz U.R., Petersen J.H., Huang R., Bings N.H., Hoffmann T.: Application of mass
35 spectrometric techniques for the trace analysis of short-lived iodine-containing volatiles emitted
36 by seaweed, *Anal. Bioanal. Chem.*, 402, 3345–3357, 2012.
- 37 Küpper F.C., Carpenter L.J., McFiggans G.B., Palmer C.J., Waite T.J., Boneberg E., Woitsch S., Weiller
38 M., Abela R., Grolimund D., Potin P., Butler A., Luther G.W., Kroneck P.M.H., Meyer-Klaucke W.,
39 Feiters M.C.: Iodide accumulation provides kelp with an inorganic antioxidant impacting
40 atmospheric chemistry, *Proc. Natl. Acad. Sci. U.S.A.*, 105, 6954–6958, 2008.

- 1 Lai S.C., Williams J., Arnold S.R., Atlas E.L., Gebhardt S., Hoffmann T.: Iodine containing species in the
2 remote marine boundary layer: A link to oceanic phytoplankton, *Geophys Res. Lett.*, 38, 2011.
- 3 Mahajan A.S., Plane J.M.C., Oetjen H., Mendes L., Saunders R.W., Saiz-Lopez A., Jones C.E., Carpenter
4 L.J., McFiggans G.B.: Measurement and modelling of tropospheric reactive halogen species over
5 the tropical Atlantic Ocean. *Atmos. Chem. Phys.*, 10, 4611–4624, 2010.
- 6 McFiggans G., Plane J.M., Allan B.J., Carpenter L.J., Coe H., O'Dowd C.: A modeling study of iodine
7 chemistry in the marine boundary layer. *J. Geophys. Res.-Atm.*, 105, 14371–14385, 2000.
- 8 Mishra S., Singh V., Jain A., Verma K.K.: Determination of iodide by derivatization to 4-iodo-N,N-
9 dimethylaniline and gas chromatography-mass spectrometry, *Analyst*, 125, 459–464, 2000.
- 10 Moore R.M., Webb M., Tokarczyk R., Wever R.: Bromoperoxidase and iodoperoxidase enzymes and
11 production of halogenated methanes in marine diatom cultures. *J. Geophys. Res.-Oc.*, 101,
12 20899–20908, 1996.
- 13 O'Dowd C.D., Hoffmann T.: Coastal new particle formation: A review of the current state-of-the-art,
14 *Environ. Chem.* 2, 245–255, 2005.
- 15 Quack B., Peeken I., Petrick G., Nachtigall K.: Oceanic distribution and sources of bromoform and
16 dibromomethane in the Mauritanian upwelling, *J. Geophys. Res.-Oc.*, 112, 2007.
- 17 [van Bergeijk S.A., Hernandez Javier L., Heyland A., Manchado M., Pedro Canavate J.: Uptake of
18 iodide in the marine haptophyte *Isochrysis sp.* \(T.ISO\) driven by iodine oxidation, *J. Phycol.*, 49,
19 640-647, 2013.](#)
- 20 von Glasow R. and Crutzen P.J. Tropospheric Halogen Chemistry. In: *Treatise on Geochemistry*
21 *Update1*, vol 4.02, pp 1–67, 2003.
- 22 Read K.A., Mahajan A.S., Carpenter L.J., Evans M.J., Faria B.V.E., Heard D.E., Hopkins J.R., Lee J.D.,
23 Moller S.J., Lewis A.C., Mendes L., McQuaid J.B., Oetjen H., Saiz-Lopez A., Pilling M.J., Plane
24 J.M.C.: Extensive halogen-mediated ozone destruction over the tropical Atlantic Ocean, *Nature*,
25 453, 1232–1235, 2008.
- 26 Roy R.: Short-term variability in halocarbons in relation to phytoplankton pigments in coastal waters
27 of the central eastern Arabian Sea, *Estuar Coast. Shelf. S.*, 88, 311–321, 2010.
- 28 Roy R., Pratihary A., Narvenkar G., Mochemadkar S., Gauns M., Naqvi S.W.A.: The relationship
29 between volatile halocarbons and phytoplankton pigments during a *Trichodesmium* bloom in
30 the coastal eastern Arabian Sea. *Estuar Coast. Shelf. S.*, 95, 110–118, 2011.
- 31 Saiz-Lopez A., Plane J.M.: Novel iodine chemistry in the marine boundary layer, *Geophys. Res. Lett.*,
32 31, 2004.
- 33 Saiz-Lopez A., Plane J.M., McFiggans G., Williams P.I., Ball S.M., Bitter M., Jones R.L., Hongwei C.,
34 Hoffmann T.: Modelling molecular iodine emissions in a coastal marine environment: the link to
35 new particle formation. *Atmos. Chem. Phys.* 6, 883–895, 2006.
- 36 Sakamoto Y., Yabushita A., Kawasaki M., Enami S.: Direct emission of I₂ molecule and IO radical from
37 the heterogeneous reactions of gaseous ozone with aqueous potassium iodide solution, *J. Phys.*
38 *Chem. A*, 113, 7707–7713, 2009.

- 1 Schall C., Heumann K.G., Kirst G.O.: Biogenic volatile organoiodine and organobromine hydrocarbons
2 in the Atlantic Ocean from 42 degrees N to 72 degrees, S. Fresen. J. Anal. Chem., 359,298–305,
3 1997.
- 4 Schwehr K.A., Santschi P.H.: Sensitive determination of iodine species, including organo-iodine for
5 freshwater and seawater samples using high performance liquid chromatography and
6 spectrophotometric detection, Anal. Chim. Acta, 482, 59–71, 2003.
- 7 Thorenz U.R., Kundel M., Müller L., Hoffmann T.: Generation of standard gas mixtures of
8 halogenated, aliphatic, and aromatic compounds and prediction of the individual output rates
9 based on molecular formula and boiling point, Anal. Bioanal. chem., 2012.
- 10 Weisse T., Grimm N., Hickel W., Martens P.: Dynamics of *Phaeocystis pouchetii* blooms in the
11 Wadden Sea of Sylt (German Bight, North Sea), Estuar. Coast. Shelf. S., 171–182, 1986
- 12 Wong G.T., Piumsomboon A.U., Dunstan W.M.: The transformation of iodate to iodide in marine
13 phytoplankton cultures. Mar. Ecol.-Prog Ser. 237, 27–39, 2002.
- 14 [Wong G.T.F., Cheng X.-H.: Dissolved organic iodine in marine waters: Determination, occurrence and](#)
15 [analytical implications. Mar.Chem. 59, 271-281, 1998.](#)
- 16

1 Table 1: Halocarbon emission rates, concentrations of chlorophyll α , iodide and iodate in the
 2 four different sample suspensions


<u>Sample</u>		<u>F/2 medium background</u>	<u><i>P. glacialis</i></u>	<u><i>M. helysia</i></u>	<u>plankton concentrate</u>
		<u>Range</u>	<u>Range</u>	<u>Range</u>	<u>Range</u>
		<u>(Mean)</u>	<u>(Mean)</u>	<u>(Mean)</u>	<u>(Mean)</u>
<u>CH₃I</u>	<u>pmol min⁻¹ m⁻²</u>	<u>0.17 – 0.72 (0.35)</u>	<u>0.21 – 0.69 (0.45)</u>	<u>0.32 – 0.82 (0.53)</u>	<u>0.08 – 0.37 (0.19)</u>
<u>CH₂Cl</u>	<u>pmol min⁻¹ m⁻²</u>	<u>0.02 – 0.22 (0.11)</u>	<u>0.02 – 0.22 (0.16)</u>	<u>0.04 – 0.22 (0.18)</u>	<u>0.02 – 0.12 (0.07)</u>
<u>CH₂I₂</u>	<u>pmol min⁻¹ m⁻²</u>	<u>0.02 – 0.08 (0.07)</u>	<u>0.27 – 0.44 (0,36)</u>	<u>0.21 – 0.50 (0.37)</u>	<u>0.04 – 0.09 (0.07)</u>
<u>CHBr₃</u>	<u>pmol min⁻¹ m⁻²</u>	<u>1.76 – 1.90(1.81)</u>	<u>1.99 – 2.17 (2.09)</u>	<u>1.75 – 2.17 (2.09)</u>	<u>1.75 – 2.33 (1.82)</u>
<u>chl α</u>	<u>$\mu\text{g L}^{-1}$</u>	<u>n.d.</u>	<u>257.27</u>	<u>926.59</u>	<u>2.53</u>
<u>Iodide</u>	<u>nmol L⁻¹</u>	<u>6.60 - 15.69 (10.35)</u>	<u>7.32 - 19.71 (12.70)</u>	<u>9.90 - 21.94 (16.84)</u>	<u>3.52 - 9.45 (6.47)</u>
<u>Iodate</u>	<u>nmol L⁻¹</u>	<u>402 - 538 (428)</u>	<u>408 - 478 (448)</u>	<u>397 - 499 (446)</u>	<u>424 - 478 (442)</u>
<u>1,3-C₃H₆Br₂ *</u>	<u>pmol min⁻¹ m⁻²</u>	<u>7.77 \pm 0.04</u>	<u>7.78 \pm 0.59</u>	<u>7.77 \pm 0.99</u>	<u>7.69 \pm 0.07</u>
<u>$\Sigma_{\text{iodocarbon}}/\text{chl } \alpha$</u>	<u>pmol/g</u>	<u>n.d.</u>	<u>19.75</u>	<u>6.06</u>	<u>694.88</u>

* evaporation standard given as

mean \pm standard deviation


chl α was measured for each sample once halocarbons, iodide, iodate mean values and ranges are calculated from 6 replicates

$\Sigma_{\text{iodocarbon}}/\text{chl } \alpha$ iodocarbon emissions were summed for the experimental conditions (time and surface area) and normalized to chl α in the watery phase




1

2 Figure 1: Experimental setup of the chamber with the phytoplankton suspension




1




1
2
3
4
5

Figure 2: Iodine emission rates normalized for the surface area of the different samples at 0 ppb and 100 ppb ozone. The error bars represent the standard deviation of the three replicates of each experiment.




1



2

3 Figure 3: Correlation of the change in the I₂ emission ~~and in dependency on~~ the iodide concentration
 4 in the ~~phytoplankton mikro algae~~ suspension



1
 2 | Figure 4: Function of the change in the total I_2 emissions ~~in dependency of~~ in relation to the amount of
 3 consumed ozone

4

5